

Risk factors for death among patients hospitalised for Covid-19 in Guinea (2020-2022)

Abstract

Introduction: The Republic of Guinea, like other countries, has recorded cases of Covid-19 in all its health districts. The aim of this study was to identify the factors associated with death in hospitalised patients.

Methods: This was a retrospective, cross-sectional, analytical study conducted from 12 March 2020 to 14 November 2022, based on records of patients admitted to treatment centres for confirmed Covid-19. Data were collected from 28 of the country's 38 health districts. A multivariate analysis was used to identify factors associated with death.

Results: The study involved 37714 confirmed cases of covid-19. The majority were male (67.33%) with an average age of 37 ± 15 years. Sixty-one percent (61%) had had contact with a covid-19 patient prior to their diagnosis. The mean time to admission to the treatment centre was 2.94 ± 0.01 days. Clinically, the main signs were dyspnoea (76.66%), cough (75.06%), anosmia (49.19%), agueusia (43.85%), fever (17.37%), headache (7.11%) and physical asthenia (5.47%). The most common comorbidities were arterial hypertension (29.41%) and diabetes (22.34%). The average hospital stay was 20 ± 0.05 days, with a case fatality rate of 1.23%. The main factors associated with death were age greater than 60 years [ORaj:6.29 and P:0.000 (CI95%:4.86-8.15)]; hospital stay greater than 2 days [ORaj: 2.67 and P:0.021 (CI95%:1.6-13.20)]; diabetes [ORaj:2.00 and P: 0.000 (CI95%:0.79-4.13)] and arterial hypertension [ORaj:9.14 and P:0.000 (CI95%:2.48-9.47)].

Conclusion: Taking better account of these factors associated with death, coupled with improving management capacity by raising the level of resuscitation facilities, will help to further reduce Covid-19-related mortality in Guinea.

Key words : Covid-19, factors, death, Guinea.

1. INTRODUCTION

Coronaviruses form a vast family of viruses, which can be pathogenic in animals or humans. In humans, several coronaviruses can cause respiratory infections ranging from the common cold to more serious illnesses, such as severe acute respiratory syndrome and Middle East respiratory syndrome [1]. In December 2019, a case of pneumonia of unknown cause was detected in China. Very quickly, the pathogenic agent involved was isolated: SARS-CoV-2. In December 2019, pneumonia of unknown cause was detected in China. The pathogen involved was very quickly isolated: SARS-CoV-2 [2]. It then spread throughout the world. Research carried out in 185 countries revealed that mortality due to COVID-19 was associated with age over 65, co-morbidities and unequal social and income conditions [3]. Other associated factors reported in the United States were socio-economic conditions and the environment/climate [4]. In addition, diabetes and chronic obstructive pulmonary disease were significantly associated with COVID-19 death in Mali in 2023.

Although 81% of cases of COVID-19 are mild, 14% of patients present with the severe form and 5% with the critical form. The mortality rate among critical cases is around 50% [5]. In West Africa in 2021, overall mortality was 5%, including 1% in patients under 40, 5% in patients aged 40 to 59 and 14% in patients aged 60 or over [6]. As in other countries, Guinea has been affected by the Covid-19 pandemic, recording its first case on 12 March 2020, in a patient returning from Europe.

Several response strategies (home confinement for uncomplicated cases, hospitalisation of serious cases, vaccination first of all of at-risk populations and then extended to the general population, compulsory compliance with barrier measures in gathering places) have been implemented by the national health security agency in collaboration with technical and financial partners under the authority of the Ministry of Health. As far as patient care is concerned, treatment centres for diseases with epidemic potential have been built throughout the country and are now operational to receive cases.

The aim of this study was to identify the risk factors associated with mortality among patients hospitalised for Covid-19 in treatment centres in Guinea between 2020 and 2022.

2. MATERIAL and METHODS

2.1. Scope of the study: The Republic of Guinea, located in West Africa, is bordered by 300 km of coastline and stretches 800 km from east to west and 500 km from north to south, with a total surface area of 245,857 km². In terms of health, the country has 33 districts plus the special region of Conakry Capital (05 districts). Treatment centres have been built throughout the country to manage diseases with epidemic potential, including Covid-19.

2.1. Type and duration of study: This was a retrospective, cross-sectional, analytical study conducted from January 4 to July 18, 2024 in 28 treatment centres in the country, using data from confirmed cases of covid-19 hospitalised from March 12, 2020 to November 14, 2022.

2.3. Selection criteria :

2.3.1. Inclusion criteria: All hospitalised patients with a confirmed diagnosis of Covid-19 by RT-PCR or rapid diagnostic test between March 2020 and November 2022.

2.3.2. Exclusion criteria: Patients with incomplete data or whose records lack information critical to the analysis.

2.3.3. Diagnosis of SARSCoV-2 infection was based on RT-PCR [7] and rapid diagnostic tests. Patients were treated symptomatically for simple forms, and severe cases were admitted to intensive care [8].

2.4. Sampling :

Sampling was non-probability and exhaustive for all Covid-19 patients hospitalised in the 28 health districts. However, the 10 other health districts were not included because of poor data management in some places and a lack of data in others.

2.5. Study variables :

The study variables were quantitative (age, length of hospitalisation, time to hospitalisation) and qualitative (occupation, sex, place of residence, mode of exposure, clinical signs, clinical classification, treatment received, site of treatment, outcome, comorbidities and factors associated with death).

The length of hospitalisation corresponded to the time elapsed in the health facility until the patient's death or discharge. Time to hospitalisation corresponded to the time between the appearance of the first symptoms and the patient's hospitalisation after confirmation.

As for comorbidities, some were self-reported by the patients themselves and others were diagnosed during complementary check-ups.

2.6. Statistical analysis tools :

The analysis was carried out using Epi-info version 7 and Stata version 11 software. The descriptive part consisted of a detailed description of the study variables.

Factors associated with death were identified using a stepwise retrograde multivariate analysis, which involved cross-tabulating death with independent variables (age, length of hospitalisation, time to hospitalisation, occupation, sex, place of residence, exposure mode, clinical signs, clinical classification, treatment received, treatment site, outcome, comorbidities).

All variables associated with the 5% threshold were retained as factors associated with death.

3. RESULTS

The study involved 37714 confirmed cases of covid-19. The majority were male (67.33%) with an average age of 37±15 years. The most common occupations were drivers (33%), followed by shopkeepers (30%) and administrators (21%). The majority lived in Conakry (83.70%) (Table 1).

Clinically, the main signs were dyspnoea (76.66%), cough (75.06%), anosmia (49.19%), agueusia (43.85%), fever (17.37%), headache (7.11%) and physical asthenia (5.47%). According to the clinical classification on admission, 69.55% of covid-19 confirmed cases were asymptomatic, 13.08% were mild cases, 11.90% were minor cases, 4.48% were severe cases and 0.99% were critical cases. The mean time to admission to the treatment centre was 2.94 ± 0.01 days. Sixty-one percent (61%) had had contact with a covid-19 patient prior to their diagnosis. The most common comorbidities were arterial hypertension (29.41%) and diabetes (22.34%). The average hospital stay was 20 ±0.05 days, with a case fatality rate of 1.23% (Table 2).

The main factors associated with death were age greater than 60 years [ORaj:6.29 and P:0.000 (IC95%:4.86-8.15)]; hospital stay greater than 2 days [ORaj: 2.67 and P:0.021 (CI95%:1.6-13.20)]; diabetes [ORaj:2.00 and P: 0.000 (CI95%:0.79-4.13)] and arterial hypertension [ORaj:9.14 and P:0.000 (CI95%:2.48-9.47)] (Table 3).

Table 1: Distribution of the 37714 confirmed cases followed up at Covid-19 treatment sites in Guinea according to their socio-demographic characteristics, from 13 March 2020 to 14 November 2022.

Socio-demographic characteristics	Number n=37714	Proportion (%)	Mean ± standard deviation
-----------------------------------	-------------------	----------------	------------------------------

Sexe		
Male	25392	67,33
Female	12322	32,67
Age (years)		37±15
Under 60	34058	90,31
Over 60	3656	9,69
Profession		
Health agent	1253	3,00
Driver	12734	33,00
Shopkeeper/Merchant	11257	30,00
Pupil/student	1008	3,00
Administrator	7942	21,00
Miner	1355	4,00
Teacher	800	2,00
Military	600	2,00
*Other	765	2,00
Residence		
Conakry (capital)	31566	83,70
Outside Conakry	6148	16,30

*Ministers (7), no profession (349), seamstresses/tailors (409)

Table 2: Breakdown of the 37714 confirmed cases followed up at Covid-19 treatment sites in Guinea according to their clinical and therapeutic aspects, from 13 March 2020 to 14 November 2022.

Clinical and therapeutic aspects	Number N=37714	Proportion (%)	Mean ± standard deviation
Exposure mode			
Contact with a Covid-19 patient	23024	61,00	
No link Epidemiological	13538	36,00	
Travel to an epidemic country	1152	3,00	
Time taken before hospitalisation		2,94 ± 0,01	
Less than 2 days	12821	34,00	
More than 2 days	24893	66,00	
Comorbidity			
Diabetes	8424	22,34	
Hypertension	11091	29,41	
HIV	7818	20,73	
Vaccination status			
Vaccinated	8062	21,38	
Non-vaccinated	29652	78,62	
Clinical signs			
Agueusia	16538	43,85	
Anosmia	18519	49,19	
Fever	6550	17,37	
Headache	2681	7,11	
Cough	28307	75,06	
Respiratory difficulties	28911	76,66	
Asthenia	2062	5,47	
Clinical classification			
Asymptomatic cases	26230	69,55	
Mild case	4934	13,08	

Minor case	4488	11,90
Severe case	1690	4,48
Critical case	372	0,99
Treatment received		
Hydroxy-chloroquine	23502	62,32
Azythromycin	12396	32,87
Dexamethasone	454	1,20
Oxygen therapy	454	1,20
Ceftriaxone inj	454	1,20
Lovenox	454	1,20
Length of hospital stay		20 ±0,05
Less than 10 days	6858	18,18
More than 10 days	30856	81,82
Patient outcome		
Cured	37250	98,77
Deceased	464	1,23

Table 3: Main factors associated with death in confirmed cases followed up at Covid-19 treatment sites in Guinea, from 13 March 2020 to 14 November 2022.

Factors linked to death	Deceased		
	P-value	ORaj	IC 95%
Age			
Under 60 years old	-	1	-
Over 60 years old	0,000	6,29	[4,86-8,15]
Time taken before hospitalisation			
Under 2 days	-	1	-
Over 2 days	0,029	2,67	[1,6-13,20]
Comorbidity			
Diabetes			
No	-	1	-
Yes	0,000	2,00	[0,79-4,13]
HTA			
No	-	1	-
Yes	0,000	9,14	[2,48-9,47]
HIV			
No	-	1	-
Yes	0,000	2,67	[1,65-4,31]
Status			
Vaccinated	-	1	-
Not vaccinated	0,000	5,30	[2,12-13,20]
Clinical classification			
Asymptomatic cases	-	1	-
Severe cases	0,000	26,45	[16-44]
Critical cases	0,000	117,6	[73,4-188,37]

ORaj: Adjusted Odds Ratio IC 95%: 95% Confidence Interval

4. DISCUSSION

Following the major Ebola epidemic between 2014 and 2016, the Republic of Guinea strengthened its health system by setting up coordination and monitoring bodies, training human resources and building and equipping diagnostic and case-management centres. A network of more than 38 centres for the treatment of diseases with epidemic potential has been built in all the country's health districts. It was against this backdrop that the great Covid-19 pandemic occurred. Guinea did not escape the explosion and transmission of cases, most of which were community-based. The aim of our study was to identify the factors associated with death in Covid-19 patients hospitalised at 28 treatment sites.

Studies on understanding the factors associated with mortality are important, as they provide a basis for decision-making in healthcare establishments [9].

The main factors found in our patients were age over 60, time to hospitalisation over 2 days, diabetes, HIV infection, non-vaccination against covid-19, arterial hypertension, and critical and severe case status.

Chronic disease increases the risk of adverse outcomes in COVID-19, including admission to intensive care and death [10].

In our series, diabetes and hypertension were associated with death.

Diabetes is responsible for an altered immune response, particularly cellular immunity. This exposes diabetics to more serious attacks from pathogenic germs [11]. In the Indian series [12] and Chinese [13], diabetes was also found. HIV-positive subjects were more likely to develop a severe form of COVID-19 and had an increased risk of death [14].

In Guinea, the prevalence of chronic diseases is increasing exponentially, with most cases being discovered by chance. The national programme to combat non-communicable diseases is working on the ground (by raising awareness and changing behaviour) to reverse this trend.

In Wuhan, China, the occurrence of death in Covid-19 positive patients was associated with an age \geq 65 years [13]. Ageing generates a number of physiological changes [15]. A younger age threshold for the risk of death has also been reported [16–18]. Kpamy [19] in Guinea reported that the risk was 24.93 times greater in patients over 60 than in those under 60.

The major presence of asymptomatic forms of covid-19 in this study can be explained by the exposure of younger people.

This observation has also been made by other authors [19,20] which found an average age of 36.86 years and affirmed that the youth of the African population went hand in hand with asymptomatic forms of covid-19 disease.

Long delays before admission to hospital were also associated with death in our series.

This can be explained, among other things, by the therapeutic itinerary of patients, most of whom first try self-medication or consultation with traditional therapists before being admitted to conventional hospitals.

Most of these patients were received in an advanced state of illness see complications was admitted to intensive care or resuscitation.

Efforts have been made by the health authorities to improve the technical facilities, but these are not available in all health districts.

In our series, non-vaccination was also identified as a factor associated with death.

In Lombardy (Italy), mortality in intensive care units and hospitals was not associated with vaccination status [21]. In the USA, severe forms of the disease were found in patients despite vaccination [22].

In the Republic of Guinea, vaccination against Covid-19 officially began on 04 March 2021. Several types of vaccine with different antigens and modes of administration have been used throughout the country. As of 11 February 2024, 12049333 doses of vaccine had been administered, giving an estimated coverage rate of 44.1%.

The study method was rigorous (triangulation of the information received to ensure the quality of the data and guarantee the validity of the results, among other things) and resulted in high-quality data and relevant results.

However, one of the limitations of this study is the possibility of under-reporting of deaths occurring in the community. In addition, the exclusion of the 10 districts could lead to a selection bias if these districts had different mortality rates.

CONCLUSION

This study identified advanced age, co-morbidities such as diabetes and hypertension, and delay before hospitalisation as major risk factors for death among Covid-19 patients in Guinea. Non-vaccination was also associated with higher mortality, highlighting the importance of increasing vaccination coverage. These results highlight the need to strengthen resuscitation infrastructures, improve rapid access to care, and develop targeted prevention strategies, including vaccination campaigns and co-morbidity management, in order to effectively reduce Covid-19-related mortality.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Generative AI technologies such as large language models (ChatGPT, COPILOT, etc.) and text-image generators were not used in the writing of this manuscript.

CONSENT

It is not applicable

ETHICAL APPROVAL

The prior agreement of the administrative and health authorities was obtained before the survey began.

Patient anonymity and confidentiality were respected.

COMPETING INTERESTS

The authors declared that there were no competing interests.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV, WHO Clinical Case Definition Working Group on Post-COVID-19 Condition. A clinical case definition of post-COVID-19 condition by a Delphi consensus. *Lancet Infect Dis.* 2022;22(4):102-7.
2. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet.* 2020;395(10229):1054-62.

3. Barbosa TP, Costa FBPD, Ramos ACV, Berra TZ, Arroyo LH, Alves YM, et al. COVID-19 morbidity and mortality associated with chronic conditions, health services, and inequities: evidence of a syndemic. *Panamerican Journal of Public Health*. 2022;46:1.
4. Luo Y, Yan J, McClure S. Distribution of the environmental and socioeconomic risk factors on COVID-19 death rate across continental USA: a spatial nonlinear analysis. *Environ Sci Pollut Res*. 2021;28(6):6587-99.
5. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239.
6. Jaspard M, Saliou Sow M, Juchet S, Dienderé E, Serra B, Kojan R, et al. Clinical presentation, survival and factors associated with mortality: a prospective study in three COVID-19 centers in West Africa. *Infect Dis Now*. 2021;51(5):59.
7. Pozzetto B, Delolme M, Rigai J, Lleres-Vadeboin M, Verhoeven P, Grattard F, et al. Virological diagnostic tests for Covid-19. *Revue de Biologie Médicale/N° 359 - MARCH-APRIL 2021*.
8. National Agency for Health Security/Ministry of Health in Guinea. National Guide for the Management of Coronavirus Disease Cases in Guinea, National Agency for Health Security, Ministry of Health, Republic of Guinea, 2020. 2nd ed.
9. Rayanne AO, Janiel CS, Livia FSS, Livia MP, Mellina Y, Floriacy SS et al. Factors associated with deaths from COVID-19 in a region of northeastern Brazil. *J Infect Dev Ctries* 2023; 17(9):1179-1187.
10. Mascarello KC, Vieira ACBC, Souza ASS, Marcarini WD, Barauna VG, Maciel ELN (2021) Hospitalization and death by COVID-19 and its relationship with social determinants of health. *Revista de Saúde Pública*. – actors related to deaths from COVID-19 in Brazil health and morbidities in Espírito Santo: a cross-sectional study.
11. Tian W, Jiang W, Yao J, Nicholson CJ, Li RH, Sigurslid HH, et al. Predictors of mortality in hospitalized COVID-19 patients: A systematic review and meta-analysis. *J Med Virol*. 2020;10.1002/jmv.26050.
12. Kumar A, Arora A, Sharma P, Anikhindi SA, Bansal N, Singla V, et al. Is diabetes mellitus associated with mortality and severity of COVID-19? A meta-analysis. *Diabetes Metab Syndr*. 2020;14(4):535-45.
13. Yu C, Lei Q, Li W, Wang X, Liu W, Fan X, et al. Clinical Characteristics, Associated Factors, and Predicting COVID-19 Mortality Risk: A Retrospective Study in Wuhan, China. *Am J Prev Med*. 2020;59(2):168-75.
14. Chanda D, Minchella PA, Kampamba D, Itoh M, Hines JZ, Fwoloshi S, et al. COVID-19 Severity and COVID-19–Associated Deaths Among Hospitalized Patients with HIV Infection — Zambia, March–December 2020. *Morb Mortal Wkly Rep*. 2021;70(22):807-10.
15. Maciel EL, Jabor P, Goncalves Júnior E, Tristão-Sá R, Lima R de CD, Reis-Santos B, et al. Factors associated with COVID-19 hospital deaths in Espírito Santo, Brazil, 2020. *Epidemiol E Serv Saude Rev Sist Unico Saude Bras*. 2020;29(4):e2020413.
16. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583 Subjects. *J Am Med Dir Assoc*. 2020;21(7):915-8.
17. Asfahan S, Deokar K, Dutt N, Niwas R, Jain P, Agarwal M. Extrapolation of mortality in COVID-19: Exploring the role of age, sex, co-morbidities and health-care related occupation. *Monaldi Arch Chest Dis Arch Monaldi Mal Torace*. 2020;90(2).

18. Smith AA, Fridling J, Ibrahim D, Porter PS. Identifying Patients at Greatest Risk of Mortality due to COVID-19: A New England Perspective. *West J Emerg Med.* 8 juill 2020;21(4):785-9.
19. DO Kpamy SK, B. Yattassaye MC, M. Barry MK, T. Camara SK, Y. Touré FC, M. Awada FT, et al. The COVID-19 pandemic in Guinea: clinical, therapeutic aspects and factors related to mortality in treatment sites. *API DPM.* 2020;67(10):509-17.
20. Etard JF, Touré A, Titiane Ndour C, Tshilolo L, Katchunga PB, Sow S, et al. Covid-19 in Africa: Do the numbers reflect reality? [Internet]. Vol. 26 July 2020. The Conversation Media Group; 2020 [cited 16 Sep 2024]. p. online [7 p.]. Available at: <https://hal.science/hal-02990361>.
21. Grasselli G, Zanella A, Carlesso E, Florio G, Canakoglu A, Bellani G, et al. Association of COVID-19 Vaccinations With Intensive Care Unit Admissions and Outcome of Critically Ill Patients With COVID-19 Pneumonia in Lombardy, Italy. *JAMA Netw Open.* 2022;5(10):e2238871.
22. Vo AD, La J, Wu JTY, Strymish JM, Ronan M, Brophy M, et al. Factors Associated With Severe COVID-19 Among Vaccinated Adults Treated in US Veterans Affairs Hospitals. *JAMA Netw Open.* 2022;5(10):e2240037.

UNDER PEER REVIEW