

Original Research Article

Phylogenetic and Gene Expression of *Norsolorinic acid* analysis of Aflatoxigenic Strains and Quantification of Aflatoxin B₁ Levels into White Rice Grains in Kenya

Abstract: The most important pathogens of global significance in food security are mycotoxigenic fungi. Aflatoxigenic strains contamination in rice grains can lead to a health risk for consumers. Under appropriate conditions, aflatoxigenic fungi can grow and subsequently produce aflatoxins. Aflatoxin poses a real threat to food safety and security and destroys approximately 25% of the world's food crops and every year 420,000 people die from consuming contaminated food causing an estimated loss of 33 million healthy life years. This study screened Norsolorinic Acid gene involved in the biosynthesis pathway, gene expression profile of aflatoxigenic strains characterized from rice grains, and assessed aflatoxin B₁ levels in white rice grains. We combine the use of molecular techniques and Enzyme-Linked Immunosorbent Assay to detect aflatoxin biosynthesis *afID* gene and quantify gene expression profile, and analyzed aflatoxin B₁ levels, respectively.

The results shown that all aflatoxigenic strains analyzed in this study are clustered within the same clade based on the *afID* gene. The targeted *afID* gene for expression was amplified in all aflatoxigenic strains with higher relative level in *Aspergillus flavus* but no relative expression shown in *Aspergillus fumigatus*. Enzyme-Linked Immunosorbent Assay showed that 43.1 % of the samples were found positive, with 15.9 % being local rice and 27.2 % imported rice. Notably, 11.3 % of analyzed samples were above the recommended aflatoxin B₁ in rice grains established by European Union with aflatoxin concentration ranging between 0 µg/kg to 3.2 µg/kg. These results are indicative of exposure of the population to aflatoxin B₁ and possible health hazard. Our study reveals high levels of aflatoxin B₁ contamination in both imported and local rice grains in Kenya. This study highlights the species diversity of aflatoxigenic fungi that have the potential to contaminate different types foodstuffs.

Keywords: Rice grains; aflatoxigenic; gene expression; aflatoxin B₁; contamination; ELISA; exposure; survey.

1. Introduction

Microbial pathogens, particularly mycotoxigenic fungi, are a significant global menace to food safety and security [1, 2]. In tropical climate conditions, aflatoxigenic fungi of *Aspergillus* species easily contaminate small grains like wheat, barley, rye, rice, triticale, and corn [3,4]. Mycotoxins are among the microbial toxins of most concern to public health and represent a significant barrier to the international trade of agricultural products and an obstacle in the face of global harmonization of regulatory standards, as discussed by [5]. At any stage of the foodstuffs production process, before and during harvesting, drying, transport, and storage, mycotoxigenic fungi can colonize grains crops and expose consumers to the risk of mycotoxin contamination directly through food consumption or indirectly through animal feed [6]. The contamination of foodstuffs and commodities with mycotoxins is a recurrent problem in Sub-Saharan Africa (SSA). It is caused mainly by post-harvest infestation of *Aspergillus* species section *flavi* during storage, holding, processing, transportation, and marketing. Among all mycotoxins, AFs are known as the most poisonous and toxic to human beings, significantly impacting the economic burden of foodstuffs and consumers [7, 8]. Aflatoxins (AFs) are secondary metabolites typically produced through certain aflatoxigenic species of *Aspergillus*, like *A. flavus*, *A. astellatus*, *A. nomius*, and *A. parasiticus* [9, 10]. AFs are genotoxic, carcinogenic, teratogenic, mutagenic, hepatotoxic, and immunosuppressive [11]. The most studied AFs among the main types are aflatoxin B₁ (AFB₁), aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁), and aflatoxin G₂ (AFG₂), while aflatoxin M₁ (AFM₁) and aflatoxin M₂ (AFM₂) have received considerable attention. AFs were first discovered following a terrible livestock poisoning occurrence in England involving turkeys [12]. Of all the AFs, AFB₁ is one of the most prevalent AFs, which represents a problem for consumers' health due to being hepatotoxic and a potent carcinogenic [13,14]. The challenge of AFs stands prominent everywhere in the world and specifically in African countries, where contamination by aflatoxins is recorded in foodstuffs with an incidence of 50 %, for instance, with 1642 µg/kg infestation in white rice grains [15]. Assessment of AFB₁ incidence in cereals is necessary to avoid and decrease the adversative health effects on the population. AFs in rice grains could be quantitatively determined by applying diverse investigative methods. Molecular techniques like PCR, Real-Time PCR, and phylogenetic analyses are commonly used to validate the morphological characterization of *Aspergillus* species section *Flavi* [5].

The biosynthesis pathway of AFs involves numerous enzymes encoded in genes located in a cluster of 75 kb in size [16]. In *Aspergillus*, the order of these genes is highly preserved [17]. There are 30 genes positioned within the same order in entirely aflatoxigenic fungi of the *Flavi* section [18, 19]. The cluster is controlled in a coordinated way, and *norsolorinic acid* is the first stable precursor of AFs [20, 21], which is then converted into *averantin* in the central of the aflatoxin biosynthetic pathway by a reductase encoded by the *aflD* gene [21, 22]. Indeed, a more profound knowledge about the elements included in the regulatory path of the biosynthetic of aflatoxin gene clusters that activate in aflatoxigenic strains is an essential prerequisite for mitigating and minimizing AFs and preventing foodstuffs and commodities contamination in the future [23, 24].

Global consumption of rice has recently increased. Rice grains were consumed around 490.27 million metric tons worldwide in the year of 2018-2019. The consumption of rice grains in Kenya is increasing at 12 % per year after wheat and maize with 4 % and 1 % respectively. Its local production is estimated to be between 100,000 and 120,000 metric tons per annum, while its consumption has increased from 400,000 to 410,000 tons per annum [25]. The contamination of rice grains with AFs has been documented in several studies, with the findings showing that rice samples surveyed contain AFs [26, 27, 28]. The European Union (EU) has set the maximum tolerable limits (MTL) for AFs in rice precisely, 4 µg/kg for total aflatoxins (B₁+B₂+G₁+G₂) and 2 µg/kg for AFB₁ [29].

Many methods are currently applied for the detection of AF contamination, including thin-layer chromatography (TLC), enzyme-linked immunosorbent assay (ELISA), high-performance liquid chromatography with a fluorescence detector (HPLC-FLD), or tandem mass spectrometry (HPLC-MS/MS) [30, 31]. TLC and ELISA are widely utilized to detect AF contamination in various foods, offering higher sensitivity around 1–20 µg/kg [32, 33].

Due to the tropical climates with extreme ranges of temperature, humidity, and rainfall levels throughout the year, rice grains sold in the markets, retail markets, and millers in Kenya are at an increased risk of contamination by AFs, posing a threat to food safety and security to the population and animal health. Aflatoxin poses a real threat to food safety and security and destroys approximately 25% of the world's food crops and every year 420,000 people die from consuming contaminated food causing an estimated loss of 33 million healthy life years. The most significant outbreaks of acute aflatoxicosis from highly contaminated food have been documented in Kenya. In July 2004, an outbreak of acute hepatotoxicity with 317 cases

and 150 deaths occurred as a result of AFs poisoning from ingestion of contaminated maize [34]. Another outbreak of aflatoxicosis occurred in 1981 from contaminated maize, with 500 acute illnesses and 200 deaths in Makueni County and other parts of Kenya [35]. From the above cases, it is clear that AFs food poisoning is a common occurrence in Kenyan cereals, a food commodity commonly used by many communities as staple food. However, no study has been performed in Kenya to assess the safety of rice grains for AFB₁ and the risk associated with consuming this staple product. Therefore, our study aimed to screen aflatoxigenic strains through Norsolorinic Acid (NOR) and quantify AFB₁ in rice grains sold in the market to evaluate the exposure levels from rice consumption and the associated liver cancer risk from this toxin.

2. Material and Methods

Sample Collection

Rice samples were collected randomly (1 kg) from local retail markets and millers in Mwea (59 samples) and at Jamhuri Market in Thika (39 samples) (Kenya) and labelled appropriately. A total of 98 samples (local rice produced in Mwea and imported rice originating from Biryani, India, Pakistan, and Thailand) were taken in line with the alternative sampling plan for the Official Control of Mycotoxins in Food [36]. Representative samples were put in sealed plastic ziplock bags and transported to the Molecular Biology and Biotechnology Laboratory at Pan African University Institute for Basic Sciences, Technology and Innovation (PAUSTI).

Isolation of Fungi

All samples of rice grains collected were analyzed using the direct plating approach defined by Pitt and Hocking [37]. Rice grains were surface sterilized by dipping in diluted commercial bleach (2.5% sodium hypochlorite) for 1 min before rinsing thrice in sterile distilled water. For every sample, 20 pieces (rice grains suspected to be infected by fungi) were positioned aseptically on a layer of moistened filter paper in the Petri dishes. The Petri dishes were incubated upright at 25 °C for 3-5 days in darkness and ventilated for 12 h on the 3rd day. Isolation of fungi from the suspected grains was performed using Ulster's method [38]. Fungi growing on distinctive seeds were isolated from emerging colonies on modified *rose Bengal chloramphenicol agar (MRBA)* [39]. The cultural characteristics described in taxonomic keys by Klich were used to identify *Aspergillus* section *Flavi* [40]. Pure cultures

were performed for subsequent studies using two different media, malt extract agar (MEA) and potato dextrose agar (PDA), according to Pitt and Hocking [39] and [41] for seven days.

Genomic DNA Extraction

Four *Aspergillus* strains (*A. flavus*, *A. parasiticus*, *A. nomius* and *A. fumigatus*) were identified for this study. The genomic DNA was extracted from seven-day-old culture grown on media according to [42] method with modification. A total of 50-100 mg of fungal mycelium was scraped and placed in a 2 mL Eppendorf tube. The tubes were frozen in liquid nitrogen and vortexed vigorously for 30 minutes with steel beads to crush the mycelial wall and release the DNA. A volume of 500 μ L of Lysis Buffer (100 mM Tris-HCl pH 8, 1 mM EDTA, 100 mM NaCl, 10 mM β -mercaptoethanol, and 1 % *Sodium dodecyl sulfate*), 5 μ L of RNase A, and 1 μ L of Proteinase K were added. Tubes were then incubated at 65 $^{\circ}$ C for 45 minutes in buffer. Thereafter, 270 μ L of Sodium/Potassium acetate (3M) was added, and samples were well-mixed and centrifuged at 13,000 rpm for 10 minutes. After centrifugation, the supernatant (700 μ L) was transferred to a fresh tube, and an equal volume of chloroform: isoamylalcohol (24:1) was added and mixed; samples were placed on a bench for 5 minutes, followed by centrifugation at 13,000 rpm for 10 minutes. The supernatant (700 μ L) was transferred into new tubes, and 80 μ L of Sodium/Potassium acetate (3M) and 587 μ L of ice-cold isopropanol were added and then mixed well by inverting the tubes and then incubated at -20 $^{\circ}$ C overnight. After that, the samples were centrifuged at 13,000 rpm for 30 minutes, and the supernatant was discarded carefully. The DNA pellets were washed with 1 mL of 70 % ethanol and centrifuged at 13,000 rpm for 10 minutes. The DNA pellets were air-dried, dissolved in 50 μ L of Tris-EDTA (TE) buffer, and stored at -20 $^{\circ}$ C until use.

Set	Primer Name	Sequences (5'-3')	Length of PCR Product (bp)
<i>afID</i>		F-ACCGCTACGCCGGCACTCTCGGCAC	400
		R-GTTGGCCGCCAGCTTCGACACTCCG	

Table 1. Oligonucleotide Primer Sets Used for the Polymerase Chain Reaction (PCR)

Abbreviations in the sequences: **F**, forward; **R**, reverse.

Sequencing and Phylogenetic Analysis

The *aflD* gene PCR products were purified using Qiagen kit for purification, and the products of *aflD* gene primers were sent to MacroGen Europe for sequencing analyses. After getting the forward and reverse primer sequences of each strain, sequences were edited with MEGA7 [43] and Geneious software (<http://www.geneious.com>). The data were contrasted in GenBank using the BLASTn algorithm. Additionally, strain information was evaluated using the reference strains of NCBI (<http://www.ncbi.nlm.nih.gov>). Using the maximum likelihood method based on the Tamura-Nei model [44], the phylogenetic relationship was found.

RNA Extraction

The four strains identified earlier were screened for gene expression to quantify the potential aflatoxigenic producers by amplifying the norsolorinic acid (NOR), *aflD* (*nor-1*) gene involved in the aflatoxin biosynthetic pathway. RNA's isolation was done using Rneasy Mini kit (Qiagen) following the manufacturer's instructions with brief modification. Briefly, 50 mg of fungal mycelium were scraped and frozen in liquid nitrogen and then ground into fine powder by vortexing in steel beads. The powder was transferred in a 2 mL Eppendorf tube, a suitable volume of Buffer RLT for a working solution was added, then centrifuged for 3 minutes at a maximum speed to disrupt and homogenize the lysate. After centrifugation, the supernatant was removed to a new tube. Additionally, 70 % of ethanol was added to the lysate and mixed well by pipetting. A volume of 700 μ L of the mixed sample was transferred to a Rneasy Mini spin column placed in a 2ml collection tube and centrifuged for 20 seconds at $\geq 8000 \times g$, the flow-through discarded. Thereafter, 700 μ L Buffer RW1 was added to the Rneasy spin column, centrifuged for 20 seconds at $\geq 8000 \times g$, and discarded the flow-through. After that, a volume of 500 μ L RPE was added to the Rneasy spin column and centrifuged, and the flow-through was discarded. This step was repeated (twice). Thereafter, the Rneasy spin column was positioned in a new 1.5 ml collection tube, 50 μ L of RNase-free water was added to the spin column membrane and centrifuged for 1 minute at $\geq 8000 \times g$ to elute the RNA. The concentration and purity of RNA were measured using a Nano-drop Spectrophotometer (Model PCR Max Lambda, Thermo Fisher Scientific, Waltham, MA, USA), and the quality of all extracted RNA was measured by taking their absorbance at 260 nm and 280 nm.

cDNA Synthesis

Reverse transcriptase was done using a cDNA synthesis kit. cDNA was prepared using 5X Hot Firepol® EvaGreen® HRM Mix (ROX) (Solis BioDyne) following the manufacturer's

instructions using random hexamers. Briefly, a master mix consisting of 1 µl of FIREScript RT, 0.5 µl of dNTP MIX (20 mM of each), 1 µl random hexamers (100 µM), 2 µl of 10x RT Reaction Buffer with DTT, 0.5 µl RiboGrip RNase Inhibitor, 5 µl of Nuclease-free H₂O was made and a 10 µl volume of total RNA was added to the master mix. Confirmation of successful cDNA synthesis was done using conventional PCR with amplification of primers for qPCR in Proflex PCR System with the following parameters: First Denaturation 95 °C for 3 minutes, second Denaturation 95 °C for 30 seconds, Primer annealing 58 °C for 45 seconds and Elongation 72 °C for 7 minutes. Then, after PCR reactions, the PCR product was confirmed by gel electrophoresis. The cDNA was then kept at -20 °C until use.

Table 2. Primer Sets Used for the Real-Time Polymerase Chain Reaction (qRT-PCR)

Set	Primer Name	Sequence (5'->3')	Product Length (bp)
aflD	Forward primer	AAT CAC AGT CCG AGG GTT CTT G	165
	Reverse primer	TGA TCA TAC CCG AGC ACA GAT	
<i>β-tubulin</i>	Forward primer	AAG ATC AAC GAG GAC GGC AC	222
	Reverse primer	CTC GAA CGA ACG ACG ACC A	

Gene Expression Profile Analysis

The gene expression in the aflatoxin biosynthetic pathway was assessed in triplicates for all aflatoxigenic strains identified. The housekeeping gene (*β-tubulin*) was used as a control during the experimentation. Two primers were designed using Genius Prime software and NCBI Gene bank for the gene of interest, *aflD* and *Beta tubulin*, a housekeeping gene Table 1. Quantifying relative gene expression by real-time quantitative PCR was conducted using the comparative of 2^{ΔΔCT} method. For gene expression analysis, *aflD* gene, which is in control for the conversion of *norsolorinic acid (NOR)* to *averantin (AVN)* in the centre of the aflatoxin biosynthetic pathway [45], was chosen.

Universal qPCR kit (5x HOT FIREPol® EvaGreen® qPCR Mix Plus) was utilized in running qPCR in qTOWER³ - Real-Time PCR Thermal Cyclers (Analytik Jena AG, Konrad-Zuse-str. 1 07745 Jena, Germany) 96-well real-time qPCR following the manufacturer's instruction. A total volume of 20 µL, including 4 µL of 5x HOT FIREPol, 13 µL of Nuclease-free H₂O, 0.5

μL of each oligonucleotide primer sets above for qPCR and 2 μL cDNA was performed in the system. The qPCR cycle parameters were set as: initial denaturation at 95 °C for 2 minutes, denaturation at 95 °C for 15 s, annealing plus elongation at 57 °C for 30 s melting curve analysis (60 -95 °C) for 10 s. Relative expression was determined using the $2^{-\Delta\Delta\text{CT}}$ method[46, 47].

Quantitative Determination of AFB₁ levels by ELISA

Local and imported white rice grains were randomly sampled from local retail markets and millers in Mwea and at Jamhuri Market in Thika, Kenya, and appropriately labelled. A total of forty-four different samples comprising homegrown rice in Mwea and imported rice originating from Biryani, India, Pakistan, and Thailand, respectively, were taken following the alternative collection plan for the Official Control of Mycotoxins in Food (OCMF) [36]. Representative samples were then sealed in plastic ziplock bags, transferred to the Laboratory of Molecular Biology and Biotechnology at Pan African University Institute for Basic Sciences, Technology and Innovation, and prepared for subsequent ELISA analyses according to OCMF.

Samples Preparation and ELISA Test Procedure

The quantitative analysis of AFB₁ from samples collected was performed based on a competitive enzyme immunoassay using RIDASCREEN® Aflatoxin B₁ 30/15 test kit (Art. No: R1211, R-Biopharm, Darmstadt, Germany). Samples collected were powdered using a moulinex blender, 5 g of the represented sample were transferred in sealed plastic bags (zip lock) and then stored at 4°C [48] until use. All samples were examined according to the instructions of the manufacturer in duplicates. Samples preparation for ELISA test were performed according to the technique described by R-Biopharm GmbH[49]. Briefly, to perform ELISA test rice grains was ground in fine powder, then 5 g of rice ground was transferred in to 50 ml of falcon tube, twenty-five milliliter methanol: water (70:30) was added in the falcon then after the samples were shaken vigorously for 3 minutes with shaker. The obtained extract was filtered through a filter paper and diluted with distilled water (1:1). At a minimum, 50 μL of the diluted filtrate was used per well in the test. Test procedure was done according to RIDASCREEN® Aflatoxin B₁ 30/15 (Art No.: 1211) test kit manual, a volume of 50 μL of the standard solution and prepared sample was added to the wells of micro titer plate in duplicate. Then, 50 μL of the enzyme conjugate was added, and this step was directly followed by adding on top 50 ml of the antibody solution to each well, then

kindly mixed and raised for 30 minutes at room temperature (20-25 °C). Later, liquid was discarded from wells by beating them upside down dynamically against the absorbent paper; followed by cleaning the wells twice with a washing buffer (250 µL). After the cleaning step, a solution of 100 µl of substrate/chromogen was added to all wells, mixed gently, and incubated for 15 minutes in the room temperature (20-25 °C). Lastly, 100 µl of the stop solution was added into all wells and the absorbance was measured at 450 nm in ELISA plate reader (ELX800 UV, Biotek, USA).

Absorbance Determination

The absorbance values of the standards, and samples tested were calculated according to the manufacture's instruction. Briefly, absorbance of the standards, and samples measured were divided by the absorbance value of the first standard (zero standard) and multiplied by 100:

$$\% \text{ absorbance} = \frac{\text{absorbance standard/sample}}{\text{absorbance zero standard}} \times 100$$

The zero standard is created equal to 100 % and the absorbance values were stated in percentages. The calculated values of all standards were typed into a system of coordinates on semilogarithmic graph paper against the concentration (µg/kg) of aflatoxin B₁. The aflatoxin B₁ concentration in µg/Kg matching to the absorbance of each sample established was read from the standardization curve correspondingly to the test preparation record for aflatoxin B₁.

Data Analysis

The data were entered in Microsoft Excel Data Analysis for statistical analysis. Phylogenetic trees based on Maximum Likelihood were constructed based on the best model using MEGA7. Graphical representation of the data was displayed in Graph Pad prism version 7.0. Analysis of variance (ANOVA) was used to compare the means and Tukey's HSD test (P≤0.05) used for mean separations, IBM® SPSS® Statistics 25.0 was used to determine the difference between levels of aflatoxin B₁ in rice grains and maximum limits levels permitted in Kenya and European Union. The difference between the aflatoxin mean in local and imported rice samples was compared by independent t-test. P<0.05 was considered as a significance.

3. Results

The Maximum Composite Likelihood (MCL) tree was constructed for the *aflD* gene sequence to describe the phylogenetic relationships among the aflatoxigenic strains, as shown in Figure 1. The tree with the highest log likelihood (-681.12) is shown. Sequencing and phylogenetic analysis of the *aflD* gene revealed that *Aspergillus flavus* strains clustered together with strains of *A. parasiticus*, *A. nomius*, and *A. fumigatus* (Figure 1). Reference strains of *A. flavus*, *A. parasiticus*, and *A. nomius* were grouped within the same clade. However, *A. fumigatus* (4A5) did not form a distinct cluster with strains of *A. parasiticus* and *A. nomius*. Overall, the aflatoxigenic strains analyzed in this study clustered within the same clade based on the *aflD* gene, which plays a critical role in converting norsolorinic acid to averantin during the intermediate stages of the aflatoxin biosynthetic pathway, rather than based on their source of isolation.




Figure 1. Molecular Phylogenetic analysis by Maximum Likelihood method showing the phylogenetic relationships among the aflatoxigenic strains based on *aflD* gene sequence. The percentage of trees in which the associated taxa clustered is shown above the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and

BioNJ algorithms to a matrix of pairwise distances estimated using the MCL approach and then selecting the topology with superior log likelihood value

Analysis of Aflatoxin Gene Expression

Four *Aspergillus* species were analyzed for the expression of the *aflD* gene, a key gene in the biosynthetic pathway of aflatoxin production. Notably, the relatively high expression levels of the *aflD* gene were observed in three strains: *A. flavus*, *A. parasiticus*, and *A. nomius*. Figure 2-A presents the mean expression levels of the *aflD* gene across these aflatoxigenic strains, with *A. flavus* showing a significantly higher mean expression, as determined by real-time PCR quantification. Figure 2- B illustrates a boxplot analysis of the relative *aflD* expression levels for all tested strains, highlighting variability among the species. Amplification curves for the *aflD* gene, as shown in Figure 2- C, confirm the specificity of the reaction by plotting fluorescence intensity against cycle number. To ensure the reaction's specificity, melting curve analysis was performed, plotting fluorescence against temperature (Figure 2- D).




Figure 2. Real-time PCR amplifications of *aflD* gene targeted in different aflatoxigenic strains: A, Graphical representation of the relative gene expression levels of the *norsolorinic acid* gene *nor-1*; B, Boxplot analysis of the relative expression of *aflD* gene; C, Relative fluorescence against cycle number; D, Melting curve analysis against temperature.

Quantitative Determination of Aflatoxin B₁ Levels in Rice Grains

Forty-four (44) samples, including local and imported rice grains (25 and 19, respectively) were analyzed for aflatoxin B₁ content (quantitatively) using an ELISA kit. Figure 3, presents the standard curve of aflatoxin B₁ generated from the absorbance values of the standards and samples in the system. An exponential correlation (Figure 3) was obtained by plotting the percentage of absorbance (y) and concentration (x) of aflatoxin B₁ ($R^2 = 0.96061$) of the standard calibration samples supplied by the manufacturer (RIDASCREEN® Aflatoxin B₁ 30/15). The determination of the aflatoxin B₁ concentration (µg/kg) corresponding to the absorbance of each sample was recited from the calibration curve according to the test preparation record. The detection limit was 1 µg/kg, and the recovery rate of 100 %. The recovery rate of less than 100 % shows more than 1 µg/kg of aflatoxin B₁. Values that exceeded the legal threshold of 2.0 µg/kg were considered unfit for consumption.




Figure 3. Standard curve of aflatoxin B₁ ($R^2 = 0.96061$).

The result revealed that 19 samples of surveyed rice grains had been contaminated by aflatoxin B₁, as shown in Table 1. Quantitative analysis reveals that 43.1 % of inspected samples were found positive, of which 31.8 % were under permitted levels. Nevertheless, 11.3 % were surpassing the accepted levels (table 3). The result of the present study demonstrates that the concentration of aflatoxin B₁ in tested rice grain samples extended from 2 µg/kg to 3.2 µg/kg using ELISA assay. The AFB₁ concentration average in the imported rice grains was higher than the findings in the local rice grains. However, no significant difference was observed in the AFB₁ concentration mean in samples collected ($P < 0.05$).

Table 3. Occurrence and level of aflatoxin B₁ in rice grains

Source of Sample	№ of Samples	Sample Status		Concentration of Aflatoxin B ₁ (µg/Kg)			Permitted levels <2µg/Kg (Commission Regulation No. 1881/2006)
		Positive	Negative	0	<2	>2	
Local	25 (56.8%)	07 (15.9%)	18 (40.9%)	17 (38.6%)	06 (13.6%)	02 (4.5%)	
Imported	19 (43.1%)	12 (27.2%)	07 (15.9%)	08 (18.2%)	08 (18.2%)	03 (6.8%)	
Total	44 (99.999%)	19 (43.1%)	25 (56.8%)	25 (56.8%)	14 (31.8%)	05 (11.3%)	

4. Discussion

In this study, genomic DNA for the norsolorinic acid gene region of aflatoxigenic strains, which is involved in the middle of the aflatoxin biosynthetic pathway, was amplified and sequenced. A comparison of *aflD* gene region sequences with aflatoxigenic strain sequences deposited in the GenBank showed a high similarity in the aflatoxigenic strains isolated. Alignment of selected isolates showed a high evolutionary divergence. A close relationship was observed between *A. flavus*, *A. parasiticus* and *A. nomius* isolates, while *A. fumigatus* was distantly related to the *A. parasiticus* and *A. nomius* isolates. The isolates of *A. flavus* strains showed close evolutionary relationships, with all isolates clustering (Figure 4). The conservation in *A. flavus*, *A. parasiticus* and *A. nomius* may be related to the functional importance of the *aflD* gene, as the first stable precursor in the aflatoxin biosynthetic pathway, encodes an enzyme responsible for catalyzing the conversion of norsolorinic acid, the initial stable intermediate, into averantin across all aflatoxigenic strains [50; 51].

Gene expression analysis showed variations, with three of four aflatoxigenic strains tested for *aflD* gene expression. We detected significantly higher gene expression stages of *aflD* in *A. flavus* than *A. parasiticus* and *A. nomius* strains. This may be linked to the aflatoxin toxicity in *A. flavus*, as reported by [52, 53]. *A. parasiticus* showed a weak expression level, followed by *A. nomius*. However, the detection of *aflD* in these strains confirms that they can equally be potential aflatoxin producers. Nevertheless, in contrast to these 3 strains in which relative gene expression levels were detected, we did not detect *aflD* in *A. fumigatus*. The relative fluorescence showed more concentration of RNA in 3 strains for each single curve observed (Figure 2-C). The amplification curves for all strains were specific and showed a single peak by melting curve analysis, as shown in Figure 2-D.

Further, we quantified aflatoxin B₁ levels in rice grains of 44 samples taken from local retail markets and millers. 43.1 % of tested samples were positive for aflatoxin B₁, with 15.9 % of contamination for local rice grains and 27.2 % of pollution in imported rice. Seven of 25 rice

samples homegrown from Mwea contained aflatoxin B₁, of which 13.6 % were below and 4.5 % above the permitted level. This observation might suggest a lack of suitable grain management or procurement of low-quality rice grains from growers who do not observe appropriate post-harvest techniques. This is in line with the research led in Turkey on the level and incidence of mycotoxin in rice [54]. Quantification of aflatoxin B₁ by ELISA assay showed a higher significance incidence in imported rice samples; 27.2 % of imported rice samples were contaminated with aflatoxin B₁ in which 6.8 % of those samples have greater levels than the 2.0 µg/kg limit established by EU. These observations agree with others [55; 56; 57] who surveyed the incidence of aflatoxin B₁ in rice grains. The highest proportion of contamination from imported rice could be explained by the storage, shipment and transport conditions along the way up to the destination point. Our findings concur with those reported by Nurshad [58], who investigated the incidence of aflatoxins in rice worldwide and communal health appearances from 1990 to 2015. In our study, the results displayed that the concentration of AFB₁ in rice grains analyzed ranges from < 2.0 µg/kg to 3.2 µg/kg. These results concur with those described by Sangare-Tigori *et al.* [56] and Anthony *et al.* [60] in West Africa (Ivory Coast and Nigeria), and by Madbouly *et al.* [61] in Egypt, which surveyed some mycotoxins and specially AFB₁ in rice grains with the range of <1.5-10.0 µg/kg, and 4.1-309.0 µg/kg, and with the limits of detection 11.0 µg/kg respectively. Here, we have established evidence of the occurrence of aflatoxin B₁ in African staple cereal, highlighting the need for new guidelines and standards to safeguard the security of consumers of rice grains. The countries in Sub-Saharan Africa must endeavour to establish policies that can guide regular empirical food quality assessments, as this would enhance and guarantee a healthy population with better standards of living and sustained development for prosperity. Acceptance of assimilated strategies across different points of the value chain to avoid human exposure to aflatoxins in consumed rice grains is important.

Conclusion

Based on molecular techniques, our analyses revealed the three aflatoxigenic strains *A. flavus*, *A. parasiticus*, and *A. nomius* are closely clustered within the same clade, suggesting the *aflD* gene plays a crucial role in regulating the biosynthetic pathway of aflatoxin production in aflatoxigenic species. The concentration of AFB₁ in the analyzed samples ranged from 0 µg/kg to 3.2 µg/kg, with 43.1% of samples containing detectable levels. Notably, 11.3% of these samples exceeded the maximum tolerable limit for aflatoxins in rice, as set by the European Union (EU), which is 2 µg/kg for AFB₁. These findings highlight the potential

exposure of humans, animals, and livestock to aflatoxin B₁, posing significant health risks and leading to indirect economic impacts on society. This underscores the importance of educating the public on proper grain handling-practices, especially during and after harvesting. Additionally, the study affirms the efficiency of ELISA method in detecting mycotoxins in cereals, offering advantages such as simplicity and reduced time requirements. Continuous monitoring of foodstuffs and agricultural products is essential to mitigate the risks associated with aflatoxin contamination. Such surveillance can help prevent critical issues in the marketing, distributing, and consuming contaminated food products.

Key Contribution: Aflatoxin contamination of cereals poses a serious concern for human health worldwide. This article highlights aflatoxigenic strains, gene expression profile and aflatoxin B₁ levels in rice grains to ensure food safety and security.

References

1. Pangga IB, Salvacion AR and Cumagun CJR. Climate change and plant diseases caused by mycotoxigenic fungi: Implications for food security. In *Climate Change and Mycotoxins; De Gruyter: Berlin, Germany*, (2015); pp. 1-28. doi: [10.1515/9783110333619-003](https://doi.org/10.1515/9783110333619-003)
2. Moretti AT, Logrieco AF and Susca A. Mycotoxin: An underhand food problem. In *Mycotoxigenic Fungi Methods and Protocols; Moretti A, Susca A, Eds.; Humana Press: New York, NY, USA. Methods in molecular biology*, (2017); pp. 3-12. doi: 10.1007/978-1-4939-6707-0_1.
3. Gacem MA and El Hadj-Khelil AO. Toxicology, biosynthesis, biocontrol of aflatoxin and new methods of detection. *Asian Pacific Journal of Tropical Biomedicine*, (2016);6, 808-814. <https://doi.org/10.1016/j.apjtb.2016.07.012>
4. Udovicki B, Audenaert K, De Saeger S and Rajkovic A. Overview on the mycotoxin's incidence in Serbia in the period 2004-2016. *Toxins*, (2018);10, 279. doi: [10.3390/toxins10070279](https://doi.org/10.3390/toxins10070279)
5. Benkerroum, N. Retrospective and prospective look at aflatoxin research and development from a practical standpoint. *International Journal of Environmental Research and Public Health*, (2019);16, 3633.
6. Winter, G. and Pereg, L. A review on the relation between soil and mycotoxins: Effects of aflatoxin on field, food and finance. *European Journal of Soil Science*, (2019);70, 882-897.

7. Mitchell NJ, Bowers E, Hurburgh C, Wu F. Potential economic losses to the US corn industry from aflatoxin contamination. *Food Addit. Contam. Part A*, 33, (2016); 540-550. doi: 10.1080/19440049.2016.1138545
8. Ostry V, Malir F, Toman J, Grosse Y. Mycotoxins as human carcinogens-the IARC Monographs classification. *Mycotoxin Res.* 33, (2017); 65-73. doi: 10.1007/s12550-016-0265-7
9. Caceres, I., Al Khoury, A., El Khoury, R., Lorber, S., Oswald I, P., El Khoury, A., et al., 2020. Aflatoxin biosynthesis and genetic regulation: a review. *Toxins* 12 (3), 150. <https://doi.org/10.3390/toxins12030150>.
10. Frisvad JC, Hubka V, Ezekiel CN, Hong SB, Nováková A., *et al.* Taxonomy of *Aspergillus* section *Flavi* and their production of aflatoxins, ochratoxins and other mycotoxins. *Stud. Mycol.* (2019); 93:1-63. doi: 10.1016/j.simyco.2018.06.001
11. Theumer, M. G., Y. Henneb, L. Khoury, S. P. Snini, S. Tadrict, C. Canlet, and M. Audebert. Genotoxicity of aflatoxins and their precursors in human cells. *Toxicol. Lett.* (2018); 287:100–107. <https://doi.org/10.1016/j.toxlet.2018.02.007>
12. Keller, N. P. Fungal secondary metabolism: regulation, function and drug discovery. *Nature Reviews Microbiology*, (2019); 17, 167-180.
13. Cao, W., Yu, P., Yang, K., Cao, D. Aflatoxin B1: metabolism, toxicology, and its involvement in oxidative stress and cancer development. *Toxicol. Mech. Methods* 32 (6), (2022); 395–419. <https://doi.org/10.1080/15376516.2021.2021339>
14. Marchese, S., Polo, A., Ariano, A., Velotto, S., Costantini, S., Severino, L. Aflatoxin B1 and M1: biological properties and their involvement in cancer development. *Toxins* 10 (2018); (6), 214. <https://doi.org/10.3390/toxins10060214>.
15. Lee HJ, Ryu D. Occurrence of Mycotoxins in Cereals and Cereal-Derived Food Products: Public Health Perspectives of Their Co-occurrence. *J. Agric. Food Chem.* (2017); doi: 10.1021/acs.jafc.6b04847
16. Bennett JW, Moore G. Mycotoxins. In: *Encyclopedia of Microbiology*. Oliver Walter, (2019); 267–273.
17. Adhikari BN, Bandyopadhyay R and Cotty P J. Degeneration of aflatoxin gene clusters in *Aspergillus flavus* from Africa and North America. *AMB Express* (2016); 6:62. doi: 10.1186/s13568-016-0228-6
18. Ehrlich KC, Yu J, Cotty PJ. Aflatoxin biosynthesis gene clusters and flanking regions. *J Appl Microbiol.* (2005); <https://doi.org/10.1111/j.1365-2672.2005.02637.x>

19. Yu J, Fedorova ND, Montalbano BG, Bhatnagar D, Cleveland TE, Bennett JW, Nierman WC. Tight control of mycotoxin bio-synthesis gene expression in *Aspergillus flavus* by temperature as revealed by RNA-Seq. *FEMS Microbiol Lett.* (2011);<https://doi.org/10.1111/j.1574-6968.2011.02345.x>
20. Bennett JW. Loss of norsolorinic acid and aflatoxin production by a mutant of *Aspergillus parasiticus*. *J Gen Microbiol.* (1981); <https://doi.org/10.1099/00221287-124-2-429>
21. Trail F, Chang PK, Cary J, Lin JE. Structural and functional analysis of the nor-1 gene involved in the biosynthesis of aflatoxins by *Aspergillus parasiticus*. *Appl Environ Microbiol*(1994); <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC201939/pdf/aem00028-0188.pdf>. Accessed 12 Jan 2018.
22. Yu J. Current understanding on aflatoxin biosynthesis and future perspective in reducing aflatoxin contamination. *Toxins.* (2012); <https://doi.org/10.3390/toxins4111024>.
23. Alkhayyat F and Yu JH. Upstream regulation of mycotoxin biosynthesis. *Advances and Applied Microbiology*, 86, (2014);251-278.doi: 10.1016/b978-0-12-800262-9.00005-6
24. Gil-Serna J, Vázquez C and Patiño B. Genetic regulation of aflatoxin, ochratoxin A, trichothecene, and fumonisin biosynthesis. *International Microbiology*, 23, (2019);89-96. doi: 10.1007/s10123-019-00084-2
25. Ministry of Agriculture. Economic Review of Agriculture in Kenya 2010. (2010);*Nairobi*.
26. Iqbal, S. Z., M. R. Asi, U. Hanif, M. Zuber, and S. Jinap. The presence of aflatoxins and ochratoxin A in rice and rice products; and evaluation of dietary intake. *Food Chem.* (2016); 210: 135–140. <https://doi.org/10.1016/j.foodchem.2016.04.104>
27. Ali, N. Aflatoxins in rice: worldwide occurrence and public health perspectives. *Toxicol. Rep.* (2019); 6:1188–1197. <https://doi.org/10.1016/j.toxrep.2019.11.007>
28. N. S. Al-Zoreky and F. A. Saleh, “Limited survey on aflatoxin contamination in rice,” *Saudi J. Biol. Sci.*, vol. 26, no. 2, (2019);pp. 225–231doi: 10.1016/j.sjbs.2017.05.010
29. European Commission. Commission regulation (EC) No. 1881/ 2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Off J Eur Union L* (2006); 364:5-24.<http://faolex.fao.org/docs/pdf/eur68134.pdf>
30. Iqbal, J., M. A. Asghar, A. Ahmed, M. A. Khan, and K. Jamil. Aflatoxins contamination in Pakistani brown rice: a comparison of TLC, HPLC, LC-MS/MS and ELISA techniques. *Toxicol. Mech.*(2014); 24:544–551. <https://doi.org/10.3109/15376516.2014.948247>.
31. Adi, P. J., and Matcha, B. Analysis of aflatoxin B₁ in contaminated feed, media, and serum samples of *Cyprinus carpio* L. by high-performance liquid chromatography. *Food Quality and Safety*,(2018);2, 199-204.

32. Shabeer S, Asad S, Jamal A, Ali A. Aflatoxin contamination, its impact and management strategies: An updated review. *Toxins (Basel)*. (2022); 14(5): 307.
33. Sakamoto, S., W. Putalun, S. Vimolmangkang, W. Phoolcharoen, Y. Shoyama, H. Tanaka, and S. Morimoto. Enzyme-linked immunosorbent assay for the quantitative/qualitative analysis of plant secondary metabolites. *J. Nat. Med.* (2018); 72:32–42. <https://doi.org/10.1007/s11418-017-1144-z>.
34. Mutiga SK, Were V, Hoffmann V, Harvey JW, Milgroom MG, Nelson RJ. Extent and drivers of mycotoxin contamination: Inferences from a survey of Kenyan maize mills. *Phytopathology*. (2014); **104**(11):1221-1231. DOI: 10.1094/phyto-01-14-0006-r.
35. Ngindu A, Johnson BK, Kenje PR, Ngira JA, Ochieng DM, Nandwa H, et al. Outbreak of acute aflatoxicosis caused by aflatoxin poisoning in Kenya. *Lancet*. (1982); **1**:1346-1348.
36. European Commission. Commission regulation (EC), No. 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Official Journal of the European Union*, (2006); L364, 5-24.
37. Pitt, J. I. and Hocking, A. D. Fungi and food spoilage; Blackie Academic and Professional, Cambridge, (1997); UK. *Food Science and Nutrition*.
38. Atlas, R. M. Handbook of Microbiological Media; CRC Press. Boca Raton, (2010); FL, USA.
39. Pitt, J. I. and Hocking, A. D. Fungi and Food Spoilage, 3rd ed, Springer: New York, NY, USA. *Advances in Microbiology*, (2009); 8, 4, 519.
40. Klich, M.A. Identification of Common *Aspergillus* Species; Centraalbureau voor Schimmelcultures: Utrecht, The Netherlands, (2002).
41. Varga, J., Frisvad, J. C. and Samson, R. A. Two new aflatoxin producing species and an overview of *Aspergillus* section *Flavi*. *Studies in Mycology*, (2011); **69**:57-80.
42. Cenis, J. L. Rapid extraction of fungal DNA for PCR amplification. *Nucleic Acids Research*, (1992); 20, 2380.
43. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* (2016); 33(7):1870-4.
44. Tamura, K. and Nei, M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, (1993); 10, 512-526.
45. Yu J, Chang PK, Ehrlich KC, Cary JW, Bhatnagar D, Cleveland TE, Gary AP, Linz JE, Woloshuk CP and Joan WB. Clustered Pathway Genes in Aflatoxin Biosynthesis. *Applied*

- and Environmental Microbiology*, 70, (2004); 1253-1262.doi: [10.1128/AEM.70.3.1253-1262.2004](https://doi.org/10.1128/AEM.70.3.1253-1262.2004)
46. Pfaffl MW. A new mathematical model for relative quantification in real time RT-PCR. *Nucleic Acids Research*, (2001); 29, 9, e45.doi: 10.1093/nar/29.9.e45
 47. Schmittgen TD and Livak KJ. Analyzing real-time PCR data by the comparative Ct method. *Nature Protocols*, 3, (2008); 1101-1108.doi: 10.1038/nprot.2008.73
 48. Mohamadi SA, Gholampour AE, Salehi AE, Rahimi K. Reduction of aflatoxin in rice by different cooking methods. *Toxicol Ind Health*(2014),30:546-50.doi: 10.1177/0748233712462466.
 49. R-Biopharm GmbH. Enzyme immunoassay for the quantitative analysis of aflatoxin B₁ Art. No.: R1211. (2010);Darmstadt, Germany: R-Biopharm GmbH.
 50. Afzaal S, Ali SW, Hameed U, Ahmad A, Akhlaq M, Haider MS Molecular probing of Aflatoxigenic fungi in rice grains collected from local markets of Lahore, Pakistan. *Adv. Life Sci.* 8(2): (2021);190-194.
 51. Kjærboelling, I. et al. A comparative genomics study of 23 *Aspergillus* species from section *Flavi*. *Nat. Commun.* (2020); 11, 1106.
 52. Shabeer, S.; Asad, S.; Jamal,A.; Ali, A. Aflatoxin Contamination,Its Impact and ManagementStrategies: An Updated Review.*Toxins* (2022); 14, 307. <https://doi.org/10.3390/toxins14050307>.
 53. Accinelli C, Abbas HK, Zablutowicz RM and Wilkinson JR. *Aspergillus flavus* aflatoxin occurrence and expression of aflatoxin biosynthesis genes in soil. *Canadian Journal of Microbiology*, 54, (2008); 371-379.doi: 10.1139/w08-018
 54. Aydin A, Aksu H and Gunsen U. Mycotoxin levels and incidence of mould in Turkish rice. *Environmental Monitoring and Assessment*, 178, (2011); 271-280.doi:10.1007/s10661-010-1688-9
 55. Alwan, N.; Bou Ghanem,H.; Dimassi, H.; Karam, L.; Hassan,H.F. Exposure Assessment ofAflatoxin B1 through Consumptionof Rice in the United Arab Emirates.*Int. J. Environ. Res. Public Health* (2022);19, 15000. <https://doi.org/10.3390/ijerph192215000>
 56. Hussein, F. H.; Rita, K.; Hani, D.; Andre, E.; Rouaa, D.; Nisreen, A.; Samar, M.; Joyce, H. And Layal, K. Aflatoxin B1 in Rice: Effects of Storage Duration, Grain Type andSize, Production Site, and Season. *Journal of Food Protection*, Vol. 85, No. 6, (2022); Pages 938–944<https://doi.org/10.4315/JFP-21-434>

57. Phan, L. T. K; De Saeger, S. ; Eeckhout, M.; Jacxsens, L. Public health risk due to aflatoxin and fumonisin contamination in rice in the Mekong Delta, Vietnam. *Food Safety and Risk* (2023); 10:4 <https://doi.org/10.1186/s40550-023-00104-0>
58. Nurshad Ali. Aflatoxins in rice: Worldwide occurrence and public health perspectives. *Toxicology Reports* 6, (2019); 1188-1197. <https://doi.org/10.1016/j.toxrep.2019.11.007>
59. Sangare-Tigori B, Moukha S, Kouadio HJ, *et al.* Co-occurrence of aflatoxin B1, fumonisin B1, ochratoxin A and zearalenone in cereals and peanuts from Cote d' Ivoire. *Food Addit Contam* (2006); 23:1000-1007. doi: 10.1080/02652030500415686
60. Anthony MH, Dutton FM, Njobeh BP *et al.* Natural multioccurrence of mycotoxins in rice from Niger State, Nigeria. *Mycotoxin Res* (2011); 27:97-104. doi: [10.1007/s12550-010-0080-5](https://doi.org/10.1007/s12550-010-0080-5)
61. Madbouly AK, Ibrahim MIM, Sehab AF, Abdel-Wahhab MA. Co-occurrence of mycoflora, aflatoxins and fumonisins in maize and rice seeds from markets of different districts in Cairo, Egypt, *Food Addit. Contam.: Part B* 5, (2012); 112-120. doi: 10.1080/19393210.2012.676078