

Review Article

An overview on *Amaranthus cruentus* L.- an alternative source of crop, its economical uses and the associated beneficial microbes

ABSTRACT

Amaranthus cruentus L. is a medicinally important locally available tropical plant belonging to the family Amaranthaceae. Now a days, this plant is designated as 'Alternative source of crop' or 'Pseudocereal' having various nutritional properties (Protein, carbohydrate, fatty acid, Vitamins, Minerals). Moreover, this plant has various medicinal properties also i.e. antioxidant, antihypertensive, antidiabetic, immune stimulation, anthelmintic, hepatoprotective, antimalarial etc. This plant has Worldwide distribution and short growing period of time. It is also regarded as Fodder crop and other industrial uses. The microbes associated with this plant help to protect the host plants from both pathogenic microorganisms and adverse climatic conditions by secreting different types of bioactive secondary metabolites i.e. phenolics, flavonoids, terpenoids, terpenes, tannins, alkaloids, quinones, anthocyanin, saponin, steroids and promote growth and development of the plant. Some unique features of this plant are also described here. This review intends to expand the understanding of taxonomic characteristics of this plant, economic uses, associated beneficial microbes and their application for sustainability of this alternative source of crop.

Keywords: *Amaranthus cruentus*, alternative source of crop, secondary metabolites, microbes.

1. INTRODUCTION

Amaranthus cruentus is a tropical, endemic, invasive, short lived perennial herbaceous plant in Africa and South America (Gresta et al., 2020). Nevertheless, now this plant is distributed all over the world and contains approximately 70 species (Gresta et al., 2020). It is cultivated mainly in tropical and temperate region (Letchamo et al., 2017). This species has greater variability (Ma et al., 2021; Gins et al., 2024). Phenotypic plasticity, Introgression and hybridization are profoundly observed in this species (Gresta et al., 2020; Prajitha & Thoppil, 2018; Jimoh et al., 2022; Amosova et al., 2024). To protect from pathogenic microorganism, this plant has some advanced physiological features (Jimoh et al., 2022). In Africa, this plant is recognized as source of food and constructive and traditional medicine (Letchamo et al., 2017). Now a days, food scarcity is a major problem and food security is necessary to protect the future generations. So, this plant is used as 'Alternative source of crop' and also considered as 'Pseudocereals' (Fletcher, 2015; Mburu, 2011; Gresta et al., 2020). Moreover, this plant is cultivated for using both the Grains and leaves in Africa and Asia (Hoidal et al., 2019). It has various medicinal principles i.e. antioxidant, anti-diabetic, anti-inflammatory properties, antihypertensive properties, stimulation of immune system, anthelmintic properties (Torane et al., 2017) etc. This plant harbor endophytic and others microflora. Endophytic microbes secrete active secondary metabolites i.e. phenolics, flavonoids, terpenoids, terpenes, tannins, alkaloids, quinones, anthocyanin, saponin, steroid etc. These metabolites are potential agents for antimicrobial, anticancerous and many other therapeutic properties which are beneficial to protect the host plant from both biotic and abiotic stresses (Mousavi & Karami, 2022). This plant is an example of C4 photosynthetic ones (Achigan et al., 2014; Gins et al., 2024) and reported as high-water using efficiency (WUE) (Wolosik & Markowska, 2019) and grow in any types of soil i.e. acidic, alkaline, saline or nutrient-poor soil (Park et al., 2014). This species is highly resistant to drought (Gresta et al., 2020; Achigan et al., 2014) and salinity (Gins et al., 2024; Emam et al., 2024) heat, oxidative stress (Jimoh et al., 2022). and UV radiation also (Castrillón & Frier, 2016 and; Omami & Hammes, 2006). It has very good nutritional properties as it contains vitamins, proteins (Lysine, Tryptophan), unsaturated fatty acids, minerals, tocopherols and tocotrienol etc. and essential antioxidant properties (Wolosik & Markowska, 2019). Light or temperature (20 -35°C) and combination of light and temperature promote higher growth and germination (Wolosik & Markowska, 2019). Its short growing period give expectancy and provide a source of employment to the farmers (Makinde & Ojieniyi, 2010). According to genomic studies, *A. cruentus* contains 17 chromosomes (Sammour & Mitra, 2012). Previous report shows that it is hybridized easily (Lanta & Ondrej, 2003). In South Africa, it is considered as potential alternative cereal having minerals, phytochemicals (Manyelo et al., 2022). Several species of *Amaranthus* have a role in CO₂ sequestration (Jimoh et al., 2022; Malik et al., 2023). This is short day plant and flowering takes place in 44 days and harvested in 102 days (Ogwu, 2020). Microbes present

Comment [U1]: Why you have chosen to make a review on this species only while they are numerous species under the genus?

Comment [U2]: The native region of that species, How many species under amaranthus genus? What is the particularity of your selected species compared to others of the same genus ?

Comment [U3]: For which techniques Africa and Asia applied to cultivate *A. cruentus* using their leaves ?

Comment [U4]: Precise the quantity?

Comment [U5]: No logic connection with previous content.

Comment [U6]: Explain

in the plant significantly help the host to increase growth, protect from both biotic and abiotic stresses and also stimulate the medicinal activities without causing any disturbance or damage. In this review, taxonomic characteristics, secondary metabolites, economic importance of this plant and lastly the microbes (PGPR, PGPF, fungal and bacterial Endophytes and Mycorrhizae) associated with this plant and how they promote the plant growth is discussed here.

2.TAXONMIC DESCRIPTION OF AMARANTHUS CRUENTUS L.

2.1Taxonomic classification: (Wolosik&Markowska,2019)

Kingdom: Plantae

Subkingdom: Tracheobionta

Subdivision: Spermatophyta

Division: Magnoliophyta

Class: Magnoliopsida

Subclass: Caryophyllidae

Order: Caryophyllales

Family: Amaranthaceae

Genus: *Amaranthus*

Species: *Amaranthuscruentus* L.

2.2Habit:

This plant is herbaceous, tropical and belongs to the family Amaranthaceae. The common name of the plant is red or blood amaranth. In Bengali, it is popularly known as 'Lal note sak'. It reproduces by seeds and have very short (4-6 weeks) growth period (Wolosik&Markowska,2019).

2.3Root: one dominant large tap root, soft, 6-7 cm long (Wolosik&Markowska,2019;Mukuwapasi et al., 2024).

2.4Stem: straight, branched, thick, ribbed, 60-65 cm in height, reddish in colour, surface minutely haired (Wolosik&Markowska,2019).

2.5Leaves: simple, reddish, 5-7.5 cm length and 2.1-4.2 cm breadth, margin entire, spiral arrangement, shape ovoid, exstipulate, leaf surface hairy. Petiolate, petiole 0.5 to 1 cm long, leaf apex retuse (Wolosik&Markowska,2019; Mukuwapasi et al., 2024).

2.6Inflorescence: Numerous concentrated spikes, axillary arrangement, ended with racemes and spikes. More than 50 cm long, reddish or pinkish colour. Branches are 40-45 cm long, perpendicular with numerous laterals. Monoecious bearing both male and female flower (Wolosik&Markowska,2019).

2.7Flower: Numerous unisexual flowers are greenish and form a terminal spike (Wolosik&Markowska,2019).

2.8Seeds: This plant produces thousands of seeds which are round or lenticular, dark brown or shiny. Campylotropus ovule (Wolosik&Markowska,2019). Endosperm present in the micropylar region (Wolosik&Markowska,2019).

Dissected parts of the plant under the simple microscope is represented in figure 1.

Comment [U7]: Summarize it in one point , for example physical characteristics / plant organs or phenological characteristics




Figure 1. Morphological characters of *Amaranthus cruentus* (where A, B, C, D, E, F, G, H represent dorsal and ventral surface of the leaf, root morphology, Dissection of female flower, male flower)

3.ECONOMICAL USES

3.1 Food: *Amaranthus cruentus* is a leafy vegetable (Torane et al., 2017; Mukuwapasiet al., 2024) and useful food supplement for human. This vegetable is similar to the well-known Barley, Wheat and Bazra crop for its nutritional properties and contain high amount of amino acids. So, this plant is considered as 'Alternative Source of Crop' or 'Pseudocereal' (Fletcher 2015; Mburu et al., 2011) and it protects the disease Malnutrition or Undernutrition. It has essential nutritional properties i.e. amino acid glutamine (Motta et al., 2019) proteins (Lysine and Tryptophan), vitamins, essential minerals, saturated and unsaturated fatty acid (Gresta et al., 2020), tocopherol, tocotrienol etc. and also contain high amount of fiber (Ali 2021). Powdery dry leaves are used in sauces during the dry weather (Aderibigbe et al., 2022; Torane et al., 2017). In Nigeria, leaves are mixed with the condiments to prepare the soups (Mepba & Banigo, 2007; Ali, 2021). Boiled leaves are mixed with groundnut flower and consumed as relish (Ali, 2021). Leaves are consumed as spinach or green vegetable (Dhellit et al., 2006). Mixture of boiled leaves and groundnut sauce are prepared for salad in West Africa and Mozambique (Kauffman and Weber, 1990). Seeds are also used as nutritious food which enriched in minerals, vitamins and proteins (Emam et al., 2024; Mukuwapasi et al., 2024).

3.2 Fodder: This plant is used not only for food but also as fodder. It has been planted as leafy vegetables for cattle, rabbit, pigs and chicken food across many countries. Dietary fiber present in the plant have positive impact on animal gut by improving behavior on digestive tract (Manyelo et al., 2022; Alegbejo 2013).

3.3 Cosmetics Industry: Amaranth oil contains high amount of unsaturated fatty acid, tocopherols, Squalene which act as hair and skin conditioner (Baraniak & Kania-Dobrowolska 2022). This seed oil also cures aging effect, wounds, skin diseases and moistening the skin. It is efficiently used as strong antioxidant and photoprotector in herbal medicine industry (Baraniak & Kania-Dobrowolska, 2022).

It is also used as lubricant in computer industry (Achigan et al., 2014).

3.4 Dye: In some countries, this plant is cultivated or grown only for making different dyes (Torane et al., 2017) Dried plant is burnt for the preparation of potash in Benin countries (Torane et al., 2017). Plants are used to extract red dye or coloring corn-based foods in South Western United states (Allemann, 2016).

3.5 Ornamental value: *A. cruentus* is cultivated as an ornamental plant in some countries (Torane et al., 2017) specially in Nigeria (Ogwu, 2020; Mukuwapasi et al., 2024) valued for its feather-like flowering plumules and also have aesthetic value also (Hricova et al., 2016). And sometimes it used as 'potherb' (Das & das, 2016). The *A. cruentus* is greatly valued as ornamental because of its bright colour flowers (Amosova et al., 2024).

Comment [U8]: You may also present a full plant with its parts

Comment [U9]: Could you search the people eat grains and how they prepare?

Comment [U10]: How used as fodder for cattles, rabbit, pigs, etc
What the contents quantity those domestic animals get from this plant?

Comment [U11]: How oil, antioxidant and others extracted from and preserved for future use?

Comment [U12]: Explore all applications of this red dye from *A. cruentus*.

3.6 Medicinal properties: It has various medicinal properties i.e. antioxidant (Osei et al., 2023), anti-inflammatory properties, antihypertensive properties stimulation of immune system, cardioprotective, antidiabetic, antimalarial, hepatoprotective (Peter&Gandhi,2017), anthelmintic (Torane et al., 2017), antiandrogenic (Alegbejo, 2013) etc. Bioactive secondary metabolites and protein hydrosylates present in the plant are potential anticancerous, antimicrobial agents (Ramkisson et al., 2020). According to report of Azab 2020, the cooked leaves of the plant have laxative properties. Seeds have some medicinal properties also i.e. hypocholesterolemic effect. Ethanolic leaf extract have protective effect such as restoring RBC, WBC and hemoglobin levels in blood (Azab, 2020). So, it is used in Anemia treatment. *A. cruentus* extract administrated to rats receive Cyclophosphamide which is an immunosuppressant to treat nephritis and arthritis and also used in Chemotherapy in combination with other drugs (Pandey et al., 2016). For infants, treatment of constipation, roots of this plant is boiled with honey and feed it (Aderibigbe et al., 2022). Its water extract is used for the treatment of limb pain and as expellant of tapeworm also for wound dressing and tumors (Aderibigbe et al., 2022). This leafy vegetable is also important for lactating women to treat kidney complaints, constipation, anemia etc. (Torane et al., 2017; Aderibigbe et al., 2022; Achigan et al., 2014) Presence of high amount of fiber reduces cholesterol levels as well as risk of cardiovascular diseases (Jimoh et al., 2022) and inhibits the intake of starchy food decreases the chances of diabetes (Adeleke&Babalola,2021). Prevention of diuretic and hypertension problem is revealed by potassium content (George, 2015). Oils or fats decreases the lipid content of blood and chances of coronary diseases have been reduced (Chatepa&Masamba,2020) Decoction of leaves is utilized as sore throat, ulcers and extraction of inflorescence swallowing by women for relieving from periodic pain (Ali et al., 2021). Ethanolic leaf extract of mixture of *Amaranthuscruentus* and *Cleomegynandra* have blood glucose lowering effect (Siwale ,2018).

Comment [U13]: Separate traditional methods from industrial preparations .

4. BIOACTIVE SECONDARY METABOLITES PRESENT IN THE PLANT

A number of different bioactive secondary metabolite present in the plant is graphically represented in Figure 2 (Peter&Gandhi,2017). which helps to promote plant growth and protect from pathogens. These metabolites also possess antioxidant, biochemical and pharmacological properties and beneficial for human health (Omale & Okafor,2008).




Figure2. Lists of Secondary Metabolites of red Amaranth (Peter&Gandhi,2017)

4.1 Betalines: Betacyanin (red violet pigment) and Betaxanthin (yellow pigment) present in different parts (root, stem, leaf, flower) of the plant (Castrillón&Frier, 2016), water soluble vacuolar pigment used in Folk medicine and also as anticancerous, anti-inflammatory and cardiovascular agent (Nana et al., 2012).

4.2 Phenolics: Anthocyanin produce excessive radicles which surpass the physiological antioxidant properties (Alvarez et al.,2010). Salicylic, cinnamic, ferulic, p-coumaric, gallic, sinapic acids are also present (Peter&Gandhi,2017;Castrillón & Frier, 2016). Premature grain of this plant contains a huge

Comment [U14]: Which quantity present in root, leaf... And how betalines tested and extracted from those parts ?

amount of phenolic acids i.e., ferulic acid, caffeic acid, galic acid, p-coumaric acid and anthocyanin) than mature grain (Manyelo et al., 2022, Osei et al., 2023). It has been reported that phenolics have beneficial role in antioxidative property serving as the free radical scavengers and lower the heart diseases and inflammation (Omale&Okafor,2008; Iwuagwu et al., 2019). Biosynthetic pathway of phenolics is diagrammatically represented in figure 3(Buchanon, 2015).

Comment [U15]: Precise the quantity of each phenolic type and the extraction method ?




Figure3. Overview of Phenylpropanoid pathway PAL- Phenylalanine ammonia lyase, C4H- Cinnamate-4- hydroxylase, 4CL- Coumarate CoA ligase. adapted from Buchanon (2015).

Comment [U16]: Explain the figure as working mechanisms of phenols

4.3Flavonoids: Rutin is one of most important flavonoids reported to have various beneficial effects on human and protective effect on memory disfunction (Koda et al., 2009). Quercetin is also reported. It serves as antimicrobial agent and showing potential antioxidant property (Castrillón&Frier, 2016) and reduce the oxidation level of triglyceride, cholesterol, low density lipoprotein (Erdtman et al., 2007).

4.4Terpenoids: Terpenoid showing antimicrobial, antifungal, antiparasitic, antiviral, anticancerous, antispasmodic, antiallergic, anti-inflammatory, immunomodulatory properties (Rabi&Bishayee,2009).

4.5Alkaloids: It is one of the important metabolites in living organisms help in development of the plant and served as useful repellent for any pathogen or parasite and possesses antimicrobial properties (Usumomena&Ngozi,2016;Cherian&Sheela,2016).

4.6Quinones: According to the result of phytochemical screening, ethanolic extract contains flavonoid, phenolics and quinones which are used in glucose tolerance test i.e., antihyperglycemia (Nofianti,2015).

4.7Saponin: It limits the growth of cancerous cells and has beneficial effects on antiinflammation (Prohp&Onoagbe,2012; Rao&Sung,1995) and stimulate release of insulin and lower blood glucose level (Siwale, 2018).

4.8Steroid: Steroid is present at vegetative and flowering stage which protect from harmful effects of microbial pathogens and also have anti-inflammatory and anti-aging properties (Okunlola et al., 2017). Amaranth grain and leaf extract contain high amount of Squalene (He&Corke,2003).

4.9Tannin: It has been reported to possess potential antiviral (Cheng& Lin, 2002) and anticancerous activity (Narayanan et al., 1999).

Comment [U17]: For all precise the quantity and the extraction methods.

5. BENEFICIAL MICROBIAL ASSOCIATION

5.1 Endophytes: Numerous studies have shown that symbiotic (mostly Endophytic) fungi enhance abiotic stress tolerance mainly water deficit i.e. Drought and protect the host plant from critical environmental condition and plant survive (Rodriguez et al., 2009; Singh et al., 2011). Endophytic microbes reside within living plant tissues and can interact with roots, stem and leaves without causing symptoms of diseases in their host plants (Schulz & Boyle, 2006). The endophyte infected plants express a range of adaptations to biotic stress (Moreli et al., 2020) and abiotic stresses mainly drought tolerance. Some previous reports demonstrated that many grass species showed vegetative growth improvement in the presence of their fungal symbionts and these endophytic fungi enlarge plant fitness by the discharge of plant growth advancing secondary metabolites (Priti et al., 2009; Hardoim et al., 2015). Endophytes have both the horizontal and vertical mode of transmission (Frank et al., 2017). Endophytic microbes are mainly bacteria and fungi and some reports revealed that it may be algae also (Rit et al., 2023). Fungal endophytes mainly belong to the Ascomycetes, Zygomycetes and Deuteromycetes group and Basidiomycetous fungi is rarely reported (Rajamanikyam et al., 2017; Stone et al., 2000). Endophytes showing mutualistic behaviour i.e., it receives nutrition and shelter from host plant and give the competitive ability to the host (Rajamanikyam et al., 2017). After isolation, pure culture and characterization of each specific endophyte, Maximum individual possessing antibiotics, anticancerous, antimycotics and immunosuppressant property (Alvin et al., 2014). Treatment of various types of endophytes upon *Amaranthus* plant showing enhancement of growth due to discharge of different plant growth regulator i.e. IAA, GA etc and increased SAR by decreasing diseases by direct antagonistic behaviour (Uppala et al., 2010). Fungi Isolated from Phyllosphere and Rhizosphere of *A. cruentus* (*Cladosporium*, *Talaromyces*, *Gibberella*, *Penicillium*, *Alternaria*, *Trichoderma*, *Rhizopus*, *Fusarium*) increase the soil quality of the plant and plants health and produce antibiotic compound against pathogens (Puszeta et al., 2015). According to Uppala et al., 2010, several bacterial endophytes promotes biometrics of Amaranth. Some endophytic bacteria isolated from plant roots and promote plant growth, survival and protect from biotic stress (Enebe & Babalola, 2018). Endophyte protects the host plant from oxidative damage by producing ROS scavenging enzymes i.e. Superoxide dismutase (SOD), Glutathione peroxidase (GR) etc. (Choudhury et al., 2023). It was reported in Red Amaranth that the application of an endophytic strain *Sphingomonas panaciterrae* not only increases the morphological parameter i.e., root length, shoot length, leaf number etc. of the host plant but also increase the chlorophyll and mineral content of the plant (Sultana et al., 2024).

Endophytic fungi are isolated (Shilpa & Shikha, 2015) from different parts of the plant *A. cruentus* is represented in figure 4.




Figure 4. Isolation of Endophyte from *A. cruentus* (Shilpa & Shikha, 2015)

Comment [U18]: Which fungi species associate with *A. cruentus*? How they symbiont

Comment [U19]: Rewrite the figure in the same writing of your manuscript

The plates containing pure culture of the isolates are represented in figure 5.




Figure 5. Fungal isolates after pure culture

According to standard protocol, leaf tissues of red Amaranth were dissolved in organic solution and used to detect endophytes under microscope after staining with Trypan blue.

5.2 PGPR and PGPF: Plant Growth Promoting Rhizobacteria (PGPR) are a group of bacteria that enhance phytoremediation (Muthukumar et al., 2017). It colonizes plant root surface efficiently and promote plant growth. PGPR may facilitate plant growth either by promoting the acquisition of nutritional resources such as Nitrogen, Phosphorus or Iron; preventing or decreasing the damage to plants by pathogenic organisms (mainly fungi and bacteria) or by directly stimulating plant growth by either providing plant hormones such as Auxin, Cytokinin or GA or lowering plant ethylene levels through the action of the enzymes 1 amino-cyclopropane 1 carboxylic acid deaminase (ACC)(del Carmen Orozco-Mosqueda et al., 2020). Other mechanisms involved i.e. solubilization of minerals, production of siderophores, increases in leaf area, chlorophyll and soluble protein content. These can also produce antioxidant, enzymes to protect plants from environmental stresses that lead to generation of ROS which cause cell damage, Rhizobacteria may reduce the level of disease by synthesizing hydrolytic enzymes, antibiotics, introduction of systemic acquired resistance (SAR) and competition for nutrients and niches (Sindhu & Sharma, 2019). PGPR significantly decrease the use of chemical fertilizer, pesticides, insecticide and herbicides in modern agricultural practices. PGPR *Pseudomonas sp.*, isolated from rhizosphere soil improve plant growth and crop production and survival of *Amaranthus* species and protect from pathogen attack(Enebe & Babalola, 2018;Ikhajiagbeetal., 2021). They have the ability to produce siderophore, hydrogen cyanide (HCN) and phosphate solubilization property (Sagar et al.,2018).

Plant Growth Promoting Fungi (PGPF) maintain plant growth through the generation of a number of significant enzymes like ACCD (1-aminocyclopropane-1-carboxylate deaminase), urease, catalase etc, phosphate solubilization, siderophores and IAA production, antagonism to pathogens and resistant to antibiotics and take a crucial role in plant growth. PGPF are non-pathogenic, soil borne fungi enhance plant growth and survival (HyAKuMAChI,1994; Debashis et al., 2020). It also enhances soil fertility, productivity, nutrient uptake and Defence mechanism (Yapaetal., 2022). It triggers induced systemic resistance (ISR) against pathogenic microorganism by modifying cell wall accumulating callose, phenol, lignin (Yapaetal., 2022). Different PGPR and PGPF strains when combined and applied to soil of Amaranth, then morphological parameters i.e., shoot height, root height, fresh weight all are increased and induced SAR (Debashis et al., 2020).

5.3 VAM: Mycorrhiza (VAM) is the symbiotic association between higher plants and fungi(Giovannini et al., 2020). The fungus grows internally inside the root cortical cells where they form specialized structures called vesicles and arbuscules(Holland& Roth, 2023). Vesicles are swollen, sac like, unbranched, modified structures of the fungal hyphae. The major function is the storage of reserve

Comment [U20]: References

Comment [U21]: They are inside *Amaranthus cruentus* ?

Comment [U22]: Search more references

Comment [U23]: References

Comment [U24]: References

food whereas arbuscules are branched, finger like projected hyphal modifications of the endomycorrhiza and the major function is the absorption of food nutrients and water from the host cytoplasm (Holland & Roth, 2023). VAM efficiently extend plant roots making the uptake of water much more efficient, better mineral nutrition especially Phosphate, alteration in root architecture, modification of some physiological and enzymatic activities, induction of the plant hormones ABA (Fengetal.,2020). VAM inoculation increases stomatal conductance under drought condition. The VAM fungi are obligatory biotrophs in nature (Giovannini et al., 2020) and also act as bioprotectants against pathogen and toxic stresses. Application of VAM as fertilizer to the Amaranth plant significantly increase the shoot diameter, root length, leaf number etc. (Dada et al., 2017). According to the study of Safeena & Sasmina, 2018, AM (*Glomus fasciculatum*) were inoculated into the pesticide treated soil of *A. cruentus*, the result revealed that growth of the plant is increased and pesticide toxicity in the soil is decreased and soil-microbe growth are not inhibited. Report of Kalaikandhan et al.,2012, revealed that when VAM (*Glomus hoi*) is inoculated in soil of *A. cruentus*, heavy metal concentration of soil is reduced because of the VAM fungi having capacity to sequester it in their mycelia and make the soil less toxic.

6.DISCUSSION

In this review, we discussed mainly the beneficial features and also the exceptional features of this plant *A. cruentus* (Red Amaranth). It distributed all over the world and can grow any type of soils. This C4 photosynthetic plant can tolerate adverse climatic conditions. Short growing period are very advantageous for agriculture and farmers are also benefitted. As India and most of the countries are suffered from Malnutrition, this Red Amaranth provide high amount of nutrition (amino acid, carbohydrate, fatty acid, vitamins, minerals, fibres, antioxidant). Financial expenditure of the maintenance of this plant is comparatively low. Food quality is more important to prevent malnutrition than food quantity (Okunlola et al., 2024). Major food crops are deficient in proper nutrient. By consuming these crops, many peoples are susceptible to diseases. This Red Amaranth is considered as 'Alternative source of Crop' or 'Pseudocereal'. It contains high percentage of nutritional compounds, bioactive secondary metabolites and medicinal properties (Manyelo et al., 2022). This plant is not only used for human food but also very efficient Fodder crop. This leafy vegetable is rich source of dietary fibre which helps in digestion of animals. In some countries, it, regarded as 'Ornamental' and also used in dye and cosmetic industry. Bioactive secondary metabolites (Phenolics, Flavonoids, Terpenoids, Saponins, Alkaloids, Quinone, Tannin) present in this plant prevent the entry of pathogenic microorganisms and have several medicinal activities i.e. antioxidant, anti-inflammatory, immunomodulatory etc. (Peter&Gandhi,2017)). Fungi isolated from this plant secret antibiotic compounds and protect plant from abiotic stresses (Pusz et al., 2015). PGPF are non-pathogenic soil borne fungi enhance nutrient uptake and soil quality of the plant have different antagonistic mechanisms (Debashis et al., 2020). VAM have the capacity to decrease the heavy metal toxicity and devastating effect of pesticides of the soil ((Kalaikandhan et al.,2012). Microbes and microbial inoculants application as biofertilizers and biocontrol agent is an indispensable component in organic farming practices. Improved understanding of 'mycorrhizosphere' interactions play pivotal role in order to preserve crop yield stability under an increasingly unstable climate change.

7.CONCLUSION

This plant has huge role in food, fodder, dye, cosmetic and medicinal industry. Secondary metabolites extracted from this plant possesses important role in pharmacy. Microbes are a noble and consistent source of unique natural medicines with a high level of biodiversity and may also yield several pharmaceutically important compounds. This symbiotic association at the molecular and genetic levels will be helpful for enhancing secondary metabolite production and can be an active area for future investigations. In recent years, there have been a large number of studies, both in the laboratory and in the field, that have provided information about the use of plant- associated microbes to the benefit of plant growth and development. Bacteria, fungi, mycorrhiza provide resistance to plants against various environmental (abiotic) stresses. In the recent time, we are hopefully determined that the purposeful use of microorganisms will be commenced on to replace the use of energy-intensive and often pollution causing chemicals in agricultural practices. Application of microbes and microbial inoculants as biofertilizers and biocontrol agent is an integral component in organic farming practices. Lastly, improved understanding of 'mycorrhizosphere' interactions is crucial in order to preserve crop yield stability under an increasingly unstable climate change. To think about the food scarcity of our near future, we should protect this plant as a Crop for alternative source

Comment [U25]: Does VAM a natural symbiont with *Amaranthus cruentus* ?

Comment [U26]: What do you discuss here? Remember that discussion is applicable to research manuscript to discuss the results?

Comment [U27]: Part of introduction

Comment [U28]: Try to include this part in conclusion

and try to increase the production of the plant by using the associated microbes and secondary metabolites extracted from this plant.

Comment [U29]: Your conclusion lacks the precision quantity of each metabolite, and other numerical characteristics and the particularities of this species compared to others.

References

Comment [U30]: Revise and rewrite correctly

Achigan-Dako, E. G., Sogbohossou, O. E., & Maundu, P. (2014). Current knowledge on *Amaranthus* spp.: research avenues for improved nutritional value and yield in leafy amaranths in sub-Saharan Africa. *Euphytica*, 197, 303-317. DOI 10.1007/s10681-014-1081-9

Adeleke, B. S., & Babalola, O. O. (2021). Pharmacological potential of fungal endophytes associated with medicinal plants: A review. *Journal of Fungi*, 7(2), 147. <https://doi.org/10.3390/jof7020147>

Aderibigbe, O. R., Ezekiel, O. O., Owolade, S. O., Korese, J. K., Sturm, B., & Hensel, O. (2022). Exploring the potentials of underutilized grain amaranth (*Amaranthus* spp.) along the value chain for food and nutrition security: A review. *Critical reviews in food science and nutrition*, 62(3), 656-669. <https://doi.org/10.1080/10408398.2020.1825323>

Alegbejo, J. O. (2013). Nutritional value and utilization of *Amaranthus* (*Amaranthus* spp.)—a review. *Bayero journal of pure and applied sciences*, 6(1), 136-143. <https://doi.org/10.4314/bajopas.v6i1.27>

Ali, M., Dahiya, S. S., Sultana, S., & Munjal, K. (2021). Fatty acid constituents from the leaves of *Amaranthus hybridus* L. *European Journal of Pharmaceutical and Medical Research*, 8(9), 613-617.

Ali, M. (2021). Determination of Proximate, Phytochemicals and Minerals Composition of *Amaranthus* spp. DOI: 10.36346/sarjps.2021.v03i05.002

Allemann, I. (2016). Influence of abiotic stress on allelopathic properties of *Amaranthus cruentus* L. <http://hdl.handle.net/11660/5825>

Alvarez-Jubete, L., Wijngaard, H., Arendt, E. K., & Gallagher, E. (2010). Polyphenol composition and in vitro antioxidant activity of amaranth, quinoa buckwheat and wheat as affected by sprouting and baking. *Food chemistry*, 119(2), 770-778. <https://doi.org/10.1016/j.foodchem.2009.07.032>

Alvin, A., Miller, K. I., & Neilan, B. A. (2014). Exploring the potential of endophytes from medicinal plants as sources of antimycobacterial compounds. *Microbiological research*, 169(7-8), 483-495. <https://doi.org/10.1016/j.micres.2013.12.009>

Amosova, A. V., Yurkevich, O. Y., Semenov, A. R., Samatadze, T. E., Sokolova, D. V., Artemyeva, A. M., ... & Muravenko, O. V. (2024). Genome Studies in *Amaranthus cruentus* L. and *A. hypochondriacus* L. Based on Repeatomic and Cytogenetic Data. *International Journal of Molecular Sciences*, 25(24), 13575. <https://doi.org/10.3390/>

Azab, A. (2020). Amaranthaceae plants of Israel and palestine: Medicinal activities and unique compounds. *EUROPEAN CHEMICAL BULLETIN*, 9(12), 366-400. <http://dx.doi.org/10.17628/ecb.2020.9.366-400>

Baraniak, J., & Kania-Dobrowolska, M. (2022). The dual nature of amaranth—Functional food and potential medicine. *Foods*, 11(4), 618. <https://doi.org/10.3390/foods11040618>

Buchanan, B. B., Grissem, W., & Jones, R. L. (2015). *Biochemistry and molecular biology of plants* (2nd ed) John wiley & sons.

Castrillón-Arbeláez, P. A., & Frier, J. P. D. (2016). Secondary metabolism in *Amaranthus* spp.—a genomic approach to understand its diversity and responsiveness to stress in marginally studied crops with high agronomic potential. *Abiotic and Biotic Stress in Plants. Recent Advances and Future Perspectives. India: In Tech*, 185-227. <https://doi.org/10.5772/61820>

Chatepa, L. E. C., & Masamba, K. G. (2020). Proximate and phytochemical

- composition of selected indigenous leafy vegetables consumed in Malawi. *African Journal of Food Science*, 14(9), 265-273. <http://dx.doi.org/10.5897/AJFS2020.1979>
- Cheng, H. Y., Lin, C. C., & Lin, T. C. (2002). Antiherpes simplex virus type 2 activity of casuarinin from the bark of *Terminalia arjuna* Linn. *Antiviral research*, 55(3), 447-455. [https://doi.org/10.1016/S0166-3542\(02\)00077-3](https://doi.org/10.1016/S0166-3542(02)00077-3)
- Cherian, P., & Sheela, D. (2016). Antimicrobial activity of Amaranth Alkaloid against pathogenic microbes. *International Journal of Herbal Medicine*, 4(5), 70-72.
- Choudhury, D., Tarafdar, S., Parvin, N., Rit, R., Roy, S., & Dutta, S. (2023). Endophytic microbes and their diverse beneficial aspects in various sectors: A critical insight. *Plant Science Today*, 10(1), 96-107. <https://doi.org/10.14719/pst.1877>
- Dada, O. A., Imade, F., & Anifowose, E. M. (2017). Growth and proximate composition of *Amaranthus cruentus* L. on poor soil amended with compost and arbuscular mycorrhiza fungi. *International Journal of Recycling of Organic Waste in Agriculture*, 6, 195-202. DOI 10.1007/s40093-017-0167-5
- Das, S., & Das, S. (2016). Amaranths: the crop of great prospect. *Amaranthus: A promising crop of future*, 13-48. https://doi.org/10.1007/978-981-10-1469-7_3
- Debasis, M., Ganeshamurthy, A. N., Komal, S., Radha, T. K., & Rupa, T. R. (2020). Plant growth promotion and biocontrol activity of some typical harsh environment rhizo-microbes and their effects on *Amaranthus cruentus* plants. *Res. J. Biotechnol. Vol*, 15, 1.
- del Carmen Orozco-Mosqueda, M., Glick, B. R., & Santoyo, G. (2020). ACC deaminase in plant growth-promoting bacteria (PGPB): An efficient mechanism to counter salt stress in crops. *Microbiological Research*, 235, 126439. <https://doi.org/10.1016/j.micres.2020.126439>
- Dhellit, J. R., Matouba, E., Maloumbi, M. G., Nzikou, J. M., Ngoma, D. S., Linder, M., ... & Parmentier, M. (2006). Extraction, chemical composition and nutritional characterization of vegetable oils: Case of *Amaranthus hybridus* (var 1 and 2) of Congo Brazzaville. *African journal of biotechnology*, 5(11). <http://www.academicjournals.org/AJB>
- Emam, T. M., Hosni, A. E. M., Ismail, A., El-Kinany, R. G., Hewidy, M., Saady, H. S., ... & El-Sayed, S. M. (2024). Physiological and Molecular Responses of Red Amaranth (*Amaranthus cruentus* L.) and Green Amaranth (*Amaranthus hypochondriacus* L.) to Salt Stress. *Journal of Soil Science and Plant Nutrition*, 1-12. <https://doi.org/10.1007/s42729-024-02125-w>
- Enebe, M. C., & Babalola, O. O. (2018). The influence of plant growth-promoting rhizobacteria in plant tolerance to abiotic stress: a survival strategy. *Applied microbiology and biotechnology*, 102, 7821-7835. <https://doi.org/10.1007/s00253-018-9214-z>
- Erdman Jr, J. W., Balentine, D., Arab, L., Beecher, G., Dwyer, J. T., Folts, J., ... & Burrowes, J. (2007). Flavonoids and Heart Health: Proceedings of the ILSI North America Flavonoids Workshop, May 31–June 1, 2005, Washington, DC1. *The Journal of nutrition*, 137(3), 718S-737S. <https://doi.org/10.1093/jn/137.3.718S>
- Feng, Z., Liu, X., Zhu, H., & Yao, Q. (2020). Responses of arbuscular mycorrhizal symbiosis to abiotic stress: a lipid-centric perspective. *Frontiers in Plant Science*, 11, 578919. <https://doi.org/10.3389/fpls.2020.578919>
- Fletcher, R. J. (2015). Pseudocereals: overview. <https://doi.org/10.1016/B978-0-12-394437-5.00039-5>
- Frank, A. C., Saldierna Guzmán, J. P., & Shay, J. E. (2017). Transmission of

- bacterial endophytes. *Microorganisms*, 5(4), 70. <https://doi.org/10.3390/microorganisms5040070>
- George, G. M. (2015). *Postharvest handling for extending shelf life of Amaranthus (Amaranthus tricolor L.)* (Doctoral dissertation, Department of Processing Technology, College of Agriculture, Vellayani). <http://krishikosh.egranth.ac.in/handle/1/5810029695>
- Gins, E. M., Baikov, A. A., Khasanova, S. D., Goryunova, S. V., Gins, V. K., Gins, M. S., & Motyleva, S. M. (2024). AMARANTHUS SPECIES ASSESSMENT FOR MORPHOLOGICAL AND BIOCHEMICAL PARAMETERS. *SABRAO Journal of Breeding & Genetics*, 56(4). <http://doi.org/10.54910/sabrao2024.56.4.6><http://sabraojournal.org/>
- Giovannini, L., Palla, M., Agnolucci, M., Avio, L., Sbrana, C., Turrini, A., & Giovannetti, M. (2020). Arbuscular mycorrhizal fungi and associated microbiota as plant biostimulants: research strategies for the selection of the best performing inocula. *Agronomy*, 10(1), 106. <https://doi.org/10.3390/agronomy10010106>
- Gresta, F., Meineri, G., Oteri, M., Santonoceto, C., Lo Presti, V., Costale, A., & Chiofalo, B. (2020). Productive and qualitative traits of *Amaranthus cruentus* L.: An unconventional healthy ingredient in animal feed. *Animals*, 10(8), 1428. <https://doi.org/10.3390/ani10081428>
- Hardoim, P. R., Van Overbeek, L. S., Berg, G., Pirttilä, A. M., Compant, S., Campisano, A., ... & Sessitsch, A. (2015). The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiology and molecular biology reviews*, 79(3), 293-320. <https://doi.org/10.1128/mmb.00050-14>
- He, H. P., & Corke, H. (2003). Oil and squalene in amaranthus grain and leaf. *Journal of Agricultural and Food Chemistry*, 51(27), 7913-7920. <https://doi.org/10.1021/jf030489g>
- Hoidal, N., Díaz Gallardo, M., Jacobsen, S. E., & Alandia, G. (2019). Amaranth as a dual-use crop for leafy greens and seeds: Stable responses to leaf harvest across genotypes and environments. *Frontiers in Plant Science*, 10, 817. <https://doi.org/10.3389/fpls.2019.00817>
- Holland, S., & Roth, R. (2023). Extracellular vesicles in the arbuscular mycorrhizal symbiosis: current understanding and future perspectives. *Molecular Plant-Microbe Interactions*, 36(4), 235-244. <https://doi.org/10.1094/MPMI-09-22-0189-FI>
- Hricova, A., Fejer, J., Libiakova, G., Szabova, M., Gazo, J., & Gajdosova, A. (2016). Characterization of phenotypic and nutritional properties of valuable *Amaranthus cruentus* L. mutants. *Turkish Journal of Agriculture and Forestry*, 40(5), 761-771. [10.3906/tar-1511-31](https://doi.org/10.3906/tar-1511-31)
- HyAKuMAcHi, M. (1994). Plant-growth-promoting fungi from turfgrass rhizosphere with potential for disease suppression. *Soil Microorganisms*, 44, 53-68. https://doi.org/10.18946/jssm.44.0_53
- Ikhajigbe, B., Ogwu, M. C., Fawehinmi, F. O., & Adekunle, I. J. (2021). Comparative growth responses of *Amaranthus* [L.] species in humus and ferruginous ultisols using plant growth promoting Rhizobacteria (*Pseudomonas* species). *South African Journal of Botany*, 137, 10-18. <https://doi.org/10.1016/j.sajb.2020.09.029>
- Iwuagwu, M. O., Ogbonna, N. C., & Okechukwu, U. H. *International Journal of Plant Science and Horticulture*. <https://doi.org/10.36811/ijpsh.2019.110008>
- Jimoh, M. O., Okaiyeto, K., Oguntibeju, O. O., & Laubscher, C. P. (2022). A systematic review on *Amaranthus*-related research. *Horticulturae*, 8(3),

239. <https://doi.org/10.3390/horticulturae8030239>

Kalaikandhan, R., Vijayarengan, P., Thamizhiniyan, P., & Sivasankar, R. (2012). Impact of the pesticides monocrotophos and quinalphos on the morphological features of red amaranth under arbuscular mycorrhizal fungus inoculation. *World J Agric Sci*, 8(2), 212-217.

Kauffman, C. S., & Weber, L. E. (1990). Grain amaranth. *Advances in new crops*. Timber Press, Portland, OR, 127-139.

Koda, T., Kuroda, Y., & Imai, H. (2009). Rutin supplementation in the diet has protective effects against toxicant-induced hippocampal injury by suppression of microglial activation and pro-inflammatory cytokines: protective effect of rutin against toxicant-induced hippocampal injury. *Cellular and molecular neurobiology*, 29, 523-531. <https://doi.org/10.1007/s10571-008-9344-4>

Lanta, V., Havranek, P., & Ondrej, V. (2003). Morphometry analysis and seed germination of *Amaranthus cruentus*, *A. retroflexus* and their hybrid (*A. x turicensis*). *Plant Soil and Environment*, 49(8), 364-369.

Letchamo, W., Hartman, T., Gosslin, A., Mamedov, N. A., & Craker, L. (2017). The accumulation of phenolic compounds in genetically selected *Amaranthus hybridus* is influenced by endophytic natural growth regulator. *International Journal of Secondary Metabolite*, 5(1), 12-19. <http://dergipark.gov.tr/ijism>

Ma, X., Vaistij, F. E., Li, Y., Jansen van Rensburg, W. S., Harvey, S., Bairu, M. W., ... & Denby, K. J. (2021). A chromosome-level *Amaranthus cruentus* genome assembly highlights gene family evolution and biosynthetic gene clusters that may underpin the nutritional value of this traditional crop. *The Plant Journal*, 107(2), 613-628. <https://doi.org/10.1111/tpj.15298>

Makinde, E. A., Ayeni, L. S., & Ojeniyi, S. O. (2010). Morphological characteristics of *amaranthus cruentus* L. as influenced by Kola Pod Husk, organomineral and NPK Fertilizers in Southwestern Nigeria. *New York Science Journal*, 3(5), 130-134. <http://www.sciencepub.net/newyork>

Malik, M., Sindhu, R., Dhull, S. B., Bou-Mitri, C., Singh, Y., Panwar, S., & Khatkar, B. S. (2023). Nutritional composition, functionality, and processing technologies for amaranth. *Journal of Food Processing and Preservation*, 2023(1), 1753029. <https://doi.org/10.1155/2023/1753029>

Manyelo, T. G., Sebola, N. A., Hassan, Z. M., Ng'ambi, J. W., Weeks, W. J., & Mabelebele, M. (2022). Chemical composition and metabolomic analysis of *Amaranthus cruentus* grains harvested at different stages. *Molecules*, 27(3), 623. <https://doi.org/10.3390/molecules27030623>

Mburu, M. W., Gikonyo, N. K., Kenji, G. M., & Mwasaru, A. M. (2011). Properties of a complementary food based on amaranth grain (*Amaranthus cruentus*) grown in Kenya. *Journal of Agriculture and Food Technology*, 1(9), 153-178.

Mepba, H. D., Eboh, L., & Banigo, D. E. B. (2007). Effects of processing treatments on the nutritive composition and consumer acceptance of some Nigerian edible leafy vegetables. *African Journal of Food, Agriculture, Nutrition and Development*, 7(1). <http://ajfand.net/AJFAND/copyrightstatement.html>

Morelli, M., Bahar, O., Papadopoulou, K. K., Hopkins, D. L., & Obradović, A. (2020). Role of endophytes in plant health and defense against pathogens. *Frontiers in Plant Science*, 11, 1312. <https://doi.org/10.3389/fpls.2020.01312>

Motta, C., Castanheira, I., Gonzales, G. B., Delgado, I., Torres, D., Santos, M., & Matos, A. S. (2019). Impact of cooking methods and malting on amino acids

content in amaranth, buckwheat and quinoa. *Journal of Food Composition and Analysis*, 76, 58-65. <https://doi.org/10.1016/j.jfca.2018.10.001>

Mousavi, S. S., & Karami, A. (2022). Application of Endophyte microbes for production of secondary metabolites. *Application of Microbes in Environmental and Microbial Biotechnology*, 1-37. https://doi.org/10.1007/978-981-16-2225-0_1

Mukuwapasi, B., Mavengahama, S., & Gerrano, A. S. (2024). Grain amaranth: A versatile untapped climate-smart crop for enhancing food and nutritional security. *Discover Agriculture*, 2(1), 44. <https://doi.org/10.1007/s44279-024-00057-8>

Muthukumar, A., Udhayakumar, R., & Naveenkumar, R. (2017). Role of bacterial endophytes in plant disease control. *Endophytes: Crop Productivity and Protection: Volume 2*, 133-161. https://doi.org/10.1007/978-3-319-66544-3_7

Nana, F. W., Hilou, A., Millogo, J. F., & Nacoulma, O. G. (2012). Phytochemical composition, antioxidant and xanthine oxidase inhibitory activities of *Amaranthus cruentus* L. and *Amaranthus hybridus* L. extracts. *Pharmaceuticals*, 5(6), 613-628. <https://doi.org/10.3390/ph5060613>

Narayanan, B. A., Geoffroy, O., Willingham, M. C., Re, G. G., & Nixon, D. W. (1999). p53/p21 (WAF1/CIP1) expression and its possible role in G1 arrest and apoptosis in ellagic acid treated cancer cells. *Cancer letters*, 136(2), 215-221. [https://doi.org/10.1016/S0304-3835\(98\)00323-1](https://doi.org/10.1016/S0304-3835(98)00323-1)

Nofianti, T. (2015). ANTIHYPERGLICEMIA ACTIVITY OF WATER FRACTION, ETHYL ACETATE FRACTION, AND N-HEXANE FRACTION OF SPINACH LEAVES EXTRACT (*Amaranthus Cruentus* L) ON MALE MICE SWISS. *JURNAL FARMASIGALENIKA*, 2(02). <https://journal.bku.ac.id/jfg/index.php/jfg/article/view/36>

Ogwu, M. C. (2020). *Value of Amaranthus [L.] species in Nigeria* (pp. 1-21). London: IntechOpen. <https://dx.doi.org/10.5772/intechopen.86990>

Okunlola, G. O., Jimoh, M. A., Olatunji, O. A., & Olowolaju, E. D. (2017). Comparative study of the phytochemical contents of *Cochorus olitorius* and *Amaranthus hybridus* at different stages of growth. *Annales of West University of Timisoara. Series of Biology*, 20(1), 43.

Okunlola, M. M., & Adejumo, S. A. (2024). Enhancing nutritional quality of *Amaranthus cruentus* (L.) with fertilisers and salicylic acid. *World Journal of Advanced Research and Reviews*, 23(1), 030-041. <https://doi.org/10.30574/wjarr.2024.23.1.1925>

Omale, J., & Okafor, P. N. (2008). Comparative antioxidant capacity, membrane stabilization, polyphenol composition and cytotoxicity of the leaf and stem of *Cissus multistriata*. *African Journal of Biotechnology*, 7(17). <http://www.academicjournals.org/AJB>

Omami, E. N., & Hammes, P. S. (2006). Interactive effects of salinity and water stress on growth, leaf water relations, and gas exchange in amaranth (*Amaranthus* spp.). *New Zealand Journal of Crop and Horticultural Science*, 34(1), 33-44. <https://doi.org/10.1080/01140671.2006.9514385>

Osei-Owusu, J., Kokro, K. B., Ofori, A., Apau, J., Dofuor, A. K., Vigbedor, B. Y., ... & Okyere, H. (2023). Evaluation of phytochemical, proximate, antioxidant, and anti-nutrient properties of *Corchorus olitorius*, *Solanum macrocarpon* and *Amaranthus cruentus* in Ghana. *International Journal of Biochemistry and Molecular*

Biology, 14(2), 17.

Pandey, S., Ganeshpurkar, A., Bansal, D., & Dubey, N. (2016). Hematopoietic effect of *Amaranthus cruentus* extract on phenylhydrazine-induced toxicity in rats. *Journal of Dietary Supplements*, 13(6), 607-615. <https://doi.org/10.3109/19390211.2016.1155685>

Park, Y. J., Nishikawa, T., Matsushima, K., Minami, M., & Nemoto, K. (2014). A rapid and reliable PCR-restriction fragment length polymorphism (RFLP) marker for the identification of *Amaranthus cruentus* species. *Breeding science*, 64(4), 422-426. <https://doi.org/10.1270/isbbs.64.422>

Peter, K., & Gandhi, P. (2017). Rediscovering the therapeutic potential of *Amaranthus* species: A review. *Egyptian journal of basic and applied sciences*, 4(3), 196-205. <https://doi.org/10.1016/j.ejbas.2017.05.001>

Prajitha, V., & Thoppil, J. E. (2018). Cytogenetic characterization of *Amaranthus caudatus* L. and *Amaranthus hybridus* subsp. *cruentus* (L.) Thell. *Cytotechnology*, 70, 95-101. <https://doi.org/10.1007/s10616-017-0100-9>

Priti, V., Ramesha, B. T., Singh, S., Ravikanth, G., Ganeshiah, K. N., Suryanarayanan, T. S., & Uma Shaanker, R. (2009). How promising are endophytic fungi as alternative sources of plant secondary metabolites?. *Current Science*, 97(4), 477-478.

Prohp, T. P., & Onoagbe, I. O. (2012). Determination of phytochemical composition of the stem bark of *Triplochiton scleroxylon* K. Schum.(sterculiaceae).

Pusz, W., Plaskowska, E., Yildirim, I., & Weber, R. (2015). Fungi occurring on the plants of the genus *Amaranthus* L. *Turkish Journal of Botany*, 39(1), 147-161. Doi:10.3906/bot-1403-106

Rabi, T., & Bishayee, A. (2009). Terpenoids and breast cancer chemoprevention. *Breast cancer research and treatment*, 115, 223-239. <https://doi.org/10.1007/s10549-008-0118-y>

Rajamanikyam, M., Vadlapudi, V., & Upadhyayula, S. M. (2017). Endophytic fungi as novel resources of natural therapeutics. *Brazilian Archives of Biology and Technology*, 60, e17160542. <https://doi.org/10.1590/1678-4324-2017160542>

Ramkisson, S., Dwarka, D., Venter, S., & Mellem, J. J. (2020). In vitro anticancer and antioxidant potential of *Amaranthus cruentus* protein and its hydrolysates. *Food Science and Technology*, 40(suppl 2), 634-639. <https://doi.org/10.1590/fst.36219>

Rao, A. V., & Sung, M. K. (1995). Saponins as anticarcinogens. *The Journal of nutrition*, 125, 717S-724S. https://doi.org/10.1093/jn/125.suppl_3.717S

Rit, R., Choudhury, D., Mukherjee, C., & Dutta, S. (2023). Investigation on microfloral association in the roots of *Macrotyloma uniflorum* (Lam.) Verdc., a medicinally important tropical pulse-crop and their possible applications for crop improvement: A review. *Plant Science Today*, 10(3), 300-311. <https://www.horizonpublishing.com/journals/index.php/PST/index>

Rodriguez, R. J., Woodward, C., Kim, Y. O., & Redman, R. S. (2009). Habitat-adapted symbiosis as a defense against abiotic and biotic stresses. In *Defensive mutualism in microbial symbiosis* (pp. 353-364). CRC Press.

Safeena, M. I. S., & Sasmina, T. (2018). Effect of biofertilizer (with mycorrhiza inoculum), cowdung, nitrogen and phosphorous on growth and yield of three varieties of amaranth (*Amaranthus tricolor* L.). <http://ir.lib.seu.ac.lk/handle/123456789/3534>

Sagar, A., Dhusiya, K., Shukla, P. K., Singh, A., Lawrence, R., & Ramteke, P. W. (2018). Comparative analysis of production of hydrogen cyanide (HCN) with production of siderophore (SD) and phosphate solubilization (PS) activity in plant growth promoting bacteria (PGPB). *Vegetos*, 31(2), 130-135. Doi:10.5958/2229-

Sammour, R. H., Radwan, S. A., & Mira, M. (2012). Genetic diversity in genus *Amaranthus*: From morphology to genomic DNA. *Res. Rev. Biosci*, 6, 351-360.

Schulz, B., & Boyle, C. (2006). What are endophytes?. In *Microbial root endophytes* (pp. 1-13). Berlin, Heidelberg: Springer Berlin Heidelberg. https://doi.org/10.1007/3-540-33526-9_1

Sharma, S., & Roy, S. (2015). Isolation and identification of a novel endophyte from a plant *Amaranthus spinosus*. *International Journal of Current Microbiology and Applied Sciences*, 4(2), 785-798. <http://www.ijcmas.com/>

Sindhu, S. S., & Sharma, R. (2019). Amelioration of biotic stress by application of rhizobacteria for agriculture sustainability. *Plant Growth Promoting Rhizobacteria for Sustainable Stress Management: Volume 2: Rhizobacteria in Biotic Stress Management*, 111-168. https://doi.org/10.1007/978-981-13-6986-5_5

Singh, L. P., Gill, S. S., & Tuteja, N. (2011). Unraveling the role of fungal symbionts in plant abiotic stress tolerance. *Plant signaling & behavior*, 6(2), 175-191. <https://doi.org/10.4161/psb.6.2.14146>

Siwale, D. R. (2018). *Blood glucose lowering effects of leaf extracts of cleome gynandra, amaranthus cruentus and their mixture in normoglycaemic and hyperglycaemic rats* (Doctoral dissertation, The University of Zambia).

Stone, J. K., Bacon, C. W., & White Jr, J. F. (2000). An overview of endophytic microbes: endophytism defined. *Microbial endophytes*, 17-44.

Sultana, R., Islam, S. M. N., Shuvo, S. B., Ehsan, G. M. A., Saha, P., Khan, M. M. R., & Rumman, N. (2024). Endophytic bacterium *Sphingomonas panaciterrae* NB5 influences soil properties and improves growth, nutrient contents, and yield of red amaranth (*Amaranthus tricolor* L.). *Current Plant Biology*, 39, 100372. <https://doi.org/10.1016/j.cpb.2024.100372>

Torane, R., Gaikwad, S., Khatiwora, E., & Adsul, V. (2017). Comparative estimation of phenol and flavonoid content of medicinally important plant- *Amaranthus currentus*. *Int J Chem Tech Res*, 10(4), 306-310.

Uppala, S., Beena, S., Chapala, M., & Bowen, K. L. (2010). Amaranth endophytes and their role in plant growth promotion. *Plant growth promotion by Rhizobacteria for sustainable agriculture*. Scientific Publishers, Jodhpur, 531-537. DOI: 10.13140/RG.2.1.3291.3441

Usunomena, U., & Ngozi, O. P. (2016). Phytochemical analysis and proximate composition of *Vernonia amygdalina*. *Int. J. Sci. World*, 4(1), 11-14. doi: 10.14419/ijsw.v4i1.5845

Wolosik, K., & Markowska, A. (2019). *Amaranthus Cruentus* taxonomy, botanical description, and review of its seed chemical composition. *Natural Product Communications*, 14(5)

1934578X19844141.<https://doi.org/10.1177/1934578X19844141>

Yapa, N., Lakmali, D., Zoysa, D., Silva, K. S., Manawadu, C., Herath, B. M., ... & Bamunuarachchige, C. (2022). Biofertilizers: An Emerging Trend in Agricultural Sustainability.<http://repo.lib.jfn.ac.lk/ujrr/handle/123456789/8984>

Chicago

UNDER PEER REVIEW