

Original Research Article

EXPLORING THE LEVELS OF HEAVY METALS IN SOIL: A CASE STUDY OF AHOADA WEST LOCAL GOVERNMENT AREA IN RIVERS STATE, NIGERIA

Abstract

Background: Heavy metal-polluted soils are becoming more widespread around the world, as geologic and anthropogenic activity grow. The risk of soil contamination by metals increases drastically with increasing industrialization, urbanization, and oil spillage. Thus, it is necessary to monitor heavy metals concentration in agricultural areas because they can pose serious risks to human health and the environment.

Objectives: the objective of the study was to investigate heavy metals concentration (lead, cadmium, chromium and iron) in soils of Ahoada West Local Government Area, specifically Akinima, Oruama, and Joinkrama 4.

Methods: Soil samples were collected from three places at varying depths (0-5, 0-10, and 0-15 cm). Concentrations of heavy metals were quantified using a Flame Atomic Absorption Spectrophotometer. ANOVA was employed to ascertain significant differences in heavy metal concentrations at $p < 0.05$ (IBM SPSS, 21 version).

Results: Results indicated that the concentrations (ppm) of lead (1.44-13.99), chromium (5.23-23.57), cadmium (0.22-2.28), and iron (3.92-13.36) were below the WHO limits. Also, it was observed that heavy metal concentrations increased with increasing soil depths, in the following order 0-15 > 0-10 > 0-5 cm. In terms of spatial distribution, Oruama had significantly higher heavy metal concentrations, followed by Joinkrama 4, and Akinima, with the exception of chromium, which was significantly higher in Joinkrama 4 (8.00; 17.29) in comparison with Oruama (7.30; 15.34), and Akinima (5.23; 12.43) for 0-5 and 0-10 cm respectively.

Conclusion: The study concludes that Oruama community generally had significantly higher levels of heavy metals, although the analyzed soil samples were below WHO permissible limits. This study, advances the understanding of heavy metal dynamics in soil health and agricultural sustainability in Ahoada West, by providing localized information on concentrations relative to depth of soil and locations. It emphasizes the need for regular monitoring to avoid possible pollution and bioaccumulation.

Key words: Akinima, environmental monitoring, heavy metals, Joinkrama 4, Oruama, soil

1.0 Introduction

Exposure to heavy metals is becoming a serious problem, particularly in emerging nations of the world¹. Although heavy metal is naturally present in the earth's crust, geogenic and human activities have caused it to contaminate the soil². Heavy metal contamination has resulted from industrialization and technological advancement, severely reducing human and animal health³. The accumulation of heavy metals in agricultural soil can lead to soil pollution and enhanced absorption by food crops, thereby jeopardizing food quality and safety¹. Thus, if accumulated in increased concentration exceeding safe thresholds, it may pose a major health risk to people and other living things⁴. Heavy metals and metalloids, including cadmium, lead, arsenic, mercury, and chromium, are environmental pollutants that can accumulate to dangerous amounts in agricultural soils, reducing crop health and productivity⁵. Lead exposure, for instance, can harm the human body's skeletal, circulating, neurological, endocrine, and immune system. Similarly, cadmium exposure can cause cancer⁶. Heavy metals can adversely affect several biological components and organelles. Metal ions can harm human DNA via interactions with both DNA and nuclear proteins. This, in turn, can induce cell cycle regulation, apoptosis, and even cancer. Heavy metals have been shown to induce site-specific damage to human DNA and nuclear proteins upon interaction. The metal causes direct damage by inducing conformational changes in the biomolecules. The generation of reactive oxygen and nitrogen species, along with other endogenous oxidants, is accountable for the indirect harm resulting from heavy metal exposure⁷.

Soil is an essential natural resource for a country's economic and agricultural growth. Its significance, which is undeniable, is evidenced by the existence of microbiological organisms as well as humans, plants, and animals. Unfortunately, a number of man-made and natural elements can cause the soil to deteriorate, which would consequently reduce its production⁸. Soil-related issues have become a significant danger to human society⁹. Heavy metal contamination induces alterations in the size, composition, and activity of microbial

populations, adversely affecting various factors associated with plant quality and production. Heavy metals interfere with important microbial processes and reduces the quantity and activity of soil microorganisms, thus having harmful impacts on the soil biota^{10,11}. Soil microorganisms are essential to soil ecosystems because they regulate nutrient cycling, sustain soil fertility, and enhance plant quality, with their activity potentially affected by heavy metal presence¹¹.

High levels of heavy metal build up are proven to considerably have detrimental effects on the ecosystem and endanger the lives of those living in it. When these heavy metals are discharged into the geoenvironment, the soil acts as a significant sink and repository¹². Wuana and Okieimen¹³ assert that the accumulation of heavy metals and metalloids in soils can occur from a variety of sources such as emissions from burgeoning industrial zones, disposal of high metal wastes, leaded gasoline and paints, fertilizers application on land, sewage sludge, pesticides, wastewater irrigation, residues from coal combustion, petrochemical spills, and atmospheric deposition. The use of soil for agricultural production and other human activities is limited by contamination¹⁴.

The physical and chemical properties of soil from the Niger Delta region of Nigeria have been compromised by petroleum hydrocarbons and heavy metals. Nigerian petroleum gotten from the Niger Delta contains various concentrations of heavy metals¹⁵. Oil spills result to heavy metal pollution of terrestrial and aquatic habitat, particularly in areas that produce oil. Crude oil extraction, processing, transportation, and storage are among the operations that contribute to oil spills; which are defined as the discharge of oil into the environment. Spills can also be caused by intentional actions (such as sabotage and oil bunkering) and accident, as well as improper maintenance of engineering equipment¹⁶.

Oil spills and illegal refining could result in unpredictable fire outbreaks that burn vegetation and leave a crust on the soil, necessitating remediation or revegetation¹⁵. Since Nigeria's crude oil activities is centered on the Niger Delta, the region frequently experience oil spills¹⁶. Consequently, the worldwide issue of heavy metal contamination necessitates cooperation across communities, governments, and scientific entities. Because governmental laws are crucial for both pollution cleanup and source management¹⁷. Hence, this study's objective was to investigate heavy metals concentration (lead, cadmium, chromium, and iron) in soils of Ahoada West Local Government Area; specifically, Akinima, Oruama, and Joinkrama 4.

2.0 Materials and Methods

Study area

The study's focus is on the communities of Joinkrama 4, Oruama, and Akinima in the Ahoada West Local Government Area of Rivers State, Nigeria. These communities are situated on the Eastern bank of the Orashi River, a prominent Niger Delta River. Akinima is situated at roughly 5.1046° N, 6.4529° E, while Joinkrama 4 is situated at roughly 4.8265° N, 6.0665° E. The Nigeria Agip Oil Company (NAOC) and Shell Petroleum Development Company (SPDC) run several hydrocarbon flow stations in the area¹⁸. The region is situated in an area where fishermen and farmers make up the majority of the population. Important operations like oil exploration, illicit bunkering, and transportation are major contaminations in that area¹⁹. Soil contamination has been an issue in the Niger Delta due to oil production and exploration. The environment has been widely contaminated by oil spills and gas flaring, adversely affecting agricultural output and jeopardizing public health²⁰⁻²².

Sample Collection

Using a clean soil auger²³, soil samples were taken from four sites in three chosen communities (Akinima, Oruama, and Joinkrama 4) in Ahoada West Local Government Area, Rivers State at surface depth of 0-5, 0-10, and 0-15 cm. Four soil samples were taken from each site and

subsequently combined to create a representative sample of that location. The samples collected were stored in an airtight container and kept in a cool, dark place until analysis.

Quantification of heavy metals using atomic absorption spectrophotometer

Sample preparation process followed the method described by Gong et al.²⁴, although slightly modified. A mixture containing 100 mL of HNO₃, H₂SO₄, and HClO₄ in the proportion of 40%:40%:20% were placed in a conical flask. Ten milliliter (10 mL) of the sample was placed into a conical flask. Two milliliter (2 mL) of the combined acid mixture was added to each of conical flask containing the sample. This was digested in a fume cupboard with hot plate until white vapors emerged. Subsequently, it was cooled and filtered into a 100 mL volumetric flask and then diluted to the 100 mL mark with distilled water. The heavy metals content was determined using flame atomic absorption spectrophotometer (mode: S4=71096) as cited by Duru *et al.*²⁵

Statistical analysis

All acquired data were statistically analysed, and the mean \pm standard deviation was used to represent the findings. One-way Analysis of Variance (ANOVA) (homogeneity of variance), followed by a post-hoc Tukey Test, was employed to ascertain whether a significant difference existed between the experimental groups. A *p-value* < 0.05 was considered statistically significant. All statistical analysis was performed using IBM SPSS software version 21.

2.0 Results

Table 1. Heavy metal concentrations in soil from Akinima

Heavy metals	mg/kg			
	0 – 5 cm	0 – 10 cm	0 – 15 cm	FAO/ WHO ^{26,27}
Pb	1.44 \pm 0.13 ^a	5.03 \pm 0.10 ^b	9.04 \pm 0.06 ^c	100-300
Cr	5.23 \pm 0.09 ^a	12.43 \pm 0.07 ^b	19.22 \pm 0.16 ^c	350

Cd	0.22±0.02 ^a	0.52±0.01 ^b	0.83±0.02 ^c	3-8
Fe	3.92±0.08 ^a	5.51±0.17 ^b	9.51±0.10 ^c	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a row with different alphabetical superscripts do differ significantly ($p < 0.05$). (a) means 0-5 cm is significantly different from other groups. (b) means 0-10 cm is significantly different from other groups, and (c) means 0-15 cm is significantly different from other groups.

Table 1 shows the heavy metal concentrations (mg/kg) in soil from Akinima, 0-15 cm was significantly higher for Pb (9.04), Cr (19.22), Cd (0.83), and Fe (9.51) in comparison with 0-10 cm (5.03,12.43,0.52, and 5.51), and 0-5 cm (1.44, 5.23, 0.22, and 3.92) respectively.

Table 2. Heavy metal concentrations in soil from Oruama

Heavy metals	mg/kg			FAO/ WHO ^{26,27}
	0 – 5 cm	0 – 10 cm	0 – 15 cm	
Pb	5.04±0.08 ^a	9.28±0.18 ^b	13.99±0.01 ^c	100-300
Cr	7.30±0.18 ^a	15.34±0.24 ^b	23.57±0.11 ^c	350
Cd	1.77±0.08 ^a	1.82±0.08 ^b	2.28±0.11 ^c	3-8
Fe	8.24±0.11 ^a	11.03±0.09 ^b	13.36±0.14 ^c	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a row with different alphabetical superscript do differ significantly ($p < 0.05$). (a) means 0-5 cm is significantly different from other groups. (b) means 0-10 cm is significantly different from other groups, and (c) means 0-15 cm is significantly different from other groups.

Table 2 shows the heavy metal concentrations (mg/kg) in soil from Oruama, 0-15 cm was significantly higher for Pb (13.99), Cr (23.57), Cd (2.28) and Fe (13.36) in comparison with 0-10 cm (9.28, 15.34,1.82, and 11.03), and 0-5 cm (5.04, 7.30, 1.77, and 8.24) respectively.

Table 3. Heavy metal concentrations in soil from Joinkrama 4(JK 4)

Heavy metals	mg/kg			FAO/ WHO ^{26,27}
	0 – 5 cm	0 – 10 cm	0 – 15 cm	
Pb	3.07±0.08 ^a	8.74±0.10 ^b	11.57±0.14 ^c	100-300

Cr	8.00±0.02 ^a	17.29±0.12 ^b	21.71±0.11 ^c	350
Cd	0.45±0.02 ^a	0.73±0.01 ^b	1.29±0.09 ^c	3-8
Fe	5.32±0.16 ^a	7.54±0.11 ^b	13.16±0.11 ^c	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a row with different alphabetical superscripts do differ significantly ($p < 0.05$). (a) means 0-5 cm is significantly different from other groups. (b) means 0-10 cm is significantly different from other groups, and (c) means 0-15 cm is significantly different from other groups.

Table 3 shows the heavy metal concentrations (mg/kg) in soil from Joinkrama 4, 0-15 cm was significantly higher for Pb (11.57), Cr (21.71), Cd (1.29) and Fe (13.16) in comparison with 0-10 cm (8.74,17.29,0.73, and 7.54), and 0-5 cm (3.07, 8.00, 0.45, and 5.32) respectively.

Table 4: Heavy metal concentrations in soil from depth 0-5 cm

Heavy metals	mg/kg			
	Akinima	Oruama	Joinkrama 4 (JK 4)	FAO/ WHO ^{26,27}
Pb	1.44±0.13 ^a	5.04±0.08 ^b	3.07±0.08 ^c	100-300
Cr	5.23±0.09 ^a	7.30±0.18 ^b	8.00±0.02 ^c	350
Cd	0.22±0.02 ^a	1.77±0.08 ^c	0.45±0.02 ^c	3-8
Fe	3.92±0.08 ^a	8.24±0.11 ^b	5.32±0.16 ^c	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a row with different alphabetical superscripts do differ significantly ($p < 0.05$). (a) means Akinima is significantly different from other groups, (b) means Oruama is significantly different from other groups, and (c) means JK4 is significantly different from other groups.

Table 4 shows the heavy metal concentrations (mg/kg) in soil from depth 0-5cm, Oruama was significantly higher for Pb (5.04), Cd (1.77), and Fe (8.24) in comparison with JK 4 (3.07,0.45, and 5.32), and Akinima (1.44, 0.22, and 3.92) respectively. While for Cr concentration, JK 4 (8.00) was significantly higher than Oruama (7.30), and Akinima (5.23).

Table 5. Heavy metal concentrations in soil from depth 0-10 cm (mg/kg)

Heavy metals	Akinima	Oruama	Joinkrama 4 (JK 4)	FAO/ WHO ^{26,27}
Pb	5.03±0.10 ^a	9.28±0.18 ^b	8.74±0.10 ^c	100-300
Cr	12.43±0.07 ^a	15.34±0.24 ^b	17.29±0.12 ^c	350
Cd	0.52±0.01 ^a	1.82±0.08 ^b	0.73±0.01 ^c	3-8
Fe	5.51±0.17 ^a	11.03±0.09 ^b	7.54±0.11 ^c	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a row with different alphabetical superscripts do differ significantly ($p < 0.05$). (a) means Akinima is significantly different from other groups, (b) means Oruama is significantly different from other groups, and (c) means JK4 is significantly different from other groups.

Table 5 shows the heavy metal concentrations(mg/kg) in soil from depth 0-10 cm, Oruama was significantly higher for Pb (9.28), Cd (1.88) and Fe (11.03) in comparison with JK 4 (8.74, 0.73, and 7.54), and Akinima (5.03, 0.52, and 5.51) respectively. While for Cr concentration, JK 4 (17.29) was significantly higher than Oruama (15.34), and Akinima (12.43).

Table 6. Heavy metal concentrations in soil from depth 0-15 cm

Heavy metals	mg/kg			
	Akinima	Oruama	Joinkrama 4 (JK 4)	FAO/ WHO ^{26,27}
Pb	9.04±0.06 ^a	13.99±0.01 ^b	11.57±0.14 ^c	100-300
Cr	19.22±0.16 ^a	23.57±0.11 ^b	21.71±0.11 ^c	350
Cd	0.83±0.02 ^a	2.28±0.11 ^b	1.29±0.09 ^c	3-8
Fe	9.51±0.10 ^a	13.36±0.14 ^b	13.16±0.11 ^b	50000

Values are expressed as Mean ± Standard Deviation (SD). Values in a column with different alphabetical superscripts do differ significantly ($p < 0.05$). (a) means Akinima is significantly different from other groups, (b) means Oruama is significantly different from other groups, and (c) means JK4 is significantly different from other groups.

Table 6 shows the heavy metal concentrations (mg/kg) in soil from depth 0-15 cm, Oruama was significantly higher for Pb (13.99), Cr (23.57), and Cd (2.28) in comparison with JK 4 (11.57, 21.71 and 1.29), and Akinima (9.04, 19.22 and 0.83 0.52) respectively. While for Fe concentration, there was no significant different between Oruama, and JK 4, although Akinima was significantly lower.

4.0 Discussion of Findings

Soil is the most valuable constituent of the farming ecosystem, and agricultural production depends on using agricultural soil sustainably so that plants can flourish. Pollutants such as heavy metals can contaminate soil, making it less suitable for use in agricultural production and other human activities^{2,28} and thus puts human health at risk. The accumulation of heavy metals in soil can degrade soil quality, diminish fertility and crop yields, hence posing risks to human and animal health²⁹. Heavy metals such as mercury (Hg), cadmium (Cd), nickel (Ni), lead (Pb), arsenic (As), and chromium (Cr) frequently serve as contaminants in soil environments. This form of contamination is extensive in the soil environment, broadly

distributed, and poses significant biological risks². This study considered the concentration of heavy metals in soil samples obtained from selected communities (Akinima, Oruama, and Joinkrama 4) in Ahoada West, Rivers State. Heavy metals such as lead (Pb), chromium (Cr), cadmium (Cd), and iron (Fe) were analysed and detected for their different concentrations.

The heavy metal analysis from soils in Akinima, Oruama, and JK4 communities revealed significant increases in lead (Pb), chromium (Cr), cadmium (Cd), and iron (Fe) concentrations with increasing depth ($p < 0.05$). Additionally, it revealed that lead, chromium, iron, and cadmium were below the FAO/WHO^{26,27} permissible limit for all the locations. The concentrations of Pb and Cd from this study was within the range of Akande and Ajayi¹ (Pb; 0.70-36.75, Cd 0.18 -0.63 mg/kg). Omeka and Igwe¹² had a similar Pb (1.15-5.6 mg/kg) value but higher Cd (1.15-3.95 mg/kg) and Fe (129.5-1304.16 mg/kg) levels when compared with this study. Additionally, there was a trend in increasing heavy metals concentration as the depth increases³⁰. This is because leaching, organic matter content, soil composition/texture and microbial activity can influence the mobility and retention of metals in soil²⁵.

The heavy metal concentration at different depths is shown on Table 4-6. For depths 0-5cm and 0-10cm, the concentration of heavy metals analyzed for the different communities was significantly in the order Oruama>Jk 4> Akinima except for Cr which had a different order of Jk4 > Oruama > Akinima respectively. Whereas for 0-15cm depth (Table 6), the concentration of all the heavy metals analysed was significantly in the order Oruama> Jk4> Akinima.

Lead (Pb) has no known biological role, and has been found to cause toxicity at quantities as low as 10 µg/kg in drinking water³¹. Lead has adverse effect on both children and adults. Lead (Pb) impairs both physical and cognitive development in children. Pregnant woman exposed to lead might result in anemia, renal impairment, cephalalgia, fatigue or irritable mood³². The lead concentration of the present study, was within the range of Adekiya *et al*³³ who had 0.7-

6.7 mg/kg. The Pb contents of the soil samples were lower than the FAO^{26,27} permissible limit. The observed Lead levels likely originated from petrochemical activities or petroleum-based products, as suggested by previous studies¹³.

Common forms of chromium in soils include Cr (III) and Cr (VI), which have different level of toxicity. Contrary to Cr (VII), which is a potent oxidizer and extremely poisonous, Cr (III) is a micronutrient and a non-toxic entity, exhibiting toxicity levels that are 10-100 times lower than those of Cr (VI). Cr (VI) is known to negatively impact microbial cell metabolism at high concentrations and alters the composition of soil microbial populations^{34,35}.

In humans, chromate compounds can induce DNA damage, and allergic dermatitis³². The chromium concentration of this study was comparable to Rani *et al.*³⁶ (9.65-127.21 mg/kg) but higher than Gogoi *et al.*³⁷ (Cr value of 0.56-0.84 mg/kg). The level of chromium in the soil from all the locations was below the FAO/WHO^{26,27} permissible limit of 300-350 mg/kg.

Cadmium is a non-essential metal, highly hazardous metal that impacts multiple organs and can, in certain instances cause death³⁸. Cadmium, present in fertilizers, pesticides, and industrial wastes, elevates the overall concentration of Cd in soils¹³, putting people at risk for cancer, anemia and kidney impairment. The level of cadmium in this study was below FAO/WHO^{26,27} limit of 5-8 mg/kg. This study aligns with the claims of Onwugbuta *et al.*³⁹ who had cadmium range of 1.91-15.04 mg/kg. Although the range, was wider than what this study had. Although, the concentration of all assessed heavy metals were below WHO standards, constant monitoring is necessary due to the disastrous impacts heavy metals have on terrestrial and aquatic life³⁴.

The research contributes greatly to the knowledge of soil health of soil health and agricultural sustainability in Ahoada West, Rivers State by emphasizing the role of soil as a resource for

agriculture and drawing attention to the risks associated by heavy metal contamination, which can hinder plant growth and pose health threats, to humans.

5.0 Conclusion

Heavy metals concentration obtained from soils of selected communities in Ahoada West, Rivers State disclosed that the concentrations of lead (Pb), cadmium (Cd), iron (Fe), and chromium (Cr) were below WHO standard. Despite the amounts of heavy metals being below WHO guidelines, regular monitoring and mitigation measures should be implemented to prevent potential future contamination. The limitations of the study include the examination of only three soil depths, which might have limited insights into changes at additional depths. The study focused on only three communities, which may distort the depiction of heavy metal poisoning in the region. Furthermore, the study focused on a single season, which might overlook seasonal variations in heavy metal concentrations. From the study, it is therefore recommended to develop a consistent program for tracking heavy metals level over time. It is necessary to encourage practices such as phytoremediation and organic amendments to improve soil health. And lastly, farmers and the communities should be educated about pollution hazards and sustainable practices.

Ethics Committee Approval: Ethical approval was granted by the Research and Ethics Committee of Rivers State University, Port Harcourt, Nigeria, with the reference number RSU-REC/2024/FS/0001.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. Chatgpt 4 was used to gain insight on how to write the limitations and recommendations.
- 2.
- 3.

References

1. Akande, F. O. and S A., Ajayi, 2017. Assessment of Heavy Metals Level in Soil and Vegetables Grown in Peri-Urban Farms around Osun State and the Associated Human Health. *International Journal of Environment, Agriculture and Biotechnology (IJEAB)*.2(6): 3250-3261. <http://dx.doi.org/10.22161/ijeab/2.6.61>
2. Angon, B. P., S. Islam, K. C. Shreejana, A. Das, N. Anjum, A. Poudel and S. A. Suchi, 2024. Sources, effects and present perspectives of heavy metals contamination: soil, plants and human food chain. *Heliyon*, 10:e28357. [Doi.org/10.1016/j.heliyon.2024.e28357](https://doi.org/10.1016/j.heliyon.2024.e28357)
3. Velayatzadeh, M., 2023. Heavy metals in surface soils and crops. *Intechopen*. [Doi:10.5772/intechopen.108824](https://doi.org/10.5772/intechopen.108824)
4. Gebeyehu, H.R. and L.D. Bayissa, 2020. Levels of heavy metals in soil and vegetables and associated health risks in Mojo area, Ethiopia. *Plos One* 15(1): 1-22. e0227883. <https://doi.org/10.1371/journal.pone.0227883>
5. Rashid, A., B.J. Schutte, A. Ulery, M. K. Deyholos, S. Sanogo, E. A. Lehnhoff and Beck, L., 2023. Heavy Metal Contamination in Agricultural Soil: Environmental Pollutants Affecting Crop Health. *Agronomy*, 13: 1521. <https://doi.org/10.3390/agronomy13061521>
6. Lemessa, F., B. Simane, A. Seyoum and G. Gebresenbet, 2022. Analysis of the concentration of heavy metals in soil, vegetables and water around the bole Lemi industry park, Ethiopia. *Heliyon* 8(e12429):1-13. <https://doi.org/10.1016/j.heliyon.2022.e12429>
7. Swain, K. C. (2024). Environmental pollution indices: a review on concentration of heavy metals in air, water, and soil near industrialization and urbanisation. *Discover Environment*. 2:5. <https://doi.org/10.1007/s44274-024-00030-8>
8. Onwuka, C., A. N. Eboatu, V. I. N. Ajiwe and E. J. Mora, 2021. Pollution Studies on Soils from Crude Oil Producing Areas of Rivers State, Niger Delta Region, Nigeria. *Open Access Library Journal*. 8: e7583:1-17. DOI: 10.4236/oalib.1107583
9. Zhao, H., Y. Wu, X. Lan, Y. Yang, X. Wu and D. Liyu, 2022 Comprehensive assessment of harmful heavy metals in contaminated soil in order to score pollution level. *Scientific Reports*, 12:3552. <https://doi.org/10.1038/s41598-022-07602-9>
10. Ren, X., G. Zeng, I. Ting, J. Wang, J. Wan, W. Jiajia, Y. Deng, Y. Liu and B. Peng, 2017. The potential impact on the biodegradation of organic pollution from composting technology for soil remediation. *Waste Management*, 72:138-149. doi: <https://doi.org/10.1016/j.wasman.2017.11.032>

11. Su, C., R. Xie, D. Liu and R. Liang, 2023. Responses of soil microbial communities to heavy metal stress in a coal-based industrial region in China. *Microorganisms*, 11(6):1392. <https://doi.org/10.3390/microorganisms11061392>
12. Omeka, M. E. and O. Igwe, 2021. Heavy metals concentration in soils and crop plants within the vicinity of abandoned mine sites in Nigeria: an integrated indexical and chemometric approach. *International Journal of Environmental Analytical Chemistry*, 1-12. <https://doi.org/10.1080/03067319.2021.1922683>
13. Wuana, R. A. and F. E. Okieimen, 2011. Heavy Metals in Contaminated Soils: A Review of Sources, Chemistry, Risks and Best Available Strategies for Remediation. *International Scholarly Research Network*. 2011:1-20. doi:10.5402/2011/402647
14. Briffa, J., E. Sinagra and R. Blundell, 2020. Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon*, 6:e04691. <https://doi.org/10.1016/j.heliyon.2020.e04691>
15. Gwary K. M., U. A. Zaria, M. S. Galadima and B. H. Diya'uddeen, 2019. Soil and Water Contamination due to Illegal Artisanal Refinery Activities: A Case Study of Okarki Community, Niger Delta Area, Nigeria. *Transactions of the VSB - Technical university of Ostrava, Safety Engineering Series*, XIV (2): 17-24, DOI 10.35182/tses-2019-0008
16. Chinedu, E. and C. K. Chukwuemeka, 2018. Oil Spillage and Heavy Metals Toxicity Risk in the Niger Delta, Nigeria. *Journal of Health & Pollution*, 8 (19):1-8. Doi:10.5636/2156-9614-8.19.180905
17. Ogbeide, O. and B. Henry, 2024. Addressing heavy metal pollution in Nigeria: Evaluating policies, assessing impacts, and enhancing remediation strategies. *Journal of Applied Sciences and Environmental Management*, 28(4):1007-1051. Doi: <https://dx.doi.org/10.4314/jasem.v28i4.5>
18. Erebi, J.L. and E.E. Davidson, 2023. Application of Vegetation Indices for Detection and Monitoring Oil Spills in Ahoada West Local Government Area of Rivers State, Nigeria. *Journal of Geographical Research*. 6(3): 29-41. DOI: <https://doi.org/10.30564/jgr.v6i3.5817>
19. Etori, E. S, O. S Etori and D. Bekee, 2021. Total Petroleum Hydrocarbons Contamination of the Surface Water and Sediments of Orashi River, Engenni, Ahoada West, Rivers State, Nigeria. *Asian Review of Environmental and Earth Sciences*, 8(1): 68-76. DOI: 10.20448/journal.506.2021.81.68.76
20. Umar, H. A., M. F. A. Khanan, S. B. Magashi, M.M. Ja'afar and M. J. Sani, 2022. fractal analysis for oil spills clustering in Ahoada communities of the Niger Delta region of Nigeria. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, XLVIII:371-377. <https://doi.org/10.5194/isprs-archives-xxviii-4-w6-2022-371-2023>
21. Bello, A. T. and T. Nwaeke, 2023. Impacts of oil exploration (oil and gas conflicts; niger delta as a case study). *Journal of Geoscience and Environmental protection*, 11(3).189-200. Doi:10.4236/gep.2023.113013
22. Ewim, D. R. E., O. F. Orikpete, T. O. Scott, C. N. Onyebuchi, A. O. Onukogu, C. G. Uzougbo and C. Onunka, 2023. Survey of wastewater issues due to oil spills and pollution in the Niger Delta area of Nigeria: a secondary data analysis. *Bulletin of the National Research Centre*, 47:116. <https://doi.org/10.1186/s42269-023-01090-1>
23. Okeke O., E. Ezeh, H. C. Okeke, C. C. Aniobi and C. I. Akagha, 2020. Comparion of soil samples from selected anthropogenic sites within Enugu metropolis for

- physiochemical paramters and heavy metal levels determination. *Journal of Environmental protection*, 11:848-861. <https://doi.org/10.4236/jep.2020.1110053>
24. Gong, I., D. Qiu, X. Yao and G. Yang, 2020. Determination of heavy metals in plant sample pre-treatment methods. *Journal of Physics: Conference Series*, 1549:022008. Doi:10.1088/1742-6596/1549/2/022008
 25. Duru, C. S., E. A. Echiegu, C. C. Anyadike, G. U. Alaneme and M. E. Okechukwu, 2024. Spatial; variability of heavy metals concentrations in soil of auto-mechanisc workshop clusters in Nsukka, Nigeria. *Scientific Reports*, 14:9681. Doi.org/10.1038/s41598-024-60044-3
 26. FAO/WHO, 2005. Regulations regarding quality standards for soil and ground . Food and agriculture organization. <https://faolex.fao.org>
 27. FAO/WHO (Food and Agriculture Organization/World Health Organization), 2016. General Standard for Contaminants and Toxins in Food and Feed (Codex Alimentarius International Food Standards.CODEX STAN, p. 193.
 28. Alengebawy, A., S. T. Abdelkhalek, S. R. Qureshi and M. Wang, 2021. Heavy Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health Implications. *Toxics*, 9(3):42. <https://doi.org/10.3390/toxics9030042>
 29. Zhyrgalovaa A., S. Yelemessova, B. Ablaikhana, G. Aitkhozhayevaa and A. Zhildikbayev, 2024. Assessment of potential ecological risk of heavy metal contamination of agricultural soils in Kazakhstan. *Brazilian Journal of Biology*, 84: e280583. <https://doi.org/10.1590/1519-6984.280583>
 30. Enya, O., C. Lin and J. Qin, 2019. Heavy metal contamination status in soil-plant system in upper Mersey Estuarine Floodplain, Northwest England. *Marine pollution Bulletin*, 146:292-304. <https://doi.org/10.1016/j.marpolbul.2019.06.026>
 31. WHO, 2016. Lead in drinking water. Background document for development of WHO Guidelines for Drinking-water Quality. World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
 32. Abdullahi, A., M. A Lawal and A. M. Salisu, 2021. Heavy Metals in Contaminated Soil: Source, Accumulation, Health Risk and Remediation Process. *Bayero Journal of Pure and Applied Sciences*, 14 (1): 1 – 12. <http://dx.doi.org/10.4314/bajopas.v14i1.1>
 33. Adekiya, O. A, A. Oloruntoba, B. S. Ewulo, T. Ogunbode and T. Sangoyomi, 2024. Heavy Metal Pollution in Soils near Ilesha Gold Mining Area, Nigeria. *The Open Agriculture Journal*, 18: e18743315290216. DOI: 10.2174/0118743315290216240123111021
 34. Zulfigar, U., F. U. Haider, M. Ahmad, S. Hussain, M. F. Maqsood, M. Ishfaq, B. Shahzad, M. M. Wagas, B. Ali, M. N. Tayyab, S. A. Ahmad, I Khan and S. M. Eldin, 2022. Chromium toxicity, speciaton and remediation strategies in soil-plant interface: A critical review. *Frontiers in Plant Science*, 13:1081624. Doi: 10.3389/fpls.2022.1081624
 35. Ullah, S., Q. Liu, S. Wang, A. U. Jan, H. M. A. Sharif, A. Ditta and G. Wang, 2023. Sources, impacts, factors affecting Cr uptake in plants, and mechanisms behind phytoremediation of Cr-contaminated soils. *Science of the total environment*. 899:165726. <https://doi.org/10.1016/j.scitotenv.2023.165726>
 36. Rani, I., Chaudhary S and T. Agarwal, 2021. Evaluation of Heavy Metals Contamination in Agricultural Soils. *Research Square*. 1-14. DOI: <https://doi.org/10.21203/rs.3.rs-327423/v1>
 37. Gogoi, N., A. Sarma, M. Choudhury, P. Samanta, M. Faizan and S. Sadhak , 2024. Soil heavy metal pollution and ecological risk assessment in disturbed and undisturbed soil

- of Morigaon, Assam. *Discover Environment* 2:13 <https://doi.org/10.1007/s44274-024-00039-z>
38. Rahimzadeh, M., R., S. Kazemi and A. Moghadamnia, 2017. Cadmium toxicity and treatment: An updatee. *Caspian Journal of Internal Medicine*, 8 (3):135-145 doi:10.22088/cjim.8.3.135
39. Onwugbuta, N., I. K. E. Ekweozor, A. P. Ugbomeh G. N. Anaero-Nweke, and K.N.O. Bobmanuel, 2022. Assessment of Heavy Metals in Soils and Manihot esculenta of Some Selected Communities in Niger Delta. *International Journal of Advanced Research in Biological Sciences*, 9(12): 74-96. <http://dx.doi.org/10.22192/ijarbs.2022.09.12.007>