

Original Research Article

Survey of parasitic nematodes associated with Cotton in Burkina Faso

ABSTRACT:

Aims: An inventory of parasitic nematodes associated with Cotton is carried out during the 2017-2018 wet season in the major production areas in Burkina Faso.

Study Design: Indeed, the cotton plant is attacked by a large number of pests, including plant parasitic nematodes, which is observed at high population levels in the country's main cotton-growing areas.

Place and Duration of Study: The study is conducted in Burkina Faso, for the Institute of Environment and Agricultural Research, Farako-Bâ station, for few months.

Methodology: 94 samples of roots and adjoining soils are taken and nematodes are extracted from 250 cubic centimeters (cc) of soil using the Seinhorst elutriator method. The nematodes present in the roots are extracted by the Seinhorst sprinkler method. Population densities are expressed in terms of Number of nematodes/dm³ of soil and Number of nematodes/g of roots.

Results: About ten (10) genera of plant-parasitic nematodes are identified and represented by *Meloidogyne*, *Pratylenchus*, *Helicotylenchus*, *Scutellonema*, *Tylenchorhynchus*, *Telotylenchus*, *Rotylenchulus*, *Xiphinema*, *Criconemella* and *Paratrichodorus*. Under the conditions of Burkina Faso, five (5) nematodes are considered as being able to cause yield losses on cotton, namely *Helicotylenchus*, *Scutellonema*, *Tylenchorhynchus*, *Pratylenchus* and *Telotylenchus* because they are frequent and abundant with frequencies between 100% and 36% for densities varying between 7,328 and 1,360 nematodes/dm³ of soil.

Conclusion: On the other hand, the nematodes *Meloidogyne* and *Rotylenchulus* as major pests of Cotton throughout the world have been observed at low frequencies but at often high levels of population abundance, Research perspectives focus on the characterization of the main parasitic nematodes of cotton and the development of control methods oriented towards resistance varieties and cultural practices which suggests that their development requires particular soil and soil moisture conditions.


Keywords: cotton; parasitic nematode; abundance; frequency; Burkina Faso

1. INTRODUCTION

Agriculture employs more than 80% of the population of Burkina Faso and contributes nearly 40% to the Gross Domestic Product (GDP). rainfed agriculture and occupy an important place because the crop employs nearly 4 million people and contributes more than 4% to GDP and about 14% of export earnings in recent years [1]. The area sown in 2016-2017 are estimated at nearly 740,000 ha, and the production was estimated at nearly 683,000 tons of seed cotton with relatively low yields of less than one ton per hectare [2]. The cotton sector has experienced a significant decline in recent years producing 407,308 tons with yields of 655 kg/ha during the 2022-2023 agricultural season [3]. This situation can be explained, among other things, by uncertain climatic conditions (rainfall and temperature) and damage due to crop pests [3]. Finally, the intensification of agricultural production in the cotton-growing area has favored the development of pests and diseases and the destruction of the many beneficial auxiliary organisms present in the soil [4]. Indeed, the cotton plant is attacked by many pests, including plant-parasitic nematodes, which is observed at high population levels in the country's main Cotton-growing areas. However, significant losses due to this group of pests, and particularly to the root-knot nematodes *Meloidogyne incognita*, is reported on cotton throughout the world ([5]; [6]). [7] have noted that the genera *Helicotylenchus*, *Pratylenchus*, *Hoplolaimus* and *Rotylenchulus* are considered as the most important and likely to cause yield losses on Cotton. *Meloidogyne incognita* and *Rotylenchulus reniformis* are considered to be the major pests limiting Cotton yield in the United States of America [8]. Cotton is known to be heavily attacked by the root-knot nematodes *Meloidogyne* spp. with yield losses of up to 60% for population densities of 1,000 nematodes/100 cm³ of soil estimated as a threshold of harmfulness ([9]; [10]; [11]). *M. incognita* is one of the world's most loss nematodes, with estimated annual losses of US\$100 billion on crops [12] and US\$283 million on Cotton in the United States of America ([13]; [8]). The aim of the inventory is to identify parasitic nematodes associated with cotton in order to develop appropriate control methods with a view to improving productivity and production of Cotton.

2. MATERIAL ET METHODS

The inventory of cotton parasitic nematodes concerns the three cotton-growing areas represented by the Société des Fibres Textiles (SOFITEX), the Société Cotonnière du Gourma (SOCOMA) and Faso Coton, during the 2017-2018 agricultural campaign (Map 1).



Map 1: Cotton-growing areas of Burkina Faso

The composite sample consisted of 10 samples of roots and adjoining soils in 0 - 20 cm horizon is collected for seed production fields at the capsulation-early maturity stage. 94 samples of roots and adjoining soils is taken and nematological analyses is carried out at the Nematology Laboratory of the Institute of Environment and Agricultural Research (INERA), Farako-Bâ/Bobo-Dioulasso Station. Nematodes is extracted from 250 cubic centimeters (cc) of soil using the [14] elutriator method. The nematodes present in the roots is extracted by the sprinkler method [15]. Population densities is expressed in terms of number of nematodes/dm³ of soil and number of nematodes/g of roots. The morpho-biometric identification of the nematodes was done according to the identification key of [16]. The data analysis focused on the frequency and abundance of parasitic nematode populations recorded. Frequency is calculated as the total Number of samples where the nematode is present divided by the total Number of samples collected multiplied by 100.

$$\text{Frequency } F = \frac{e}{n} \times 100$$

e = Number of samples where the nematode considered is present
n = Total Number of samples

Abundance is calculated by the sum of the samples where the nematode divide by the number of samples.

$$\text{Abundance A} = \frac{\sum X_i}{n}$$

X_i = Number of individuals of the nematode per dm³ of soil or per gram of roots

n = Number of samples where the nematode under consideration is present

The importance of the main genera of nematodes parasitic on Cotton is determined according to the method of [17], which proposed that a nematode is said to be abundant in the soil, if the abundance is ≥ 200 individuals/dm³ of soil and in the roots if the abundance is ≥ 20 individuals/g of roots. A nematode is said to be share in soil or roots, if it is observing in at least 30% of the samples. Statistical analyses is performed with the XLSTAT 2016 software and the separation of the means according to the Newman Keuls test.

3. RESULTS AND DISCUSSION

3.1. RESULTS

3.1.1. Frequency and abundance of observed nematodes

About ten genera of parasitic nematodes are associated with the cotton plant and represented by *Meloidogyne*, *Pratylenchus*, *Helicotylenchus*, *Scutellonema*, *Tylenchorhynchus*, *Telotylenchus*, *Rotylenchulus*, *Xiphinema*, *Criconemella* and *Paratrichodorus* (Table 1).

3.1.2. Nematodes extracted from soil samples

Helicotylenchus, *Scutellonema*, *Tylenchorhynchus*, *Pratylenchus* and *Telotylenchus* are frequent and abundant with frequencies between 100% and 36% for densities varying between 7,328 and 1,360 nematodes/dm³ of soil. According to the method of [17], this group of nematodes can be capitalized as the most important on Cotton and likely to cause significant damage (Frequency $>30\%$ and Abundance <200 nematodes/dm³ of soil). *Xiphinema*, *Rotylenchulus* and *Meloidogyne* are infrequent ($<30\%$) but abundant (<200 nematodes/soil dm³) and their presence may be associated with particular environmental conditions (soil texture, humidity, etc.). *Paratrichodorus Minor* and *Criconemella onoensis* are uncommon ($<30\%$) and not very abundant (<200 nematodes/dm³ of soil) and can be considered as little damage to Cotton.

3.1.3. Nematodes extracted from the roots

Pratylenchus is observed in 73% of the samples with root population densities of 30 nematodes/g of roots; This nematode is viewed to be frequent and abundant (frequency >30 and abundance $>30\%$) and can be considered as a parasite of Cotton according to the method of [16]. The nematodes *Scutellonema* and *Helicotylenchus* are infrequent ($>30\%$) and scarce (<20 nematodes/g

of roots). This group of nematodes is considered mainly as ectoparasites and therefore little observed in the roots.

Table 1: Frequency and densities of nematode communities observed on cotton

Nematode genera	Frequency	Minimum	Maximum	Average	standard error
Extracted from soils	(%)	Nber N/dm ³	Nber N/dm ³	Nber N/dm ³	Nber N/dm ³
<i>Helicotylenchus</i>	100	40	37,040	7,328	± 781
<i>Scutellonema</i>	100	60	4,800	1,133	± 107
<i>Tylenchorhynchus</i>	75	0	7,560	464	± 117
<i>Pratylenchus</i>	71	0	4,940	584	± 93
<i>Telotylenchus</i>	36	0	1,360	121	± 27
<i>Xiphinema</i>	26	0	380	25	± 781
<i>Rotylenchulus</i>	19	0	1,460	41	± 18
<i>Meloidogyne</i>	13	0	200	7	± 3
<i>Paratrichodorus</i>	9	0	120	6	± 2
<i>Criconemella</i>	3	0	20	1	± 0
Extracted from roots	(%)	Nber N/g	Nber N/g	Nber N/g	Nber N/g
<i>Pratylenchus</i>	73	0	30	3	± 1
<i>Scutellonema</i>	14	0	7	1	± 0
<i>Helicotylenchus</i>	27	0	14	1	± 0

Legend: Nber/dm³: number of nematodes/dm³ of soil; Nber/g: number of nematodes/g of roots

3.1.4. Nematode community densities by cotton-growing areas and prospecting localities

The nematode samples is taken in the 3 cotton-growing areas represented by the cotton companies SOFITEX (18 sites), SOCOMA (7 sites) and Faso Coton (7 sites).

The high population densities of *Pratylenchus* is observed in the SOFITEX zone with 1,029 nematodes/dm³ of soil with the highest densities observed on the Sarba site with 2,960 nematodes/dm³ of soil (P<0.05). Respective population densities of 350 and 187 individuals/dm³ of soil are observed, respectively in the SOCOMA and Faso Coton zones. The cotton-growing area of SOCOMA appears to be more infested with *Helicotylenchus* (1,0408 individuals/dm³ soil), *Scutellonema* (1,546/dm³ soil), *Tylenchorhynchus* (243/dm³ soil) and *Telotylenchus* (264/dm³ soil) (P<0.05), representing, with *Pratylenchus*, the group of nematodes likely to cause significant damage to cotton. The SOFITEX zone appears to be less infested for these nematodes, compared to the Faso Coton zone, except for *Helicotylenchus*, with an average density of 7,239 individuals/dm³ of soil. The highest population densities of *Helicotylenchus* were observed at the Dédougou site with 29,600 individuals/dm³ of soil in the SOFITEX zone and at the Gongongwana/Pama site with 25,280 individuals/dm³ of soil in the Faso cotton zone. The total

population densities of all parasitic nematodes observed in soil samples are higher in the SOCOMA zone, with 12,911 individuals/dm³ of soil (P<0.05). In the SOFITEX zone, the total soil populations are estimated at 9,583 nematodes/dm³ of soil and 6,624/dm³ of soil in the Faso Coton zone. Total populations of root-extracted nematodes are relatively low across cotton-growing areas with less than 10 nematodes/g of roots.

Table 2: Population densities of the leading parasitic nematodes according to cotton-growing areas and sampling sites

Cotton-Zone	Location	Pray/dm ³	Heli/dm ³	Scute/dm ³	Tyle/dm ³	Telo/dm ³	Tot/dm ³	Tot/g rac.
SOFITEX	Satiri	4,680	3,360	680	560	0	9,280	12
	Sara	1,160	10,030	650	180	0	12,070	5
	Bondokuy	1,180	3,755	1,030	55	0	6,140	7
	Ouarkoye	1,193	4,267	587	67	0	6,307	5
	Dédougou	0	29,600	800	80	0	30,560	1
	Dara	847	3,167	733	207	0	4,960	1
	Boron	700	10,640	1,180	180	0	12,940	1
	Tcheriba	900	6,040	340	340	0	7,640	2
	Bognounou	380	3,180	880	100	0	4,620	3
	Lon	1,345	4,250	915	90	25	6,675	7
	Tabou	1,090	3,390	1,273	170	203	6,243	2
	Sarba	2,960	10,920	1,720	40	280	16,600	4
	Bafor	45	6,290	1,280	170	0	7,810	4
	Klesso	1,320	3,140	420	0	120	5,160	4
	Soumousso	260	5,380	1,820	180	260	7,940	7
	Gnafogo	150	8,720	1,030	60	110	10,110	1
Noumoudara	193	5,527	733	53	20	6,600	3	
Tapoko	120	8,640	1,860	20	0	10,840	1	
Average		1,029	7,239	996	142	57	9,583	4
SOCOMA	Kouaré	749	7,832	335	886	349	10,194	7
	Niendouga	270	7,410	590	80	460	8,830	26
	Gongongwana	25	25,280	2,205	45	435	28,560	1
	Diapangou	80	17,100	1,330	260	20	18,830	4
	Dourtenga	327	1,660	1,393	147	60	3,593	2
	Tangonko	7	973	3,800	173	433	5,400	9
	Koghin	990	12,600	1,170	110	90	14,970	3
Average		350	10,408	1,546	243	264	12,911	7
Faso coton	Gonbloré V6	67	1,567	247	4,793	0	6,680	1
	Gonbloré V7	0	1,560	100	4,640	0	6,300	0
	Gnangdin	0	7,707	2,107	7	0	9,820	1
	Bittou	100	3,090	870	150	70	4,280	1
	Zekeze	900	5,967	1,740	20	0	8,647	2
	Ouadav1	200	260	3,560	0	0	4,020	1
	Kaïbo Sud	45	4835	1,085	648	8	6,620	1

Everage	187	3,569	1,387	1,465	11	6,624	1
---------	-----	-------	-------	-------	----	-------	---

Legend : Praty. : *Pratylenchus* ; Heli : *Helicotylenchus* ; Scute : *Scutellonema*

Tyle : *Tylenchorhynchus* ; Telo : *Telotylenchus* ; Tot : Total in soil and roots

UNDER PEER REVIEW

3.2. DISCUSSION

The results of this study, which covered all major areas of Burkina Faso, are in line with the work of [7] which had identified *Helicotylenchus*, *Pratylenchus*, *Hoplolaimus*, *Tylenchorhynchus*, *Rotylenchulus* and *Meloidogyne* as the main nematodes associated with the cotton-maize-sorghum cropping system in the cotton basin of the Houndé zone in western Burkina Faso. This study did not make it possible to rule on the pathogenicity of the root-knot nematodes *Meloidogyne* spp. on cotton, unlike several studies conducted in the primary production areas of the United States and South Africa where *Meloidogyne incognita* race 3, is known to cause significant damage [18]. *M. incognita* (Cophoid and white), Chitwood, and *Rotylenchulus reniformis* (Linford and Oliveira) are identified as major yield-limiting pests of upland cotton [19]. *M. incognita* is considered to be the nematode causing yield losses on a global scale on crop plants [12]. Our research in the main cotton-growing areas of Burkina Faso shows that the root-knot nematodes *Meloidogyne* spp. do not constitute a significant problem because they are observed at low frequencies (13%) and at average densities of 7 nematodes/dm³ of soil. Studies on the pathovars of *Meloidogyne incognita*, recognized as the dominant species, would make it possible to decide on its pathogenicity. However, *M. acronea*, known to be a parasite on cotton and, present in South Africa and Malawi, is not present in Burkina Faso [20]. The present study showed a predominance of the lesion nematode *Pratylenchus brachyurus* in the SOFITEX zone, a former cotton production area in western Burkina Faso (P<0.05) where cotton is generally in rotation with maize, the preferred host plant of this group of nematodes. Its abundance in this area confirms its adaptation to this culture. The lesion nematode *Pratylenchus* is known to cause significant damage to cotton and [21] have shown that the presence of *P. brachyurus* would induce a decrease in *Meloidogyne incognita* populations. Similar competition has been observed between *Meloidogyne incognita* and *Rotylenchulus reniformis*. As for the kidney-shaped nematode *Rotylenchulus reniformis*, it is known as an important pest of cotton throughout the world mainly, in soils with a high silt/clay content ([18]; [22]). This species may pose a danger to cotton in Burkina Faso where it is observed in high populations but limited to a few sites with 1,460 nematodes/dm³ of soil. The species has experienced a substantial expansion in the southern United States, causing significant damage and losses to cotton ([24]; [8]). Several ectoparasitic nematodes is identified on Cotton, mainly in the Central, Cotton-producing states of the United States of America, but their pathogenicity has not been demonstrated. These are mostly *Helicotylenchus*, *Paratrichodorus minor*, *Tylenchorhynchus* spp. but with very low population levels (less than 100 nematodes/100cm³ of soil or less than 10/dm³ of soil) ([24]; [25]). On the other hand, other studies have shown high densities of these groups of nematodes, which are often considered secondary but likely to cause yield losses on cotton ([26]; [27]). The high populations of the nematodes

Helicotylenchus, *Scutellonema*, *Tylenchorhynchus* and *Telotylenchus annulatus* observed on cotton can cause significant damage to this crop under the conditions of Burkina Faso.

4. Conclusion

About ten genera of parasitic nematodes are associated with cotton in Burkina Faso, seven (7) of which are likely to cause yield losses on cotton in Burkina Faso. *Meloidogyne*, *Pratylenchus*, *Helicotylenchus*, *Scutellonema*, *Rotylenchulus*, *Tylenchorhynchus* and *Telotylenchus*. The nematodes *Meloidogyne* and *Rotylenchulus*, known as major cotton pests throughout the world, have been observed at low frequencies but often at high population levels, suggesting their development in particular soil and soil moisture conditions. In perspective, it will be a question of identifying the existence of races of nematodes belonging to the species *Meloidogyne javanica* and *M. incognita* by molecular characterization that can attack cotton in Burkina Faso.

5. REFERENCES

- [1] - **AICB. 2023a.** Rapport du Burkina Faso. 81^{ème} Assemblée Plénière du Comité Consultatif International du Coton. Mumbai, 02 au 05 décembre 2023. 8p.
- [3] - **AICB. 2023b.** Campagne cotonnière 2023-2024 au Burkina Faso. Ministère de l'Industrie, du Commerce et de l'Artisanat. <https://www.commerce.gov.bf/accueil/actualites>.
- [2] **AICB. 2008.** Note d'information sur la filière coton du Burkina. 10p.
- [4] - **Berimey M. A., 2012.** Contrôle des nématodes et des symphytes, parasites des racines de l'ananas par l'utilisation de plantes de service. Master En Sciences – Technologies –Sante, Université Des Antilles Et De La Guyane. 30p
- [5] - **Taylor AL., Sasser J.N. and Nelson L.A., 1982.** Relationship of climatic and soil characteristics to geographical distribution of nematode species in agricultural soils, Raleigh, North Carolina State University and USAID, NC., 65 p.
- [6] - **Pages S.L.J., 1983.** Biological studies of the African cotton root nematode, *Meloidogyne acrona*. Ph.D thesis, University of London., 294 p.
- [7] - **Sawadogo A., Dakouo D., Konaté Y. A. & Thio B. 1998-1999.** Les nématodes parasites associés au système de culture coton-maïs-sorgho dans la zone de Houndé (Ouest du Burkina Faso). *Sci. et Tech., Série Sciences Naturelles*, Vol. 23, n° 2 - juillet-décembre 1998 - Janvier - juin 1999.

- [8] - **Lawrence K., 2022.** Reniform nematode (*Rotylenchulus reniformis*) and its interactions with cotton (*Gossypium hirsutum*) Pp. 94-99 in R. A. Sikora, J. Desaeger, and L. Molendijk, eds. Integrated nematode management: state of-the-art and visions for the future. Oxfordshire, UK: CAB International. doi:10.1079/9781789247541.0014.
- [9] - **Blasingame D., Gazaway W., Goodell P., Kemerait R., Kirkpatrick T., Konning S., Lawrence G. W., McClure M., Mueller J., Newman M., Overstreet C., Phipps P., Rich J., Thomas S., Wheeler T. and Wrather A. 2002.** Cotton nematodes: your hidden enemies National Cotton Council, Memphis, TN.
- [10] - **Doshi R. A., King R. L. and Lawrence G. W.. 2010.** Classification of *Rotylenchulus reniformis* numbers in cotton using remotely sensed hyperspectral data on self-organizing maps. *Journal of Nematology* 42:179–193.
- [11] - **Moore S. R. and Lawrence K. S. 2012.** *Rotylenchulus reniformis* in cotton: current methods of management and the future of site-specific management. *Nematropica* 42:227–236.
- [12] - **Wram, C. L., and Zasada, I. A. 2019.** Shortterm effects of sublethal doses of nematicides on *Meloidogyne incognita*. *Phytopathology* 109:1605- 1613. doi:10.1094/phyto-11-18-0420.
- [13] - **Forghani, F., and Hajihassani, A. 2020.** Recent advances in the development of environmentally benign treatments to control root-knot nematodes. *Frontiers in Plant Science* 11:1125. doi:10.3389/fpls.2020.01125.
- [14] - **Seinhorst J.W., 1962.** Modifications of the elutriation method for extracting nematodes from soils. *Nematologica*. 8 : 117- 28.
- [15] - **Seinhorst J.W., 1950.** De betekenis van de grond voor het optreden van aanstasting door het stengelaaltje (*Ditylenchus dipsaci*) (Kühn Filipjev). Tijdschr. *Plantenziekten*, 56: 289-348.
- [16] **Mai W.F. & Lyon H.H. 1975.** Pictorial Key to Genera of Plant Parasitic Nematodes. 4th edition Revised. Ithaca and London, Comstock Publishing Associates, Cornell University Press. pp219.
- [17] - **Fortuner R., Merny G., 1973.** Les nématodes parasites des racines associés au riz en Basse Casamance (Sénégal) et en Gambie. *Cah. ORSTOM, Sér. Biol.*, 21: 3-20.
- [18] - **Starr J.L., Carneiro R.G. et Ruano O. 2005.** Nematode Parasites of Cotton and other Tropical Fibre Crops. In Luc M., Sikora R.A. and Bridge J., eds. *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*. Wallingford, U.K: CAB International, 733-750.

- [19] - **Moore S. R. and Lawrence K. S. 2013.** The Effect of Soil Texture and Irrigation on *Rotylenchulus reniformis* and Cotton. *J. Nematol*, 45(2):99–105.
- [20] - **CABI, 2019.** *Meloidogyne acronea* (African cotton root nematode). datasheet on *Meloidogyne acronea* covers Identity, Overview, Distribution, Dispersal, Hosts/Species Affected, Diagnosis, Biology & Ecology, Natural Enemies. <https://www.cabidigitallibrary.org> ›.
- [21] - **Gay, C.M. and Bird, G.W. 1973.** Influence of concomitant populations of *Pratylenchus brachyurus* and *Meloidogyne incognita* on root penetration and population dynamics. *Journal of Nematology* 5,212–217.
- [22] - **Gordon, K. L., Schrimsher, D. W., and Lawrence, K. S. 2022.** Additional fertilizer and nematicide combinations on upland cotton to manage *Rotylenchulus reniformis* and *Meloidogyne incognita* in Alabama. *Journal of Nematology* 54:1-15.
- [23] - **McLean, K.S. and Lawrence, G.W. 2003.** Efficacy of aldicarb to *Rotylenchulus reniformis* and biodegradation in cotton field soils. *Journal of Nematology* 35, 65–72.
- [24] - **Bridge, J. 1992.** Nematodes.: Cotton Diseases. R. J. Hillocks, ed. CAB International. Wallingford, UK. 331-353.
- [25] - **Wrather, J. A., Niblack, T. L., and Milam, M.R. 1992.** Survey of plant-parasitic nematodes in Missouri cotton fields. Suppl. *J. Nematol.* 24:779-782.
- [26] - **Singh A.U. and Mishra, S.D. 2016.** Nematode problems in cotton crop and their management Archana. *Current Nematology*, 27 (1): 103–113.
- [27] - **Schumacher, L.A.; Grabau, Z.J.; Wright, D.L.; Small, I.M.; and Liao, H.L. 2020.** Nematicide influence on cotton yield and plant-parasitic nematodes in conventional and sod-based crop rotation. *Journal of Nematology* 52: e2020-34. DOI: 10.21307/jofnem-2020-034.