

EFFECT OF OVIRICH® ON BLOOD CHEMISTRY IN WHITE LEGHORN LAYER

ABSTRACT

Present experiment was conducted to discern the effect of Ovirich® in White Leghorn birds (90 in numbers; 22 weeks age) which were divided into three treatment groups using completely randomized design (CRD). The control group (T1) received the basal diet without any supplementation, while in treatment groups T2 and T3; the basal diet was supplemented with 0.75 kg and 1.0 kg of supplement per ton of basal feed, respectively. Blood chemistry was assessed at the end of feeding trial. Significant differences ($P \leq 0.05$) were observed in haemoglobin, PCV, TEC, TLC, MCV, MCH, MCHC, SGPT, SGOT, total protein, albumin, globulin, glucose, total cholesterol, triglycerides, serum calcium and phosphorus in supplemented groups. It was concluded that Ovirich® in the basal diet can affect blood biochemistry in White Leghorn layers.

Keywords: blood chemistry, Ovirich®, white leghorn

1.Introduction: Ovirich® supplement has beneficial phytochemical molecules from Capparis spinosa (Caper bush), Terminalia arjuna (Arjuna tree), Cichorium intybus (Chicory), Solanum nigrum (Black nightshade) Tamarix gallica, Achillea millefolium and Andrographis paniculata. It also contains some essential minerals like Organic Chromium, Zinc fortified with other minerals viz. (Calcium, Phosphorus, Manganese, Cobalt, Selenium) and Yeast complex. In this experiment effect of supplementing Ovirich® on blood parameters were studied in white leghorn.

2.Materials and methods: White Leghorn birds (90 in number, 22 weeks age) kept at Instructional Poultry Farm, Nagla were selected and divided into three treatment groups using completely randomized design (CRD) for feeding trial of 14 weeks. The control group (T1) received the basal diet without any supplementation, while in treatment groups T2 and T3, the basal diet was supplemented with 0.75 kg and 1.0 kg of supplement per ton of basal feed, respectively. Standard basal diets for laying birds was prepared by mixing the feed ingredients to meet the nutrient requirements of birds as per specification recommendations of BIS (2007). Proximate composition of experimental feed was analysed using AOAC

(2003). Phytochemical analysis of Ovirich was conducted at the Department of Biochemistry at GBPUA&T, Pantnagar. Ovirich sample was subjected to analysis to determine their nutritional parameters like phenolic compound, amino acid content (methionine, arginine, tryptophan and lysine), total content of flavonoid, ascorbic Acid. Antinutritional factors like phytic acid and tannin were also analysed. Standard procedure was followed during biochemical analysis. Arginine was estimated by Greenstein and Winitz (1961), methionine by Horn *et al.*, (1946), Lysine by Sadasivam and Manickam (1992), tryptophan by Spies and Chambers (1949), Phenol by Bray and Thorpe (1954) Total flavonoids by Jagadish *et al.*, (2009), Tannins by Kavitha and Indira (2016), Phytic acid by Haug and Lantzsch (1983) and Total anti-oxidants by Prieto *et al.* (1999) method.

At the end of the experiment, around 2 ml of blood sample was collected into an EDTA-coated vial for the assessment of haematological parameters. Additionally, approximately 3 ml of blood was collected into plain vials (devoid of anticoagulant) from the wing vein. These plain vials were kept in a slanting alignment at room temperature for a period of 3 to 4 hours to facilitate the natural clotting of the blood. Once the blood had clotted, serum samples were separated by subjecting them to centrifugation at 3000 rpm for duration of 10-15 minutes. The separated serum was carefully collected into eppendorf tubes and subsequently stored at a temperature of -20°C within a deep freeze. These samples were labelled with dates and sample numbers, and were kept in this frozen state until the completion of biochemical estimations.

Haematological parameters

Haemoglobin: The concentration of hemoglobin (g/dl) was determined using the cyanomethemoglobin method (Drabkin and Austin, 1932).

Packed cell volume (PCV) : The packed cell volume (PCV) was determined using the Microhematocrit method, following the procedure given by Martha *et al.* (2012).

Total erythrocytes count and total leucocyte count : The total erythrocyte count (TEC) and total leukocyte count (TLC) of the blood samples were assessed using the Hemocytometer (Neubauer's counting chamber) method, as outlined in the procedure by the author Natt and Herrick (1952).

Mean corpuscular volume : Mean corpuscular volume (MCV) was expressed in femtolitre (fl) and calculated as per following formula.

$$MCV = \frac{PCV (\%)}{TEC (10^6/\mu l)} \times 10$$

Mean corpuscular haemoglobin (MCH) :The mean corpuscular haemoglobin (MCH) was expressed in terms of picogram and calculated as per formula prescribed below.

$$MCH = \frac{\text{Hb (g/dl)}}{\text{TEC (106/\mu l)}} \times 10$$

Mean corpuscular haemoglobin concentration (MCHC):The mean corpuscular haemoglobin concentration (g/dl) was calculated as per formula given below.

$$MCHC = \frac{\text{Hb (g/dl)}}{\text{PCV\%}} \times 100$$

Serum biochemical parameters :Serum Glucose, Uric Acid, Serum Glutamic Oxaloacetic Transaminase (SGOT), Serum Glutamate Pyruvate Transaminase(SGPT) and Lactate dehydrogenase(LDH) were estimated using Erba diagnostic kit using GOD-POD Method, Uricase - PAP method, IFCC method, IFCC method and DGKC method.

Serum protein profile:

The total protein and albumin concentrations in the serum were assessed through spectrophotometric analysis using the Erba Diagnostics kits. Total protein was quantified using the bromocresol green (BCG) method at a wavelength of 540 nm, and albumin was determined using the bromocresol purple (BCP) method at 630 nm.

Serum lipid profile :Cholesterol and triglycerides were measured using specific methods: CHOD-PAP for cholesterol, and GPO-PAP method for triglycerides using Diagnostic kit.

Serum calcium and serum phosphorous:The concentrations of serum calcium and phosphorus were determined using the Arsenazo III method and Ammonium Molybdate method, respectively, at 630 nm and 340 nm wavelengths, respectively, as per the manufacturer's instructions, using a standard kit supplied by ERBA Mannheim Diagnostic Centre.

Albumin Globulin ratio (A:G) :Albumin globulin ratio was calculated by comparing their ratios

A:G = Concentration Of Albumin / Concentration Of Globulin

Serum LH: A CLIA (chemi luminescence immunoassay) test for LeutinisingHormone was used for estimation.

3.Results and discussion

Thecomposition of ingredientsof standard basal diets fed to white leghorn during feeding trial are shown in the Table 1 and proximate and phytochemical composition of Ovirich® is given in Table 2.

HaematologicalParameters of the white leghorn laying birds

The average values of haematological parameters presented in laying birds fed diet supplemented with Ovirich® in different treatment groups (T1, T2 and T3), at the end of supplementation period are presented in Table 3.

Table.1 Composition ofBasalDietfedtolayingbirdsduringthetrial period

S No.	Ingredients	% Composition
1.	Yellow Maize	57.05
2.	GNC-SolventExtracted	8.50
3.	SoyabeanMeal	16.50
4.	De-oiledRicebran	7.0
5.	Rice Polish	5.0
6.	MarblePowder	3.0
7.	TraceMineralMixture	0.10
8.	Dicalcium Phosphate	1.0
9.	DL-Methionine	0.35
10.	CholineChloride	0.05
11.	Hepatocare	0.10
12.	CommonSalt	0.40
13.	Vitamin Premix	0.10
14.	Toxin Binder	0.05
15.	Multi-Enzyme	0.05
16.	Coccidio-stat	0.05
17.	Lysine	0.70
		100.00

% CrudeProteinand Metabolizable energy in the basal diet was 18.13 and 2743.55 Kcal/kg respectively.

a) Haemoglobin (g/dl)

The average blood haemoglobin(g/dl) values showed significant differences ($P \leq 0.05$) among the control(T1) and supplemented (T2 and T3) groups. Numerically, T3 had a higher value compared to T1 and T2, but no significant difference ($P \geq 0.05$) was observed between T2 and T3 group.

b) Packed cell volume (PCV %)

The average packed cell volume values showed significant differences ($P \leq 0.05$) among the control (T1) and treatment(T2 and T3) groups. Specifically, T3 had a higher value compared to T1 and T2 group.

c) Total erythrocyte count ($10^6/\text{mm}^3$)

The average total erythrocyte count of T1, T2 and T3 treatment groups at the end of the supplementation period showed significant differences ($P \leq 0.05$) among the control(T1) and treatment(T2 and T3) groups. Specifically, T3 had a higher value compared to T1 and T2 group.

d) Total leucocyte count ($10^3/\text{mm}^3$)

The average total leucocyte count of T1, T2 and T3 treatment groups showed significant differences ($P \leq 0.05$) among the control(T1) and treatment(T2 and T3) groups. Numerically, T3 had a lower value compared to T1 and T2 group.

Table 2: Proximate and phytochemical composition of Ovirich® supplement on % dry matter basis.

S.No.	Nutrient Composition	Values
1	Dry Matter(%)	96.84
2	Crude Protein(%)	8.03
3	Ether Extract(%)	3.08
4	Crude fibre(%)	15.96
5	Total Ash(%)	37.04
6	Nitrogen free Extract(%)	35.98
7	Acid Insoluble Ash(%)	4.08
8	Total Phenol (mg GAE/g)	101.95

9	Total Flavonoid (mg QE/g)	45.80
10	Total Tannin (mg TAE/g)	80.90
11	Ascorbic Acid (mg/100gm)	115.94
12	Phytic Acid (g/100gm)	0.408
13	Methionine (g/100g)	3.624
14	Arginine (g/100g)	2.02
15	Tryptophan (g/100g)	3.22
16	Lysine (g/100g)	2.86

e) Mean corpuscular volume (fl)

The average mean corpuscular volume These values showed significant differences ($P \leq 0.05$) among the control(T1) and treatment(T2 and T3) groups.T3 had a higher value compared to T1 and T2 group.

f) Mean corpuscular haemoglobin (pg) and Mean corpuscular haemoglobin concentration (g/dl)

Significant differences ($P \leq 0.05$) were found in Mean corpuscular haemoglobin (pg) and Mean corpuscular haemoglobin concentration (g/dl) values among all treatment groups.

Increase in haemoglobin, Packed cell volume(PCV), Total erythrocyte count(TEC) and Mean corpuscular volume(MCV) values in Uttara chicken have been reported by Nayalet *al.* (2023), due to presence of different minerals like Calcium, Zinc, Aluminium, Copper, Magnesium. Haematinic property of Arjun bark and minerals have important role in the synthesis of haemoglobin. The bark contains compounds that promote the formation of red blood cells, leading to higher haemoglobin concentrations. Decrease in Mean corpuscular haemoglobin (MCH) and Mean corpuscular haemoglobin concentration (MCHC) values have also been reported by Nayalet *al.* (2023). Suelyet *al.* (2016)also reported that supplementation of Arjun increases MCV, decreases MCH and MCHC.

Contrary to our results increased total leucocyte count (TLC) values have been reported by Nayalet *al.* (2023).

Table 3. Average Haematological test results of the whiteleghorn laying birds at the end of supplementation period of Ovirich®

Parameter	T1	T2	T3	P-Value
Hb (g/dl)	8.23 ^b ±0.008	8.29 ^a ±0.011	8.32 ^a ±0.005	0.001
PCV (%)	29.38 ^c ±0.04	30.24 ^b ±0.04	30.96 ^a ±0.008	0.00
TEC (10 ⁶ /mm ³)	2.33 ^c ±0.005	2.36 ^b ±0.005	2.38 ^a ±0.005	0.003
TLC(10 ³ /mm ³)	4.13 ^a ±0.005	4.10 ^b ±0.005	4.08 ^c ±0.005	0.003
MCV (fl)	125.75 ^c ±0.005	126.95 ^b ±0.33	130.15 ^a ±0.005	0.00
MCH (pg)	35.22 ^a ±0.005	35.10 ^b ±0.011	35.0 ^c ±0.023	0.00
MCHC (g/dl)	28.02 ^a ±0.011	27.49 ^b ±0.023	26.88 ^c ±0.023	0.00

Mean values bearing different superscripts (a,b,c) within a row differ significantly from each other

Serological parameters of laying birds

The average values of serological parameters in White Leghorn layers fed with diet supplemented with Ovirich® in different treatment groups (T1, T2 and T3), are presented in Table 4.

a) SGOT (U/L)

Significant differences ($P \leq 0.05$) existed among the control(T1) and treatment(T2 and T3) groups. T3 had a lower value compared to T1 and T2 group.

Decrease in SGOT values have also been reported in Uttara chicken due to supplementation of Arjun bark (Nayalet *et al.*, 2023).

b) SGPT (U/L)

Significant differences ($P \leq 0.05$) were found among the control(T1) and treatment(T2 and T3) groups in average serum SGPT level.

Solanum nigrum extract significantly reduced the level of SGOT and SGPT, while also lowering the generation of superoxide and hydroxyl radicals (Raju *et al.*, 2003). Similar results have also been reported by (Malisornet *et al.*, 2020 and Nayal *et al.*, 2023).

c) Serum Protein (g/dl)

The average serum protein values differed significantly ($P \leq 0.05$) among the control(T1) and treatment(T2 and T3) groups.

Sharma *et al.* (2023)alsoreported that supplementation of Arjun bark powder and Sahjan leaf powder at different levels to basal diet increases total protein (g/dl) and albumin (g/dl).Inulin supplementation enhances globulin and total protein concentration (Mateovaet *et al.*, 2008).

d) Serum Albumin (g/dl)

Significant differences ($P \leq 0.05$) were noted among the control(T1) and treatment(T2 and T3) groups. Numerically, T3 had a higher value compared to T1 and T2 group.

Dietary supplementation of Arjun and Sahjan increased serum albumin values in Uttara chicken (Sharma *et al.*, 2023).

e) Serum Globulin (g/dl)

The average serum globulin values differed significantly ($P \leq 0.05$) among the control (T1) and treatment (T2 and T3) groups. Specifically, T3 had a higher value compared to T1 and T2 group.

Globulin fractions are increased due to increased protein synthesis in liver during the peak laying period (Swathi and Sudhamayee, 2005).

f) Serum Albumin Globulin Ratio (A:G)

Non-significant differences ($P \geq 0.05$) were noted among the control (T1) and treatment (T2 and T3) groups.

g) Serum Triglycerides (mg/dl)

Significant differences ($P \leq 0.05$) were noted among all treatment groups. Specifically, T3 had a lower value compared to T1 and T2 group.

The average serum triglycerides level of the different treatment groups T1, T2 and T3 at the end of the withdrawal period were recorded as 62.18 ± 0.08 , 58.92 ± 0.44 and 56.99 ± 0.21 , respectively. Significant differences ($P \leq 0.05$) among the control (T1) and treatment (T2 and T3) groups.

Decrease in serum triglycerides have been reported by (Malisornet *et al.*, 2020) on feeding mixed herbal extract in hybrid ducks. Nayalet *et al.* (2023) **also** reported decrease in serum triglycerides on feeding Arjun bark powder in Uttara layer birds.

Methanolic and phenolic extract of *Tamarix gallica* showed dose dependent decrease in the levels of cholesterol, triglyceride in hyperlipidemic rats. Phenolic extract 500 mg/kg body weight had antihyperlipidemic activity (Naveed *et al.*, 2015).

h) Serum Cholesterol (mg/dl)

Significant differences ($P \leq 0.05$) were noted among the control (T1) and treatment (T2 and T3) groups. Numerically, T3 had a lower value compared to T1 and T2 groups.

Decrease in serum cholesterol level have been reported by Nayalet *et al.* (2023). *C. spinosa* may be attributed to its ability to reduce the activity of 3-hydroxy-3-methyl-glutaryl coenzyme A reductase (HMG-CoA reductase), a key enzyme in cholesterol biosynthesis (Ness *et al.*, 2015).

i) Serum Lactate Dehydrogenase (LDH) (U/L)

No significant differences ($P \geq 0.05$) was noted among different treatment group both on supplementation of supplement.

j) Uric Acid (mg/dl)

The average serum uric acid level values showed significant differences ($P \leq 0.05$) among the control (T1) and treatment (T2 and T3) groups. Specifically, T3 had lower value compared to T1 and T2 group.

Sharma et al. (2023) reported similar results that supplementation of Arjun bark powder and Sahjan leaf powder at different levels to basal diet decreases serum uric acid levels in Uttara chicken.

k) Glucose (mg/dl)

Significant differences ($P \leq 0.05$) were noted among different treatment groups in glucose values.

Nayalet al. (2023) reported similar results which indicated that supplementation of Arjun bark powder at 1g, 2g, and 4g/100gm, respectively to basal diet decreases the serum glucose in Uttara birds. Arjun bark may be responsible for stimulating B cells and elevating insulin, which increases glucose utilization in tissues and lowers glucose levels. Ragavan and Kumari, (2006) reported that oral administration of *Terminalia arjuna* in rats for 30 days significantly reduce the glucose level.

C. spinosa reduces carbohydrate absorption rates and exerts a postprandial hypoglycemic effect in the gastrointestinal tract (Lemhadriet al., 2007).

l) Calcium (mg/dl)

The average serum calcium level values showed significant differences ($P \leq 0.05$) among the control (T1) and treatment (T2 and T3) groups. Specifically, T3 had a higher value compared to T1 and T2 group.

Frost and Roland (1991) reported that the level of plasma ionized Ca significantly increases in a linear manner by increasing dietary Ca levels from 2.75% to 4.25%, but not plasma total calcium.

m) Phosphorus (mg/dl)

The average serum phosphorus level values showed significant differences ($P \leq 0.05$) among the control (T1) and treatment (T2 and T3) groups. Specifically, T3 had a higher value compared to T1 and T2 group.

Miles et al. (1982) reported increased in plasma phosphorus as the dietary phosphorus increased in White Leghorn Layers.

Dietary Calcium has no effect on serum phosphorus level in Hy-Line Brown layers (Anet al., 2016).

Table 4 Average Serological test results of the laying birds at the end of supplementation period of Ovirich®

Parameters	T1	T2	T3	P-Value
SGOT (U/L)	182.64 ^a ±0.50	175.88 ^b ±1.16	171.40 ^c ±0.12	0.00
SGPT (U/L)	21.6 ^a ±0.19	18.67 ^b ±0.13	15.77 ^c ±0.04	0.00
Protein (g/dl)	4.88 ^c ±0.005	4.91 ^b ±0.008	4.95 ^a ±0.008	0.001
Albumin (g/dl)	2.68 ^c ±0.003	2.70 ^b ±0.004	2.72 ^a ±0.004	0.001
Globulin (g/dl)	2.196 ^c ±0.002	2.21 ^b ±0.003	2.23 ^a ±0.003	0.001
Albumin:Globulin	1.22±0.0005	1.221±0.0004	1.219±0.0005	0.125
Triglycerides (mg/dl)	63.03 ^a ±0.38	57.36 ^b ±0.18	55.44 ^c ±0.28	0.00
Cholesterol (mg/dl)	120.27 ^a ±0.03	118.36 ^b ±0.04	115.44 ^c ±0.11	0.00
LDH (U/L)	220.50±2.37	218.02±2.48	217.22±3.10	0.490
Uric acid (mg/dl)	8.62 ^a ±0.03	7.79 ^b ±0.24	7.91 ^c ±0.01	0.012
Glucose (mg/dl)	223.66 ^a ±1.20	218.33 ^b ±0.88	215.33 ^c ±0.88	0.003
Calcium (mg/dl)	12.48 ^c ±0.20	14.62 ^b ±0.07	15.62 ^a ±0.30	0.00
Phosphorus (mg/dl)	5.53 ^c ±0.01	5.64 ^b ±0.008	5.75 ^a ±0.01	0.00
LH (mIU/ml)	30.28±0.57	30.33±0.60	30.35±0.54	0.996

Mean values bearing different superscripts (a,b,c) within a row differ significantly from each other

n) Leutinizinghormone(LH)

The average LH level of the different treatment groups T1, T2 and T3 at the end of the supplementation period were non significantly different among all the treatment groups.

Due to heat stress no effect in plasma LH of White Leghoirn bird have been reported by (Rozenboimet *al.*, 2007).

4.Conclusion: It can be concluded that Ovirich can affect haemato-biochemical parameters in white leghorn layer bird.

References

- An, S. H.; Kim, D. W. and An, A. B.(2016).Effects of dietary calcium levels on productive performance, eggshell quality and overall calcium status in aged laying hens. *Asian-Australasian Journal of Animal Science*, 29(10): 1477.
- AOAC. (2003). Official methods of Analysis. *Association of Official Analytical Chemists*. 17thEdn. Washington, DC.

- BIS. (2007). Indian Standard: Poultry Feeds- Specification, IS-1374 *Bureau of Indian Standards*, 5th rev., Bahadur Sah Zafar Marg, Manaka Bhawan, New Delhi.
- Bray, H.G. and Thorpe, W.V.,(1954). Analysis of phenolic compounds of interest in metabolism. In: *Methods Biochem. Anal.*, Glick, D. (Ed.), Interscience Publishers, Inc., Hoboken, NJ: 27-52.
- Drabkin, D.L. and Austin, J.H.,(1932) Spectrometric studies: Spectrometric constant for common hemoglobin derivatives in human, dog, and rabbit blood. *Journal of Biological chemistry.*, 98(2): 719-733.
- Frost, T.J. and Roland, D.A., Sr.,(1991). The influence of various calcium and phosphorus levels on tibia strength and eggshell quality of pullets during peak production. *Poultry Science. Sci.*, 70: 963-969.
- Greenstein, J.P. and Winitz, M.,(1961). Chemistry of the amino acids. John Wiley & Sons, New York, 763p.
- Horn, J.M.; Jones, O.B. and Blum, A.E. (1946) Calorimetric determination of methionine in protein and foods. *Journal of Biological Chemistry*, 1 (16) : 313.
- Haug, W. and Latzsch, H.J.,(1983). Sensitive method for the rapid determination of phytate in cereal and cereal products. *J. Sci. Food Agric.*, 34(12): 1423-1426.
- Jagadish, L.K.; Krishnan, V.V.; Shenbhagaraman, R. and Kaviyarasan, V.,(2009). Comparative study on the antioxidant, anticancer, and antimicrobial property of *Agaricusbisporus* (J.E. Lange) Imbach before and after boiling. *Afr. J. Biotechnol.*, 8(4).
- Kavitha Chandran, C.I. and Indira, G.,(2016). Wound healing activity of ethanolic extract of roots of *Morindapubescens*. *J. Pharmacogn. Phytochem.*, 5(3): 43-46.
- Lemhadri, A.; Eddouks, M.;Sulpice, T. and Burcelin, R. (2007). Anti-hyperglycaemic and anti-obesity effects of *Capparis spinosa* and *Chamaemelumnobile* aqueous extracts in HFD mice. *Am. J. Pharmacol. Toxicol.*, 2: 106–110.
- Malisorn, M.; Akkaramadhurakul, P. O.; Songserm, T.;Wannachart, S. and Paraksa, N.,(2020). Effects of dietary mixed herbal extracted product supplementation on fatty liver hemorrhagic syndrome protection and productive performances of broilers. *Agric. Nat. Resour.*, 54(5): 485-490.
- Mátéová, S.; Šály, J.; Tučková, M.; Koscova, J.; Nemcova, Radomira;Gaálová, M. and Baranová, D. (2008.) Effect of probiotics, prebiotics and herb oil on performance and metabolic parameters of broiler chickens. *Med. Weter.*, 64: 294-297.

- Martha ,D.O.;Adetokundu,S.A.;Olabanji;O.S.;TakpejewhoE.G.andSunday,O.T(2012).The effect of supplementation of enzyme on performance and some blood chemistry parameters in broiler finisher chickens fed ginger byproduct meal(*Zingiberofficinale*).Int.J.Biol.Sci.,2(7):59-65.
- Miles, R.D.; Costa, P.T. and Harms, R.H.,(1983). The influence of dietary phosphorus level on laying hen performance, eggshell quality, and various blood parameters. *Poult. Sci.*, 62(6): 1033-1037.
- Natt,M.P and Harrick ,C A.1952 A new blood diluent for counting the erythrocytes and leukocytes of chicken.Poultry Science,31:735-738.
- Naveed, S.A.; Reddy, M.S.; Kumar, C.H.P.;Suhasini, B. and Dontamalla, S.K., (2015).Anti-hyperlipidemicactivity of *Tamarix gallica* extracts in Triton X-100 induced hyperlipidemic rats. *Int. J. Pharm.*, 6(4): 7880-7895.
- Nayal, K.; Kumar, A.; Gururaj, V.K. and Prasad, S.,(2023). Effect of Arjun (*Terminalia arjuna*) bark powder supplementation on haemato-biochemical parameters in Uttara layers. *Pharma Innov. J.*, 12(8): 906-911.
- Ness, G. C.(2015). Physiological feedback regulation of cholesterol biosynthesis: Role of translational control of hepatic HMG-CoA reductase and possible involvement of oxylanosterols. *Biochim. Biophys. Acta*, 1851(5): 667–673.
- Prieto, P.; Pineda, M. and Aguilar, M.,(1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal. Biochem.*, 269(2): 337-341.
- Raghavan, B. and Kumari, S.K.,(2006). Effect of *Terminalia arjuna* stem bark on antioxidant status in the liver and kidney of diabetic rats. *Indian J. Physiol. Pharmacol.*, 50: 133-142.
- Raju, K., Anbuganapathi, G., Gokulakrishnan, V., Rajkapoor, B., Jayakar, B. and Manian, S. (2003). Effect of dried fruits of *Solanum nigrum* L. against CCl4 induced hepatic damage in rats. *Biol. Pharm. Bull.*, 26(11): 1618-1619.
- Sadasivam, S. and A. Manickam, (1992). Biochemical Methods for Agricultural Science. Willey Eastern Ltd., New Delhi, p. 8-21.
- Rozenboim, I., Tako, E., Gal-Garber, O., Proudman, J.A. and Uni, Z.,(2007). The effect of heat stress on ovarian function of laying hens. *Poult. Sci.*, 86(8): 1760-1765.
- Sharma, A., Kumar, A., Sharma, R.K., Singh, B., Mondal, B.C., Prakash, O., Gangwar, S., Azad, A.K. and Nayal, K.,(2023). Effect of dietary supplementation of Arjuna (*Terminalia arjuna*) bark and Sahjan (*Moringa oleifera*) leaf powder on haematology and serum biochemicals of Uttara chicken. *Pharma Innov. J.*, 12(8):117-121.

Spies, J.R. and Chambers, D.C.,(1949). Chemical determination of tryptophan in proteins. *Anal. Chem.*, 21(10): 1249-1266.

Suely, A.; Zabed, H.; Ahmed, A.B., Mohamad, J., Nasiruddin, M., Sahu, J.N. and Ganesan, P. (2016). Toxicological and hematological effect of *Terminalia arjuna* bark extract on a freshwater catfish, *Heteropneustes fossilis*. *Fish PhysiolBiochem.* 42(2): 431-444.

Swathi, B. and Sudhamayee, K.G.,(2005). Studies on haematological parameters of ducks during prelaying and peak production periods. *Indian J. Poult. Sci.*, 40(2): 146-149.

UNDER PEER REVIEW