

# OVERWEIGHT AND OBESITY AMONG HEALTHCARE PROFESSIONALS IN TUNISIA: PREVALENCE AND ASSOCIATED FACTORS

## Abstract

### INTRODUCTION:

Obesity has become a major public health problem because of its potential impact on health and its alarming increase worldwide.

The objectives were to estimate the prevalence of overweight and obesity among healthcare professionals (HCPs) and to examine their associated factors.

### METHODS

We conducted a cross-sectional study during the period August 2022–November 2022 among HCPs in the two-university hospital Habib Bourguiba and Hedi Chaker of Sfax governorate, Southern Tunisia. To screen for EDs, we opted for the validated French version of the Eating Attitude Test (EAT-26).

### RESULTS

The median age was 30 years (Interquartile range (IQR)=[26–40]). In this study, 253 HCPs were aged 35 years or more (68.4%). Females represented 74.6% of participants (n=276). The mean Body Mass Index (BMI) was  $24.79 \pm 3.74$  kg/m<sup>2</sup>. The prevalence of overweight was 40.8%.

Among the participants, 24.6% were at high risk of developing eating disorders. In multivariate analysis, factors independently associated with the prevalence of overweight in HCP were a number of working years  $\geq 10$  years (adjusted odds ratio (AOR) = 5.7;  $p < 0.001$ ), large weight variation  $\geq 25$  Kg during adulthood (AOR=3.5 ;  $p = 0.038$ ), distortion in body image perception (AOR=2.2;  $p = 0.01$ ), dissatisfaction with the current weight (AOR=5.3;  $p < 0.001$ ) and high risk of developing eating disorders (ED) according to the EAT-26 score (AOR= 1.8;  $p = 0.04$ ).

### CONCLUSION

The alarming figures for overweight have been illustrated, justifying the introduction of preventive measures in the workplace in order to avoid the deleterious consequences.

*Keywords: [obesity; overweight; healthcare professionals; epidemiology; associated factors]*

## **INTRODUCTION**

Obesity is defined by the World Health Organisation (WHO) as an abnormal or excessive accumulation of body fat that can damage health. The notion of risk to an individual's health is part of the very definition of obesity (1). Overweight and obesity are one of the major lifestyle diseases that lead to other health problems and contribute to many chronic diseases, including cancers, diabetes, metabolic syndrome and cardiovascular disease (2). Obesity has become a major public health problem because of its potential impact on health and its alarming rise worldwide (3).

Tunisia does not seem to be spared from this affliction. Indeed, the prevalence of obesity in Tunisia increased between 2005 and 2016 among people aged 35-70, rising from 27.2% to 34.1% (4). The prevalence of overweight and obesity among healthcare professionals (HCPs) in Tunisia is still underestimated, and there is a lack of statistical data on the subject.

An accurate assessment of the prevalence of obesity among HCPs is necessary for several reasons. Firstly, obesity increases the likelihood of musculoskeletal disorders (5) and mental health problems (6), which are the main causes of work-related illness and accidents in the workplace for HCPs. As well as being implicated in the onset of chronic disease, these conditions and their associated absenteeism rates pose a potential problem for the efficiency and sustainability of the healthcare system by potentially reducing the performance of the healthy workforce (7).

Comparing the rate of obesity in HCPs with that in the general population will help to identify the possible contribution of unfavourable factors in the workplace, such as a lack of access to healthy food options (8,9), night work (10,11) and work with high demand but little control over working conditions. These working conditions can lead to increased stress and even burnout, which may increase the risk of overweight in this sub-group (12).

In a recent study (13) published in 2023, involving 709 HCPs in Saudi Arabia, 21% of participants were overweight and 15.5% were obese. These results are not negligible, and effective and early preventive measures should be established in the workplace to prevent overweight among HCPs and avoid its deleterious consequences.

In this perspective, this study aimed to estimate the prevalence of overweight and obesity among HCPs and to examine the factors associated with it.

## **METHODS**

### **1. Study design and settings**

This was a cross-sectional study, conducted over a 4-month period: from August 2022 to November 2022, among nursing staff at the two-university hospital in the governorate of Sfax, Habib Bourguiba and Hedi Chaker, southern Tunisia.

### **2. Sampling procedure**

A representative sample of healthcare professionals from both medical and paramedical cadres working in the two teaching hospitals was randomly selected. Calculation of the minimum sample size based on an expected overweight prevalence of 30.9% from a previous study conducted among HCPs in Italy (14), a confidence level of 95% and a 5%

margin of error, giving a sample size of 328 subjects. An additional 10% was required to allow adjustment for some missing data. Finally, from an exhaustive list of all eligible HCP, 370 participants were randomly recruited.

## 2. Study population and inclusion criteria

All healthcare professionals (doctors, interns, residents, seniors, nurses, anaesthetists, instrument technicians, physiotherapists, etc...) from the two university hospitals were eligible for the study. We did not include people who did not agree to complete the form because they refused or were mentally incapable of answering, or trainees. We excluded people who did not return the questionnaire and those who gave incomplete responses or missing data.

## 3. Data collection

The study was conducted as follows: data were collected by means of a self-report questionnaire, including questions on socio-demographic characteristics, personal and family history, EAT-26, exercise and sleep habits, exposure to recent stress, as well as position and number of shifts per week. An explanatory note for participants was appended to the questionnaire. We took the time to explain to the participants who agreed to take part in the survey the interest and purpose of this study and to ask them to complete a questionnaire. Each participant was then weighed and tapped by the investigating doctor to calculate their body mass index (BMI). This was defined as the ratio of weight (P) in kg to the square of height (T) in metres. To screen for eating disorders, we opted for the validated French version of the Eating Attitude Test (EAT-26) (15), which is the language of instruction of our population.

The EAT-26 questionnaire, a tool developed by Garner and Garfinkel in 1979, consists of 26 items and participants were asked to rate their responses on a 6-point Likert scale (always = 3, usually = 2, often = 1, sometimes rarely and never = 0 for items 1-25, and the reverse direction of scoring for item 26) (16,17).

The responses to the items were added together and a score was calculated with extremes from 0 to 78.

- A score of less than 20 indicated that the participant was at low risk of developing EDs.
- A score of 20 or more indicated that the participant was at high risk of developing EDs.

## 4. Statistical analysis

Data entry was performed using SPSS.20 software. For continuous variables, we calculated the means and standard deviations in case of Gaussian distribution. In the case of a non-Gaussian distribution, medians and interquartile intervals (IQR) were calculated. For the categorical variables, we calculated the frequencies and the percentages. The comparison of frequencies on independent series was done by the Pearson's chi-square test or Fisher's exact test if the conditions of validity were not met. The results of the risk associated with EDs were expressed by the Crude Odds Ratio (COR) with their 95% confidence intervals using the univariate logistic univariate logistic regression method. To compare the means for independent samples, we opted for the student's t-test for variables with a Gaussian distribution, and for non-Gaussian variables, we opted for non-parametric statistical tests. We opted for a multivariate analysis by binary logistic regression analysis (Adjusted odds

ratio (AOR), 95% CI) to identify independent factors of EDs. A p value <0.05 was statistically significant.

## RESULTS

The median age was 30 years (IQR= [26-40]). Two hundred and fifty-three HCPs were aged 35 years or more (68.4%) and 117 HCPs (31.6%) were aged less than 35 years. Our population was predominantly female (n=276; 74.6%). More than half of the HCPs who took part in our study belonged to the paramedical profession (n=206; 55.7%). The mean BMI of the HCPs was 24.79±3.74 kg/m<sup>2</sup>. Of the HCPs who took part in our study, 40.8% were overweight. Ninety-one participants (24.6%) were at high risk of developing AUD. The median value of the score calculated was 10 (IQR= [5-19]) (Table 1).

**Table 1: Socio-demographic and clinical characteristics of HCPs**

		NUMBER	%
<b>GENDER</b>	Male	94	25.4
	Female	276	74.6
<b>AGE GROUPS</b>	≥ 35 years	253	68.4
	< 35 years	117	31.6
<b>LIVING SITUATION</b>	With family	324	87.6
	Alone	28	7.6
	With friends	18	4.9
<b>MARITAL STATUS</b>	Married	194	52.4
	Engaged	33	8.9
	Single	134	36.2
	Divorced	9	2.4
<b>HAVING CHILDREN</b>		169	45.7
<b>PROFESSIONAL CATEGORY</b>	Medical	164	44.3
	Paramedical	206	55.7
<b>OCCUPATION</b>	interns	51	13.8
	Residents	79	21.4
	Seniors	28	7.6
	Nurses	90	24.3
	Anesthetists	14	3.8
	Instrumentalists	7	1.9
	other	101	27.3
<b>NUMBER OF YEARS OF WORK</b>	< 10 years	242	65.4
	≥ 10 years	128	34.6
<b>NIGHTSHIFTS</b>	>1 nightshift/week	89	24.1
	1 nightshift/week	281	75.9
<b>PHYSICAL ACTIVITY</b>	Light-intensity	232	62.7
	Moderate-intensity	55	14.9
	high-intensity	32	8.6
<b>SLEEP DURATION</b>	<5 hours	25	6.8
	5-8 hours	325	87.8
	>8 hours	20	5.4
<b>PRESENCE OF RECENT STRESSORS CURRENTLY</b>		191	51.6
<b>SMOKING</b>		42	11.4
<b>ALCOHOL CONSUMPTION</b>		14	3.8
<b>CHRONIC DISEASE</b>		63	17

<b>ALREADY CONSULTED A PSYCHIATRIST</b>		311	84.1
		59	15.9
<b>CURRENTLY TAKING A PSYCHIATRIC MEDICATION</b>		16	4.3
<b>FAMILY HISTORY OF OTHER MENTAL HEALTH DIAGNOSIS</b>		50	13.5
<b>FAMILY HISTORY OF EATING DISORDER</b>		46	12.4
<b>WEIGHT STATUS</b>	Underweight	8	2.2
	Normal	211	57
	Overweight	114	30.7
	Obese	37	10
<b>WEIGHT VARIATION DURING ADULTHOOD</b>	<25kg	347	93.8
	≥25kg	23	6.2
<b>PERCEPTION</b>	Normal	252	68.1
	Thin	28	7.6
	Obese	90	24.3
<b>SATISFACTION</b>	Satisfied	165	44.6
	Want to reduce weight	173	8.6
	Want to gain weight	32	46.8

N: Number; %: percentage

Univariate analysis showed that the prevalence of overweight and obesity in HCPs was significantly associated with the age group '< 35 years' (COR=2.6 ; p<0.001), marital status 'in couple' (COR=2.7 ; p<0.001), having children (COR=1.8 ; p=0.004), paramedical profession (COR=1.8 ; p=0.004) and number of years working ≥ 10 years (COR=2.8 ; p<0.001), the presence of warning signs of EDs in participants (COR=1.6; p=0.046), the presence of a chronic illness (COR=1.7 ; p=0.047), wide variation in weight ≥25Kg during adulthood (COR=5.7; p<0.0001), abnormal perception of body image (COR=4.7; p<0.001) and dissatisfaction with current weight (COR=5.2; p<0.001). Also, the prevalence of overweight was significantly higher in POC at risk of developing EDs according to the EAT-26 score (35.5% vs 58.2%; COR=2.5; p<0.001) (**table 2**).

**Table 2: Association of socio-demographic and clinical characteristics with overweight in HCPs**

Variable		Not Overweight n(%) N=218	Overweight n(%) N=152	COR (CI 95%)	p
<b>Gender</b>	Male	58 (61,7)	36 (38,3)	1	0,5
	Female	160 (58)	116 (42)	1,8 (0,7-4,6)	
<b>Age group</b>	≥ 35 years	168 (66,4)	85 (33,6)	1	<0,001
	< 35 years	50 (42,7)	67 (57,3)	2,6 (1,7-4,1)	

<b>Marital status</b>	No	103 (72)	40 (28)	1	<b>&lt;0,001</b>
	Engaged	115 (50,7)	112 (49,3)	2,7 (1,1-5,6)	
<b>Having Children</b>	No	132 (65,7)	69 (34,3)	1	<b>0,004</b>
	Yes	86 (50,9)	83 (49,1)	1,8 (1,2-2,8)	
<b>Living situation</b>	No	203 (59,4)	139 (40,6)	1	0,5
	Alone	15 (53,6)	13 (46,4)	1,2 (0,6-2,7)	
<b>Origin</b>	No	178 (59,7)	120 (40,3)	1	0,5
	yes	40 (55,6)	32 (44,4)	1,2 (0,7-1,9)	
<b>Position</b>	Médical	110 (67,1)	54 (32,9)	1	<b>0,004</b>
	Paramédical	108 (52,4)	98 (47,6)	1,8 (1,2-2,8)	
<b>Time of work</b>	Dayshift	120 (62,5)	72 (37,5)	1	0,1
	Day and nightshift	98 (55,1)	80 (44,9)	1,4 (0,9-2,1)	
<b>Nightshift</b>	≤ 1 nightshift/week	58 (65,2)	31 (34,8)	1	0,1
	>2 nightshift/week	160 (56,9)	121 (43,1)	1,7 (0,7-2,3)	
<b>Number of years of work</b>	< 10 years	167 (69)	75 (31)	1	<b>&lt;0,001</b>
	≥ 10 years	51 (39,8)	77 (60,2)	2,8 (1,4-5,5)	
<b>Smoking</b>	No	25 (59,5)	17 (40,5)	1	0,9
	Yes	193 (58,8)	135 (41,2)	1,1 (0,5-1,9)	
<b>Number of pack-years</b>	< 10 PA	206 (59,7)	139 (40,3)	1	0,2
	≥ 10 PA	12 (48)	13 (52)	1,6 (0,7-3,6)	
<b>Alcohol consumption</b>	Yes	9 (64,3)	5 (35,7)	1	0,6
	No	209 (58,7)	147 (41,3)	1,3 (0,4-3,8)	
<b>Already consulted a psychiatrist</b>	No	187 (60,1)	124 (39,9)	1	0,2
	Yes	31 (52,5)	28 (47,5)	1,4 (0,8-2,4)	
<b>Currently taking a psychiatric medication</b>	No	210 (59,3)	144 (40,7)	1	0,4
	Yes	8 (50)	8 (50)	1,5 (0,5-3,9)	
<b>Chronic disease</b>	No	188 (61,2)	119 (38,8)	1	<b>0,047</b>
	Yes	30 (47,6)	33 (52,4)	1,7 (1,1-2,9)	
<b>Family history of other mental health diagnosis</b>	No	192 (60)	128 (40)	1	0,2
	Yes	26 (52)	24 (48)	1,4 (0,8-2,5)	
<b>Family history of</b>	No	194 (60,1)	129 (39,9)	1	0,1

<b>eating disorder</b>	Yes	23 (50)	23 (50)	1,5 (0,8-2,8)	
<b>Weight variation during adulthood</b>	<25Kg	213 (61,4)	134 (38,6)	1	<b>&lt;0,001</b>
	≥25Kg	5 (21,7)	18 (78,3)	5,7 (2,1-15,8)	
<b>Perception</b>	Normal	178 (70,6)	74 (29,4)	1	<b>&lt;0,001</b>
	Thin or obese	40 (33,9)	78 (66,1)	4,7 (2,9-7,5)	
<b>Satisfaction</b>	Satisfied	131 (79,4)	34(20,6)	1	<b>&lt;0,001</b>
	No	87 (42,4)	118 (57,6)	5,2 (3,2-8,3)	
<b>Physical activity</b>	Yes	147 (59,5)	100 (40,5)	1	0,7
	No	71 (57,7)	52 (42,3)	1,1 (0,7-1,7)	
<b>Sleep duration</b>	Sufficient	192 (59,1)	133 (40,9)	1	0,8
	Insufficient or excessive	26 (57,8)	19 (42,2)	1,1 (0,6-1,9)	
<b>Presence of recent stressors Currently</b>	Yes	117 (61,3)	74 (38,7)	1	0,3
	No	101 (56,4)	78 (43,6)	1,2 (0,8-1,8)	

N: Number ; n : Number ; COR: Crude Odds Ratio; 95% CI: 95% confidence interval;

UNDER PEER REVIEW

In multivariate analysis, the factors independently associated with the prevalence of overweight in the HCPs who participated in our study were a number of working years  $\geq 10$  years (AOR=5.7;  $p<0.001$ ), large weight variation  $\geq 25$ Kg during adulthood (AOR=3.5 ;  $p=0.038$ ), abnormal perception of body image (AOR=2.2;  $p=0.01$ ), dissatisfaction with current weight (AOR=5.3;  $p<0.001$ ) and high risk of developing EDs according to the EAT-26 score (AOR= 1.8;  $p=0.041$ ) (table 3).

**Table 3: Multivariate analysis: factors associated with overweight among healthcare professionals**

Variable		AOR	CI 95%	p
Number of years of work	< 10 years	1		<0,001
	$\geq 10$ years	5,7	3,2-10,1	
Weight variation during adulthood	<25Kg	1		0,038
	$\geq 25$ Kg	3,5	1,1-11,4	
Perception	Normal	1		0,01
	Thin or obese	2,2	1,2-3,9	
Satisfaction	Satisfied	1		<0,001
	No	5,3	2,8-9,9	
High risk of developing EDs	No	1		0,041
	Yes	1,8	1,1-3,3	

N: Number ; n : Number ; AOR: Adjusted Odds Ratio; 95% CI: 95% confidence interval;

## DISCUSSION

In our study, the mean BMI of HCPs was  $24.79 \pm 3.74$  kg/m<sup>2</sup>. Of the HCPs in our study, 115 (30.8%) were overweight and 37 (10%) were obese. A recent study carried out in Spain (18) in 2022 also revealed overweight and obesity rates of 33.7% and 7.3%, respectively. Although overweight and obesity rates differ considerably from region to region and country to country, these significantly high rates are in line with those observed in similar studies (6,9). Rates of overweight and obesity were particularly high in the United States (19) and Brazil (20).

Differences in obesity prevalence between countries may result from a complex combination of factors, including eating habits, levels of physical activity, public health policies and socio-economic disparities.

In our study, a quarter of the participants (24.6%) were at high risk of developing EDs. These results are similar to those found in the Tunisian (21), Lebanese (22) and Bangladeshi (23) studies. However, higher figures were found in Morocco (24) and Egypt (25). We suggest



that the variations in prevalence are due to differences in the means of study used to screen for eating disorders, which may lead to heterogeneity of results, to the variety of the population studied and to the cultural context, which may have a direct impact on the risk of eating disorders in the population.

We found a significant association between the number of working years and the risk of developing overweight in HCPs. This result was consistent with that of Samhat et al (26). In that study, 56% of nurses reported having gained weight since starting work. After taking into account the age and sex of the population studied, they found that BMI and waist circumference increased with the cumulative number of years worked throughout the professional history. However, it is important to note that this result may be modulated by the influence of age and menopause, which have been identified in previous studies (3,27) as having a significant effect on the prevalence of overweight.

We found that wide weight variation during adulthood was an independent factor associated with the risk of overweight. This result was well confirmed by the study by Marleen A et al (28). In this study, it was shown that variations induced by weight loss, including cellular stress, remodeling of the extracellular matrix, inflammatory responses, secretion of adipokines and lipolysis, appear to be associated with weight regain after successful weight loss. The increased risk of recurrence of overweight could therefore, at least in part, be explained by a combination of these factors (28). Another study in 2023 (29) suggests that adipose tissue immune cells; hormonal and neural factors influencing hunger, satiety and reward; resting energy expenditure and adaptive thermogenesis; and lipid metabolism (lipolysis and lipid oxidation) play an important role in weight regain.

We found that abnormal body image perception was an independent factor associated with overweight. This result is consistent with those of other authors (30,31), who have shown that increasing BMI is a key point influencing body image perception. Indeed, overweight and obesity are often associated with serious disturbances in the perception of body shape and weight, as well as a desire to become thinner and obsessive fears of becoming fat (32). A negative body image increases the risk of suffering from eating disorders and developing overweight (33).

In our study, dissatisfaction with current body weight was an independent factor associated with the risk of overweight and obesity in SP. This result is consistent with studies by Toselli et al (34). Indeed, the results of the multiple linear regression analysis of this study also showed that BMI was an independent factor associated with weight dissatisfaction for both sexes, and that its increase led to an increase in the level of body dissatisfaction. This was also well confirmed in the study by Ku et al (19), where it was shown that obese participants, compared with overweight participants, reported lower body satisfaction and a more negative body perception. For both men and women, body dissatisfaction increased as BMI increased with a desire for smaller body sizes and obese individuals expressed a stronger desire for change to achieve their ideal body.

We found that eating disorders were an independent factor associated with the risk of overweight and obesity in HCPs. Several studies (24,25) have examined the relationship between eating disorders and the risk of obesity in students and healthcare professionals. This may be explained by a greater preoccupation with food and a tendency to follow inappropriate diets in patients with AUDs (25). Indeed, it has been shown that the weight control strategies used by overweight people are often ineffective and actually lead to weight gain and eating disorders such as binge eating, further increasing the risk of obesity. This suggests that high BMI and disturbed eating attitudes may be mutually reinforcing (35).

To our knowledge, this study provides the first data on the prevalence of overweight among practitioners of different health professions in Tunisia, as well as the predisposing factors. However, our study has certain limitations, including the cross-sectional nature of the study, which limits the interpretation of the causality of the risk factors studied, and the collection of data from a self-administered questionnaire; participants may intentionally under- or overestimate their attitudes and dietary behaviours in the questionnaire, which may introduce a bias in the interpretation of the responses.

## **CONCLUSION**

Our study has highlighted an underestimated problem. Overweight has complex and interdependent etiologies and risk factors, and is associated with a high mortality rate. These findings are not negligible, and effective, early preventive measures need to be put in place in the workplace to prevent overweight and obesity and avoid their deleterious consequences. It is therefore recommended that medical and paramedical staff be made aware of the importance of screening for overweight, particularly in vulnerable patients, and referring them to a qualified specialist for further assessment.

## **CONSENT (WHERE EVER APPLICABLE)**

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

## **ETHICAL APPROVAL (WHERE EVER APPLICABLE)**

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

## **REFERENCES**

1. Organisation mondiale de santé. Obésité et surpoids, 2020 [En ligne]. OMS [cité le 07/09/2022] ; [environ 12 écrans]. Disponible à l'URL : <https://www.who.int/fr/news-room/fact-sheets/detail/obesity-and-overweight>.
2. Safaei M, Sundararajan EA, Driss M, Boulila W, Shapi'i A. A systematic literature review on obesity: Understanding the causes & consequences of obesity and reviewing various machine learning approaches used to predict obesity. *Comput Biol Med.* 2021 Sep;136:104754.
3. Opoku AA, Abushama M, Konje JC. Obesity and menopause. *Best Pract Res Clin Obstet Gynaecol.* 2023 Jun;88:102348.
4. Tunisienne R. Indicateurs clefs de la santé des Tunisiens: Résultats de l'enquête "Tunisian Health Examination Survey-2016". Organisation Mondiale de la Santé. 2019.
5. Anandacoomarasamy A, Caterson I, Sambrook P, Fransen M, March L. The impact of obesity on the musculoskeletal system. *Int J Obes (Lond).* 2008 Feb;32(2):211-22.
6. Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BW, Zitman FG. Overweight,

- obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch Gen Psychiatry*. 2010 Mar;67(3):220-9.
7. Kyle RG, Wills J, Mahoney C, Hoyle L, Kelly M, Atherton IM. Obesity prevalence among healthcare professionals in England: a cross-sectional study using the Health Survey for England. *BMJ Open*. 2017 Dec;7(12):e018498.
  8. Nicholls R, Perry L, Duffield C, Gallagher R, Pierce H. Barriers and facilitators to healthy eating for nurses in the workplace: an integrative review. *J Adv Nurs*. 2017 May;73(5):1051-1065.
  9. Cheong ZY, Lopez V, Tam WSW. Barriers to healthy eating among nurses working in hospitals: A meta-synthesis. *J Adv Nurs*. 2022 Feb;78(2):314-331.
  10. Saulle R, Bernardi M, Chiarini M, Backhaus I, La Torre G. Shift work, overweight and obesity in health professionals: a systematic review and meta-analysis. *Clin Ter*. 2018 Jul-Aug;169(4):e189-e197.
  11. Tavares Amaro MG, Conde de Almeida RA, Marques Donalsonso B, Mazzo A, Negrato CA. Prevalence of overweight and obesity among health professionals with shift work schedules: A scoping review. *Chronobiol Int*. 2023 Mar;40(3):343-352.
  12. Fujishiro K, Lawson CC, Hibert EL, Chavarro JE, Rich-Edwards JW. Job strain and changes in the body mass index among working women: a prospective study. *Int J Obes (Lond)*. 2015 Sep;39(9):1395-400.
  13. Elsafi SH, Al-Dossari RH, Al-Shaqi RA, Fakirah WE, Al-Dossari RF, Al-Sharif OJ, Maawadh RM, Al Musallam LD, Alaohali A, Abu Hassan AM, Alfahad OA, Al Naam YA, Al Zahrani EM. Obesity-Related Knowledge and Practice Among the Healthcare Professions Students in Saudi Arabia. *Diabetes Metab Syndr Obes*. 2024 Jan ;17:427-434.
  14. Latina R, Petruzzo A, Vignally P, Cattaruzza MS, Vetri Buratti C, Mitello L, Giannarelli D, D'Angelo D. The prevalence of musculoskeletal disorders and low back pain among Italian nurses: An observational study. *Acta Biomed*. 2020 Nov; 30:91.
  15. Leichner P, Steiger H, Puentes-Neuman G, Perreault M, Gottheil N. Validation d'une échelle d'attitudes alimentaires auprès d'une population québécoise francophone [Validation of an eating attitude scale in a French-speaking Quebec population]. *Can J Psychiatry*. 1994 Feb;39(1):49-54.
  16. Garner DM, Olmsted MP, Bohr Y, Garfinkel PE. The eating attitudes test: psychometric features and clinical correlates. *Psychol Med*. 1982;12(4):871-8.
  17. Al-Jumayan AA, Al-Eid NA, AlShamlan NA, AlOmar RS. Prevalence and associated factors of eating disorders in patrons of sport centers in Saudi Arabia. *J Family Community Med*. 2021;28(2):94-102.
  18. Forcada-Parrilla I, Reig-Garcia G, Serra L, Juvinyà-Canal D. The Influence of Doing Shift Work on the Lifestyle Habits of Primary Care Nurses. *Nurs Rep*. 2022 Apr;12(2):291-303.
  19. Ku B, Phillips KE, Fitzpatrick JJ. The relationship of body mass index (BMI) to job performance, absenteeism and risk of eating disorder among hospital-based nurses. *Appl Nurs Res*. 2019 Oct ;49:77-9.

20. Guedes, F. R., Navarro, F. M., Nakao, R. Y., Franco, I. P., & Rodrigues, L. C. L. The prevalence of low back pain in nurses at a university hospital in the eastern area of Sao Paulo. *Coluna/Columna*. 2022 Jul; 21(21): e262474.
21. Jaweher M, Sonda T, Uta O, Inès F, Rim S, Imene B, et al. Troubles des conduites alimentaires et tempérament cyclothymique : étude transversale à propos de 107 étudiants Tunisiens. *Pan Afr Med J* . 2014 Jun;18:117.
22. Hoteit M, Mohsen H, Bookari K, Moussa G, Jurdi N, Yazbeck N. Prevalence, correlates, and gender disparities related to eating disordered behaviors among health science students and healthcare practitioners in Lebanon: Findings of a national cross sectional study. *Front Nutr*. 2022 Jul ;9:956310.
23. Banna MHA, Dewan MF, Tariq MR, Sayeed A, Kundu S, Disu TR, Akter S, Sahrin S, Khan MSI. Prevalence and determinants of eating disorder risk among Bangladeshi public university students: A cross-sectional study. *Health Psychol Res*. 2021 Jun ;9(1):24837.
24. Azzouzi N, Ahid S, Bragazzi NL, Berhili N, Aarab C, Aalouane R, Boujraf S, Rammouz I. Eating disorders among Moroccan medical students: cognition and behavior. *Psychol Res Behav Manag*. 2019 Mar;12:129-135.
25. Ali E, Shehata WM. Eating Disorder Risk among Medical Students at Tanta University, Egypt. *Egypt J Community Med*. 2020 Oct ;38(4):17–23.
26. Samhat Z, Attieh R, Sacre Y. Relationship between night shift work, eating habits and BMI among nurses in Lebanon. *BMC Nurs*. 2020 Apr;19(1):1–6.
27. Bogossian FE, Hepworth J, Leong GM, Flaws DF, Gibbons KS, Benefer CA, et al. A cross-sectional analysis of patterns of obesity in a cohort of working nurses and midwives in Australia, New Zealand, and the United Kingdom. *Int J Nurs Stud*. 2012 Jun;49(6):727–38.
28. van Baak MA, Mariman ECM. Mechanisms of weight regain after weight loss - the role of adipose tissue. *Nat Rev Endocrinol*. 2019 May;15(5):274-287.
29. van Baak MA, Mariman ECM. Obesity-induced and weight-loss-induced physiological factors affecting weight regain. *Nat Rev Endocrinol*. 2023 Nov;19(11):655–70.
30. Valerio G, Balsamo A, Baroni MG, Brufani C, Forziato C, Grugni G, Licenziati MR, Maffei C, Miraglia Del Giudice E, Morandi A, Pacifico L, Sartorio A, Manco M; on the behalf of the Childhood Obesity Group of the Italian Society of Pediatric Endocrinology and Diabetology. Childhood obesity classification systems and cardiometabolic risk factors: a comparison of the Italian, World Health Organization and International Obesity Task Force references. *Ital J Pediatr*. 2017 Feb;43(1):19.
31. Stagi S, Ibáñez-Zamacona ME, Jelenkovic A, Marini E, Rebato E. Association between self-perceived body image and body composition between the sexes and different age classes. *Nutrition*. 2021 Feb;82:111030.
32. McAdams CJ, Smith W. Neural correlates of eating disorders: translational potential. *Neurosci Neuroecon*. 2015 Sep;4:35-49.
33. Rounsefell K, Gibson S, McLean S, Blair M, Molenaar A, Brennan L, et al. Social media, body image and food choices in healthy young adults: A mixed methods systematic review. *Nutr Diet*.

2020 Feb ;77(1):19–40.

34. Toselli AL, Villani S, Ferro AM, Verri A, Cucurullo L, Marinoni A. Eating disorders and their correlates in high school adolescents of Northern Italy. *Epidemiol Psichiatr Soc.* 2005 Apr-Jun;14(2):91-9.
35. Chaudhari, B., Tewari, A., Vanka, J., Kumar, S., & Saldanha, D. The Relationship of Eating Disorders Risk with Body Mass Index, Body Image and Self-Esteem among Medical Students. *Annals of Medical and Health Sciences Research.* 2017; 7:144-149.

UNDER PEER REVIEW