

# ANTHROPOMETRIC STUDY OF ANGLE OF HUMERAL TORSION OF SOUTH-SOUTH, SOUTH-EAST OF NIGERIA

## ABSTRACT

**Aims:** Torsion is defined as a state of longitudinal twisting or spiraling of shaft of a long bone and can be measured as the difference between joint axis of proximal and distal ends of the bones. Angle of humeral torsion entails the reorientation of head of humerus bone relative to its shaft, the head being directed medially rather than caudally. Torsion occurs between proximal and distal extremities of the humerus, with humeral head facing postero-medially in anatomical position. The anthropometric study of angle of humeral torsion aims to understand the variation in the angle of humeral torsion among individuals and populations; however, this study focused on South-South and South East, Nigeria. The results concluded that there was no significant difference in mean values of angle of humeral torsion on right and left side bones. **Study design:** Descriptive and cross-sectional study. **Place and Duration of Study:** University of Port Harcourt, South-South, Nigeria.

**Methodology:** Two hundred humeri belonging to skeletons of unknown age, sex, and stature were collected from Medical Colleges of various Nigeria universities in South-South and South East, Nigeria. The torsion angles of these bones were then measured with the goniometer. Out of 200 humeri, 112 were of right side and 88 were of left side. **Results:** The result of the study showed the right torsion ( $\phi$ )  $73.87 \pm 5.67$  and left torsion ( $\phi$ )  $73.78 \pm 5.48$ .

**Conclusion:** There was no significant difference in mean values of angle of humeral torsion on right and left side bones.

**Keywords:** [Anthropometric study, Angle, humeral torsion, South-South and South East, Nigeria.]

**(Note:** 1. Case Reports should follow the structure of Abstract, Introduction, Presentation of Case, Discussion, Conclusion, Acknowledgements, Competing Interests, Authors' Contributions, Consent (where applicable), Ethical approval (where applicable), and References plus figures and/or tables. Abstract (not more than 250 words) of the Case reports should have the following sections: Aims, Presentation of Case, Discussion and Conclusion. Only Case Reports have word limits: Papers should not exceed 2000 words, 20 references or 5 figures. Other Type of papers have no word limits.

2. Review papers may have different headings of the sections and are exempted from following these suggestions.

3. Research Papers and Short Notes should follow the structure of Abstract, Introduction, Methodology, Results and Discussion, Conclusion, Acknowledgements, Competing Interests, Authors' Contributions, Consent (where applicable), Ethical approval (where applicable), and References plus figures and/or tables.)

## 1. INTRODUCTION

The anthropometric study of the angle of humeral torsion is a branch of research that focuses on the measurement and analysis of the rotational orientation of the humerus bone in the human body. This angle, also known as the humeral torsion angle, plays a crucial role in the biomechanics of the shoulder joint and affects the range of motion and functionality of the upper limb (Krishnan et al, 2019).

The angle of humeral torsion is formed by a line drawn through the center of longitudinal axis of the neck and head of humerus meeting a line drawn along the transverse axis of the condyle, when the base is viewed from above, looking straight down from above the head of the humerus; the normal angle of humeral torsion is between 20 and 40 degree (Stedman's medical dictionary. 2006). The average angle of the humeral torsion is 67.57° with a range of variation from 38° to 98°(Shahet al. 2006).

Anthropometric study aims to understand the variation in the angle of humeral torsion among individuals and populations, as well as its implications musculoskeletal health and performance. Through precise measurements and statistical analyses, anthropometric studies provide valuable insights into the anatomical variability and functional significance of the angle of humeral torsion, thus contributing to advancements in orthopedics, sports medicine and rehabilitation practices (Fadhil & Mubarak, 2020).

Thus, the aim of this study is to ascertain and document scientifically the humeral torsion angle of Nigerians in the South South and South Eastern region. This study also aims to observed the prevalence of humeral torsion angle in the South South and South Eastern region of Nigeria and to compare their humeral torsion angle to other races.

UNDER PEER REVIEW

## 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The following materials were used as the cause of this project:

### 2.1 Materials:

Humerus Bone, Plasticin, Cardboard Paper, Goniometer, Osteometric Board, Meter Ruler, Compass and Protractor, Calculator, Pencil, Pin, Pen, Exercise Book, SPSS 17 software.



Fig 1: Humeri Bone



Fig 2: Goniometer

### 2.2 Study Areas

Two hundred humeri belonging to skeletons of unknown age, sex, and stature were collected from the medical college of the following universities: University of Port Harcourt, Rivers State, University of Nigeria, Enugu State, University of Calabar, Cross River State, University of Uyo, Akwa Ibom State, University of Benin, Edo State, Delta State University, Abraka, Ambros Ali University, Ekpoma. Edo State, Nnamdi Azikiwe University, Awka. Anambra State, Niger Delta University, Amasoma. Bayelsa State.

## 2.3 Methodology

For this study two hundred humeri belonging to skeletons of unknown age, sex, and stature were collected from Medical Colleges of various Nigeria universities (already mentioned in Study Area above). Out of 200 humeri, 112 were of right side and 88 were of left side. The mechanical axis of the head was taken as the line joining the two points namely center of the articular surface of head where transverse diameter of articular surface is maximum and at the greater tuberosity approximately between the insertions of supraspinatus and of the infraspinatus muscles. A pin was stuck with plasticin along the line of these two points. For the distal end axis, a point was marked on the anterior surface of capitulum along the center of its vertical diameter. Similarly, a second point was marked on the trochlea. A pin was stuck with plasticin along the line of these two points. The angle formed by the crossing of these two reference lines was then measured with the goniometer; this angle is called the torsion angle.

## 2.4 Precautions

- i. A pin was fixed along the reference line employed for the upper and lower ends of humerus.
- ii. A line was drawn along the fixed reference line for the upper and lower ends of humerus with a sharp pencil and meter ruler.
- iii. Care was taken to ensure that the goniometer was properly placed to avoid error due to parallax


## 3. RESULTS AND DISCUSSION

### Data of Torsion Angle

Table 1: Descriptive Statistics

	N	Range	Minimum	Maximum	Mean		Std.	Varianc
	Statistic	Statistic	Statistic	Statistic	Statistic	Std. Error	Deviation	e
							Statistic	Statistic
Right humerus (φ)	111	34.00	50.00	84.00	73.8739	.53770	5.66507	32.093
Left humerus (φ)	87	28.00	55.00	83.00	73.7816	.58800	5.48449	30.080
Average (φ)					73.82			

Fig 1: Chart Showing the Descriptive Statistics



**Table 2: Z-Test for Hypothesis**

	Mean	variance	STD Deviation	Z-critical	Z-Value	Level of significance
Right torsion ( $\phi$ )	73.87	32.093	5.665	1.96	0.10	Not significant
Left torsion ( $\phi$ )	73.78	30.080	5.484			

Table 2 shows sample test to assess the difference in the means, variance and standard deviation between the right and left side

### 3. DISCUSSION

Torsion and rotation are two different phenomena. The angle made by the crossing of the axes of the two opposite ends, when measured obtusely includes 90-degree rotation, which the entire upper limb undergoes during its development in the embryo. In this study the mean angle reveals that there is variation on the angle of humeral torsion which is 73.83 degree. The range of variation in this study was 50 degrees to 84 degrees.

A comparison of torsion angle reported by different studies show that there is a considerable racial variation. Among all the investigators who have studied torsion in different races, highest average degree of torsion is found to occur in white races. It would be interesting to note that with lower degree of torsion the incidence of recurrent anterior dislocation may be less common in Nigeria but so far no clinical data is available to support this hypothesis.

Studies from Patil *et al* (2016) and Skaria & Kulkarni (2022) have found positive relationship between increased humeral torsion and recurrent anterior dislocation of shoulder (RADS), in which there was significant increase in the angle of humeral torsion in patients with RADS as compared to persons without any history of RADS or shoulder trauma. However, Larson (2007) reported no difference in torsion between normal shoulders and those with recurrent anterior dislocation of shoulder (RADS).

Furthermore, studies of individuals, whether participants in selected sports, show that humeral torsion, rather than being a static attribute that encodes only phylogenetic information, actually is developmentally dynamic in its morphological expression. Because the lateral rotators insert within the proximal epiphysis whereas most of the medial rotators act distally on the shaft, forces working in opposite directions during development normally add approximately 32° of secondary torsion. The extent of this secondary (developmental) torsion depends on a variety of factors.

#### 4. CONCLUSION

The result of the study showed the right torsion ( $\phi$ )  $73.87 \pm 5.67$  and left torsion ( $\phi$ )  $73.78 \pm 5.48$ . There was no significant difference in mean values of angle of humeral torsion on right and left side bones. From this study it has been established that torsion angle is very important for movement of upper limb, upright and posture therefore, care should be taken in other not to fracture proximal and distal part of humerus. Hence public enlightenment and awareness should be created to avoid deformity.

#### COMPETING INTERESTS

The authors declare that there is no competing interest in this study.

#### CONSENT (WHEREEVER APPLICABLE)

Not applicable

#### ETHICAL APPROVAL (WHEREEVER APPLICABLE)

Ethical clearance was obtained from the research ethics committee of the University of Port Harcourt before commencement of the study.

#### REFERENCES

- Fadhil, S. A. N., & Mubarak, H. J. (2020). A Morphometric Evaluation of the Humerus in relation to Shoulder Arthroplasty. *Research Journal of Pharmacy and Technology*, 13(10), 4817-4822.
- Krishnan, R., Björzell, N., Gutierrez-Farewik, E. M., & Smith, C. (2019). A survey of human shoulder functional kinematic representations. *Medical & biological engineering & computing*, 57, 339-367.
- Larson, S. G. (2007). The definition of humeral torsion: a comment on Rhodes (2006). *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 133(2), 819-820.
- Patil, S., Sethi, M., & Vasudeva, N. (2016). Determining angle of humeral torsion using image software technique. *Journal of Clinical and Diagnostic Research: JCDR*, 10(10), AC06.
- Shah R K, Trivdei B D, Patel J P, Shah G V, Nirvan A B;(2006): angle of humeral Torsion of India, *Journal of Anatomy Society of India*, vol 55 (2) 43-47.
- Skaria, S., & Kulkarni, M. (2022). Morphometric study of torsion of humerus and its applications in total shoulder arthroplasty. *International Journal of Health Sciences*, 6(S1), 8314-8322.
- Stedman's Medical Dictionary, 2006. Lippincott Williams and Wilkin, 28th edition, pg 90 and 92.