

Evaluation and production of Enriched (Bio-fertilizers) Vermi-compost on Bio-waste by earthworms (*Eisenia fetida* sp.)

ABSTRACT

This investigation was associated with different materials and techniques on bio-fertilizer application in soybean stover (dry matter) and fresh Cow dung for effect of bio fertilizer and earthworms (*Eisenia fetida* sp.) on bio waste decomposition. This study was analyzed by RBD (Randomized Block Design) with 6 treatments have to bio fertilizers combination *i.e.* (E₁-control, E₂- Rhizobium, E₃-Rhizobium+PSB, E₄-Rhizobium+KSB, E₅-Rhizobium+PSB+KSB, and E₆-Rhizobium+PSB+KSB+Trichoderma) and earthworms with 4 replications. The research was conducted at MRPC in Department of Soil Science and Agricultural Chemistry, JNKVV, Jabalpur during the Rabi season of 2021-22 and 2022-23. The result interpreted by partial decomposition, duration of completion, conversion rate and recovery percentage of vermicomposting. The result revealed that partial decomposition (kg) found in enriched vermicompost higher to lower sequence *i.e.* E₅ (7.11 kg) > E₃ (7.18 kg) > E₂ (7.19 kg) > E₄ (7.20 kg) = E₆ (7.20 kg) > E₁ (7.28 kg) from initial weight (10 kg). The production of enriched vermicompost maximum found in E₆ (5.58, 5.63 and 5.61 kg pot⁻¹) within respective duration (42, 40 and 41 days) of vermicomposting in sequent years and statically pooled analysis. Conversion rate and recovery percentage of enriched vermicompost were increase with the combination of bio-agents. This investigation useful to making vermicompost using of agricultural waste through bio-fertilizer.

Keyword: *Rhizobium, Trichoderma, PSB, KSB* and microorganisms *etc.*

1. INTRODUCTION

Waste management (solid, liquid and gaseous waste) is a major global challenge with increasing demand for protecting human health and the environment. In this research review, the classification and overview of research will help us identify the most important research areas for waste management. Furthermore, to promote the transition from a linear to a circular economy, waste management should be supported by policy-based initiatives and management policies (Maqsoodi *et al.*, 2023). The case of vermicomposting, or with mixed agricultural and eco-vermicomposting, the earthworms can assist in detoxifying commercial and ecological wastes and their very last product is useful for plant increase as natural fertilizers (Raza *et al.*, 2024). Millions of tons of agricultural and industrial waste are discarded every year at considerable financial and environmental cost. Instead of discarding the food scraps and waste, we can recycle these with the help of earth worms (Fayaz *et al.*, 2016). The Vermicompost is an eco-friendly, low cost, and effective way to recycle agricultural and kitchen waste. It is a mixture of earthworm castings, organic materials, humus, and other organisms. It has been advocated in integrated nutrient management systems in field crops due to its rich source of macro and micronutrients, vitamins, enzymes, antibiotics, and growth hormones. It also improves fertilizer and water use efficiency even better than FYM (Singh and Agarwal, 2005). Organics, inorganic and bio-fertilizers are essential to raise the crop yield. Vermicomposting of non-toxic biodegradable matter produces a stabilized humus like product known as vermicompost, which has a great potential as soil amendment. Vermicompost is a good soil conditioner that is rich in NPK, micronutrients, and growth hormones. Vermicompost application to soil also increases microbial populations and activities that further influence nutrient cycling, production of plant growth-regulating materials, and build up plant resistance to pathogen and nematode attacks (Gopal *et al.*, 2009). Composting waste and using it in agriculture is the most economical way to deal with it. Worldwide, there is a systematic improvement in the methods of treating waste and then recycling it for use in technological processes. This approach can be applied in a circular economy in which the value of raw materials and finished products can be preserved for as long as possible, minimizing the amount of waste (Szulc *et al.*, 2021). In agriculture, production is massive every year, but millions of tons of agricultural products are lost during the agricultural process (Serpil *et al.*, 2012). To achieve maximum and quality production, fertilizers are used. The necessary elements are absorbed by plants from fertilizers. This is called plant nutrition. In permanent agriculture, the nutrient content is reduced to compensate for this excessive use of artificial fertilizers [6]. With excessive use, soils are salinized, heavy metals come into contact with humans, and nitrates accumulate in water sources. They have harmful effects on the environment. Instead of artificial fertilizers, bio-fertilizers are used for mass and quality food production. In this case, agricultural waste produces bio-fertilizer and energy. Composting is a biological process in which microorganisms decompose organic matter and lower the carbon-nitrogen ratio of the substrate. It is generally prepared

57 from organic waste material such as crop residue, household waste etc. this research focus the using
58 of different bio agent on agricultural waste.

59 2. METHOD AND MATERIAL

60 The experiment was conducted in the Department of soil science and agriculture chemistry,
61 JNKVV, Jabalpur. The Enriched vermicompost made through biowaste of soybean stover (dry matter)
62 and fresh cow dung with earthworms (*Eisenia fetida* sp.) under 6 treatments of biofertilizer combination
63 with 4 replications and the data was statically analysis by RBD design (Gomez and Gomez, 1984).
64 Vermicompost made by continue two years (2022 or 2023) completely in *Rabi* season. This
65 experimental trial have using different materials and technology that is following heading listed below.

66 2.1 Collection of bio wastes, bio agents and earthworms

67 Soybean stover and cow dung were gathered from study campus Breeder Seed Research Unit
68 and Dairy Research Farm respectively, for use in the current study as vermicomposting substrates.
69 However, bio fertilizers and earthworms were obtained from the Microbes Research Production Centre
70 and Dairy Research Farm of the vermicompost unit at JNKVV, Jabalpur, Madhya Pradesh respectively.
71 The dry matter contents of soybean stover and fresh cow dung were determined using the oven dry
72 method at 105 °C and were 67.8, 22.3%, respectively.

73 2.2 Partial decomposition after adding of bio fertilizers and earthworms in pot⁻¹

74 The earthworms have a completely unique capacity to transform degradable bio wastes into
75 precious composts. However, those wastes need to be partly decomposed bio inoculate and launch of
76 earthworms throughout the start of decomposition technique can be survival and improvement of
77 earthworms. Before pre decomposition, dung was turned into delivered as additive in reputable
78 substrata in an identical proportion (1:1) which allowed to decompose for 28 days. After pre-
79 decomposition technique Rhizobium, PSB, KSB and Trichoderma have been injected discretely @ 10
80 g kg⁻¹ every pots. Desired moisture stage of 70-80% turned into maintained with inside the decomposing
81 bio-waste with normal watering on the price of 4 liter water for 8 kg of substrata at 7 days periods
82 throughout the partial decomposition (28 days) duration the partial decomposition was calculated
83 through following formula.

84
$$\text{Partial Decomposition (kg)} = \text{Fresh weight of substrata (at initial)} - \text{Oven dry weight of substrata (after 28 days)}$$

85 2.4 Duration of vermicomposting

86 The total number of days required to complete the composting process, including 28-days for
87 partial decomposition, were recorded as vermicompost duration for every treatment.

88 2.5 Conversion rate of vermicompost

89 The vermicomposting rate is the daily conversion of bio waste into decomposed organic matter
90 by bio inoculants and earthworms were recorded in the final stage of vermicomposting. In fact, the
91 amount of vermicompost produced during a treatment period is known as the conversion rate and is
92 determined for each treatment according to a given method.

93
$$\text{Conversion rate (g/day)} = \frac{\text{Quantity of vermicompost (weight)}}{\text{Number of days it took to convert the substrate into vermicompost}}$$

94 2.6 Recovery of vermicompost

95 Vermicompost recovery was calculated based on the final dry matter obtained from each
96 treatment from the total dry matter used for decomposition according to the following formula:

97
$$\text{Recovery of vermicompost (\%)} = \frac{\text{Dry weight of vermicompost}}{\text{Dry weight of substrata}} \times 100$$

98 3. RESULT AND DISCUSSION

99 3.1 Partial decomposition of bio-waste

100 The data table 1 revealed that the partial decomposition of bio-waste shows non-significantly
101 in the both year and statistical pooled data. However, The partial decomposition maximum observed
102 in treatment E₄ (Rhizo.+PSB+KSB) *i.e.* 7.11 kg found in both year and pooled data followed by E₃ -
103 Rhizobium + PSB (7.12, 7.24 and 7.18 kg) letter on E₄-Rhizobium + PSB (7.13, 7.27 and 7.20 kg) and
104 the minimum partial decomposition found in E₁ - control *i.e.* 7.25, 7.31 and 7.28 in the year 2021,2022
105 and pooled data respectively. Wiharyanto *et.al.*, 2018 reported that the partial decomposition in addition
106 of local microorganisms mixture of food waste can accelerate the process of compost maturity.

107 **Table 1. Partial Decomposition (kg) of bio-waste under different bio agent with earthworm after**
108 **28 days**

Treatments	2021	2022	Pooled
E ₁ (Control)	7.25	7.31	7.28
E ₂ (Rhizobium)	7.16	7.22	7.19
E ₃ (Rhizobium + PSB)	7.12	7.24	7.18
E ₄ (Rhizobium + KSB)	7.13	7.27	7.20

E₅ (Rhizo. + PSB + KSB)	7.11	7.11	7.11
E₆ (Rhizo.+PSB+KSB+Tricho)	7.18	7.21	7.20
SEm±	0.23	0.13	0.19
CD (P=0.05)	0.69	0.37	0.55


3.2 Production of vermicompost under different treatments with duration of completion of vermicomposting

Production of vermicompost and vermicompost completion under different treatments were presented in Table 2 and Figure 1. The results were found significantly on production and duration of completion of vermicompost. The production of vermicompost found in E₆ - Rhizo. + PSB+ KSB+ Tricho (5.58, 5.63 and 5.61 kg plot⁻¹) within respective duration of vermicomposting (42, 40 and 41 days) followed by E₅ - Rhizo. + PSB + KSB (5.20, 5.25 and 5.23 kg plot⁻¹) within 52, 54 and 53 days after completion of decomposition of bio- waste significantly superior to E₂ (Rhizobium) *i.e.* 4.39, 4.45 and 4.42 kg plot⁻¹ within 66, 63 and 64.50 days and E₁ (Control) *i.e.* 4.23, 4.25 and 4.24 kg plot⁻¹ within 79, 74 and 76.50 days of bio-waste decomposition. While, the treatments E₃-Rhizobium + PSB (4.90, 4.95 and 4.93 kg plot⁻¹) and E₄-Rhizobium + KSB (4.92, 4.87 and 4.90 kg plot⁻¹) within E₃ (55, 57 and 56 days) and E₃ (57, 55 and 56 days) respective duration of vermicomposting. Addition of the bio-fertilizers altered the soil physicochemical properties due to the microbial activity from the bio-fertilizers. Zea Mays showed an enhanced growth and reproduction rate upon application of the bio-fertilizers. Vermicomposting can be used as a waste corn pulp management strategy and at the same time obtain bio-fertilizers reported by Manyuchi, 2013. Under paddy straw based vermicomposting reported by Vijaya *et al.*, 2008 and similarly observation obtained in different agricultural crops reported by researcher that is Kalantari *et al.*, 2010, Kumari *et al.*, 2011 and Ansari and Sukhraj, 2010.

Table 2. Production of Vermicompost (Kg pot⁻¹) under various treatments.

Treatments	Vermicompost Production (kg pot ⁻¹)			Duration (days) of completion vermicomposting		
	2021	2022	Pooled	2021	2022	Pooled
E₁ (Control)	4.23	4.25	4.24	79.00	74.00	76.50
E₂ (Rhizobium)	4.39	4.45	4.42	66.00	63.00	64.50
E₃ (Rhizobium + PSB)	4.90	4.95	4.93	55.00	57.00	56.00
E₄ (Rhizobium + KSB)	4.92	4.87	4.90	57.00	55.00	56.00
E₅ (Rhizo. + PSB + KSB)	5.20	5.25	5.23	52.00	54.00	53.00
E₆ (Rhizo.+PSB+KSB+Tricho)	5.58	5.63	5.61	42.00	40.00	41.00
SEm±	0.28	0.26	0.27	2.62	2.55	2.58
CD (P=0.05)	0.82	0.77	0.79	7.72	7.51	7.62

Figure 1. Production of Vermicompost (Kg pot⁻¹) and Duration (days) of completion vermicomposting under various treatments.



3.3 Vermicompost recovery percentage under various treatments

The data Table 3 and Figure 2 shows that vermicompost recovery percentage on E₆- Rhizo.+ PSB+ KSB+ Tricho (69.75, 70.38 and 70.06 %) and E₅ - Rhizo. + PSB + KSB (65.00, 65.63 and 65.31%) were significantly superior to E₂ –Rhizobium (54.88, 55.63 and 55.25 %) and E₁-Control (52.88, 53.13 and 53.00 %). While, the treatments E₃ -Rhizobium + PSB (61.25, 61.88 and 61.56 %) and E₄ - Rhizobium + KSB (61.50, 60.88 and 61.19 %) were found partly in the year of 2021 and 2022 as well as statically pooled analysis. The recovery percentage depend on dry vermicompost production *i.e.* increase with the increasing the different bio-fertilizer numbers.

Table 2. Vermicompost recovery percentage and Conversion rate of bio-waste (g day⁻¹) under various treatments.


Treatments	Vermicompost Recovery (%)			Conversion rate of waste (g day ⁻¹)		
	2021	2022	Pooled	2021	2022	Pooled
E ₁ (Control)	52.88	53.13	53.00	55.00	59.15	57.07
E ₂ (Rhizobium)	54.88	55.63	55.25	69.80	73.76	71.78
E ₃ (Rhizobium + PSB)	61.25	61.88	61.56	92.34	89.70	91.02
E ₄ (Rhizobium + KSB)	61.50	60.88	61.19	90.97	93.67	92.32
E ₅ (Rhizo. + PSB + KSB)	65.00	65.63	65.31	103.16	99.85	101.50
E ₆ (Rhizo.+PSB+KSB+Tricho)	69.75	70.38	70.06	134.90	142.88	138.89
SEm±	3.47	3.26	3.37	6.77	6.51	6.64
CD (P=0.05)	10.24	9.62	9.93	19.96	19.20	19.58

3.4 Conversion rate of bio-waste (g day⁻¹) under different treatments

The data have present in Table 3 and Figure 2 Bio-waste Conversion rate estimated that significantly in the both year and pooled analysis. The heights conversion rate computed in E₆- Rhizo.+PSB+KSB+Tricho (134.90, 142.88 and 138.89 g day⁻¹) *i.e.* was significantly superior to all bio agent based treatments. While, E₅ - Rhizo. + PSB + KSB (103.16, 99.85 and 101.50 g day⁻¹), E₄ - Rhizo.+ KSB (90.97, 93.67 and 92.32 g day⁻¹) E₃- Rhizo.+ PSB (92.34, 89.70 and 91.02 g day⁻¹) were found significant to E₂- Rhizo. (69.80, 73.76 and 71.78 g day⁻¹) and E₁- control (55.00, 59.15 and 57.07 g day⁻¹) in the year of 2021, 2022 and statically pooled analysis data. The conversation rate might be dependent on the bio-fertilizer effected to deteriorate the cellulose tissue in soybean straw so that earthworm easy to digest the soybean straw. Similar finding reported by Tsai *et al.*,2007, Yong *et al.*,2021, Dikko *et al.*,2019, Srivastava *et al.*,2024.

156
157

Figure 2. Vermicompost recovery percentage and Conversion rate of bio-waste (g day⁻¹) under various treatments.



158
159

ACKNOWLEDGEMENTS

I express my gratitude and honors to my chairmen and my advisory committee for his able guidance, constant inspiration, and valuable suggestion and critical during entire course of present studies and preparation manuscript. I extend my heartiest thanks to HOD sir Dr. P.S. Kulhare, Dr. B. K. Dikxit, Dr. H. K. Rai, Dr. Y. M. Sharma, Dr. S. S. Baghel, Dr. B. S. Dubedi, Dr. R. K. Sahu, Dr. A. K. Upadhyay, Dr. G. S. Tegore and all department members soil science and agricultural chemistry, CoA, JNKVV, Jabalpur. For provide valuable resources for vermicompost production, field works, research works and laboratory works. There is no funding agency for support this research.

AUTHORS' CONTRIBUTIONS

The K.K.P. (1st author) carried out the experimental during the research period, D.S.S.B. (2nd author) given guidance for above investigation, the D.S.B.A. (3rd author), D.H.K.R. (4th author), D.R.K.S. (5th author), and D.U.S. (6th author) supporting for laboratory guidance and valuable suggestion for the research after that A.K.S. (7th author) and P. (8th author) were supported in data collection, vermicompost testing analysis and lot of moral supported for research.

Conflicts of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

- Ansari, A. A., and Jaikishun, S., (2010) "An investigation into the vermicomposting of sugarcane bagasse and rice straw and its subsequent utilization in cultivation of *Phaseolus vulgaris* L. In Guyana", *American-Eurasian J. Agriculture and Environmental Science*, 8 (6), 666-671.
- Dikko, A., Mahmud, Mustapha, Y., Abubakar, D., and Yusuf, A., (2019) STUDIES ON CONVERSION OF SOLID WASTE TO BIOFERTILIZER BY VERMICOMPOSTING. *FUW Trends in Science & Technology Journal*, www.ftstjournal.com e-ISSN: 24085162; p-ISSN: 20485170; 4 (1): 137 – 139.
- Fayaz, A., Ahmad, L., and Singh, P. (2016). Vermi-composting: an effective technique to recycle wastes into valuable organic fertilizer: a review. *Journal of Pure and Applied Microbiology*, 10(2), 1109. <https://link.gale.com/apps/doc/A481650197/AONE?u=anon~5c8ff683&sid=googleScholar&xid=6b493c7a>
- Gomez, K.A., and Gomez, A.A., (1984) *Statistical Procedures for Agricultural Research*. 2nd Edition, John Wiley and Sons, New York, 680 p.
- Gopal, M., Gupta, A., Sunil, E., Thomas, V.G., (2009) Amplification of plant beneficial microbial communities during conversion of coconut leaf substrate to vermicompost by *Eudrilus* sp. *Curr Microbiol.* 59:15–20. doi: 10.1007/s00284-009-9388-9.

193

194 Kalantari, S., Hatami, S., Ardalan, M. M., Alikhani, H. A., and Shorafa, M., (2010) "The effect of compost
 195 and vermicompost of yard leaf manure on growth of corn", *African Journal of Agricultural*
 196 *Research*, 5 (11), 1317-1323.

197 Kumari, M., Kumar, S., Chauhan, R.S., and Ravikanth, K., (2011) "Bioconversion of herbal industry
 198 waste into vermicompost using an epigeic earthworm *Eudrilus Eugeniae*", *Waste Management*
 199 *and Research*, 29 (11), 1205-1212.

200 Maghsoudi, M.S., Shokouhyar, S., Khanizadeh, S., and Shokoohyar, (2023). Towards a taxonomy of
 201 waste management research: An application of community detection in keyword network J.
 202 *Clean. Prod.*, 401 Article 136587.

203 Manyuchi, M., (2013). Production of Bio-Fertilizers from Vermicomposting of Waste Corn Pulp Blended
 204 with Cow Dung as a Solid Waste Management Approach.

205 Raza, S.T., Zhu, B., Yao, Z.Y., Wu, J.P., Chen, Z., Ali, Z., and Tang, J.L., (2023) Impacts of
 206 vermicompost application on crop yield, ammonia volatilization and greenhouse gases
 207 emission on upland in Southwest China *Sci. Total. Environ.*, 860 Article 160479.

208 Serpil S., and Scribe B., (2012) *International Journal of Environmental Science and Development*, 3 (1)
 209 An Agricultural Pollutant: Chemical.

210 Singh, R., and Agarwal, S.K., (2005). Effect of levels of farmyard manure and nitrogen fertilization on
 211 grain yield and use efficiency of nutrients in wheat (*Triticum aestivum*). *Indian Journal of*
 212 *Agricultural Sciences*. 75. 408-413.

213 Srivastava, S., Singh, P., Barbora, L., Baruah, D., Saikia, R., Mohanty, K., and Kalita, P., (2024)
 214 Performance assessment of innovative waste management system developed for the
 215 production of bio-fertilizer, *Sustainable Chemistry for the Environment*, 7, 100148, ISSN 2949-
 216 8392, <https://doi.org/10.1016/j.scenv.2024.100148>. (<https://www.sciencedirect.com/science/article/pii/S2949839224000919>)
 217

218 Szulc, W., Rutkowska, B., Gawroński, S., Wszelaczyńska, E., (2021) Possibilities of Using Organic
 219 Waste after Biological and Physical Processing-An Overview. *Processes*, 9, 1501.
 220 <https://doi.org/10.3390/pr9091501>

221 Tsai, S., Liu, C., and Yang, S., (2007) Microbial conversion of food wastes for biofertilizer production
 222 with thermophilic lipolytic microbes, *Renewable Energy*, 32 (6): 904-915, ISSN 0960-1481,
 223 <https://doi.org/10.1016/j.renene.2006.04.019>. (<https://www.sciencedirect.com/science/article/pii/S0960148106001030>)
 224

225 Vijaya, D., Padmadevi, S.N., Vasandha, S., Meerabhai, R.S., and Chellapandi, P., (2008) "Effect of
 226 vermicomposting coirpith on the growth of *Andrographis Paniculata*", *Journal of Organic*
 227 *Systems*, 3 (2), 51-56.

228 Wiharyanto, O., Mochtar, H., Purwono, P., (2018) Decomposition of food waste using bulking agent and
 229 bio-drying technology. *E3S Web of Conferences*. 73. 05013. 10.1051/e3sconf/20187305013.

230 Yong, Z. J., Bashir, Mohammed J.K., and Hassan, M. S., (2021) Biogas and biofertilizer production from
 231 organic fraction municipal solid waste for sustainable circular economy and environmental
 232 protection in Malaysia, *Science of The Total Environment*, 776, 145961, ISSN 0048-9697,
 233 <https://doi.org/10.1016/j.scitotenv.2021.145961>. (<https://www.sciencedirect.com/science/article/pii/S0048969721010287>)
 234

235 ABBREVIATIONS

236 Rhizo= Rhizobium, PSB= Phosphorus solubilizing bacteria, KSB= Potassium solubilizing
 237 bacteria, Tricho.= Trichoderma, *i.e.* = that is