

# ANTIPLASMODIAL PROPERTIES OF MEDICINAL PLANTS FROM EAST SINGHBHUM DISTRICT OF JHARKHAND IN INDIA

## ABSTRACT

It has been reported that malaria is one of the most important vectors borne disease in Jharkhand with significant morbidity and mortality. The National Vector Borne Disease Control Programme has prepared new treatment guidelines for malaria that recommend phytochemical constituents based therapy as the first-line drugs for malaria. It is being observed in the treatment of malaria that the cases of drug resistant are increasing day by day. Finding of new antiplasmodial or antimalarial plant species to develop effective malarial drugs are the necessity of time to avoid the problem of drug resistant. The tribal people have been using traditional medicinal plants since ancient time. They are using the medicinal plants found in their area for the treatment of fever, loss of appetite, digestive disorders, diabetes, jaundice, joint pain, skin and various other diseases. A community level study was conducted to document the traditional knowledge and identify the antimalarial plant species with the help of traditional healers (vaidya/hakeem) as well as by involving Sabar, Santal, Ho tribes and other communities of the area. The present study is focused on the exploration of antiplasmodial and antimalarial medicinal plants which are traditionally used by the ethnic or tribal people of East Singhbhum district in Jharkhand. Further the review of scientific literatures were conducted to validate the documented knowledge and related antimalarial plant species. A total of 29 medicinal plants of East Singhbhum have been identified and reviewed which have good antiplasmodial properties. These medicinal plants are rich in bioactive compounds such as flavanoids, alkanoids, terpenoids, tannins, essential oils, anthraquinones and pyrroles etc. The present study deals with the details of those active compounds and concludes the best drug for control of *Plasmodium vivax* and *P. falciparum*. However, more advanced research is needed to establish the effectiveness of bioactive compounds in vitro and in vivo, understand their basic mechanisms of action and begin the process of clinical research.

**KEYWORDS:** Medicinal Plants, Traditional Medicines, Antimalarial, Antiplasmodial Properties, Bioactive Compounds, East Singhbhum, Jharkhand, India

## 1. INTRODUCTION

India has been center to varied range of medicinal plants for centuries which are used by the tribal and local communities in the treatment various ailments. These plants are not only used in human diseases but also applied in veterinary treatment and even insect-pests management of different agricultural crops. There are about 18000 flowering plant species in India of which around 7000 are medicinally important. These plants are used in folk or traditional and documented systems of medicines including Ayurveda, Unani, Siddha and Homeopathy. These documented plants and their associated traditional knowledge are key resource base for traditional healing systems [1]. Jharkhand is one of the biodiversity rich states of India because of its diverse geographical and ecological conditions. The state is land of different tribes, tropical dry deciduous forests, flora and fauna and hence called cultural and herbal heritage and natural

sanctuary of the country. The total forest cover in Jharkhand is 23,605 km<sup>2</sup> which is 29.62% of the geographical area of Jharkhand [2]. The rich biodiversity of Jharkhand has more than 600 species of medicinally important plants. About 70% of the state population relies on traditional and herbal systems of medicines. East Singhbhum, the focused area of the present study is one of the biodiversity rich districts of Jharkhand. The district is dominated by ethnic and tribal populations (28.5%), which include Santal, Ho, Bhumij, Munda and Oraon [3]. These people have a treasure of traditional knowledge in ethno-medicines and herbal formulations. Forest cover and biodiversity of the district is important source livelihood, food and health of the local communities. The community used medicinal plants available in their area as folk medicines for centuries. These medicinal plants contain various bioactive compounds, mainly secondary metabolites and essential oils which are applied as the potential cure to different diseases or disorders including malaria in human beings. Malaria is a vector born disease widely observed in different climatic zones of Jharkhand. Changing in climatic conditions is one of the major factors spreading malaria. The warm and humid conditions favor the breeding of vectors at large scale. Hence, the transmission of malaria is a dynamic process influenced by the changes in ecological and meteorological conditions [4, 5]. It has been reported that at present the malarial patients in Jharkhand is about 31,600 [6]. Malaria has been a major health problem in the East Singhbhum region of Jharkhand particularly in rainy season. [7]. The cases of malaria in East Singhbhum region of Jharkhand are about 3000. Most of the tribal peoples and vulnerable tribal groups (PVTG) of the region are suffering from malaria. These communities are living in inaccessible forests, hills, valleys, and perennial streams. This topography and environment are favorable for rapid vector growth and the propagation of malaria parasite that results in a strong transmission of fatal form of malaria parasite *Plasmodium falciparum*. The disease is a big cause of death, disability and financial loss of the rural and tribal peoples of the area. The community has limited access to timely treatment and social securities. Experts found that malaria is the reason of maternal death, stillbirth, low birth weight in infants and weak immunity [8, 9]. The government of India has adopted a framework for the elimination of malaria covering the period 2016 to 2030 as per the Global Technical Strategy. The aim set by the government is to reach zero Malaria cases by 2027 before the certification for Malaria free status by WHO [4]. Medicinal plants traditionally used by tribal people can play effective role to achieve the aim set by the government to eliminate Malaria. The rural and tribal people of the Jharkhand use different plant based traditional medicines in the treatment of Malaria and high fever. It was found in the reviewing of different scientific literatures that there are 29 medicinal plants which are mostly used by the people of East Singhbhum and other regions of Jharkhand in the treatment of Malaria and intermittent fever. These medicinal plants are rich in bioactive compounds such as flavanoids, alkanoids, terpenoids, tannins, essential oils, anthraquinones and pyrroles etc. The present study deals with the review of all the 28 plant species to know about the antimalarial properties of the plants and their active compounds for control of *Plasmodium vivax* and *P. falciparum*. This review work will provide updated information for further research on the effectiveness of bioactive compounds and development of antiplasmodial drugs.

## 2. ANTIPLASMODIAL PLANT SPECIES

East Singhbhum region of Jharkhand has a rich diversity of medicinally important plant species along with traditional healing knowledge. It has been reported in several studies conducted by ethnobotanists in and around the region that there are a large number of plant species used by

ethnic people and traditional healers in fever, malaria, body pain, shivering and headache. A total of 29 highly effective antiplasmodial or antimalarial plant species of the region identified and documented in the present study are mentioned below.

Table 1 : Antiplasmodial Medicinal Plants of East Singhbhum

S. No.	Plant Species	Common Name	Part used	Traditional Uses
1.	<i>Aegle marmelos</i>	Wood Apple, Bael	Leaf, Seed	Antipyretic, Antidiabetic [10], Analgesic, Antiinflammatory, Antihepatotoxic, Antibacterial [11]
2.	<i>Andrographis paniculata</i>	King of Bitters, Kalmegh	Whole plant	Antimalarial, Antipyretic, Antiinflammatory, Antidiabetic, Antihypertensive, Antihepatotoxic [12, 13]
3.	<i>Aristolochia indica</i>	Birthwort, Snakeroot, Ishwarmul	Leaf, Root	Snakebites, Fever, Malaria, Skin Diseases, Helminthes, Intestinal Disorders [14]
4.	<i>Artemisia indica</i>	Wormwood, Majtari	Whole plant	Malarial Fever, Tuberculosis, Lice, Wounds, Scabies, Dysentery [15]
5.	<i>Azadirachta indica</i>	Neem	Leaf, Stem, Bark, Roots, Fruits	Fever, Malaria, Headache, Ulcers, Diabetes, Leprosy, Chickenpox, Skin Disease [16]
6.	<i>Berberis aristata</i>	Barberry, Daru, Haldi	Root	Malaria, Skin Diseases, Diarrhea, Eye Problem, Inflammation, Wound Healing, [17]
7.	<i>Cassia alata</i>	Ringworm Shrub, Senna,	Leaf, Seeds, Stem, Flower	Eczema, Ringworm, Malaria, Constipation [18]
8.	<i>Cassia occidentalis</i>	Coffee Senna, Chakod	Root bark	Fever, Laxative, Whooping Cough, Skin Diseases, Asthma, Joint pains [19]
9.	<i>Cissampelos pareira</i>	Velvetleaf, Harjori or Akanadi	Root, Stem	Malaria, Ulcer, Wound, Rheumatism, Asthma, Cholera, Diarrhoea, Snakebite, Rabies [20]
10.	<i>Curcuma longa</i>	Turmeric, Haldi	Rhizome	Malaria, Cancer, Skin Diseases, Joint Pain [21]
11.	<i>Cyclea barbata</i>	Hairy Cyclea, Raj Patha	Root, Leaf	Malarial Fever, Stomach Trouble [22, 23, 24]
12.	<i>Cyperus rotundus</i>	Purple Nutsedge, Motha	Root, Tuber	Anti-Malarial, Anti-Pyretic, Anti-Inflammatory, Hepatoprotective, Anti-Obesity, Anti-Diabetic [25, 26]
13.	<i>Datura metel</i>	Thornapple, Datura	Fruit	Malaria, Dandruff, Arthritis, Liver Problem, Viral Fever,

				Antifungal [27]
14.	<i>Diospyros melanoxyton</i>	Coromandel ebony, Tendu	Bark	Antiplasmodial, Antiprotozoal Antimicrobial, Anti-Inflammatory, Antidiabetic [28]
15.	<i>Eclipta alba</i>	False Daisy, Bhringaraj	Leaf	Malaria, Fever, Hair loss, Jaundice, Skin Diseases, [30, 31, 32]
16.	<i>Ficus sur</i>	Bush Fig	Stem Bark, Leaf	Malarial Fever, Leukoderma, Leprosy, Wounds, Respiratory Problems, Diarrhoea, Tuberculosis, Anaemia, Epilepsy [33]
17.	<i>Holarrhena pubescens</i>	Fever pod, Indrajau	Root, Bark, Seed	Malarial Fever, Diabetes, Piles, Leprosy, Diarrhea [34, 35]
18.	<i>Lantana camara</i>	Lantana, Putoos	Leaf	Fevers, Colds, Rheumatism, Asthma, Cuts, Intestinal Worms [36, 37, 38]
19.	<i>Moringa oleifera</i>	Drumstick Tree, Sahjan	Leaf, Bark, Flower, Fruit, Seed	Malaria, Malnutrition, Water Purifier, Joint Pain, Diabetes, Hypertension, Wound, Paralysis [39, 40, 41, 42, 43]
20.	<i>Ocimum sanctum</i>	Holy Basil, Tulsi	Bark, Leaves, Flowers, Roots, Fruits, Seeds	Malarial Fever, Cold, Headache, Stomach Disorders, Inflammation, Heart Disease [47, 48]
21.	<i>Picrorhiza kurroa</i>	Bitter-root, Kutki	Root, Rhizome	High Fever, Stomachache, Cough, Cold, Jaundice, Diarrhea, Dysentery [49, 50, 51]
22.	<i>Plumbago zeylanica</i>	Ceylon leadwort, Chitrak	Root, Aerial Part	Anti-Plasmodial, Anti-Tumour, Anti-Bacterial, Hepatoprotective, Anti-Fungal, Anti-Inflammatory [52, 53, 54]
23.	<i>Pongamia pinnata</i>	Indian Beech, Karanj	Leaf, stem, bark, flower, fruit, kernel, latex	Malaria, Fever, Wound, Skin Diseases, Diabetes, Head Lice [55, 56, 57, 58, 59]
24.	<i>Solanum incanum</i>	Bitter Apple, Jangli Bhata	Leaf, Fruit, Root	Malaria, Sore Throat, Stomach-ache, Head-ache, Painful Menstruation, Liver pain, Diabetes, Common Cold, Snake Bites [60, 61]
25.	<i>Solanum nigrum</i>	Black Nightshade, Makoi	Leaf, Fruit, Root	Malarial Fever, Eczema, Urinary Tract Infections, Dysentery, Prostate [62, 63]
26.	<i>Stachytarpheta cayennensis</i>	Snakeweed, Barahmasi	Leaf, Root	Malarial Fever, Eyewash, Laxative Inflammation, Hepatic and Renal Disorder, Helminthiasis, Diabetes [64,

				65]
27.	<i>Swertia chirayita</i>	Bitter stick, Chiraita	Whole Plant	Malarial Fever, Blood Purification, Laxative, Inflammation, Tuberculosis, Diabetes [66]
28.	<i>Swietenia mahagoni</i>	Mahogany	Seed, Bark	Malaria, Diabetes, Diarrhea, Wound [68, 69]

### 3. ANTIPLASMODIAL BIOACTIVE COMPOUNDS OF THE ENLISTED PLANT SPECIES

#### 3.1 *Aegle marmelos*

*Aegle marmelos* is a valuable medicinal tree. All parts of the tree are used in traditional system of medicine. The bioactive compounds extracted from the fruits, leaves, stem, stem bark, root, and root bark of *A. marmelos* have different pharmacological properties including antiplasmodial efficacy [69]. Terpenoids and Fagarine are two important bioactive compounds that have antimalarial or antiplasmodial properties [70]. *In vitro* antimalarial activity of *A. marmelos* leaf methanol extract has showed highest activity against *Plasmodium falciparum* [70, 71].

#### 3.2 *Andrographis paniculata*

Andrographolide and diterpene lactone compound are derived from *Andrographis paniculata* as a potent antiplasmodial phytochemicals. These compounds are synergistic with curcumin and additively interactive with artesunate. It has been reported that andrographolide-curcumin exhibited better antimalarial activity. These are nontoxic compounds can be used as template molecule for designing new derivatives with improved antimalarial properties [72, 73].

#### 3.3 *Aristolochia indica*

The antimalarial bioactive compounds isolated from different parts of *A. indica* are aristolochic acid, ceryl alcohol, -sitosterol, stigmast-4-en-3-one, friedelin, cycloeucaenol, and rutin [74]. Some of the compounds are extracted from essential oil of the plants; these are trans-pinocarveol (24.2%),  $\alpha$ -pinene (16.4%) and pinocarvone (14.2%) [75].

#### 3.4 *Artemisia indica*

Some of the chemical constituents extracted from *Artemisia indica* are effective against *Plasmodium falciparum*. These are exiguaflavone A, exiguaflavone B, maackiain, and 2-(2, 4-dihydroxyphenyl)-5,6-methylenedioxybenzofuran. Exiguaflavones A and B exhibit *in vitro* antimalarial activities of  $4.60 \times 10^{-6}$  and  $7.05 \times 10^{-6}$  g/mL, respectively, against *Plasmodium falciparum* [76]. Artemisinin and polymethoxyflavonoid are also antiprotozoal potential extracted from essential oil [77].

#### 3.5 *Azadirachta indica*

*A. indica* has been reported to be effective against *P. falciparum* in humans [78]. It has been reported that *in vitro* and *in vivo* models show that aqueous, methanolic, or ethanolic extracts of neem bark, stems and leaves have significant activity against the strains of *Plasmodium falciparum* and *Plasmodium berghei* [79]. Some of the researchers have found that limonoid

deacetylnimbin extracted from neem seeds is effective against early sporogony stages of *P. bergehi* [81]. Azadirachtin is one of the important phytochemicals isolated from *A. indica* is effective in the treatment for cerebral malaria [80].

### **3.6 *Berberis aristata***

The roots and other parts of *Berberis aristata* contain alkaloids such as berberine and berbamine, which have antimalarial properties. The berberine is an isoquinoline which are used as a substitute of quinine in the treatment of malaria. [81].

### **3.7 *Cassia alata***

The fresh leaf extract of *Cassia alata* was observed as antimalarial properties against one of the strains of *P. falciparum* parasites. This action was generated by some of the phytochemicals of the plant leaves, such as anthraquinones, alkaloids, tannins, flavonoids, terpenoids and steroids. The dichloromethane/methane (1:1, v/v) extract of *Cassia alata* was also reported effective against *P. falciparum* [82, 83].

### **3.8 *Cassia occidentalis***

The leaf extract of *Cassia occidentalis* has reported outstanding antiplasmodial activity *in vitro*. The important phytochemicals including Anthraquinone, Emodin, Rhein Crysophanol, Pinselín, Physcion and Aloe emodin are effective against the malaria [84, 85].

### **3.9 *Cissampelos pareira***

The roots and stem of *Cissampelos pareira* contain alkaloids hayatinine and isoquinoline which are effective antimalarial agents. There are scientific evidences for the traditional use of this plant in the treatment of malaria [86].

### **3.10 *Curcuma longa***

*Curcuma longa* has different kinds of bioactive compounds effective against a large range of ailments. Curcumin is one of the most important phytochemicals of *C. longa* has been in malaria, skin diseases and joint pains. It has been observed that curcumin, a polyphenolic organic molecule restrain chloroquine-resistant *P. falciparum* growth in culture in a dose dependent mode with an IC<sub>50</sub> of ~5 µM. Moreover, oral administration of curcumin to mice infected with malaria parasite (*P. berghei*) reduces blood parasitemia by 80–90% and enhances their survival significantly. Therefore, curcumin may represent a novel treatment for malarial infection [87, 88].

### **3.11 *Cyclea barbata***

The fresh leaf of *Cyclea barbata* has been used traditionally in various ailments including intermittent fever, intestinal problems, bacterial diseases, inflammation and wound healing. It has been reported that the alkaloids and flavonoids of the plant species present in leaf, stem and root are effective in malaria [89].

### **3.12 *Cyperus rotundus***

It was investigated that the crude hexane extracts of the air-dried tubers of *Cyperus Rotundus* is having high potency in the *in vitro* test against *Plasmodium falciparum* (EC<sub>50</sub> = 0.66 µg ml<sup>-1</sup>);

the result showed the significant anti-malarial activity of *Cyperus Rotundus*. the anti-malarial activity of *C. rotundus*. Among tested compounds, 10, 12-peroxycalamenene had a maximum effect with an EC<sub>50</sub> of  $2.33 \times 10^{-6}$  M and dichloromethane extract with an IC<sub>50</sub> between 10 and 50 µg/mL [90, 91, 92].

### **3.13 *Datura metel***

*Datura metel* is one of the plant species used in the traditional and modern systems of medicines for different diseases. The leaf and seed of *D. metel* contain alkaloids, flavonoids, glycosides, steroids and saponins which could be effective against larvicidal activities [93, 94].

### **3.14 *Diospyros melanoxylon***

The bark of *Diospyros melanoxylon* is used in the treatment of malarial fever. Plumbagin and diospyrin are the two significant bioactive compounds of *D. melanoxylon* which have potent activities against *P. falciparum* [95].

### **3.15 *Eclipta alba***

*Eclipta alba* is one of the highly valuable medicinal plants which has different important pharmacological activities such as antimalarial, analgesic, hair growth promoting, hepatoprotective, antibacterial and antidiabetic [98]. The fresh tender leaf extract of the species is against *P. berghei* as it contains various phytoconstituents such as β-amirin, phytosterols, flavones (luteolin), coumarin (wedelolactone) and triterpenes (ecalbatin, ursolic acid and echinocystic acid), Sesquiterpene lactones. Other biochemicals present in the plant parts are 5-hydroxymethyl-(2,2':5',2'')-terthienyl tiglate, 5-hydroxymethyl-(2,2':5',2'')-terthienyl agelate, 5-hydroxymethyl-(2,2':5',2'')-terthienyl acetate [96].

### **3.16 *Ficus sur***

The review of current status of *Ficus sur* has shown that the plant species has antiplasmodial activity. *F. sur* contains pentacyclic triterpenoids of oleanane and ursine structures as well as flavonoids and coumarins which are effective against antiplasmodial activities [97].

### **3.17 *Holarrhena pubescens***

*Holarrhena pubescens* is one of the important medicinal plant species of Indian subcontinent. The species is used in traditional systems of medicine by the tribal people. The bark, seed and root of the plant are used in diabetes and malarial fever. The species contains several bioactive compounds including holarrhimine, conarrhimine, conessine, isoconessine, antidysentericine, conessimine, conimine, conessidine, mokluangin A-D and konkurchicine. Chloroform extracts of the plant bark reported significant in vitro and in vivo anti-malarial activity [98, 99].

### **3.18 *Lantana camara***

*Lantana camara* is a common plant species in East Singbhum region of Jharkhand which is used by the tribal communities for various uses. The aqueous and ethanolic leaf extracts of the plant leaf have shown antimalarial activity close to that of the standard drug chloroquine. The leaf extracts of *L. camara* contains alkaloids, flavonoids, terpenoids, saponins, carbohydrates, steroids and tannins in the different extracts [100].

### **3.19 *Moringa oleifera***

*Moringa oleifera* is a very common plant in India including Jharkhand. The leaves and seeds of this plant are the main parts used in the formulation of medicine for the treatment of different ailments including malaria. It has been confirmed that the bioactive compounds extracted from the leaves and seeds are effective against *P. falciparum*, *P. berghei*, *P. chabaudi* and *P. yoelii*. Flavonoids such as apigenin, kaempferol, rutin, and quercetin extracted from the leaf and seeds have shown antimalarial properties [101, 102, 103].

### **3.20 *Ocimum sanctum***

Extracts of leaves and roots of *Ocimum sanctum* Linn are used as antimalarial against *Plasmodium berghei* in vivo. The antiplasmodial bioactive compounds of *O. sanctum* include eugenol, apigenin, ursolic acid, rosmarinic acid [104, 105]

### **3.21 *Picrorhiza kurroa***

The dried roots of *Picrorhiza kurroa* are traditionally used in the treatment malaria and intermittent fever. Kutkin is the important bioactive compound of the plant species which includes kutkin is the active principle of *P. kurroa* and is comprised of kutkoside and the iridoid glycoside picrosides. Other bioactive compounds are apocynin, drosin, and nine cucurbitacin glycosides [106, 107].

### **3.22 *Plumbago zeylanica***

The root extract of *Plumbago zeylanica* contains plumbagin which was evaluated for chemical constituent and antimalarial effect against *Anopheles stephensi* Liston (Diptera). Plumbagin compound of *P. zeylanica* can be considered as a new source of natural larvicide for the control of malarial vector [108].

### **3.23 *Pongamia pinnata***

*Pongamia pinnata* has been reported to have antiplasmodial or antimalarial properties. The methanol extract of the bark of the tree contains alkaloids, flavonoids, triterpenes, tannins, carbohydrates, phenols, coumarins, saponins, phlobatannins and steroids. The isolated bioactive compounds are effective against *Plasmodium falciparum* and *Plasmodium berghei* parasites [109].

### **3.24 *Solanum incanum***

The leaves of *Solanum incanum* have prophylactic antimalarial properties. The ethyl acetate fraction of the leaves contain an alkane compound which play a significant role in the inhibition of enzymatic activity [110].

### **3.25 *Solanum nigrum***

*Solanum nigrum* is an important medicinal plant used in the treatment of malaria in traditional medicine. The bioactive compounds of the herb are alkaloids, flavonoids, phenolic, diterpenes and saponin isolated from the leaf ethanol extract of the plant used in malaria and other diseases [111].



### **3.26 *Stachytarpheta cayennensis***

The ethanolic leaf extract of *Stachytarpheta cayennensis* contains antimalarial properties as observed in its characteristic to suppress *Plasmodium berghei* infection. The fresh leaf extract of *S. cayennensis* constitutes antiplasmodial bioactive compounds such as alkaloids, terpenes and flavonoids [112].

### **3.27 *Swertia chirayita***

*Swertia chirayita*) is an important herb used traditionally in the treatment of several ailments including malaria, diabetes and liver problems. The herb is rich source of flavonoids, xanthenes, terpenoids, iridoid and alkaloids. The chemical components such as swerchirin, swertinin, decussatin, 1-hydroxy-3,5,8-trimethoxyxanthen-9-one and chiritol of *S. chirayita* have effective binding interactions to control malaria [113].

### **3.28 *Swietenia mahagoni***

In traditional system of medicine, *Swietenia mahagoni* is an important medicinal plant used in the treatment of malaria, diarrhea and diabetes. Leaf, bark and seed of the plant consist of secondary metabolites which is effective in malarial fever, bacterial and fungal infection. The methanol extract of seed of the plant was reported with antimalarial properties [115, 116].

## **4. SAFETY OF THE DRUGS**

The above mentioned medicinal plants and bioactive compounds are valuable sources for the development of novel drugs. The extracts of medicinal plants used in the treatment of malaria are common as an alternative to modern antimalarial drugs. The recommended doses of the antiplasmodial bioactive compounds were found safe and non-toxic for human beings. However, the efficacy and safety of most of the herbal medicines has not yet been established.

## **5. CONCLUSION**

Malaria is one of the lives threatening parasitic disease affecting the whole developing and poor countries causing huge social and economic loss. The situation is harsh as the malarial parasites have developed resistant against majority of the modern drugs. To control and eradicate the devastating disease and overcome of the drug resistance, novel medicinal plant based bioactive compounds are required. There are many plant species in East Singhbhum region of Jharkhand which have been left unexplored with its properties to cure malaria. Further studies are required to find out the phytochemical constituents effective in the treatment of malarial.

## **6. FUTURE REMARKS**

Development of new antiplasmodial drugs using the bioactive compounds extracted from medicinal plants will play a big role in solving the problems of developing countries to combat this formidable disease. The pace of research progress must be high to develop novel antiplasmodial medicines.

**DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Artificial Intelligence (AI) and AI-assisted technologies have not been used in the writing process of this paper.

**CONSENT AND ETHICAL APPROVAL**

Not required

**COMPETING INTERESTS**

I declare that no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the position presented in, or the review of, the manuscript entitled.

UNDER PEER REVIEW

Plate 1 : Photographs of the plants



*Aegle marmelos*



*Andrographis paniculata*



*Aristolochia indica*



*Artemisia indica*



*Azadirachta indica*



*Berberis aristata*



*Cassia alata*



*Cassia occidentalis*



*Cissampelos pareira*



*Curcuma longa*



*Cyclea barbata*



*Cyperus rotundus*



*Datura metel*



*Diospyros melanoxylon*



*Eclipta alba*



*Ficus sur*



*Holarrhena pubescens*



*Lantana camara*



*Moringa oleifera*



*Ocimum sanctum*



*Picrorhiza kurroa*



*Plumbago zeylanica*



*Pongamia pinnata*



*Solanum incanum*



*Solanum nigrum*



*Stachytarpheta cayennensis*



*Swertia chirayita*

## REFERENCES

1. <https://nmpb.nic.in/content/medicinal-plants-fact-sheet>
2. Anonymous. Forest Survey of India, Ministry of Environment, Forest and Climate Change, Dehradun, Uttarakhand, India.
3. District Census Handbook, Purbi Singhbhum, Jharkhand. 2011
4. World Health Organization, editor. Global Technical Strategy for Malaria 2016–2030, monograph on the Internet. Geneva; 2016.
5. Editor. National Framework for Malaria Elimination in India 2016–2030, Government of India, New Delhi. 2016.
6. National Vector-Borne Disease Control Programme, editor. Annual Report 2018, Monograph on the Internet. New Delhi; 2018.
7. Malaria Annual Report. National Centre for Vector Borne Diseases Control, 22-Shamnath Marg, Delhi-110054. 2023.
8. Singh N., Singh O.P. and Sharma V.P. Dynamics of malaria transmission in forested and deforested region of mandla district, Central India, Madhya Pradesh. *J Am Mosq Contr Assoc* 1996;12:225-234.
9. Schantz-Dunn J, Nour NM. Malaria and pregnancy: a global health perspective. *Rev Obstet Gynecol*. Summer. 2009; 2(3):186-92.
10. Lampronti I, Martello N, Bianchi N, Borgatti M, Lambertini E, Piva R, Jabbar S, Choudhuri MSK, Khan MTH, Gambari R. In vitro antiproliferative effects on human tumor cell lines of extracts from the Bangladeshi medicinal plant *Aegle marmelos* Correa. *Phytomedicine*. 2003; 10:300-08.
11. Singh J. Anjana Shukla A. S., Singh S. C., Jain S. P., Ashok Sharma A. S., and Khanuja S. P. S.. Traditional uses of *Aegle marmelos* (Bilva). 2007.
12. Dua V.K., Ojha V.P., Roy R., Joshi B.C., Valecha N., Usha C., Devi M.C., Bhatnagar V.P., Sharma S.K. and Subbarao. Anti-malarial activity of some xanthenes isolated from the roots of *Andrographis paniculata*, *Journal of Ethnopharmacology*. 2004; 95:2–3:247-251.
13. Kabir M. H., Hasan N., Rahman M. M., et al. A survey of medicinal plants used by the Deb barma clan of the Tripura tribe of Moulvibazar district, Bangladesh. *Journal of Ethnobiology and Ethnomedicine*. 2014;10
14. Heinrich M. et al. Local uses of *Aristolochia* species and contents of nephrotoxic aristolochic acid 1 and 2—a global assessment based on bibliographic sources, *Journal of Ethnopharmacology*. 2009.
15. Nurlybekova, A., Kudaibergen A., Kazymbetova A., Amangeldi M., Baiseitova A., Ospanov M., Aisa H.A. Ye Y. Ibrahim M.A. Jenis J. Traditional Use, Phytochemical Profiles and Pharmacological Properties of *Artemisia* Genus from Central Asia. *Molecules*. **2022**; 27:5128.
16. Pankaj S, Lokeshwar T, Mukesh B, Vishnu B. Review on neem (*Azadirachta indica*): Thousand problems one solution. *IRJP*. 2011;2(12):97-102.
17. Chander V., Aswal J. S., Dobhal R., Uniyal D. P. A review on Pharmacological potential of Berberine; an active component of Himalayan *Berberis aristata*. *Journal of Phytopharmacology*. 2017.6(1):53-58.
18. Palanichamy S., Nagarajan S. Analgesic activity of *Cassia alata* leaf extract and kaempferol 3-O-sophoroside. *Journal of Ethnopharmacology*. 1990; 29:73-78.
19. Yadav J.P., Vedpriya A., Yadav S., Panghal M., Kumar S., Dhankhar S. *Cassia occidentalis* L.: A review on its ethnobotany, phytochemical and pharmacological profile, *Fitoterapia*. 2010;81(4): 223-230.
20. Kumari S., Vinod B.A., Shivprasad P., Upendra S.S., *Cissampelos pareira* L.: A review of its traditional uses, phytochemistry and pharmacology, *Journal of Ethnopharmacology*. 2021;274: 113850.
21. Reddy R. C., Vathsala P. G., Keshamouni V. G., Padmanaban G., Rangarajan P. N. Curcumin for malaria therapy. *Biochem. Biophys. Res. Commun*. 2005;326:472-474.
22. Nandakumar D.N., Nagaraj V.A., Vathsala P.G., Rangarajan P., Padmanaban G. Curcumin-artemisinin combination therapy for malaria. *Antimicrob Agents Chemother*. 2006; 50(5):1859-60.

23. Arba P. R., Paloque L., Belda H., Anshory H., Masriani T., Jean-Michel A., J., Valentin A., Agus W.A., Mustofa, Benoit-Vical F. Antiprotozoal properties of Indonesian medicinal plant extracts, *Journal of Herbal Medicine*. 2018;11:46-52.
24. Manilal K.S., Sabu T. *Cyclea barbata* miers (menispermaceae): a new record of a medicinal plant from South India. *Anc Sci Life*. 1985;4(4):229-31.
25. Kirtikar & Basu. *Indian Medicinal Plants*. International Book distributors & publishers, Dehradun. 1(III):2638-2640.
26. Nadkarni A.K., *Indian Materia Medica*, Popular Prakashan, Mumbai. 1976;(1):428-430.
27. Alam W., Khan H., Khan S.A., Nazir S., Akkol E.K. *Datura metel*: A Review on Chemical Constituents, Traditional Uses and Pharmacological Activities. *Curr Pharm Des*. 2021;27(22):2545-2557.
28. Kantamreddi, Venkata & Wright, Colin. Investigation of Indian Diospyros Species for Antiplasmodial Properties. *Evidence-based complementary and alternative medicine:eCAM*. 2008;(5):187-90.
29. Harun M.A. Rashid I., Majumder S., Mandal V., Subhash C. M., Amirthalingam R. T. In search of suitable extraction technique for large scale commercial production of bioactive fraction for the treatment of diabetes: The case *Diospyros melanoxylon* Roxb., *Journal of Traditional and Complementary Medicine*. 2019;9(2):106-118.
30. Pani M., Nahak G., Sahu R.K. Review on ethnomedicinal plants of Odisha for the treatment of malaria. *International Journal of Pharmacognosy and Phytochemical Research*. 2014;7:156–165.
31. Jahan R., Al-Nahain A., Majumder S., Rahmatullah M. Ethnopharmacological Significance of *Eclipta alba* (L.) Hassk. (Asteraceae) *Int. Sch. Res. Not*. 2014;385-969.
32. Dalal S., Kataria S., Sastry K., Rana S.V.S. Phytochemical Screening of Methanolic Extract and Antibacterial Activity of Active Principles of Hepatoprotective Herb, *Eclipta alba*. *Ethnobot. Leafl*. 2010;3.
33. Clark A.M. Natural products as a resource for new drugs. *Pharm Res*. 1996;13:1133–1144.
34. Yetein M.H., Houessou L.G., Loubégnon T.O., Teka O., Tente B. Ethnobotanical study of medicinal plants used for the treatment of malaria in plateau of Allada, Benin (West Africa). *J Ethnopharmacol*. 2013.
35. Wyk V.B.E., Gericke N. *People's plants. A guide to useful plants of Southern Africa*. Pretoria, South Africa: Briza Publications. 2000.
36. Sharma V.S., Kaul K.N. *Chem. Abstr*. 1959;53:652.
37. Sastri B.N. *The Wealth of India*, New Delhi, Council of Scientific and Industrial Res. 1962;5.
38. Verma R.K., Verma S.K. Phytochemical and termiticidal studies of *Lantana camara* var *aculeata* leaves. *Fitoterapia*. 2006;(77):466-468.
39. Adekunle M.F. Indigenous uses of plant leaves to treat malaria fever at Omo Forest reserve (OFR) Ogun state, Nigeria. *Ethiop J Environ Stud Manag*. 2008;1(1):31–35.
40. Dube F.F., Tadesse K., Birgersson G., Seyoum E., Tekie H., Ignell R., Hill S.R. Fresh, dried or smoked? Repellent properties of volatiles emitted from ethnomedicinal plant leaves against malaria and yellow fever vectors in Ethiopia. *Malar J*. 2011;10:375.
41. Alebie G., Urga B., Worku A. Systematic review on traditional medicinal plants used for the treatment of malaria in Ethiopia: trends and perspectives. *Malar J*. 2017;16:307.
42. Dhawan B.N., Dubey M.P., Mehrotra B.N. Screening of Indian plants for biological activity: part IX, *Indian J Exp Biol*. 1980;18:594–606.
43. Kulkarni A.R. Comparative studies on nutritive values of tender foliage of seedlings and mature plants of *Moringa oleifera* Lam *J Econ Taxon Bot*. 1993;17:479–85.
44. Biswas N.P., Biswas A.K. Evaluation of some leaf dusts as grain protectant against rice weevil *Sitophilus oryzae* (Linn.) *Environ Ecol*. 2005;23:485–8.
45. Arya D., Bhatt D., Kumar R., Tewari L.M., Kishor K., Joshi G.C. Studies on natural resources, trade and conservation of Kutki (*Picrorhiza kurroa* Royle ex Benth., Scrophulariaceae) from Kumaun Himalaya. *Scientific Res Essays*. 2013; 8(14): 575-580.
46. Khan A.H., Zaidi S.H. Propagation and Regeneration Technology of Pharmacopeial Medicinal Plants of the Temperate Regions of Pakistan. *Biological Sciences Research Division, Pakistan Forest Institute, Peshawar, Pakistan*. 1989;8.
47. Arya D., Bhatt D., Kumar R., Tewari L.M., Kishor K., Joshi G.C. Studies on natural resources, trade and conservation of Kutki (*Picrorhiza kurroa* Royle ex Benth., Scrophulariaceae) from Kumaun Himalaya. *Scientific Res Essays*. 2013;8(14):575-580.
48. Maniafu B.M., Wilber L., Ndiege I.O., Wanjala C.C., Akenga T.A. Larvicidal activity of extracts from three *Plumbago* spp against *A. gambiae*. *Mem Inst Oswaldo Cruz*. 2009;104(6):813-817.

49. Patil C.D., Patil S.V., Salunke B.K., Salunkhe R.B. Bioefficacy of *Plumbago zeylanica* (Plumbaginaceae) and *Cestrum nocturnum* (Solanaceae) plant Extracts against *Aedes aegypti* (Diptera: Culicidae) and *Poeciliareticulata*. Parasitol. Res. 2011;108(5):1253-1263.
50. Kumar R., Kumar S., Patra A., Jayalakshmi S. Hepatoprotective activity of aerial parts of *Plumbago zeylanica* linn against carbon tetrachloride induced hepatotoxicity in rats. International Journal of Pharmacy and Pharmaceutical Sciences. 2009;1:171-175.
51. Simonsen H.T., Nordskjold J.B., Smitt U.W., Nyman U., Palpu P., Joshi P. In vitro screening of Indian medicinal plants for antiplasmodial activity. Journal of Ethnopharmacology. 2001;74:195-204.
52. Srinivasan K., Muruganandan S., Lal J., Chandra S., Tandan S.K., Ravi Prakash V., Kumar D. Antinociceptive and Antipyretic Activities of *Pongamia pinnata* Leaves. Phytotherapy Research. 2003;17:259–264.
53. Dwivedi D., Dwivedi M., Malviya S., Singh V. Evaluation of wound healing, anti-microbial and antioxidant potential of *Pongamia pinnata* in wistar rats. Journal of Traditional and Complementary Medicine. 2016;7(2017):79-85.
54. Punitha, R., Vasudevan K., Manoharan S. Effect of *Pongamia pinnata* flowers on blood glucose and oxidative stress in alloxan induced diabetic rats. Indian Journal of Pharmacology 2006;38:62–63.
55. Mumcuoglu K.Y. Prevention and treatment of head lice in Children. PEDIATR DRUGS. 1999; 1:211-218.
56. Mirutse G., Tilahun T. Ethnobotanical Study of Medicinal Plants used by People in Zegie Peninsula, Northwestern Ethiopia, Journal of Ethnobiology Ethnomedicine. 2007;3:12.
57. Hyde M., Wursten B. Flora of Zimbabwe, Species Information of *Solanum incanum*. September 2006.
58. Gao S.H., Su Z. Z., Yang L. J., Li Z. Y. Chemical Components From Stems of *Solanum nigrum* by LC-MS and NMR. Chin. Tradit. Herb. Drugs. 2021;52 (5):1263–1273.
59. Aburjai T. A., Oun I. M., Auzi A. A., Hudaib M. M. Volatile Oil Constituents of Fruits and Leaves of *Solanum nigrum* L. Growing in Libya. J. Essent. Oil Bear. Plants. 2014;17(3):397–404.
60. Uphof J.J. Th. Dictionary of Economic Plants. Weinheim Publisher. 1959.
61. Cano J.H., Volpato G. Herbal mixture in the traditional medicine of Eastern Cuba. J Ethnopharmacol. 2004;90:293–316.
62. Rahman S., Husen A. Antimalarial Response, Traditional and Other Potential Uses of *Swertia* Genus. Antimalarial Medicinal Plants. Taylor & Francis. 2024;294-300.
63. Yelaware P. Naveen G., Rupini D, Ahmed F, Urooj A. Pharmacological effects and active phytoconstituents of *Swietenia mahagoni*: a review, Journal of Integrative Medicine. 2014;12(2): 86-93.
64. Angajala G., Ramya R., Subashini R. In-vitro anti-inflammatory and mosquito larvicidal efficacy of nickel nanoparticles phytofabricated from aqueous leaf extracts of *Aegle marmelos* Correa. Acta Tropica. 2014;135:19-26.
65. Misra P., Pal N.L., Guru P.Y., Katiyar J.C., Tandon J.S. Antimalarial Activity of Traditional Plant against Erythrocyte Stages of *Plasmodium berghei*. International Journal of Pharmacognosy. 1991;29:19-23.
66. Kamaraj C., Kaushik N.K., Rahuman A.A., Mohanakrishnan D., Bagavan A., Elango G. Antimalarial activities of medicinal plants traditionally used in the villages of Dharmapuri regions of South India. J. Ethnopharmacol. 2012;141(3):796–802.
67. Angajala G., Pavan P., Subashini R. One-step biofabrication of copper nanoparticles from *Aegle marmelos* Correa aqueous leaf extract and evaluation of its anti-inflammatory and mosquito larvicidal efficacy. RSC Adv. 2014;4(93):51459–51470.
68. Moghadamtousi S.Z., Goh B.H., Chan C.K., Shabab T., Kadir H.A. Biological activities and phytochemicals of *Swietenia macrophylla* King. Molecules. 2013;18(9):10465-8.
69. Dua V.K., Ojha V.P., Roy R. Anti-malarial activity of some xanthenes isolated from the roots of *Andrographis paniculata*. Journal of Ethnopharmacology. 2004;95(2-3):247–251.
70. Sati H., Sati B., Saklani S., Bhatt P.C., Mishra A. P. Phytochemical and pharmacological potential of *Aristolochia indica*: a review. Research journal of pharmaceutical, biological and chemical sciences. 2011;2(4):647-654.

71. Kanjilal P.B., Kotoky R., Couladis M. Chemical composition of the stem oil of *Aristolochia indica* L. *Journal of Essential Oil Research*. 2009; 21(1): 24-25.
72. Chanphen R., Thebtaranonth Y., Wanauppathamkul S., Yuthavong Y. Antimalarial principles from *Artemisia indica*. *J Nat Prod*. 1998;61(9):1146-7.
73. Chatterjee A., Pakraõshi S.C. *The treatise on Indian medicinal plants*. New Delhi: National Institute of Science Communication CSIR. 1997:142–143.
74. Adebayo J.O., Krettli A.U. Potential antimalarials from Nigerian plants: a review. *J Ethnopharmacol*. 2011;133: 289–302.
75. Tepongning R.N., Mbah J.N., Avoulou F. L., Jerme M. M., Ndanga E. K., Fekam F. B. Hydroethanolic Extracts of *Erigeron Floribundus* and *Azadirachta indica* Reduced *Plasmodium Berghei* Parasitemia in Balb/c Mice. *Evid. Based Complement. Altern. Med*. 2018;156:710.
76. Tapanelli S., Chianese G., Lucantoni L., Yerbanga R. S., Habluetzel A., Tagliatela-Scafati O. Transmission Blocking Effects of Neem (*Azadirachta indica*) Seed Kernel Limonoids on *Plasmodium Berghei* Early Sporogonic Development. *Fitoterapia*. 2016;114:122–126.
77. Okoh M.P., Singla R.K., Madu C., Soremekun O., Adejoh J., Alli L. A. Phytomedicine in Disease Management: In-Silico Analysis of the Binding Affinity of Artesunate and Azadirachtin for Malaria Treatment. *Front. Pharmacol*. 2021;12:751032.
78. Bhanu M., Giriraj T. K., Neerupma D., Joshi D.D., Chander S., Kharkwal A., Arun K. S., Kharkwal H. Recent advances on *Berberis aristata* emphasizing berberine alkaloid including phytochemistry, pharmacology and drug delivery system, *Journal of Herbal Medicine*. 2021;27: :100433:ISSN 2210-8033.
79. Vigbedor Y.B., Osafo Acquah S., Adu Gyan B.B., Lotsi B. In vitro antimalarial activity of the ethanol extracts of *Azela africana* and *Cassia alata* commonly used as herbal remedies for malaria in Ghana. *Int J Nov Res Life Sci*. 2015;2(6):10–16.
80. Atanu F.O., Rotimi D., Ilesanmi O.B., Al Malki J.S., Batiha G.E., Idakwoji P.A. Hydroethanolic extracts of *Senna alata* Leaves possess antimalarial effects and reverses haematological and biochemical perturbation in plasmodium berghei infected mice. *J Evidence-Based Integr Med*. 2022;27:2515690X2211164.
81. Daskum A. M., Godly C., Qadeer M. A., Ling L. Y. Effect of *Senna occidentalis* (Fabaceae) leaves extract on the formation of  $\beta$ -hematin and evaluation of in vitro antimalarial activity. *International Journal of Herbal Medicine*. 2019;7(3):46–51.
82. Aliyu U. M., Ramgopal D., Pankaj K.T., Umesh G., Umar Aliyu A., S. Saminu M., Khalipha A. J., Fatima I. S. In-silico evaluation of *Cassia occidentalis* phytochemicals for *Plasmodium falciparum* plasmepsin V inhibition: Revealing antimalarial potential, *South African Journal of Botany*. 2025;177:527-541.
83. Bhatt V., Kumari S., Upadhyay P., Agrawal P. A., Sahal D., Sharma U. Chemical profiling and quantification of potential active constituents responsible for the antiplasmodial activity of *Cissampelos pareira*. *J Ethnopharmacol*. 2020;15(262):113185.
84. Hatcher H., Planalp R., Cho J., Torti F.M., Torti S.V. Curcumin: From ancient medicine to current clinical trials. *Cell. Mol. Life Sci*. 2008;65:1631–1652.
85. Raju C. R., Palakkodu G. V., Venkateshwar G. K., Govindarajan P., Pundi N. Rangarajan, Curcumin for malaria therapy, *Biochemical and Biophysical Research Communications*. 2005; 326(2):472-474.
86. Lin L. Z., Shieh H. L., Angerhofer C. K., Pezzuto J. M., Cordell G. A., Xue L., Johnson M. E., Ruangrungsi N. Cytotoxic and antimalarial bisbenzylisoquinoline alkaloids from *Cyclea barbata*, *Journal of Natural Products*. 1993;56(1):22–29.
87. Thebtaranonth C., Thebtaranonth Y., Wanauppathamkul T.S., Yuthavong Y. Antimalarial Sesquiterpenes from Tubers of *Cyperus rotundus*: Structure of 10, 12-Peroxy calamene, A Sesquiterpene Endoperoxide, *Phytochemistry*. 1995;40(1):125-128.
88. Weenen H., Nkonya M.H., Bray D.H., Mwasumbi L.B., Kinabo L.S., Kilimali V.A. Antimalarial activity of Tanzanian medicinal plants. *Planta Med*. 1990;56(4):368-70.



89. Kamala A., Middha S.K., Karigar C.S. Plants in traditional medicine with special reference to *Cyperus rotundus* L.: A review of Biotech. 2018;8(7):309.
90. Berkov S., Zayed R. Comparison of Trophane alkaloid spectra between *Datura innoxia* grown in Egypt and Bulgaria. Zeitschrift for Nature Ferschung. 2004;89:184-186.
91. Gopieshkhanna V.K., Kannabiran K. Larvicidal effect of *Hemidesmus indicus*, *Gymnema Sylvestre*, and *Eclipta prostrate* against *Culex quinquefastiatus* mosquito larvae. African Journal of Biotechnology. 2007;6(3):307-311.
92. Kantamreddi V., Wright C. Investigation of Indian *Diospyros* Species for Antiplasmodial Properties. Evidence-based complementary and alternative medicine : eCAM. 2008;5:187-90.
93. Bapna S., Adsule S., Jadhav S., Patil L.S., Deshmukh R.A. Anti-malarial activity of *Eclipta alba* against *Plasmodium berghei* infection in mice, J. Commun. Dis. 2007;39(2):91-94.
94. Jahan R., Al-Nahain A., Majumder S., Rahmatullah M. Ethnopharmacological Significance of *Eclipta alba* (L.) Hassk. (Asteraceae), Int. Sch. Res. Not. 2014;1-22.
95. Govindarajan M. Evaluation of indigenous plant extracts against the malarial vector, *Anopheles stephensi* (Liston) (Diptera: Culicidae) Parasitology Research. 2011;109(1):93–103.
96. Baah K., Ameyaw A. A., Isaac K. A., Reinhard I. N. Antiplasmodial Activity and Phytochemical Evaluation of the Stems of *Albizia Coriaria* and *Ficus Sur*. European Journal of Medicinal Plants. 2024;35(2):1-10.
97. Verma G., Dua V.K., Agarwal D.D., Atul P.K. Anti-malarial activity of *Holarrhena antidysenterica* and *Viola canescens*, plants traditionally used against malaria in the Garhwal region of north-west Himalaya. Malar. J. 2011;10:20.
98. Simonsen H.T., Nordskjold J.B., Smitt U.W., Nyman U., Palpu P., Joshi P., Varughese G. In vitro screening of Indian medicinal plants for anti-plasmodial activity, J. Pharmacol. 2001;74:195–204.
99. Baba G., Adewumi A.A.J., Aina V.O. Phytochemical Characterization and in-vivo Anti- Malaria Activity of *Lantana camara* Leaf Extract. Br. J. Pharmacol. Toxicol. 2011;2:277–282.
100. Fallatah O., Georges E. Apigenin-induced ABCC1-mediated efflux of glutathione from mature erythrocytes inhibits the proliferation of *Plasmodium falciparum* Int J Antimicrob Agents. 2017;50(5):673-77.
101. Somsak V., Damkaew A., Onrak P. Antimalarial activity of kaempferol and its combination with chloroquine in *Plasmodium berghei* infection in mice. J Pathog. 2018; 2:39:12090.
102. Bhatt D., Kumar S., Kumar P., Bisht S., Kumar A., Maurya A.K., Pal A., Bawankule D.U. *Rutin ameliorates* malaria pathogenesis by modulating inflammatory mechanism: an *in vitro* and *in vivo* study. Inflammo pharmacology. 2022;30(1):159-71.
103. Banya H.S. Sharma R., Devi N. Antimalarial effect of *Ocimum sanctum* Linn. and *Bauhinia variegata* Linn. on *Plasmodium berghei*, IOSR Journal of Pharmacy and Biological Sciences. 2015;10:70-72.
104. Baliga M.S., Jimmy R., Thilakchand K.R., Sunitha V., Bhat N.R., Saldanha E., Rao S., Rao P., Arora R., Palatty P.L. *Ocimum sanctum* L (Holy Basil or Tulsi) and its phytochemicals in the prevention and treatment of cancer. Nutr Cancer. 2013;65(1):26-35.
105. Idrisi M.S., Badola H.K., Singh R. Indigenous knowledge and medicinal use of plants by local communities in Rangit Valley, South Sikkim, India. NeBIO. 2010;1(2):34-45.
106. Stuppner H., Wagner H. New cucurbitacin glycosides from *Picrorhiza kurrooa*. Planta Med. 1989;55(6):559-63.
107. Pradeepa V., Sathish-Narayanan S., Kirubakaran S.A., Senthil-Nathan S. Antimalarial efficacy of dynamic compound of plumbagin chemical constituent from *Plumbago zeylanica* Linn (Plumbaginaceae) against the malarial vector *Anopheles stephensi* Liston (Diptera: Culicidae). Parasitol Res. 2014;113(8):3105-9.
108. Satish P.V.V., Sunita K. Antimalarial efficacy of *Pongamia pinnata* (L) Pierre against *Plasmodium falciparum* (3D7 strain) and *Plasmodium berghei* (ANKA). BMC Complement Altern Med. 2017;17(1):458. Retraction in: BMC Complement Med Ther. 2021;21(1):139.
109. Chiamah O.C., Atieno D., Karani L., Chepng'etich J., Osano M., Gachie B., Kipkoech G., Jepkorir M., Ndungu J.W., Kuria J., Kimani F., Njeru S.N., Gathirwa J.W. Evaluation of the antimalarial properties of *Solanum incanum* L. leaf extract fractions and its ability to down regulate delta aminolevulinic acid dehydratase to prevent the establishment of malaria infection. J Ethnopharmacol. 2024;323:117613.
110. Melese Birru E., Adela Alemu M., Tsegaw Weredekal A., Belay Asrie A., Geta Mengistie M. Antimalarial Activity of Crude Extract and Solvent Fractions of Leaves of *Solanum nigrum* L.

(*Solanaceae*) against *Plasmodium berghei* in Mice. Evid Based Complement Alternat Med. 2022;34:261-75.

111. Christensen S.B., Kharazmi A., Tringali C. Antimalarial natural products. Isolation, characterization and biological properties Bioactive compounds from natural sources: Isolation, characterization and biological properties, London Taylor and Francis. 2001;379–432.
112. Lal Swagat Shrestha, R., Marasini, B.P., Adhikari Subin J. Phytochemicals of *Swertia chirayita* Roxb. ex Fleming against malarial dihydroorotate dehydrogenase: an in silico study. Discov Mol. 2024;1:3.
113. Yelaware Puttaswamy Naveen, Gunashekar Divya Rupini, Faiyaz Ahmed, Asna Urooj, Pharmacological effects and active phytoconstituents of *Swietenia mahagoni*: a review, Journal of Integrative Medicine. 2014;(12):2:86-93.
114. Soediro I., Padmawinata K., Wattimena J.R., Rekita S. Study of the active antimalarial methanolic extract of *Swietenia macrophylla* king (meliaceae) Acta Pharm. Indones. 1990;15:1–13.

UNDER PEER REVIEW