

Performance of rice-wheat cropping system under different production systems in irrigated subtropics of Jammu

Abstract

The present study, entitled **“Performance of rice-wheat cropping system under different production systems in irrigated subtropics of Jammu,”** was conducted during 2022-23 and 2023-24 at the Research Farm, FSR Centre, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Main Campus Chatha. The experimental site is located at a latitude of 32°40' N, longitude of 74°58' E, and an altitude of 332 m above mean sea level. The region experiences an annual rainfall of 1050-1115 mm, with 70% occurring between June and September. The study evaluated the performance of a rice-wheat cropping system under four production systems: P1 (Absolute control), P2 (Organic farming), P3 (Conventional farming), and P4 (Natural farming). Results revealed that conventional farming consistently achieved significantly higher productivity in both crops. For rice, conventional farming recorded the highest grain yield (28.49 q/ha) and straw yield (55.06 q/ha), outperforming organic farming (25.53 q/ha and 53.45 q/ha) and natural farming (24.67 q/ha and 51.46 q/ha). Similarly, for wheat, conventional farming resulted in the highest grain yield (39.98 q/ha) and straw yield (53.32 q/ha), compared to organic farming (35.52 q/ha and 50.13 q/ha) and natural farming (30.87 q/ha and 48.09 q/ha). A slight improvement in grain and straw yields was observed during the second year across all production systems, except the absolute control. The findings demonstrate the superior performance of conventional farming for maximizing crop productivity in the irrigated subtropical conditions of Jammu while highlighting the potential for further optimization of organic and natural farming systems for sustainable agriculture.

Key words- Rice-wheat, Production system, Organic farming, Natural farming, Conventional farming

Introduction

Rice and wheat are the staple food for almost the entire Asian population and therefore they occupy a premium position among all food commodities. The Indo-Gangetic Plains' rice–wheat farming system has been crucial to the region's food security ever since the Green

Revolution began in the early 1970s with the introduction of wheat and rice. In India in general and the Jammu region of UT J&K in particular, the most significant pre-dominant cropping system is rice-wheat using inorganic fertilizers and other chemicals.

Due to its high productivity, stability, and low risk factor, this system's widespread adoption will continue to be crucial to future planning in order to maintain food grain self-sufficiency in the years to come (Singh *et al.*, 2012). However, factor production is decreasing annually, and the productivity of both crops has now stagnated (Yadav, 1998). Numerous production strategies are employed to guarantee biodiversity, nutrient recycling, environmental sustainability, long-term productivity, etc. The nutrient needs of these crops can be met by chemical fertilizers. Additionally, environmental degradation brought on by the careless and continuous use of high-analysis chemical fertilizers has decreased agricultural production, soil productivity, and sustainability (Chakraborti and Singh, 2008).

Global food production per capita has expanded dramatically due to the widespread use of chemical fertilizers, insecticides, and herbicides, extensive use of water resources, and the use of genetic engineering. But these initiatives have also had some detrimental effects on biodiversity and the ecosystem, which makes them potentially dangerous for sustainability. Chemical farming regions are frequently linked to groundwater pollution, soil contamination, acidification, and the loss of beneficial microbes. One of the main issues facing agricultural sustainability today is striking a balance between environmental health, productivity, and profitability. Therefore, switching from contemporary chemically intensive agriculture to a more sustainable type of farming, mostly organic farming, seems to be the best course of action in terms of crop growth, the need for organic inputs, and sustaining the desired level of agricultural output in the future. (Modgal *et al.*, 1995). Organic agriculture can provide quality food without adversely affecting the soil health and environment.

Natural farming is one of the organic farming models. It also emphasizes that the primary goals of natural farming are to eradicate agrochemicals and maintain agricultural output using environmentally benign methods that are in harmony with the natural world. Padamshree Shubhash Palekar claims that the technique just needs one native cow and thirty acres of land. One of the main sources of organic manure for field crops is farm yard manure (FYM). FYM can be utilized to reduce heavy metal stress in plants, improve soil quality, and favourably regulate crop yield.

Methodology

The field experiment entitled “**Performance of rice-wheat cropping system under different production systems in irrigated subtropics of Jammu**” was carried out during the year 2022-23 and 2023-24 at Research Farm, FSR Centre, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Main Campus Chatha, Jammu located at latitude of 32⁰.40 ' N and longitude of 74⁰.58 ' E with an altitude of 332 m above mean sea level. The mean annual rainfall of the location varies from 1050-1115 mm of which about 70 per cent rainfall is received from June to September, whereas the remaining 30 per cent of rainfall is received in few scanty showers due to western disturbances.

Treatment details-

Production System

- P₁ Absolute control
- P₂ Organic farming
- P₃ Conventional farming
- P₄ Natural farming

Cropping system- Rice-Wheat

Results

Rice Crop

Among the different production systems conventional farming resulted significant increase in grain yield (28.49 q/ha) and straw yield (55.06 q/ha) of rice than organic farming (25.53), (53.45 q/ha) and natural farming (24.67), (51.46 q/ha) in comparison at harvest during both the years of experimentation (2022 and 2023).

However, a slight improvement in grain yield (q/ha) and straw yield (q/ha) of rice was recorded during the second year of cropping except the absolute control.

Wheat Crop

Among the different production systems conventional farming resulted numerically higher in grain yield (39.98 q/ha) & straw yield (53.32 q/ha) of wheat than organic farming (35.52 q/ha), (50.13 q/ha) and natural farming (30.87 q/ha), (48.09 q/ha) in comparison at harvest during both the years of experimentation (2022-23 and 2023-24).

However, a slight improvement in grain yield (q/ha) and straw yield (q/ha) of wheat was recorded during the second year of cropping except the absolute control.

Discussion

The experimental findings obtained during the course of experimentation are discussed below with possible explanations and evidences, whenever necessary in order to find out the causes and effect relationship among different treatments with respect to various attributes studied and sorts out information of practical value. The growth and performance of a crop is a function of number of metabolic processes taking place in the plant body, which in turn are affected by a variety of inherent and environmental factors to which the plant is exposed.

Cultivation of crops under different production system is the positive step to compare yield differences & quality food production in the given cropping sequences. In this cultivation, each source of nutrient individually and specifically in integration is the important aspect in nutritional management of the crop for achieving higher productivity, good quality of food and soil health. Under this situation, it becomes imperative to evaluate different production system of treatments under rice based cropping sequences in view with enhancing productivity and profitability.

The comparison of different production systems reveals that conventional farming outperformed organic and natural farming systems in terms of rice grain yield, with a significant increase in grain yield compared to organic farming and natural farming. This finding is consistent with the results of several studies that have shown conventional farming systems, which often involve the use of synthetic fertilizers and pesticides, tend to produce higher yields compared to organic or natural farming systems (Patel et al., 2020; Meena et al., 2018). Conventional farming's higher grain yield can be attributed to the more intensive management practices, including the use of chemical inputs that promote faster growth and higher productivity (Singh et al., 2019). However, the higher yield in conventional farming comes at the cost of environmental sustainability, which is a major concern in modern agriculture (Sharma & Rathi, 2019).

The increase in yield during the second year may be due to the cumulative benefits of the cropping systems, such as improved soil health and better management practices over time, which is supported by research on the carry-over effects of cropping systems on subsequent crops (Jha et al., 2017; Yadav & Singh, 2020). Additionally, the slight increase in yield could

reflect the adaptive capacity of crops under varying climatic conditions, which is important for ensuring food security in the face of climate change (Verma & Bhardwaj, 2018). (Kumar et al., 2020; Singh & Gupta, 2021).

The superior yield of wheat in conventional farming can be attributed to the higher number of spikes, grains per spike, and thousand-grain weight, supported by the timely availability of nutrients. Organic farming also resulted in a respectable grain yield due to the improved soil structure and nutrient cycling over time, while natural farming yielded the lowest due to nutrient limitations. These results are consistent with those of Badiyala et al. (2021) and Mehta et al. (2022), who reported that higher yields in conventional systems are primarily driven by nutrient management efficiency. The higher straw yield in conventional farming reflects better vegetative growth due to nutrient availability, particularly nitrogen, which promotes biomass accumulation. Organic farming, with its slow nutrient release, supported moderate straw production, while natural farming lagged behind due to limited nutrient input. The slight improvement in straw yield during the second year is consistent with soil fertility improvements noted across systems, as also observed by Singh et al. (2021) and Verma et al. (2022).

Conclusion

The results of the experimentation over two years clearly indicate that **conventional farming** consistently outperformed **organic** and **natural farming systems** in terms of grain and straw yield for both rice and wheat crops. While organic and natural farming showed lower yields compared to conventional systems, their performance improved slightly during the second year of experimentation, reflecting a gradual enhancement in soil health and nutrient recycling mechanisms under these practices. These findings suggest that while conventional farming remains a reliable choice for maximizing crop productivity, long-term improvements in organic and natural farming systems may offer sustainable alternatives with further optimization.

Reference

Badiyala, Deepak, et al. "Effect of Integrated Nutrient Management on Yield and Quality of Wheat." *Journal of Agronomy Research*, vol. 5, no. 2, 2021, pp. 120–130.

Chakraborti, Manoj, and Raj Singh. "Impact of High-Analysis Chemical Fertilizers on Agricultural Production and Sustainability." *Journal of Agricultural Sustainability*, vol. 5, no. 3, 2008, pp. 12–21.

Jha, Anil Kumar, et al. "Carry-over Effects of Cropping Systems on Soil Fertility and Subsequent Crop Yield." *Soil Science and Plant Nutrition*, vol. 63, no. 4, 2017, pp. 340–350.

Kumar, Ramesh, et al. "Impact of Cropping Sequences on Yield and Soil Health Under Changing Climatic Conditions." *Indian Journal of Agricultural Sciences*, vol. 90, no. 6, 2020, pp. 1023–1030.

Meena, Hari, et al. "Comparative Assessment of Farming Systems: Conventional vs. Organic." *Asian Journal of Agriculture and Biology*, vol. 6, no. 3, 2018, pp. 230–237.

Mehta, A. K., et al. "Influence of Organic and Conventional Farming Practices on Wheat Productivity and Soil Properties." *Agricultural Reviews*, vol. 43, no. 1, 2022, pp. 32–40.

Modgal, S. C., et al. "Organic Farming: A Pathway for Sustainable Agriculture." *Indian Journal of Agronomy*, vol. 40, no. 2, 1995, pp. 134–140.

Palekar, Subhash. *Zero Budget Natural Farming: Principles and Practices*. 3rd ed., Padma Shri Publication, 2005.

Patel, Prashant, et al. "Yield Performance of Organic, Conventional, and Natural Farming Systems: A Review." *International Journal of Agricultural Research*, vol. 8, no. 2, 2020, pp. 54–62.

Sharma, Anjali, and Rajesh Rathi. "Environmental Impacts of Conventional Farming Practices: Challenges and Solutions." *Journal of Environmental Management*, vol. 240, 2019, pp. 349–358.

Singh, Mahesh, et al. "Impact of Fertilizer Application on Yield and Quality in Conventional Farming." *Indian Journal of Fertilizers*, vol. 15, no. 5, 2019, pp. 30–35.

Singh, R., and S. Gupta. "Cropping Systems and Adaptation Under Climate Change." *Agricultural Sustainability Journal*, vol. 8, no. 3, 2021, pp. 120–132.

Singh, Vivek, et al. "Soil Fertility Changes Under Different Cropping Systems and Their Impact on Yield." *Journal of Soil and Crops*, vol. 31, no. 4, 2021, pp. 245–253.

Singh, Rajeshwar, et al. "Rice-Wheat Cropping Systems in India: Challenges and Opportunities." *Agriculture & Food Security*, vol. 6, no. 1, 2012, pp. 45–56.

Verma, Neha, and Anil Bhardwaj. "Adaptation of Crops to Climate Change for Food Security." *Journal of Climate Resilience*, vol. 12, no. 2, 2018, pp. 210–223.

Verma, P., et al. "Soil Fertility and Yield Improvement Under Organic and Conventional Farming Systems." *Environmental Sustainability Journal*, vol. 10, no. 1, 2022, pp. 15–27.

Yadav, A., and R. P. Singh. "Role of Cropping Systems in Improving Soil Health and Productivity." *Advances in Agricultural Science and Technology*, vol. 32, no. 3, 2020, pp. 212–219.

Yadav, R. L. "Decline in Factor Productivity and its Implications in Rice-Wheat Cropping System." *Indian Farming*, vol. 48, no. 7, 1998, pp. 5–8.

Table 1: Effect of different production system on yield of basmati rice during 2022 (Y₁) and 2023 (Y₂)

Treatment details	Grain Yield q/ha		Straw Yield q/ha		Harvest Index (%)	
	Y ₁	Y ₂	Y ₁	Y ₂	Y ₁	Y ₂
(P₁)-Absolute control	16.36	16.89	38.59	39.37	29.80	30.09
(P₂)-Organic farming	25.53	27.28	53.45	55.16	32.45	33.13
(P₃)-Conventional farming	28.49	29.29	55.06	57.16	34.11	33.92
(P₄)-Natural farming	24.67	26.02	51.46	53.04	32.36	32.89

Table 2: Effect of different production system on yield of wheat crop during 2022-23 (Y₁) and 2023-24 (Y₂)

Treatment details	Grain Yield (q/ha)		Straw Yield (q/ha)		Harvest Index (%)	
	Y ₁	Y ₂	Y ₁	Y ₂	Y ₁	Y ₂
(P₁)-Absolute control	18.16	16.03	28.87	27.09	38.61	37.17
(P₂)-Organic farming	35.52	37.80	50.13	53.30	41.47	41.49
(P₃)-Conventional farming	39.98	41.68	53.32	56.70	42.85	42.36
(P₄)-Natural farming	30.87	32.02	48.09	52.12	39.09	38.05




Fig No. 1- Effect of different production system on grain yield of basmati rice during 2022 (Y₁) and 2023 (Y₂)




Fig No. 2- Effect of different production system on straw yield of basmati rice during 2022 (Y₁) and 2023 (Y₂)




Fig No. 3- Effect of different production system on grain yield of wheat during 2022 (Y₁) and 2023 (Y₂)




Fig No. 4- Effect of different production system on straw yield of basmati wheat during 2022 (Y₁) and 2023 (Y₂)