

## **Original Research Article**

# **Impact of Nitrogen Dosage through various sources on Growth of Dragon Fruit in mid-hill condition of Nagaland**

### **ABSTRACT**

**Aims:** To evaluate optimum dose of nitrogen through various sources for growth and development of dragon fruit grown under mid-hill condition of Nagaland

**Study design:** Randomized Block Design

**Place and Duration of Study:** The present experiment was carried out during the year of 2021 and 2022 has been conducted in the research experimental block of Department of Horticulture, School of Agricultural Sciences, Medziphema Campus, Nagaland.

**Methodology:** Different levels of nitrogen with and without organic manures were applied to the one-year-old plants to evaluate the optimum dose for growth and development of dragon fruit plants. Ten treatments of different doses of nitrogen and combinations of organic manures (Farm Yard Manure (FYM), Pig manure), common doses of phosphorous and potassium were applied to plants. [ $N_0$  Control;  $N_{100g}$ /plant;  $N_{125g}$ /plant;  $N_{150g}$ /plant; 75% of  $N_{100g}$ +FYM<sub>1.0 kg</sub>/plant; 75% of  $N_{125g}$ +FYM<sub>1.25 kg</sub>/plant; 75% of  $N_{150g}$ +FYM<sub>1.5kg</sub>/plant; 75% of  $N_{100g}$ +Pig manure<sub>1.66kg</sub>/plant; 75% of  $N_{125g}$ +Pig manure<sub>2.1kg</sub>/plant; 75% of  $N_{150g}$ +Pig manure<sub>2.5kg</sub>/plant.]

**Results:** The data of two-year study revealed that the maximum cladode length (232.04 cm) and number of areoles/cladode (79.32) were recorded in the treatment with 75% of  $N_{150g}$ +Pig manure<sub>2.5kg</sub>/plant ( $T_{10}$ ) whereas maximum cladode diameter (5.02 cm), cladode girth (15.78 cm), distance between areoles (3.30 cm) and area of areoles (0.18 mm<sup>2</sup>) were recorded in the treatment 75% of  $N_{125g}$ +Pig manure<sub>2.1kg</sub>/plant ( $T_9$ ) and minimum values were recorded mostly in treatments supplied with only inorganic manures ( $T_1$  and  $T_2$ )

**Conclusion:** The combined application of both organic and inorganic manures ( $T_9$  and  $T_{10}$ ) found to be most superior treatment combination with regard to vegetative growth in *Hylocereus polyrhizus*.

### **1. INTRODUCTION**

Dragon fruit (*Hylocereus* spp.), a climbing cactus vine native to tropical regions of Mexico, Central and South America (Mizrahi *et al.*, 2010) is an emerging super crop among farmers for its economic value and rich in nutrient content. It is commonly known as kamalam, strawberry pear, pithaya, night-blooming cereus, Belle of the night, Jesus in the cradle etc. It is a fast-growing, herbaceous, perennial, epiphytic, cacti crop which has been introduced into India in late 90's and is considered as a promising remunerative fruit crop and future crop of India (Arivalagan *et al.*, 2019). With its high nutritional value, resilience against pests and diseases, minimal orchard maintenance, low water needs, frequent yields throughout the year and the potential for high output lasting up to 20 years, this crop has enticed numerous growers to cultivate it. This surge in cultivation has also amplified the fruit's export potential, catering to the growing market demand. The growing importance of dragon fruit cultivation in India, major emphasis should be given to standardize the fertilizer doses in different agro-climatic conditions in order to help farmers take up its cultivation on commercial scale and gain economic rewards as it is a nutrient loving plant.

Plant nutrition is one of the most important factors that plays crucial role in growth, yield and quality attributes in dragon fruit. Indiscriminate and prolonged use of chemical fertilizers without adding organic fertilizers deteriorated the health of soil and

increase soil pollution by decreasing the microbial activity (Singh and Kallo, 2000). This calls for a holistic approach to maintain soil health as well as obtain high productivity in the crop grown. A new farming strategy therefore entails a plant nutritional package which would provide all elements through both organic, inorganic and biofertilizers which would not only reduce the soil pollution but also produces quality produce with greater production and productivity and also keep the production cost at bearable level to the average farmer. Several reports show that the integration of organic and inorganic fertilizers proved superior to the sole application (Abusaleha and Shanmugavelu, 1988). Use of biofertilizers along with organic and inorganic fertilizers created a lot of impact in horticulture. Most soils in the North East region of India are acidic in nature where phosphorous content in soil is high but is not readily available to the plant. Organic acids secreted during organic matter decomposition and phosphobacteria makes the insoluble form of phosphorous to available form and saves up to 30-50 kg of superphosphate (Chen *et al.*, 2006). Nitrogen occupies a conspicuous place in plant metabolism system and it plays a key role in agriculture by increasing crop yield. It is a part of protein, important constituent of protoplasm, enzymes, the biological catalytic agents which speed up life processes. Nitrogen is also present as a part of nucleoprotein, amino acids, amines, amino sugar, polypeptides and other organic compounds in plants.

North Eastern India is considered as a hub of organic products because there is minimal or no use of chemical fertilizers by most farmers. The fertile soil and congenial climatic conditions in the region contribute as positive factor in the growing of several exotic crops including dragon fruit. Although dragon fruit belongs to cactaceae family, it requires water during critical periods of its growth because unlike other cacti it is from tropical rainforest. According to the studies conducted on dragon fruit for adaptability and production aspects by Karunakaran *et al.* (2014) it was reported that dragon fruit prefers dry tropical climate with optimum temperature of 20-29°C and can withstand temperatures up to 38-40°C and as low as 0°C for shorter periods. Heavy rainfall during flowering season may lead to flower and fruit drop and high temperatures exceeding 40°C may lead to flower burn. These conditions make the north eastern regions even more suitable for dragon fruit cultivation except during high rainfall season which may negatively impact flowering and fruit setting. Considering the importance of this emerging crop with high potential in commercialization it was a felt need to conduct the present study which would augment dragon fruit crop cultivation and add fillip to the economy of average farmers.

## 2. MATERIAL AND METHODS

The present investigation was carried out during the year of 2021 and 2022 at research experimental block of Department of Horticulture, School of Agricultural Sciences, Medziphema Campus, Nagaland situated at 25° 45' 53" N latitude and 93° 53' 04" E longitudes at an elevation of 310 m above sea level. The established field of dragon fruit with three-year-old plants were used for the phenological studies in the present experiment. The nutrients *i.e.*, nitrogen (N), phosphorus (P) and potassium (K) were applied in the form of urea, single super phosphate and muriate of potassium, respectively. The different doses of organic fertilizers (Farm yard manure and Pig manure) were applied in the month of March followed by inorganic fertilizers in the month of May. Nitrogen was applied in split doses. Doses of FYM and Pig manure were calculated on basis of N content in the recommended dose of fertilizers (RDF) (N: P: K) - 135:78:63g/plant. Common doses of P<sub>50g</sub> and K<sub>75g</sub>/plant was applied to every plant in the fields. PSB @ 30 g/pillar was supplied in equal doses to all treatments. The experiment was laid out in randomized complete block design with three replications.

Various treatments consisting of organic (FYM, Pig manure) and inorganic manures (NPK) were applied. Treatment details: N<sub>0</sub> Control; N<sub>100g</sub>/plant; N<sub>125g</sub>/plant; N<sub>150g</sub>/plant; 75% of N<sub>100g</sub>+FYM<sub>1.0 kg</sub>/plant; 75% of N<sub>125g</sub>+FYM<sub>1.25 kg</sub>/plant; 75% of N<sub>150g</sub>+FYM<sub>1.5kg</sub>/plant; 75% of N<sub>100g</sub>+Pig manure<sub>1.66kg</sub>/plant; 75% of N<sub>125g</sub>+Pig manure<sub>2.1kg</sub>/plant; 75% of N<sub>150g</sub>+Pig manure<sub>2.5kg</sub>/plant. Growth parameters like cladode length, diameter, girth,

number of areoles, spines, distance between areoles, area of areoles, fresh weight and dry weight of cladode were observed. The data collected and computed was completed by using SPSS software (Gomez and Gomez, 2010).

### 3.RESULTS AND DISCUSSION

Marked variations were observed with respect to morphological characteristics among the different treatments by the application of nitrogen through various sources. The data recorded revealed that the treatment 75% of N<sub>150g</sub>+Pig manure<sub>2.5 kg</sub>/plant (T<sub>10</sub>) recorded the maximum cladode length (232.04 cm) and was found highly significant whereas the minimum was recorded in N<sub>0</sub> Control (T<sub>1</sub>) (82.16 cm) (Table 1). According to Verma *et al.* (2019) maximum plant height (129.30 cm) in dragon fruit was recorded in the treatment supplied with FYM+NPK+ Azotobactor + PSB whereas in the present investigation pig manure recorded significantly better results compared to the above treatment. This might be due to the high nitrogen content in the pig manure compared to FYM and vermicompost. The combination of inorganic and organic fertilizers might have improved the soil health thus resulting in creating favourable nutrient environment for plant growth (Lodhi *et al.*, 2017).

**Table 1. Effect of various nitrogen sources on physical cladode characteristics of dragon fruit**

Treatments	Cladode length (cm)	Cladode diameter (cm)	Cladode girth (cm)	No. of spines/ areoles
N <sub>0</sub> Control	82.16	3.86	12.15	4.00
N <sub>100g</sub> / plant	105.33	3.80	12.12	4.00
N <sub>125g</sub> /plant	106.06	4.00	12.59	3.50
N <sub>150g</sub> /plant	117.46	4.35	13.66	4.16
75% of N <sub>100g</sub> + FYM <sub>1kg</sub> /plant	130.20	4.44	13.97	3.66
75% of N <sub>125g</sub> +FYM <sub>1.25 kg</sub> /plant	135.27	4.51	14.17	3.50
75% of N <sub>150g</sub> + FYM <sub>1.5 kg</sub> /plant	154.04	4.59	14.43	3.50
75% of N <sub>100g</sub> + Pig manure <sub>1.66 kg</sub> /plant	178.69	4.87	15.32	3.50
75% of N <sub>125g</sub> + Pig manure <sub>2.1 kg</sub> /plant	215.65	5.02	15.78	3.50
75% of N <sub>150g</sub> + Pig manure <sub>2.5 kg</sub> /plant	232.04	4.87	15.33	3.16
S Em±	2.44	0.05	0.50	0.23
CD at 5%	7.32	0.15	1.51	NS

Olusegun (2014) supported these findings stating that pig manure in combination with NPK significantly increased the plant height.

The maximum cladode diameter (5.02 cm) and minimum in treatment N<sub>100g</sub>/ plant (T<sub>2</sub>) (3.80 cm). Ringphawon (2018) reported that 25% pig manure gave best results regarding the diameter of dragon fruit compared to mineral fertilizers. This might be because of availability of more nutrients due to the addition of organic and inorganic fertilizers (Lodhi *et al.*, 2017).

**Table 2. Effect of various nitrogen sources on areole characteristics of dragon fruit**

Treatments	No. of areoles/cladode (cm)	Distance between areoles (cm)	Area of areoles (mm <sup>2</sup> )
N <sub>0</sub> Control	53.32	1.71	0.07
N <sub>100g</sub> /plant	66.49	1.80	0.07
N <sub>125g</sub> /plant	41.45	2.83	0.08
N <sub>150g</sub> /plant	43.34	3.11	0.09
75% of N <sub>100g</sub> + FYM <sub>1kg</sub> /plant	49.13	3.14	0.12
75% of N <sub>125g</sub> +FYM <sub>1.25 kg</sub> /plant	62.14	2.26	0.13
75% of N <sub>150g</sub> + FYM <sub>1.5 kg</sub> /plant	72.46	2.28	0.13
75% of N <sub>100g</sub> + Pig manure <sub>1.66 kg</sub> /plant	68.02	3.41	0.14
75% of N <sub>125g</sub> + Pig manure <sub>2.1 kg</sub> /plant	67.00	3.53	0.18
75% of N <sub>150g</sub> + Pig manure <sub>2.5 kg</sub> /plant	79.32	3.26	0.15
S Em±	2.76	0.31	0.01
CD at 5%	8.27	0.94	0.03

Cladode girth (15.78 cm) was recorded maximum in 75% of N<sub>125g</sub>+Pig manure<sub>2.1 kg</sub>/plant (T<sub>9</sub>) and treatment 75% of N<sub>150g</sub>+Pig manure<sub>2.5 kg</sub>/plant (T<sub>10</sub>) was found statistically at par with it whereas minimum was recorded in the treatment N<sub>100g</sub>/plant (T<sub>2</sub>) (12.12 cm). The results are in agreement with Awosika *et al.* (2014) and Iren *et al.* (2016) where the combination of NPK and pig manure significantly increased the stem girth. This might be due to the increase in soil nutrients because of addition of organic manures, which helped in modifying the soil properties and boosted the soil health. The maximum number of spines (4.16) were recorded in treatment N<sub>150g</sub>/plant (T<sub>4</sub>) and minimum in 75% of N<sub>150g</sub>+Pig manure<sub>2.5 kg</sub>/plant (T<sub>10</sub>) (3.16). Rawat *et al.* (2022) reported similar results, where number of spines were found statistically non-significant with the different nutrient sources.

The distance between areoles (0.53 cm) and area of areoles (0.18 mm<sup>2</sup>) (Table 2) were recorded highest in treatment supplied with 75% of N<sub>125g</sub>+Pig manure<sub>2.1 kg</sub>/plant (T<sub>9</sub>) and lowest in N<sub>0</sub> Control (T<sub>1</sub>) (1.17 cm and 0.07 mm<sup>2</sup>). The results showed that the dragon fruit responded well to the application of pig manure when combined with chemical fertilizers. Abo Sedera *et al.* (2009) stated that nitrogen fertilizer when used along with compost had enhanced the vegetative growth. This might also be due to the availability of chemical fertilizers in the initial stages and organic manures in latter stages which might have helped in improving the soil organic matter, total nitrogen and soil carbon (Kumar *et al.*, 2016 and Reza and Jafar, 2007). The least supply of N, P and K in control might have resulted in limited plant growth and development altering photosynthesis and carbohydrate production (Zekri and Obreza, 2003).

#### 4.CONCLUSION

From the above experiment it can be stated that pig manure with highest amount of nitrogen along with inorganic manures resulted in rapid growth of cladode length, diameter, girth, distance between areoles, area of areoles stating that the combined application of pig manure with inorganic fertilizers (NPK) found to be most superior treatment combination with regard to vegetative growth in *H. polyrhizus*.

## REFERENCES

- Abo Sedera, F.A., Shafshak, S.N., Eid, M.S., & Mohamed, M.H.M. (2009). Improving productivity and quality of strawberry via organic fertilization and application of some natural growth stimulants. *Indian Journal of Horticulture*, 1:481-485.
- Abusaleha & Shanmugavelu, K.G. (1988). Studies on the effect of organic vs inorganic source of nitrogen on growth, yield and quality of okra (*Abelmoschus esculentus*). *Indian Journal of Horticulture*, 45 (4): 312-318.
- Arivalagan, M., Sriram, S., & Karunakaran, G. (2019). *Dragon fruit country report from India. FFTC Agricultural Policy Platform (FFTC-AP)*. 1-8.
- Awosika, O.E., Awodun, M.A., & Ojeniyi, S.O. (2014). Comparative effect of pig manure and NPK fertilizer on agronomic performance of tomato (*Lycopersicon esculentum mill*). *American Journal of Experimental Agriculture*, 4 (11): 1330-1338.
- Chen, Y. P., Rekha, P. D., Arun, A. B., Shen, F. T., Lai, W. A., & Young, C. (2006). Phosphate solubilizing bacteria from subtropical soil and their tricalcium phosphate solubilizing abilities. *Applied Soil Ecology*, 34: 33–41.
- Gomez, A. K. & Gomez, A. A. (2010). *Statistical procedures for agricultural research* (2<sup>nd</sup> ed.). Wiley India Private Limited. New Delhi. pp 134-138.
- Iren, O.B., Ijah, C.J., Asawalam, D.O., & Osodeke, V.E. (2016). Comparative effect of pig manure, urea fertilizer and their combinations on the performance of *Amaranthus cruentus* in a rainforest ultisol, Nigeria. *Journal of Agricultural Science and Practice*, 1: 52-57.
- Karunakaran, G., Tripathi, P.C., Sankar, V., Sakthivel, T. & Senthilkumar, R.( 2014). Dragon Fruit - A new introduction crop to India: A potential market with promising future. In: *Proceeding of National Seminar on Strategies for conservation, Improvement and utilization of underutilized fruits*, 138-139.
- Karunakaran, G., Tripathi, P.C., Sankar, V., Sakthivel, T., & Senthilkumar, R. (2014). Dragon Fruit - A new introduction crop to India: A potential market with promising future. In: *Proceeding of National Seminar on Strategies for conservation, Improvement and utilization of underutilized fruits*, 138-139.
- Kumar, N., Attar, S.K., & Patel, S.N. (2016). Effect of organic and inorganic sources of nitrogen on growth, yield and quality of mango. *Annals of Plant and Soil Research*, 18 (1): 29-32.
- Lodhi, P., Singh, D., & Tiwari, A. (2017). Effect of inorganic and organic fertilizers on yield and economics of Broccoli (*Brassica oleracea var. italica*). *International Journal of Current Microbiology and Applied Sciences*, 6: 562-566.
- Mizrahi, Y., Nerd, A., & Nobel, P. S. (2010). Cacti as a crops. *Horticultural Reviews*, 18: 291-320.
- Olusegun, O.S. (2014). Influence of NPK 15-15-15 fertilizer and pig manure on nutrient dynamics and production of cowpea, *Vigna unguiculata L.* Walp. *American Journal of Agriculture and Forestry*, 2(6): 267-273.

- Reza, M.T., & Jafar, K. (2007). Influence of organic and chemical fertilizers on growth and yield of tomato (*Lycopersicon esculentum* L.) and soil chemical properties. *Annual Report*, Tehran, Iran: Shahid Beheshti University.
- Ringphawon. H. (2018). Effect of various nutrient sources on the vegetative growth of dragon fruit. *M.Sc. Thesis*, Nagaland University, School of Agricultural Sciences and Rural Development, Medziphema Campus, India.
- Singh, K.P., & Kalloo, G. (2000). Nutrient management in vegetable crops. *Fertilizer News*, 45: 77-81.

UNDER PEER REVIEW