

On Certain Results On The Diophantine Equation:

$$\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3 \left(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2 \right)$$

**ORIGINAL
RESEARCH
ARTICLE**

Abstract

Consider a sequence w_r in arithmetic progression with a common difference d . The exploration of Diophantine equations, which are polynomial equations seeking integer solutions, has been a fascinating endeavor in number theory. These equations have historically intrigued mathematicians due to their inherent complexities and their importance in understanding the properties of integers. In this study, we investigate a Diophantine equation that relates the sum of squares of integers from specific sequences to a variable d . Specifically, we extend existing results on the Diophantine equation: $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3 \left(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2 \right)$. We aim to determine the conditions under which integer solutions for w_r and d exist within this equation. Our methodology involves decomposing and factoring polynomials and exploring the solution set of the given equation.

Keywords: Sequences, Diophantine equation, Integer, Polynomial, Factorization

1 Introduction

Diophantine equations, tracing their origins back to the era of the ancient Greek mathematician Diophantus, continue to pose captivating challenges within number theory. These equations, which seek integer solutions, hold significant importance due to their real-life applications. Despite the extensive exploration of various Diophantine equations, including renowned challenges like Fermat's Last Theorem, the Ramanujan-Nagell equation, and the Lebesgue-Nagell equation, as well as studies focusing on polynomials of degree less than five, specific examinations of the Diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3\left(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2\right)$ remain largely uncharted. Recent research has delved into the intricacies of polynomials with degrees less than five [5, 1, 3, 15, 9, 13]. For a comprehensive understanding of studies related to Fermat's Last Theorem and the Ramanujan-Nagell equations, readers are encouraged to explore [4, 7, 10, 12, 8, 14, 2, 11, 16]. Within the existing body of work, the literature concerning the Diophantine equation remains largely unexplored. This study aims to contribute to this knowledge gap by extending existing results on the Diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3\left(\frac{nd^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{3r-1}^2\right)$ first introduced by Lao [12, 14], thus seeking to enhance our comprehension of this specific Diophantine equation within the broader landscape of mathematical exploration.

2 Main Results:

Theorem 1.1: Consider equation 1 satisfying condition $(n, w_1, w_2, \dots, w_{15}, 9d) = (27, w_1, w_2, \dots, w_{27}, 9d)$

Then, the diophantine equation:

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + 9d^2 = 3(9d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2)$$

has the solution in integers if $w_{27} - w_{26} = w_{26} - w_{25} = w_{25} - w_{24} = w_{24} - w_{23} = w_{23} - w_{22} = w_{22} - w_{21} = w_{21} - w_{20} = w_{20} - w_{19} = w_{19} - w_{18} = w_{18} - w_{17} = w_{17} - w_{16} = w_{16} - w_{15} = w_{15} - w_{14} = w_{14} - w_{13} = w_{13} - w_{12} = w_{12} - w_{11} = w_{11} - w_{10} = w_{10} - w_9 = w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$

Proof: Consider the equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + 9d^2 = 3(9d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2)$$

And suppose that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d, w_{10} = w_1 + 9d, w_{11} = w_1 + 10d, w_{12} = w_1 + 11d, w_{13} = w_1 + 12d, w_{14} = w_1 + 13d, w_{15} = w_1 + 14d, w_{16} = w_1 + 15d, w_{17} = w_1 + 16d, w_{18} = w_1 + 17d, w_{19} = w_1 + 18d, w_{20} = w_1 + 19d, w_{21} = w_1 + 20d, w_{22} = w_1 + 21d, w_{23} = w_1 + 22d, w_{24} = w_1 + 23d, w_{25} = w_1 + 24d, w_{26} = w_1 + 25d, w_{27} = w_1 + 26d,$

The left hand side expressed as:

And suppose that $w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + (w_1 + 7d)^2 + (w_1 + 8d)^2 + (w_1 + 9d)^2 + (w_1 + 10d)^2 + (w_1 + 11d)^2 + (w_1 + 12d)^2 + (w_1 + 13d)^2 +$

$$(w_1 + 14d)^2 + (w_1 + 15d)^2 + (w_1 + 16d)^2 + (w_1 + 17d)^2 + (w_1 + 18d)^2 + (w_1 + 19d)^2 + (w_1 + 20d)^2 + (w_1 + 21d)^2 + (w_1 + 22d)^2 + (w_1 + 23d)^2 + (w_1 + 24d)^2 + (w_1 + 25d)^2 + (w_1 + 26d)^2 + 9d^2$$

Simplifies to

$$27w_1^2 + 702w_1d + 6210d^2 = 3(9w_1^2 + 234w_1d + 2070d^2) \dots(1.1)$$

Splitting equation (2.2) into thrice sums of squares we obtain:

$$3(9d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2) + (w_1^2 + 20w_1d + 100d^2) + (w_1^2 + 26w_1d + 169d^2) + (w_1^2 + 32w_1d + 256d^2) + (w_1^2 + 38w_1d + 361d^2) + (w_1^2 + 44w_1d + 484d^2) + (w_1^2 + 50w_1d + 625d^2))$$

$$= 3(9d^2) + (w_1 + d)^2 + (w_1 + 4d)^2 + (w_1 + 7d)^2 + (w_1 + 10d)^2 + (w_1 + 13d)^2 + (w_1 + 16d)^2 + (w_1 + 19d)^2 + (w_1 + 22d)^2 + (w_1 + 25d)^2$$

$$= 3(9d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2)$$

This completes the proof \square .

Theorem 1.2: Consider equation 1 satisfying condition $(n, w_1, w_2, \dots, w_{30}, 10d) = (30, w_1, w_2, \dots, w_{30}, 10d)$

Then, the diophantine equation:

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + \dots + w_{28}^2 + w_{29}^2 + w_{30}^2 + 10d^2 = 3(10d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2)$$

has the solution in integers if $w_{30} - w_{29} = w_{29} - w_{28} = w_{28} - w_{27} = w_{27} - w_{26} = w_{26} - w_{25} = w_{25} - w_{24} = w_{24} - w_{23} = w_{23} - w_{22} = w_{22} - w_{21} = w_{21} - w_{20} = w_{19} - w_{18} = w_{18} - w_{17} = w_{17} - w_{16} = w_{16} - w_{15} = w_{15} - w_{14} = w_{14} - w_{13} = w_{13} - w_{12} = w_{12} - w_{11} = w_{11} - w_{10} = w_{10} - w_9 = w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$

Proof: Consider the equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + \dots + w_{28}^2 + w_{29}^2 + w_{30}^2 + 10d^2 = 3(6d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2)$$

And suppose that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d, w_{10} = w_1 + 9d, w_{11} = w_1 + 10d, w_{12} = w_1 + 11d, w_{13} = w_1 + 12d, w_{14} = w_1 + 13d, w_{15} = w_1 + 14d, w_{16} = w_1 + 15d, w_{17} = w_1 + 16d, w_{18} = w_1 + 17d, w_{19} = w_1 + 18d, w_{20} = w_1 + 19d, w_{21} = w_1 + 20d, w_{22} = w_1 + 21d, w_{23} = w_1 + 22d, w_{24} = w_1 + 23d, w_{25} = w_1 + 24d, w_{26} = w_1 + 25d, w_{27} = w_1 + 26d, w_{28} = w_1 + 27d, w_{29} = w_1 + 28d, w_{30} = w_1 + 29d$

The left hand side expressed as:

$$\text{And suppose that } w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + (w_1 + 7d)^2 + (w_1 + 8d)^2 + (w_1 + 9d)^2 + (w_1 + 10d)^2 + (w_1 + 11d)^2 + (w_1 + 12d)^2 + (w_1 + 13d)^2 + (w_1 + 14d)^2 + (w_1 + 15d)^2 + (w_1 + 16d)^2 + (w_1 + 17d)^2 + (w_1 + 18d)^2 + (w_1 + 19d)^2 + (w_1 + 20d)^2 + (w_1 + 21d)^2 + (w_1 + 22d)^2 +$$

$$(w_1 + 23d)^2 + (w_1 + 24d)^2 + (w_1 + 25d)^2 + (w_1 + 26d)^2 + (w_1 + 27d)^2 + (w_1 + 28d)^2 + (w_1 + 29d)^2 + 10d^2$$

Simplifies to

$$30w_1^2 + 870w_1d + 8565d^2 = 3(10w_1^2 + 290w_1d + 2855d^2) \dots(1.2)$$

Splitting equation (1.2) into thrice sums of squares we obtain:

$$\begin{aligned} & 3(10d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2) + (w_1^2 + 20w_1d + 100d^2) + \\ & (w_1^2 + 26w_1d + 169d^2 + (w_1^2 + 32w_1d + 256d^2) + (w_1^2 + 38w_1d + 361d^2) + (w_1^2 + 44w_1d + 484d^2) + \\ & (w_1^2 + 50w_1d + 625d^2) + (w_1^2 + 56w_1d + 784d^2)) \\ & = 3(10d^2) + (w_1 + d)^2 + (w_1 + 4d)^2 + (w_1 + 7d)^2 + (w_1 + 10d)^2 + (w_1 + 13d)^2 + (w_1 + 16d)^2 + (w_1 + \\ & 19d)^2 + (w_1 + 22d)^2 + (w_1 + 25d)^2 + (w_1 + 28d)^2 \\ & = 3(6d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2) \end{aligned}$$

This completes the proof \square .

Theorem 1.3: Consider equation 1 satisfying condition $(n, w_1, w_2, \dots, w_{33}, 11d) = (33, w_1, w_2, \dots, w_{33}, 11d)$

Then, the diophantine equation:

$$\begin{aligned} & w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + \\ & w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + w_{28}^2 + w_{29}^2 + w_{30}^2 + w_{31}^2 + w_{32}^2 + w_{33}^2 + 11d^2 \\ & = 3(11d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2) \end{aligned}$$

has the solution in integers if $w_{33} - w_{32} = w_{32} - w_{31} = w_{31} - w_{30} = w_{30} - w_{29} = w_{29} - w_{28} = w_{28} - w_{27} = w_{27} - w_{26} = w_{26} - w_{25} = w_{25} - w_{24} = w_{24} - w_{23} = w_{23} - w_{22} = w_{22} - w_{21} = w_{21} - w_{20} = w_{20} - w_{19} = w_{19} - w_{18} = w_{18} - w_{17} = w_{17} - w_{16} = w_{16} - w_{15} = w_{15} - w_{14} = w_{14} - w_{13} = w_{13} - w_{12} = w_{12} - w_{11} = w_{11} - w_{10} = w_{10} - w_9 = w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$

Proof: Consider the equation

$$\begin{aligned} & w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + \\ & w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + w_{28}^2 + w_{29}^2 + w_{30}^2 + w_{31}^2 + w_{32}^2 + w_{33}^2 + 11d^2 \\ & = 3(11d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2) \end{aligned}$$

And suppose that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d, w_{10} = w_1 + 9d, w_{11} = w_1 + 10d, w_{12} = w_1 + 11d, w_{13} = w_1 + 12d, w_{14} = w_1 + 13d, w_{15} = w_1 + 14d, w_{16} = w_1 + 15d, w_{17} = w_1 + 16d, w_{18} = w_1 + 17d, w_{19} = w_1 + 18d, w_{20} = w_1 + 19d, w_{21} = w_1 + 20d, w_{22} = w_1 + 21d, w_{23} = w_1 + 22d, w_{24} = w_1 + 23d, w_{25} = w_1 + 24d, w_{26} = w_1 + 25d, w_{27} = w_1 + 26d, w_{28} = w_1 + 27d, w_{29} = w_1 + 28d, w_{30} = w_1 + 29d, w_{31} = w_1 + 30d, w_{32} = w_1 + 31d, w_{33} = w_1 + 32d$

The left hand side expressed as:

$$\begin{aligned} & \text{And suppose that } w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + \\ & (w_1 + 7d)^2 + (w_1 + 8d)^2 + (w_1 + 9d)^2 + (w_1 + 10d)^2 + (w_1 + 11d)^2 + (w_1 + 12d)^2 + (w_1 + 13d)^2 + \\ & (w_1 + 14d)^2 + (w_1 + 15d)^2 + (w_1 + 16d)^2 + (w_1 + 17d)^2 + (w_1 + 18d)^2 + (w_1 + 19d)^2 + (w_1 + 20d)^2 + \end{aligned}$$

$$(w_1 + 21d)^2 + (w_1 + 22d)^2 + (w_1 + 23d)^2 + (w_1 + 24d)^2 + (w_1 + 25d)^2 + (w_1 + 26d)^2 + (w_1 + 27d)^2 + (w_1 + 28d)^2 + (w_1 + 29d)^2 + (w_1 + 30d)^2 + (w_1 + 31d)^2 + (w_1 + 32d)^2 + 11d^2$$

Simplifies to

$$33w_1^2 + 1056w_1d + 11451d^2 = 3(11w_1^2 + 352w_1d + 3817d^2) \dots(1.3)$$

Splitting equation (1.3) into thrice sums of squares we obtain:

$$3(11d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2) + (w_1^2 + 20w_1d + 100d^2) + (w_1^2 + 26w_1d + 169d^2) + (w_1^2 + 32w_1d + 256d^2) + (w_1^2 + 38w_1d + 361d^2) + (w_1^2 + 44w_1d + 484d^2) + (w_1^2 + 50w_1d + 625d^2) + (w_1^2 + 56w_1d + 784d^2) + (w_1^2 + 62w_1d + 961d^2))$$

$$= 3(11d^2) + (w_1 + d)^2 + (w_1 + 4d)^2 + (w_1 + 7d)^2 + (w_1 + 10d)^2 + (w_1 + 13d)^2 + (w_1 + 16d)^2 + (w_1 + 19d)^2 + (w_1 + 22d)^2 + (w_1 + 25d)^2 + (w_1 + 28d)^2 + (w_1 + 31d)^2$$

$$= 3(11d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2)$$

This completes the proof \square .

Theorem 1.4: Consider equation 1 satisfying condition $(n, w_1, w_2, \dots, w_{36}, 12d) = (36, w_1, w_2, \dots, w_{36}, 12d)$

Then, the diophantine equation:

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + w_{28}^2 + w_{29}^2 + w_{30}^2 + w_{31}^2 + w_{32}^2 + w_{33}^2 + w_{34}^2 + w_{35}^2 + w_{36}^2 + 12d^2 = 3(12d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2 + w_{35}^2)$$

has the solution in integers if $w_{36} - w_{35} = w_{35} - w_{34} = w_{34} - w_{33} = w_{33} - w_{32} = w_{32} - w_{31} = w_{31} - w_{30} = w_{30} - w_{29} = w_{29} - w_{28} = w_{28} - w_{27} = w_{27} - w_{26} = w_{26} - w_{25} = w_{25} - w_{24} = w_{24} - w_{23} = w_{23} - w_{22} = w_{22} - w_{21} = w_{21} - w_{20} = w_{20} - w_{19} = w_{19} - w_{18} = w_{18} - w_{17} = w_{17} - w_{16} = w_{16} - w_{15} = w_{15} - w_{14} = w_{14} - w_{13} = w_{13} - w_{12} = w_{12} - w_{11} = w_{11} - w_{10} = w_{10} - w_9 = w_9 - w_8 = w_8 - w_7 = w_7 - w_6 = w_6 - w_5 = w_5 - w_4 = w_4 - w_3 = w_3 - w_2 = w_2 - w_1 = d$

Proof: Consider the equation

$$w_1^2 + w_2^2 + w_3^2 + w_4^2 + w_5^2 + w_6^2 + w_7^2 + w_8^2 + w_9^2 + w_{10}^2 + w_{11}^2 + w_{12}^2 + w_{13}^2 + w_{14}^2 + w_{15}^2 + w_{16}^2 + w_{17}^2 + w_{18}^2 + w_{19}^2 + w_{20}^2 + w_{21}^2 + w_{22}^2 + w_{23}^2 + w_{24}^2 + w_{25}^2 + w_{26}^2 + w_{27}^2 + w_{28}^2 + w_{29}^2 + w_{30}^2 + w_{31}^2 + w_{32}^2 + w_{33}^2 + w_{34}^2 + w_{35}^2 + w_{36}^2 + 12d^2 = 3(12d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2 + w_{35}^2)$$

And suppose that $w_2 = w_1 + d, w_3 = w_1 + 2d, w_4 = w_1 + 3d, w_5 = w_1 + 4d, w_6 = w_1 + 5d, w_7 = w_1 + 6d, w_8 = w_1 + 7d, w_9 = w_1 + 8d, w_{10} = w_1 + 9d, w_{11} = w_1 + 10d, w_{12} = w_1 + 11d, w_{13} = w_1 + 12d, w_{14} = w_1 + 13d, w_{15} = w_1 + 14d, w_{16} = w_1 + 15d, w_{17} = w_1 + 16d, w_{18} = w_1 + 17d, w_{19} = w_1 + 18d, w_{20} = w_1 + 19d, w_{21} = w_1 + 20d, w_{22} = w_1 + 21d, w_{23} = w_1 + 22d, w_{24} = w_1 + 23d, w_{25} = w_1 + 24d, w_{26} = w_1 + 25d, w_{27} = w_1 + 26d, w_{28} = w_1 + 27d, w_{29} = w_1 + 28d, w_{30} = w_1 + 29d, w_{31} = w_1 + 30d, w_{32} = w_1 + 31d, w_{33} = w_1 + 32d, w_{34} = w_1 + 23d, w_{35} = w_1 + 34d, w_{36} = w_1 + 35d$

The left hand side expressed as:

$$\text{And suppose that } w_1^2 + (w_1 + d)^2 + (w_1 + 2d)^2 + (w_1 + 3d)^2 + (w_1 + 4d)^2 + (w_1 + 5d)^2 + (w_1 + 6d)^2 + (w_1 + 7d)^2 + (w_1 + 8d)^2 + (w_1 + 9d)^2 + (w_1 + 10d)^2 + (w_1 + 11d)^2 + (w_1 + 12d)^2 + (w_1 + 13d)^2 + (w_1 + 14d)^2 +$$

$$(w_1 + 15d)^2 + (w_1 + 16d)^2 + (w_1 + 17d)^2 + (w_1 + 18d)^2 + (w_1 + 19d)^2 + (w_1 + 20d)^2 + (w_1 + 21d)^2 + (w_1 + 22d)^2 + (w_1 + 23d)^2 + (w_1 + 24d)^2 + (w_1 + 25d)^2 + (w_1 + 26d)^2 + (w_1 + 27d)^2 + (w_1 + 28d)^2 + (w_1 + 29d)^2 + (w_1 + 30d)^2 + (w_1 + 31d)^2 + (w_1 + 32d)^2 + (w_1 + 33d)^2 + (w_1 + 34d)^2 + (w_1 + 35d)^2 + 12d^2$$

Simplifies to

$$36w_1^2 + 1260w_1d + 14922d^2 = 3(12w_1^2 + 420w_1d + 4974d^2) \dots(1.4)$$

Splitting equation (1.3) into thrice sums of squares we obtain:

$$3(12d^2 + (w_1^2 + 2w_1d + d^2) + (w_1^2 + 8w_1d + 16d^2) + (w_1^2 + 14w_1d + 49d^2) + (w_1^2 + 20w_1d + 100d^2) + (w_1^2 + 26w_1d + 169d^2) + (w_1^2 + 32w_1d + 256d^2) + (w_1^2 + 38w_1d + 361d^2) + (w_1^2 + 44w_1d + 484d^2) + (w_1^2 + 50w_1d + 625d^2) + (w_1^2 + 56w_1d + 784d^2) + (w_1^2 + 62w_1d + 961d^2) + (w_1^2 + 68w_1d + 1156d^2))$$

$$= 3(12d^2 + w_2^2 + w_5^2 + w_8^2 + w_{11}^2 + w_{14}^2 + w_{17}^2 + w_{20}^2 + w_{23}^2 + w_{26}^2 + w_{29}^2 + w_{32}^2 + w_{35}^2)$$

This completes the proof \square .

3 Conclusion

In summary, the solution of the diophantine equation $\sum_{r=1}^n w_r^2 + \frac{n}{3}d^2 = 3(\frac{nd^2}{3} + \sum_{n=1}^{\frac{n}{3}} w_{3r-1}^2)$, under the specified conditions of a common difference d between consecutive terms w_n, w_{n1}, w_2, w_1 where $w_n w_{n1} = w_{n1} w_{n2} = w_2 w_1 = d$ has been achieved for some cases. This solution provides valuable insights into the relation among the sequence terms, enhancing our understanding of the inherent patterns and structures within the equation. For future investigations, it is recommended to explore extensions of this diophantine equation by proving conjecture (1).

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

References

- [1] Bombieri, E. and Bourgain, J. (2015). A problem on sums of two squares. *International Mathematics Research Notices*, 2015(11):3343–3407.
- [2] Cai, Y. (2016). Waring-goldbach problem: two squares and higher powers. *Journal de théorie des nombres de Bordeaux*, 28(3):791–810.
- [3] Cavallo, A. (2019). An elementary computation of the galois groups of symmetric sextic trinomials. *arXiv preprint arXiv:1902.00965*.

-
- [4] Christopher, A. D. (2016). A partition-theoretic proof of fermat's two squares theorem. *Discrete Mathematics*, 339(4):1410–1411.
- [5] Fathi, A., Mobadersany, P., and Fathi, R. (2012). A simple method to solve quartic equations. *Australian Journal of Basic and Applied Sciences*, 6(6):331–336.
- [6] Kimtai, B. S. and Mude, L. H. (2023). On generalized sums of six, seven and nine cube. *Science Mundi*, 3(1):135–142.
- [7] Kouropoulos, G. P. (2021). A combined methodology for the approximate estimation of the roots of the general sextic polynomial equation.
- [8] Lao, H. M., Zachary, K. K., and Kinyanjui, J. N. (2023). Some generalized formula for sums of cube. *British Journal of Mathematics & Computer Science*.
- [9] Mochimaru, Y. (2005). Solution of sextic equations. *International Journal of Pure and Applied Mathematics*, 23(4):577.
- [10] Mude, L. H. (2022). Some formulae for integer sums of two squares. *Journal of Advances in Mathematics and Computer Science*, 37(4):53–57.
- [11] Mude, L. H. (2024). On some mixed polynomial exponential diophantine equation: $(\alpha^n + \beta^n + a(\alpha^s \pm \beta^s)^m + d = r(u^k + v^k + w^k))$ with (α) and (β) consecutive. *Journal of Advances in Mathematics and Computer Science*, 39(10):11–17.
- [12] Mude, L. H., Ndung'u, K. J., and Kayiita, Z. K. (2024). On sums of squares involving integer sequence: $(\sum_{r=1}^n w_r^2 + \frac{n}{3} d^2 = 3 \left(\frac{n^2}{3} + \sum_{r=1}^{\frac{n}{3}} w_{r-1}^2 \right))$. *Journal of Advances in Mathematics and Computer Science*, 39(7):1–6.
- [13] Najman, F. (2010a). The diophantine equation $x^4 \pm y^4 = iz^2$ in gaussian integers. *The American Mathematical Monthly*, 117(7):637–641.
- [14] Najman, F. (2010b). Torsion of elliptic curves over quadratic cyclotomic fields. *arXiv preprint arXiv:1005.0558*.
- [15] Simatwo, K. B. (2024). Equal sums of four even powers. *Asian Research Journal of Mathematics*, 20(5):50–54.
- [16] Tignol, J.-P. (2015). *Galois' theory of algebraic equations*. World Scientific Publishing Company.