

Locally attractivity results for fractional order quadratic functional integral equations

Abstract: we prove using hybridfixed point theorem in Banach algebra an existence of solutions to fractional order quadratic functional integral equation in \mathcal{R}_+ . Also locally attractivity results and solutions for fractional order quadratic functional integral equations is proved. Also, one example is considered.

Keywords: Fixed point theorem, Banach algebra, Quadratic functional integral equation, existence result, locally attractive solution, Extremal Solution.

1. INTRODUCTION:

Fractional Calculus is a generalization of ordinary differentiation and integration to arbitrary order. The subject has its origin in 16th century. During three centuries, the theory of fractional calculus developed as pure theoretical field, useful only for Mathematicians [22] Quadratic functional integral equation has newly received a lot of attention and establishes a meaningful branch of nonlinear analysis[18-29]. For examples, quadratic integral equations are often applicable in the theory of radioactive transfer, kinetic theory of gases, in the theory of neutron transport and in the traffic theory. Numerous research papers and monographs devoted to quadratic differential and integral equations of fractional order have appeared (see [1-3, 5-7, 9-17]. These papers contain different types of existence results for equations of fractional order. Here we are concerned with the existence of solution for fractional order quadratic functional integral equation also locally attractive solutions of following QFIE will be proved.

2. Statement of the Problem:

Let $\alpha, \zeta \in (0,1)$ and \mathcal{R} denote the real numbers whereas \mathcal{R}_+ be the set of nonnegative numbers i.e. $\mathcal{R}_+ = [0, \infty) \subset \mathcal{R}$.

Consider the following nonlinear functional quadratic integral equations of fractional order

$$x(t) = \left[\frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t)x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \right] \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds \right]$$

$$\forall t \in \mathcal{R}_+ \quad (2.1)$$

Where $q: \mathcal{R}_+ \rightarrow \mathcal{R}, f(t, x(t)x(\gamma(s))) = f: \mathcal{R}_+ \times \mathcal{R} \rightarrow \mathcal{R}, g(t, x(s)x(\beta(s))) = g: \mathcal{R}_+ \times \mathcal{R} \rightarrow \mathcal{R}$ and By a solution of the (2.1) we mean a function $x \in \mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ that satisfies (2.1) on \mathcal{R}_+ . Where $\mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ is the space of continuous and bounded real-valued functions defined on \mathcal{R}_+ .

In this paper, we prove the locally attractive of the solutions for QFIE (2.1) employing a classical hybrid fixed point theorem of B.C. Dhage [5]. In the next section, we collect some preliminary definitions and auxiliary results that will be used in the follows.

3.Preliminaries:

Let $X = \mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ be Banach algebra with norm $\|\cdot\|$ and let Ω be a subset of X . Let a mapping $\mathcal{A}: X \rightarrow X$ be an operator and consider the following operator equation in X , namely,

$$x(t) = (\mathcal{A}x)(t) \quad t \in \mathcal{R}_+ \quad (3.1)$$

Below we give different characterizations of the solutions for operator equation (3.1) on \mathcal{R}_+ .

We need the following definitions in the sequel.

Definition 3.1[29]: The solution $x(t)$ of the equation (3.1) is said to be locally attractive if there exists an closed ball $B_r[0]$ in $\mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ such that for arbitrary solutions $x = x(t)$ and $y = y(t)$ of equation (3.1) belonging to $B_r[0] \cap \Omega$ such that

$$\lim_{t \rightarrow \infty} (x(t) - y(t)) = 0 \quad (3.2)$$

Definition 3.2[29]: Let X be a Banach space. A mapping $\mathcal{A}: X \rightarrow X$ is called Lipschitz if there is a constant $\alpha > 0$ such that $\|\mathcal{A}x - \mathcal{A}y\| \leq \alpha \|x - y\|$ for all $x, y \in X$. If $\alpha < 1$ then \mathcal{A} is called a contraction on X with the contraction constant α .

Definition 3.3: (Dugundji and Granas [9]). An operator \mathcal{A} on a Banach space X into itself is called Compact if for any bounded subset S of X , $\mathcal{A}(S)$ is a relatively compact subset of X . If \mathcal{A} is continuous and compact, then it is called completely continuous on X .

Let X be a Banach space with the norm $\|\cdot\|$ and Let $\mathcal{A}: X \rightarrow X$ be an operator (in general nonlinear). Then \mathcal{A} is called

- (i) Compact if $\mathcal{A}(X)$ is relatively compact subset of X ;
- (ii) totally bounded if $\mathcal{A}(S)$ is a totally bounded subset of X for any bounded subset S of X .
- (iii) Completely continuous if it is continuous and totally bounded operator on X .

It is clear that every compact operator is totally bounded but the converse need not be true.

The solutions of (2.1) in the space $\mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ of continuous and bounded real-valued functions defined on \mathcal{R}_+ . Define a standard supremum norm $\|\cdot\|$ and a multiplication “.” in

$$\mathcal{BC}(\mathcal{R}_+, \mathcal{R}) \text{ by } \|x\| = \sup\{|x(t)| : t \in \mathcal{R}_+\} \quad (3.3)$$

$$(xy)(t) = x(t)y(t) \quad t \in \mathcal{R}_+ \quad (3.4)$$

Clearly, $\mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ becomes a Banach space with respect to the above norm and the multiplication in it. By $\mathcal{L}^1(\mathcal{R}_+, \mathcal{R})$ we denote the space of Lebesgue integrable functions on \mathcal{R}_+ with the norm $\|\cdot\|_{\mathcal{L}^1}$ defined by

$$\|X\|_{\mathcal{L}^1} = \int_0^\infty |x(t)| dt \quad (3.5)$$

Denote by $\mathcal{L}^1(a, b)$ be the space of Lebesgue integrable functions on the interval (a, b) , which is equipped with the standard norm. Let $x \in \mathcal{L}^1(a, b)$ and let $\beta > 0$ be a fixed number.

Definition 3.4[3]: The Riemann-Liouville fractional integral of order β of the function $f(t)$ is defined by the formula $I^\beta f(t) = \frac{1}{\Gamma(\beta)} \int_0^t \frac{f(s)}{(t-s)^{1-\beta}} ds \quad t \in (a, b) \quad (3.6)$

Where $\Gamma(\beta)$ denote the gamma function.

It may be shown that the fractional integral operator I^β transforms the space $\mathcal{L}^1(a, b)$ into itself and has some other properties (see [12-19])

Definition 3.5[5]: A set $A \subseteq [a, b]$ is said to be measurable if $m^* A = m_* A$. In this case we define mA , the measure of A as $mA = m^* A = m_* A$

If A_1 and A_2 are measurable subsets of $[a, b]$ then their union and their intersection is also measurable.

Clearly every open or closed set in \mathbb{R} is measurable.

Definition 3.7[12]: Let f be a function defined on $[a, b]$. Then f is measurable function if for each $\alpha \in \mathbb{R}$, the set $\{x : f(x) > \alpha\}$ is measurable set.

i.e. f is measurable function if for every real number α the inverse image of (α, ∞) is an open set.

As (α, ∞) is an open set and if f is continuous, then inverse image under f of (α, ∞) is open. Open sets being measurable, hence every continuous function is measurable.

Definition 3.8[7]: A sequence of functions $\{f_n\}$ is said to converge uniformly on an interval $[a, b]$ to a function f if for any $\epsilon > 0$ and for all $x \in [a, b]$ there exists an integer N (dependent only on ϵ) such that for all $x \in [a, b]$

$$|f_n(x) - f(x)| < \epsilon \quad \forall n \geq N$$

Definition 3.7[7]: The Family F is Equi continuous at a point $x_0 \in X$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that $d(f(x_0), f(x)) < \epsilon$ for all $f \in F$ and all x that $d(x_0, x) < \delta$.

The family is point wise Equi continuous if it is Equi continuous at each point of X .

The family is uniformly Equi continuous if for every $\varepsilon > 0$ there exists $\delta > 0$ such that $d(f(x_1), f(x_2)) < \varepsilon$ for all $f \in F$ and all $x_1, x_2 \in X$ such that $d(x_1, x_2) < \delta$.

Theorem 3.1: (Arzela-Ascoli theorem (7)): If every uniformly bounded and Equi-continuous sequence $\{f_n\}$ of functions in $\mathcal{C}(J, \mathcal{R})$, then it has a convergent subsequence.

Theorem 3.2[7]: A metric space X is compact iff every sequence in X has a convergent subsequence.

We employ a hybrid fixed point theorem of Dhage [5] for proving the existence result.

Theorem 3.3 : (Dhage [5]). Let S be a closed-convex and bounded subset of the Banach space X and let $\mathcal{A}, \mathcal{B}: S \rightarrow S$ be two operators satisfying:

- (a) \mathcal{A} is Lipschitz with the Lipschitz constant k ,
- (b) \mathcal{B} is completely continuous,
- (c) $\mathcal{A}\mathcal{B}x \in S$ for all $x \in S$ and
- (d) $Mk < 1$ Where $M = \|\mathcal{B}(S)\| = \sup \{\|\mathcal{B}x\|: x \in S\}$

Then the operator equation $\mathcal{A}\mathcal{B}x = x$ has a solution and the set of all solutions is compact in S

4. Existence results:

Definition 4.1.: A mapping $g: \mathcal{R}_+ \times \mathcal{R} \rightarrow \mathcal{R}$ is said to be Caratheodory if

1. $t \rightarrow g(t, x)$ is measurable for all $x \in \mathcal{R}$, and
2. $t \rightarrow g(t, x)$ is continuous almost everywhere for $t \in \mathcal{R}_+$

Again a caratheodory function g is called \mathcal{L}^1 -Caratheodory if

3. for each real number $r > 0$ there exists a function $h_r \in \mathcal{L}^1(\mathcal{R}_+, \mathcal{R})$ such that

$$\left| g\left(s, x(s)x(\beta(s))\right) \right| \leq h_r(t) \quad \text{a.e. } t \in \mathcal{R}_+ \text{ for all } x \in \mathcal{R} \text{ with } |x| \leq r$$

Finally, a Caratheodory function $g(t, x)$ is called $\mathcal{L}^1_{\mathcal{R}}$ - Caratheodory if

4. there exists a function $h \in \mathcal{L}^1(\mathcal{R}_+, \mathcal{R})$ such that $\left| g\left(s, x(s)x(\beta(s))\right) \right| \leq h(t)$
a.e. $t \in \mathcal{R}_+$ for all $x \in \mathcal{R}$

Throughout this paper, we assume the following Hypothesis

(H_1) The function $f: \mathcal{R}_+ \times \mathcal{R} \rightarrow \mathcal{R}$ is continuous and bounded with bound

(H_1) $F = \sup_{(t,x) \in \mathcal{R}_+ \times \mathcal{R}} |f(t, x)|$ there exists a bounded function $l: \mathcal{R}_+ \rightarrow \mathcal{R}_+$ with bound L Satisfying

$$\left| f\left(t, x(t)x(\gamma(s))\right) - f\left(t, y(t)y(\gamma(s))\right) \right| \leq l(t) \frac{r(\alpha+1)}{(t)^\alpha} |x - y| \quad t \in \mathcal{R}_+ \text{ for all } x, y \in \mathcal{R}$$

(H_2) $q: \mathcal{R}_+ \rightarrow \mathcal{R}$ is continuous function on \mathcal{R}_+ ; also $\lim_{t \rightarrow \infty} q(t) = 0$

(H₃) The functions $f, g: \mathcal{R}_+ \times \mathcal{R} \rightarrow \mathcal{R}$ satisfy caratheodory condition (i.e. measurable in t for all $x \in \mathcal{R}$ and continuous in x for all $t \in \mathcal{R}_+$) and there exist function $h_1, h_2 \in \mathcal{L}^1(\mathcal{R}_+, \mathcal{R})$ Such that $f(t, x(t)x(\gamma(s))) \leq h_1(t)$ and $g(s, x(s)x(\beta(s))) \leq h_2(t) \quad \forall (t, x) \in \mathcal{R}_+ \times \mathcal{R}$

(H₄) The uniform continuous function $v: \mathcal{R}_+ \rightarrow \mathcal{R}_+$ defined by the formulas

$$v_1(t) = \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds, v_2(t) = \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds$$

is bounded on \mathcal{R}_+ and vanishes at infinity, that is, $\lim_{t \rightarrow \infty} v_i(t) = 0$

Remark 4.1: Note that if the hypothesis (H₂) and (H₃) hold, then there exist constants $K_1 > 0$ and $K_2 > 0$ such that: $K_1 = \sup\{q(t): t \in \mathcal{R}_+\}$,

$$K_2 = \sup_{t \geq 0} \frac{1}{\Gamma(\alpha)} \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds, \quad K_3 = \sup_{t \geq 0} \frac{1}{\Gamma(\beta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds$$

Theorem 4.1: Let the assumptions [(H₁)-(H₅)] are satisfied. Furthermore if $L(K_1 + K_3) < 1$, where K_1, K_2 and K_3 are defined remark (4.1), Then the equation (2.1) has a solution in the space $\mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ and solutions of the equation (2.1) are locally attractive on \mathcal{R}_+ . Moreover (2.1) has attractive solutions.

Proof: By a solution of the (2.1) we mean a continuous function $x: \mathcal{R}_+ \rightarrow \mathcal{R}$ that satisfies (2.1) on \mathcal{R}_+ .

Let $X = \mathcal{BC}(\mathcal{R}_+, \mathcal{R})$ be Banach Algebras of all continuous and bounded real valued function on \mathcal{R}_+ with the norm $\|x\| = \sup_{t \in \mathcal{R}_+} |x(t)|$ (4.1)

We show that existence of solution for (2.1) under some suitable conditions on the functions involved in (2.1).

Consider the closed ball $B_r[0]$ in X centered at origin 0 and of radius r , where

$$r = K_2[K_1 + K_3] > 0$$

Let us define two operators \mathcal{A} and \mathcal{B} on $B_r[0]$ by

$$\mathcal{A}x(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t)x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \quad (4.2)$$

$$\text{and } \mathcal{B}x(t) = \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \quad \forall t \in \mathcal{R}_+ \quad (4.3)$$

The function q is continuous on \mathcal{R}_+ , the function $\mathcal{B}x$ is also continuous and bounded in view of hypotheses $(H_2 - H_3)$. The mapping \mathcal{A} is well defined (Since the hypotheses (H_1) holds) and the function $\mathcal{A}x$ is continuous and bounded on \mathcal{R}_+ .

Therefore \mathcal{A} and \mathcal{B} define the operators $\mathcal{A}, \mathcal{B}: B_r[0] \rightarrow X$. we shall show that \mathcal{A} and \mathcal{B} satisfy all the requirements of theorem (3.3) on $B_r[0]$.

Step I: Firstly, we show that \mathcal{A} is Lipschitz on $B_r[0]$. let $x, y \in B_r[0]$ be arbitrary, and then by hypothesis (H_2) , we get

$$\begin{aligned} |\mathcal{A}x(t) - \mathcal{A}y(t)| &= \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(s), x(\gamma(s)))}{(t-s)^{1-\alpha}} ds - \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, y(s), y(\gamma(s)))}{(t-s)^{1-\alpha}} ds \right| \\ &\leq \frac{1}{\Gamma(\alpha)} \int_0^t \frac{1}{(t-s)^{1-\alpha}} |f(t, x(s), x(\gamma(s))) - f(t, y(s), y(\gamma(s)))| ds \\ &\leq \frac{1}{\Gamma(\alpha)} \int_0^t \frac{1}{(t-s)^{1-\alpha}} |x(s) - y(s)| \\ &\leq l(t) \frac{\Gamma(\alpha+1)}{(t)^\alpha} |x-y| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{1}{(t-s)^{1-\alpha}} ds \\ &\leq l(t) \frac{\Gamma(\alpha+1)}{(t)^\alpha} |x-y| \frac{1}{\Gamma(\alpha)} \left[\frac{(t-s)^\alpha}{\alpha} \right]_0^t \\ &\leq l(t) \frac{\Gamma(\alpha+1)}{t^\alpha} |x-y| \frac{t^\alpha}{\Gamma(\alpha+1)} \end{aligned}$$

$$\leq L \|x - y\| \text{ for all } t \in \mathcal{R}_+ \quad (4.4)$$

Taking supremum over t

$$\|\mathcal{A}x - \mathcal{A}y\| \leq L \|x - y\| \text{ for all } x, y \in B_r[0] \quad (4.5)$$

Gives that \mathcal{A} is Lipschitz on $B_r[0]$ with the Lipschitz constant L .

Step II: Now we show that \mathcal{B} is completely continuous operator on $B_r[0]$.

Firstly we show that \mathcal{B} is continuous on $B_r[0]$.

Case I: let us assume that, $t \in [0, T]$ then evaluating we obtain the following estimate

$$\begin{aligned}
& |(Bx)t - (By)t| \\
& \leq \left| q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds - q(t) - \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right| \\
& \leq \left| \frac{1}{\Gamma(\beta)} \int_0^T \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds - \frac{1}{\Gamma(\beta)} \int_0^T \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right| \\
& \leq \frac{1}{\Gamma(\zeta)} \left[\int_0^T \frac{|g(s, x(s)x(\beta(s))) - g(s, y(s)y(\beta(s)))|}{(t-s)^{1-\zeta}} ds \right] \\
& \leq \frac{1}{\Gamma(\zeta)} \left[\int_0^T \frac{w_r^T(h_2, \epsilon)}{(t-s)^\zeta} ds \right] \\
& \leq \frac{1}{\Gamma(\zeta)} \left[\frac{w_r^T(h_2, \epsilon)}{\Gamma(\beta)\beta} T^\zeta ds \right] \\
& \leq \left[\frac{w_r^T(h_2, \epsilon)}{\Gamma(\zeta + 1)} T^\zeta ds \right] \tag{4.6}
\end{aligned}$$

Where $w_r^T(g, \epsilon) = \sup \{|g(s, x) - g(s, y)| : s \in [0, T]; x, y \in [-r, r], |x - y| \leq \epsilon\}$

Therefore, from the uniform continuity of the function $g(t, x)$ on the set $[0, T] \times [-r, r]$.we derive that $w_r^T(g, \epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

Case II: Suppose that $t \geq T$ there exist $T > 0$ and let us fix arbitrary $\epsilon > 0$ and take $x, y \in B_r[0]$ such that $\|x - y\| \leq \epsilon$ Then

$$\begin{aligned}
& |(Bx)t - (By)t| \\
& \leq \left| q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds - q(t) - \frac{1}{\Gamma(\beta)} \int_0^t \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right| \\
& \leq \left| \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\beta}} ds - \frac{1}{\Gamma(\beta)} \int_0^t \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{\Gamma(\zeta)} \left[\int_0^t \frac{|g(s, x(s)x(\beta(s)))|}{(t-s)^{1-\beta}} ds + \int_0^t \frac{|g(s, y(s)y(\beta(s)))|}{(t-s)^{1-\zeta}} ds \right] \\
&\leq \frac{1}{\Gamma(\zeta)} \left[\int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds + \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right] \\
&\leq \frac{2}{\Gamma(\zeta)} \left[\int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right] \\
&\leq \frac{2v_2(t)}{\Gamma(\zeta)} \tag{4.6}
\end{aligned}$$

Hence we see that There exists $T > 0$ s.t.

$$v_2(t) \leq \frac{\varepsilon \Gamma(\zeta)}{2} \text{ for } t > T$$

Since ε is an arbitrary,

$$\text{from (4.7) we derive that } |(Bx)t - (By)t| \leq \varepsilon \tag{4.8}$$

Now combining the case I and II, we conclude that the operator B is continuous operator on closed ball $B_r[0]$ in to itself.

Step III: Next we show that B is compact on $B_r[0]$.

(A) First prove that every sequence $\{Bx_n\}$ in $B(B_r[0])$ has a uniformly bounded sequence in $B(B_r[0])$. Now by $(H_2) - (H_3) - (H_4)$

$$\begin{aligned}
|(Bx_n)t| &= \left| q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x_n(\beta(s)))}{(t-s)^{1-\zeta}} ds \right| \\
|(Bx_n)t| &\leq |q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{|g(s, x(s)x_n(\beta(s)))|}{(t-s)^{1-\zeta}} ds \\
|(Bx_n)t| &\leq |q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \\
|(Bx_n)t| &\leq |q(t)| + \frac{v_2(t)}{\Gamma(\zeta)}
\end{aligned}$$

$$|(Bx_n)t| \leq K_1 + K_3 \forall t \in \mathcal{R}_+ \tag{4.9}$$

Taking supremum over t , we obtain $\|Bx_n\| \leq K_1 + K_3 \quad \forall n \in N$

This shows that $\{Bx_n\}$ is a uniformly bounded sequence in $B(B_r[0])$.

(B) Now we proceed to show that sequence $\{Bx_n\}$ is also equicontinuous.

Let $\varepsilon > 0$ be given. Since $\lim_{t \rightarrow \infty} q(t) = 0$ there is constant $T > 0$ such that

$$|q(t)| < \varepsilon/2 \text{ for all } t \geq T$$

Case I: If $t_1, t_2 \in [0, T]$ then we have

$$\begin{aligned} & |(Bx_n)_{t_2} - (Bx_n)_{t_1}| \\ & \leq \left| q(t_2) + \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{g(s, x(s)x_n(\beta(s)))}{(t_2 - s)^{1-\zeta}} ds - q(t_1) - \frac{1}{\Gamma(\zeta)} \int_0^{t_1} \frac{g(s, x(s)x_n(\beta(s)))}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \left| \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_2 - s)^{1-\zeta}} ds - \frac{1}{\Gamma(\beta)} \int_0^{t_1} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \left| \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{h_2(s)}{(t_2 - s)^{1-\zeta}} ds - \frac{1}{\Gamma(\beta)} \int_0^{t_1} \frac{h_2(s)}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \frac{1}{\Gamma(\zeta)} \left| \int_0^{t_2} \frac{h_2(s)}{(t_2 - s)^{1-\zeta}} ds - \int_0^{t_1} \frac{h_2(s)}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \frac{1}{\Gamma(\zeta)} |v_2(t_2) - v_2(t_1)| \quad (4.10) \end{aligned}$$

from the uniform continuity of the function $q(t), v(t)$ on $[0, T]$, we get

$$|(Bx_n)_{t_2} - (Bx_n)_{t_1}| \rightarrow 0 \text{ as } t_1 \rightarrow t_2$$

Case II: If $t_1, t_2 \geq T$ then we have

$$\begin{aligned} & |(Bx_n)_{t_2} - (Bx_n)_{t_1}| \\ & \leq \left| q(t_2) + \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{g(s, x(s)x_n(\beta(s)))}{(t_2 - s)^{1-\zeta}} ds - q(t_1) - \frac{1}{\Gamma(\zeta)} \int_0^{t_1} \frac{g(s, x(s)x_n(\beta(s)))}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \left| \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_2 - s)^{1-\zeta}} ds - \frac{1}{\Gamma(\beta)} \int_0^{t_1} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_1 - s)^{1-\zeta}} ds \right| \\ & \leq |q(t_2) - q(t_1)| + \left| \frac{1}{\Gamma(\zeta)} \int_0^{t_2} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_2 - s)^{1-\zeta}} ds \right| + \left| \frac{1}{\Gamma(\beta)} \int_0^{t_1} \frac{|g(s, x(s)x_n(\beta(s)))|}{(t_1 - s)^{1-\zeta}} ds \right| \end{aligned}$$

$$\leq |q(t_2) - q(t_1)| + \frac{v_2(t_2)}{\Gamma(\zeta)} + \frac{v_2(t_1)}{\Gamma(\zeta)} \leq 0 + \frac{\epsilon}{2} + \frac{\epsilon}{2}$$

$$\leq \epsilon \text{ as } t_1 \rightarrow t_2. \quad (4.11)$$

Case III: If $t_1, t_2 \in \mathcal{R}_+$ With $t_1 < T < t_2$ then we have

$$|(Bx_n)_{t_2} - (Bx_n)_{t_1}| \leq |Bx_n(t_2) - Bx_n(T)| + |Bx_n(T) - Bx_n(t_1)| \quad (4.12)$$

Now if $t_1 \rightarrow t_2$ then $t_1 \rightarrow T$ and $T \rightarrow t_2$

Therefore, $|Bx_n(t_2) - Bx_n(T)| \rightarrow 0, |Bx_n(T) - Bx_n(t_1)| \rightarrow 0$

and so $|(Bx_n)_{t_2} - (Bx_n)_{t_1}| \rightarrow 0$ as $t_1 \rightarrow t_2$ for all $t_1, t_2 \in \mathcal{R}_+$ (4.13)

Hence $\{Bx_n\}$ is an equicontinuous sequence of functions in $\mathcal{B}(B_r[0])$.

So applying Arzela-Ascoli theorem (see[23]) we say that $\{Bx_n\}$ has a uniformly convergent subsequence in $\mathcal{B}(B_r[0])$ and consequently $\mathcal{B}(B_r[0])$ is a relatively compact subset of X . This shows that \mathcal{B} is compact operator on $B_r[0]$. Hence by Dugungi \mathcal{B} is completely continuous on $B_r[0]$.

Step IV: Next we show that $\mathcal{A}x\mathcal{B}x \in B_r[0]$ for all $x \in B_r[0]$ is arbitrary, then

$$|\mathcal{A}x(t)\mathcal{B}x(t)| \leq |\mathcal{A}x(t)| |\mathcal{B}x(t)|$$

$$|\mathcal{A}x(t)\mathcal{B}x(t)| \leq \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t), x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \right| \left| \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, y(s), y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \right|$$

$$\leq \left[\frac{1}{\Gamma(\alpha)} \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds \right] \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right]$$

$$\leq \frac{v_1(t)}{\Gamma(\alpha)} \left[|q(t)| + \frac{v_2(t)}{\Gamma(\zeta)} \right]$$

$$\leq K_2[K_1 + K_3] = r \text{ for all } t \text{ in } \mathcal{R}_+ \quad (4.14)$$

Taking the supremum over t , we obtain $\|\mathcal{A}x\mathcal{B}x\| \leq r$ for all $x \in B_r[0]$.

Hence $\mathcal{A}x\mathcal{B}x \in B_r[0]$

Hence hypothesis (c) of Theorem (3.3) holds.

Also we have $M = \| \mathcal{B}(B_r[0]) \| = \sup \{ \| \mathcal{B}x \| : x \in B_r[0] \}$

$$\begin{aligned}
&= \sup \{ \underbrace{\sup_{t \geq 0}}_{t \geq 0} \{ |q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{|g(s, y(s)y(\beta(s)))|}{(t-s)^{1-\zeta}} ds \} : x \in B_r[0] \} \\
&= \sup \{ \underbrace{\sup_{t \geq 0}}_{t \geq 0} \{ |q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \} : x \in B_r[0] \} \\
&\leq \underbrace{\sup_{t \geq 0}}_{t \geq 0} \{ |q(t)| + \underbrace{\sup_{t \geq 0}}_{t \geq 0} \frac{v_2(t)}{\Gamma(\zeta)} \} \leq K_1 + K_3 \tag{4.15}
\end{aligned}$$

Therefore $Mk = L(K_1 + K_3) < 1$

Now Applying Theorem 3.3 to shows that (2.1) has a solution on \mathcal{R}_+ .

Step V. In this step we show the local attractivity of the solutions for (2.1).

Let x and y be any two solutions of the (2.1) in $B_r[0]$ defined on \mathcal{R}_+ . Then we have,

$$\begin{aligned}
|x(t) - y(t)| &= \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t)x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, y(t)y(\gamma(s)))}{(t-s)^{1-\alpha}} ds \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \right| \\
|x(t) - y(t)| &\leq \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t)x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \left[q(t) + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, x(s)x(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \right| \\
&\quad + \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, y(t)y(\gamma(s)))}{(t-s)^{1-\alpha}} ds \left[q(t) \right. \right. \\
&\quad \left. \left. + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{g(s, y(s)y(\beta(s)))}{(t-s)^{1-\zeta}} ds \right] \right| \\
|x(t) - y(t)| &\leq \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, x(t)x(\gamma(s)))}{(t-s)^{1-\alpha}} ds \right| \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{|g(s, x(s))|}{(t-s)^{1-\zeta}} ds \right]
\end{aligned}$$

$$\begin{aligned}
& + \left| \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(t, y(t), y(s))}{(t-s)^{1-\alpha}} ds \right| \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{|g(s, y(s))|}{(t-s)^{1-\zeta}} ds \right] \\
& |x(t) - y(t)| \leq \frac{1}{\Gamma(\alpha)} \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right] \\
& \quad + \frac{1}{\Gamma(\alpha)} \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right] \\
& |x(t) - y(t)| \leq 2 \frac{1}{\Gamma(\alpha)} \int_0^t \frac{h_1(s)}{(t-s)^{1-\alpha}} ds \left[|q(t)| + \frac{1}{\Gamma(\zeta)} \int_0^t \frac{h_2(s)}{(t-s)^{1-\zeta}} ds \right] \\
& |x(t) - y(t)| \leq 2 \frac{v_1(t)}{\Gamma(\alpha)} \left[|q(t)| + \frac{v_2(t)}{\Gamma(\zeta)} \right] \tag{4.16}
\end{aligned}$$

For all $t \in \mathcal{R}_+$. Since $\lim_{t \rightarrow \infty} q(t) = 0$ and $\lim_{t \rightarrow \infty} v_i(t) = 0$ this gives that,

$\lim_{t \rightarrow \infty} \sup |x(t) - y(t)| = 0$. Thus the (2.1) has a solution and all the solutions are locally attractive on \mathcal{R}_+ .

5.Conculsion: In this paper we have studied that the attractivity results and existence solutions of quadratic type integral equation of fractional order. However, the method can be applied to other more than nonlinear quadratic integral equation involving integers or Riemann-Louville type fractional order of integration. The research in this direction forms the further scope for the work and some of the results of attractivity and extremal solution of maximal and minimal solutions.

7.References:

- [1] A.A.Kilbas, J.J.Trujillo, Differential equations of fractional order: Methods, results, Problems, I.Appl.Anal. Vol.78 (2001),153-192..
- [2] A.A.Kilbas, Hari M. Srivastava and Juan J.Trujillo, Theory and Applications Fractional Differential equations , North-Holland Mathematics Studies,204, Elsevier Science B.V., Amsterdam ,2006, MR2218073 (2007a:34002).Zbl 1092.45003.
- [3] I.Podlubny, Fractional differential equations, Mathematics in science and engineering, volume 198.
- [4] A.Babakhani, V.Daftardar-Gejii, Existence of positive solutions of nonlinear fractional differential equations, J.Math.Appl. Vol.278 (2003).434-442.
- [5] B.C,Dhage, A Fixed point theorem in Banach algebras involving three operators with applications, Kyungpook Math J. Vol.44 (2004),pp.145-155.

- [6] B.D.Karande, Existence of uniform global locally attractive solutions for fractional order nonlinear random integral equation, Journal of Global Research in Mathematical Archives , Vol.1 (8), (2013) ,34-43.
- [7] B.D.Karande, Fractional Order Functional Integro-Differential Equation in Banach Algebras, Malaysian Journal of Mathematical Sciences, Volume 8(S),(2014), 1-16.
- [8] B.D.Karande, Global attractivity of solutions for a nonlinear functional integral equation of fractional order in Banach Space, AIP Conf.Proc. "10th international Conference on Mathematical Problems in Engineering, Aerospace and Sciences"1637 (2014), 469-478.
- [9] Dugungi, A.Granas, Fixed point Theory, MonographieMath.,Warsaw, 1982.
- [10] H.M.Srivastava,R.K.Saxena, Operators of fractional integration and applications, Appl.Math.Comput. Vol.118 (2006),.147-156.
- [11] I.Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1993.
- [12] J.Banas,B.Rzepka, An application of measures of noncompactness in the study of asymptotic stability, Appl. Math. Lett. Vol.16 (2003),1-6.
- [13] K.S.Miller, B.Ross, An Introduction to the Fractional Calculus and Differential Equations, John Wiley, New York, 1993.
- [14] Lakshmikantham and A.S.Vatsala, Basic theory of fractional differential equations, Nonlinear Analysis, 69(2008),2677-2682.
- [15] M.M.El-Borai and M.I.Abbas, on some integro-differential equations of fractional orders involving caratheodory nonlinearities, Int.J.Modern Math, Vol.2(1) (2007)41-52.
- [16] M.M.El-Borai and M.I.Abbas, Solvability of an infinite system of singular integral equations, SerdicaMath.J. Vol.33 (2007), pp.241-252.
- [17] M.M.El-Borai, W.G.El-Sayed and M.I.Abbas,Monotonic solutions of a class of quadratic singular integral equations of Voltra type, Int.J.Contemp.Math.Sci. Vol.2 (2007),89-102.
- [18] Mohamed I. Abbas, on the existence of locally attractive solutions of a nonlinear quadratic voltra integral equation of fractional order, Advances in difference equations, (2010), 1-11.
- [19] S.Samko, A.A.Kilbas, O.Marichev, Fractional Integrals and Derivatives: Theory and Applications, Gordon and Breach, Amsterdam (1993)
- [20] X.Hu,J.Yan, The global attractivity and asymptotic stability of solution of a nonlinear integral equation, J.Math..Anal. Appl., 321(2006),147-156.
- [21] Shanti Swarup, Integral equations, Krishnaprakashan media(P) Ltd. Meerut, Fourteenth Edition (2006).
- [22] Moulay Rchid Sidi Ammi, El Hassan El Kinani, Delfim F.M. Torres.,Existence and Uniqueness of solutions to a functional Integr0-Differential Equation,Electronic Journal of Differential equation(2012)

- [23]** D.J.Guo and V. Lakshmikantham, Nonlinear problems in Abstract cones, Notes and Reports in Mathematics in Science and engineering , vol.5, Academic press, Massachusetts,1988.
- [24]** S.Heikkila and V. Lakshmikantham, Monotone iterative Techniques for discontinuous Nonlinear Differential Equations, Monographs and Textbooks in Pure and applied mathematics, vol.181, Marcel Dekker, New York, 1994
- [25]** Dhage B.C., A Non-Linear alternative in Banach Algebras with applications to functional differential equations, Non-linear functional Analysis Appl 8, (2004)563-575
- [26]** Dhage B.C., Fixed Point theorems in ordered Banach Algebras and applications Panam Math J 9, (1999) 93-102
- [27]** Hongling Lu, Shurong Sun, Dianwuyamand Houshan Teng , Theory of fractional hybrid differential equations with linear perturbations of second type , Springer
- [28]** R.P. Agarwal, Y. Zhou, Y. He, Existence of fractional neutral functional differential equations, Comput. Math App. 59(3), (2010)1095-1100
- [29]** J. Banas, B.C. Dhage, Globally Asymptotic Stability of solutions of a functional integral equations, Non-linear functional Analysis 69 (7) ,(2008),1945-1952
- [30]** M.A. Krasnosel'skii , Topological Methods in the theory of Nonlinear integral equations, Pergamon Press Book, The Macmillan , new York 1964.
- [31]** B.C.Dhage , On Existence of Extremal solutions of Nonlinear functional Integral equations in Banach Algebras, Journal of applied mathematics and stochastic Analysis 2004:3(2004)271-282.