

Effect Of Rosemary Leaf Ethanol Extract (*Rosmarinus Officinalis L.*) On Arginase-1 And Tumor Suppressor P53 Expression Of Hepatocellular Carcinoma In Male Wistar Strain Rats (*Rattus Norvegicus L.*) Induced By p-Dimethylamino benzaldehyde (DMBA)

ABSTRACT

Aims: Tumor Suppressor p53 is a frequent research target to further understand the mechanisms of cancer development and the pathogenesis of cancer. Arginase-1 may also be a potential prognosis predictor for HCC patients, and also be a new target for HCC treatment. This research aims to scientifically test whether the administration of ethanol extract from Rosemary leaves (*Rosmarinus officinalis L.*) inhibits the increase in Arginase-1 and inhibits the increase in Tumor Suppressor p53 from Hepatocellular Carcinoma in Rats (*Rattus norvegicus L.*) Male Wistar induced by p - Dimethylamino Benzaldehyde (DMBA).

Study design: This study used an in vivo laboratory test method

Place and Duration of Study: Department of Pharmacology and Therapy and Department of Histology, Faculty of Medicine, Udayana University, between December 2023 to July 2024.

Methodology: This study employed true experimental laboratories in vivo test method. Wistar rats induced by p -Dimethylamino Benzaldehyde (DMBA) treatment. Evaluation of normal and negative controls, using ethanol extract of Rosemary leaves for testing with varying doses of 200 mg/kg BW, 400 mg/kg BW, and 800 mg/kg BW with replication carried out 5 times. Observation of Hepatocellular Carcinoma in rat liver tissue samples using immunohistochemical expression of Arginase-1 test and the Tumor Suppressor p53. Interpretation of immunoreactivity was based on Histochemical scoring (H-score) assessment.

Results: The results of the study showed that ethanol extract of rosemary leaves can inhibit the decrease in the expression of tumor suppressor p53 in the group treated with DMBA, with an increase seen at doses of 400 mg/kg BW and 800 mg/kg BW. The results of the hypothesis test study on the immunohistochemical test of arginase-1 and tumor suppressor p53 with One-way ANOVA showed a significant difference between groups ($p < 0.05$).

Conclusion: Ethanol extract of rosemary leaves could not inhibit the increase in arginase-1 expression and inhibit the decrease in tumor suppressor p53 of hepatocellular carcinoma in male Wistar rats induced by p-Dimethylamino benzaldehyde (DMBA).

Keywords: Tumor Suppressor p53, Arginase-1, DMBA, Rosemary leaf

1. INTRODUCTION

The death rate from liver cancer in the United States appears to have increased by about 40% from 1990 to 2004, in contrast to the overall cancer death rate, which decreased by about 18% over the same period [1]. The worldwide occurrence of primary liver cancer is steadily rising, with projections indicating that over one million individuals will be diagnosed with the disease each year by 2025 [2]. Hepatocellular carcinoma (HCC) ranks as the fifth most prevalent cancer globally and the fourth most common among men in South

Korea, where chronic hepatitis B infection is widespread among middle-aged and older individuals [3]. Hepatocellular carcinomas vary widely and are not uniform tumors, they exhibit diverse layers of heterogeneity [4]. The discovery of effective biomarkers for monitoring and early diagnosis of HCC remains inadequate. Current serum biomarkers exhibit low sensitivity and inconsistent specificity, even when evaluated over time or used in combination with other markers, despite employing varying cut-off points [5]. Early detection is crucial for

hepatocellular carcinoma (HCC) because an early diagnosis can greatly enhance the prognosis, which is closely linked to the tumor stage[6]. Liver cancer shows extensive diversity in its development, histopathological traits, and biological behavior. This variation in causative factors significantly contributes to the generally unfavorable prognosis seen in liver cancer patients[7]. The classification of hepatocellular carcinoma (HCC) is mainly based on its morphological features, with histopathological evaluation serving as the gold standard for diagnosing HCC and differentiating it from other possible conditions[1].

L-arginine is a positively charged amino acid that plays a role in various metabolic processes[8]. Arginase 1 (ARG1) and arginase 2 (ARG2), the two mammalian arginase isoforms, are significantly overexpressed in various cancers. They contribute to tumor growth and metastasis through several mechanisms, such as altering L-arginine metabolism and influencing the tumor immune microenvironment[9]. Arginase 1 has a significant role in the cancer microenvironment (tumor microenvironment) and is also a new target for the treatment of HCC[10].

The normal function of tumor suppressor p53 is a powerful barrier to cancer. The tumor suppressor protein p53 plays a key role in regulating the cell cycle and responding to cellular stress, such as exposure to carcinogens (carcinogenic agents)[11]. Loss of normal p53 function due to mutations can allow uncontrolled growth of cancer cells and is a major contributing factor in the development of cancer. Tumor Suppressor p53 is a frequent target of research in an effort to better understand the mechanisms of cancer development and potential cancer therapies. Disruption of the p53 tumor suppressor signaling pathway overcomes the apoptosis checkpoint and causes further cell division in hepatocytes with already shortened telomeres until the telomeres become very short. At this point, the cells enter a crisis checkpoint, characterized by massive cell death. Insufficient telomerase activity leads to accelerated telomere shortening in proliferating liver cells and as a result, genomic instability. Telomere shortening is an important risk factor for tumor development in liver carcinogenesis. The risk of tumor formation increases dramatically in the cirrhosis stage, which is characterized by increased apoptosis in hepatocytes, and further cell division in the crisis stage through hepatocyte apoptosis[7].

Anticancer therapy for hepatocellular carcinoma is currently not as much as other anticancer therapies. One of the recommended first-line anticancer drugs for hepatocellular carcinoma is Sorafenib and Lenvatinib. These drugs have various limitations in the side effects they cause and drug resistance often occurs. Side effects occur because the selectivity of the drug works at many drug target sites[12]. This problem underlines the importance of developing anticancer drugs for therapy in hepatocellular carcinoma.

Medicinal plants that are efficacious in cancer therapy are rosemary. Rosmarinic acid has also been studied in animal models of Hepatocarcinoma with biomarkers TNF- α , TGF- β and IFN- γ [13]. In addition, there has been no testing and development of this compound as an antitumor drug on the expression of Arginase-1 and tumor suppressor p53.

Potential as a useful material as an anticancer and phytochemicals taken for example phenol and flavonoids. Rosmarinic acid has solubility can dissolve in ethanol so this study uses ethanol extract as a material in its testing. Rosmarinic acid is a compound that can be found in nature especially in rosemary leaves. Several studies have confirmed the various therapeutic benefits of rosmarinic acid in various diseases, including cancer, diabetes, inflammatory disorders, neurodegenerative disorders, and liver disease. Rosmarinic acid is a bioactive phenolic compound commonly found in plants of the Lamiaceae and Boraginaceae families. Rosmarinic acid is biosynthesized using the amino acids tyrosine and phenylalanine through reactions catalyzed by enzymes.

This study aims to scientifically test the administration of rosemary leaf ethanol extract in inhibiting the increase in arginase-1 expression and inhibiting the decrease in tumor suppressor p53 from hepatocellular carcinoma in male Wistar rats (*Rattus norvegicus* L.) induced by p-Dimethylamino benzaldehyde (DMBA).

2. MATERIAL AND METHODS

2.1 MATERIALS

The tools used in this study include analytical scales, stainless spoons, ovens, *object glasses*, *cover glasses*, droppers, mouse cages, mouse drinking bottles, husks, Hardware (Asus Vivo Book Flip 14 Laptop with

Intel Celeron N4020 processor specifications, 4GB RAM, 64-bit operating system, window 2019), SPSS 26 software , mouse food, gloves, masks, 1 cc and 3 cc syringes, probes, mouse fixation tools, ketamine, data loggers, cameras, microscopes , styrofoam boxes, plastic tissue pots, plastic bottles, scalpels, tweezers, strainers, tissue cassettes, automatic processor machines, vacuum machines, blocking machines, microtome machines, microtome knives, 46 ° C water baths, object glasses, cover slips, special staining racks, 60 ° C ovens.

2.2 METHODOLOGY

The type of research used is true experimental laboratories in vivo. The research design applied for oral administration of rosemary leaf ethanol extract (*Rosmarinus officinalis* L.) inhibits hepatocellular carcinoma levels in this study is a post test only control group design.

2.3 SAMPLE

In this study, specimens were collected from rat that met the inclusion and exclusion criteria with the following provisions:

2.3.1 INCLUSION CRITERIA

- Male Wistar Rat (*Rattus norvegicus* L.)
- Rat weight 100-250 grams
- Healthy rats aged 2-3 months.

2.3.2 EXCLUSION CRITERIA

- Rats don't want to eat or drink
- Rats died and liver tissue could not be taken.

2.4 PROCEDURES

This study is a true experimental laboratories in vivo. This research protocol has met the ethical principles of research, the research protocol was reviewed and approved by the Udayana University ethics committee with a letter or Ethical Clearance document Number: B / 259 / UN14.2.9 / PT.01.04 / 2023. This research protocol has received approval

for the implementation of research from the Integrated Biomedical Laboratory with a letter or document Number: 1568 / UN14.2.2.VII.6 / LT / 2023.

- group of rats were induced with DMBA twice a week for 5 weeks at a dose of 25 mg/kg BW. DMBA was administered orally using a probe. DMBA was dissolved in corn oil in a Bio Safety Cabinet for the carcinogenic induction process.
- Normal rat group and DMBA-induced rat group were given pellets and drink ad libitum during the treatment process. The waiting period for cancer growth was 2 weeks.
- Group of rats that met the inclusion criteria with DMBA induced treatment of 25 mg/kg BW for 2 times a week for 5 weeks. Waited for 2 weeks for tumor formation with treatment given food and water ad lib during the study. Continued by giving rosemary leaf ethanol extract at a dose of 200 mg/kg BW , 400 mg/kg BW, 800 mg/kg BW once a day for 20 days
- Final treatment all rats (*Rattus norvegicus* L.) were sacrificed with ketamine. Liver organ was taken for Immunohistochemistry measurement of Tumor Suppressor p53 expression.

3. RESULTS AND DISCUSSION

The organoleptic properties of rosemary leaf ethanol extract were identified visually. The observation results showed that the rosemary leaf ethanol extract had a blackish brown color. If the extract is open and at room temperature and exposed to light, the extract can absorb oxygen from the air so that it can produce a darker color. Phytochemical screening tests were conducted with the aim of determining secondary metabolite compounds and bioactive components contained in the ethanol extract of rosemary leaves.

Table 1. Results of the Normality Test for Body Weight of Research Rats

Sampel	Fenol (mg/100g)	Flavonoid (mg/100g)	IC 50 (ppm)	Antioxidant capacity (mg/L GAEAC)
Ekstrak Rosemary	4880,70	51348,04	216,0088	4452,90




Fig 1. Arginase-1 expression and p53 wild type expression compare to normal liver and control negative sample. A. Negative Arginase-1 expression from normal liver. B. Negative p53 wild type expression from normal liver. C. Negative Arginase-1 expression from control negative sample. D. Negative p53 wild type expression from control negative sample. E. Negative arginase-1 expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 200 mg/kg BW. F. Negative p53 expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 200 mg/kg BW. G. Negative arginase-1 expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 400 mg/kg BW. H. Negative p53 wild type expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 400 mg/kg BW. I. Positive arginase-

1 expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 800 mg/kg BW. J. Positive p53 wild type expression from liver DMBA induced with treatment rosemary leaf ethanol extract at a dose of 800 mg/kg BW.




Fig. 2 Descriptive analysis results of Arginase-1 Immunohistochemistry H-score Data (indicates negative arginase-1 expression and abnormal data from group control normal, control negative, Rosemary 200mg/kg BW, Rosemary 400mg/kg BW with P-value < 0.001, indicates positive arginase-1 expression and normal data from rosemary 800mg/kg BW group with P-value > 0.01).

Table 2. Results of the Homogeneity Test and One-way ANOVA Test Results of Arginase-1 Immunohistochemistry H-score Data

Variable	Levene statistic	p	Result	One-way Anova p	Result
ARG 1 Expression	2,083	0,247	Homogen	0,02	Significant differences

Table 3. Post Hoc Test Results with Least Significantly Different (LSD) H-score Data of Arginase-1 Immunohistochemistry

Variable	p	Result	
Normal Control	Negative Control	1,000	Not Significant
Normal Control	Rosemary 200mg/kg BW	1,000	Not Significant
Normal Control	Rosemary 400mg/kg BW	1,000	Not Significant
Normal Control	Rosemary 800mg/kg BW	0,006	Significant differences
Negative Control	Rosemary 200mg/kg BW	1,000	Not Significant
Negative Control	Rosemary 400mg/kg BW	1,000	Not Significant
Negative Control	Rosemary 800mg/kg BW	0,006	Significant differences
Rosemary 200mg/kg BW	Rosemary 400mg/kg BW	1,000	Not Significant
Rosemary 200mg/kg BW	Rosemary 800mg/kg BW	0,006	Significant differences
Rosemary 400mg/kg BW	Rosemary 800mg/kg BW	0,006	Significant differences




Fig. 3 Descriptive analysis results of p53 Wild Type Immunohistochemistry H-score Data (indicates negative p53 Wild Type expression and abnormal data from group control normal with P-value < 0.001, indicates positive p53 Wild Type expression and normal data from control negative, Rosemary 200mg/kg BW, Rosemary 400mg/kg BW, Rosemary 800mg/kg BW group with P-value > 0.01).

Table 4. Results of the Homogeneity Test and One-way ANOVA Test Result of p53 wild type Immunohistochemistry H-score Data

Variable	Levene statistic	p	Result	One-way Anova p	Result
P53 wild type Expression	4,935	0,053	Homogen	0,000	Significant differences

Table 5. Post Hoc Test Results with Least Significantly Different (LSD) H-score Data of p53 wild type Immunohistochemistry

Variable	p	Result	
Normal Control	Negative Control	0,837	Not Significant
Normal Control	Rosemary 200mg/kg BW	0,415	Not Significant
Normal Control	Rosemary 400mg/kg BW	0,000	Significant differences
Normal Control	Rosemary 800mg/kg BW	0,000	Significant differences
Negative Control	Rosemary 200mg/kg BW	0,539	Not Significant
Negative Control	Rosemary 400mg/kg BW	0,001	Significant differences
Negative Control	Rosemary 800mg/kg BW	0,000	Significant differences
Rosemary 200mg/kg BW	Rosemary 400mg/kg BW	0,003	Significant differences
Rosemary 200mg/kg BW	Rosemary 800mg/kg BW	0,000	Significant differences
Rosemary 400mg/kg BW	Rosemary 800mg/kg BW	0,229	Not Significant

Anticancer therapy for hepatocellular carcinoma is currently not as much as other anticancer therapies. Rosemary for biological activities, such as antibacterial, anticancer, antidiabetic, anti-inflammatory and antinociceptive, antioxidant, antithrombotic, antiulcerative, improving cognitive deficiency, antidiuretic, and hepatoprotective effects are the prospects for the use of rosemary as an agent in the treatment of various diseases. Rosemary has long been considered as a herb and occupies a special place in traditional medicine. Phenolic diterpenes, triterpenes, phenolic acids, such as carnosic acid (CA), carnosol, rosmanol, ursolic acid, betulinic acid, and rosmarinic acid (RA), and nepitrin are the pharmacologically active constituents identified in Rosemary. Isolated phenolic compounds in rosemary, carnosic acid and rosmarinic acid have been shown to have the most common pharmacological effects and interact with several molecular targets[14].

Literature review study of existing in vitro and in vivo studies focused on the anticancer effects of rosemary extract and rosemary extract polyphenols namely carnosic acid and rosmarinic acid, and their effects on key signaling molecules. The main polyphenols found in rosemary extract include the diterpenes carnosic acid and rosmarinic acid. In vitro and in vivo studies on the effects of rosemary and its major polyphenols have been summarized and sorted by cancer cell type, in chronological order from earliest to latest. The anticancer effects of rosemary extract include several in vitro studies using colon cancer cell lines have shown rosemary to exhibit anticancer properties. Rosemary extract is summarized to have effects in increasing the tumor suppressor p53 as a proapoptotic[15].

The selection of multilevel maceration method in this study was to separate the antioxidant compounds contained in the rosemary leaf sample from other substances according to their polarity level. The main constituents of rosemary leaves, namely phenol and flavonoids, have solubility that can dissolve in ethanol solvents although they are insoluble in n-hexane and slightly soluble in ethyl acetate. The results of the analysis of rosemary leaf ethanol extract using the multilevel maceration method in this study were organoleptically in the form of a thick liquid, reddish brown in color and a distinctive odor of rosemary leaves. Testing of flavonoid, phenol, IC50 levels, and antioxidant capacity of rosemary leaf ethanol extract was carried out as confirmation of the content of secondary

metabolite compounds from the rosemary leaf ethanol extract made.

The antioxidant capacity value in the ethanol extract of rosemary leaves obtained results of 4452.90 mg/L GAEAC and the IC50 value was found to be 216.0088 ppm. The IC50 value shows a concentration of 216.0088 ppm where the extract inhibits 50% of free radicals with an antioxidant capacity of 4452.90 mg/L GAEAC. The IC50 value in this study shows that a large concentration of extract is needed to achieve 50% free radical inhibition. Although, the antioxidant capacity measured in mg/L GAEAC (Antioxidant Capacity Equivalent to Gallic Acid) has a very high value, indicating strong overall antioxidant activity. The antioxidant content is increased by administering 3 high dose variants. This study used a dose of rosemary ethanol extract of 200 mg/kg BW, 400 mg/kg BW and 800 mg/kg BW.

Albino rats (*Rattus norvegicus* L.), better known as laboratory rats, are used as a research model in biomedicine. Rats can represent mammalian biological systems, making them very suitable for preclinical research. The reproductive period of rats can be determined by observing their various life stages and behaviors. The liver of Wistar rats has a structure and function similar to the liver of other mammals, used in biomedical research to study liver function, metabolism, and the effects of various compounds or drugs.

a. Effect of Rosemary Leaf Ethanol Extract on Arginase-1 Expression of Hepatocellular Carcinoma in Male Wistar Rats Induced by p-Dimethylamino benzaldehyde (DMBA).

ARG1 contributes to the development of hepatocellular carcinoma by promoting carcinogenesis. Elevated ARG1 expression is associated with more aggressive tumor growth, larger tumor size, and increased levels of alanine aminotransferase (ALT) and glutamyl transferase (GGT)[9]. Immunosuppressive macrophages are traditionally identified by the expression of the enzyme Arginase 1 (ARG1), which we observed to be highly expressed in pancreatic tumor-associated macrophages in both human patients and mouse models[16].

Arginase 1 also plays a role in modulating the immune system, primarily through its

effects on arginine metabolism. Arginase 1 is expressed by alternatively activated macrophages (M2) and myeloid suppressor cells (MDSCs). This enzyme catalyzes the depletion of arginine in the microenvironment, which can suppress T cell activity and facilitate anti-inflammatory or immunosuppressive responses. High expression of arginase 1 by tumor cells or stromal cells in the tumor microenvironment can lead to local immunosuppression, allowing tumors to evade detection by the immune system. Arginase 1 is considered a potential target for immune cancer therapy in this study. Decreased Arginase 1 function in chronic liver diseases, such as cirrhosis or hepatitis, may impair ammonia detoxification and slow hyperammonemia. Further study of Arginase 1 may pave the way for novel therapies in these conditions.

Arginase 1 is often used as a marker to identify specific cell types and enzymatic activities in various tissues, especially in the liver. Arginase 1 is highly expressed in hepatocytes, the primary cells of the liver. In liver histology, immunohistochemistry (IHC) staining for arginase 1 will show a broad and intense distribution throughout the liver tissue. Immunohistochemistry is a technique used to detect arginase 1 in tissue samples. This technique uses specific antibodies that bind to arginase 1, which is then visualized through staining. In IHC, arginase 1 is often stained using monoclonal or polyclonal antibodies that impart a specific color (usually brown or red) at the location of the enzyme in the tissue. Arginase 1 is used as a marker to diagnose cellular hepatocarcinoma (HCC), a liver cancer. High expression of arginase 1 in a liver biopsy can help distinguish HCC from liver metastases or other liver tumors. Other histological staining techniques such as H&E (Hematoxylin and Eosin) staining may not specifically demonstrate arginase 1, although they can be used in conjunction with IHC to provide a general picture of tissue morphology and structure.

In rats treated with DMBA and rosemary 800 mg/kg BW with arginase-1 expression, the effect of the combination treatment was accompanied by an H-score value and there was a significant difference with other groups. In other

samples, no arginase-1 expression was found and there was no significant difference between the normal control group, negative control, rats treated with DMBA with rosemary 200 mg/kg BW and rosemary 400 mg/kg BW. The phenomenon in this study explains that it can damage cell structure with 10 doses of DMBA and a longer exposure time is needed for the development of hepatocellular carcinoma formation. The assessment by immunohistochemical test is still qualitative, so it requires quantitative testing such as ELISA for further research. Arginase-1 immunohistochemistry in this study cannot be concluded because of weaknesses in this study, namely the method of cutting the preparations only once so that it does not ensure the formation of cancer.

b. Effect of Rosemary Leaf Ethanol Extract on the Expression of Tumor Suppressor p53 of Hepatocellular Carcinoma in Male Wistar Rats Induced by p-Dimethylamino benzaldehyde (DMBA).

The tumor suppressor gene P53 is one of the most frequently inactivated tumor suppressors in human cancers. The function of p53 during cancer development has been associated with a variety of transcriptional and non-transcriptional activities that lead to tight control of cell proliferation, senescence, DNA repair, and cell death. However, evidence suggests that the tumor suppressor p53 also plays a major role in normal and cancer cell metabolism[17].

Cancer cells often show various abnormalities or damage to the nucleus, cytoplasmic organelles, and cytoskeleton. The nucleus tends to be enlarged and irregularly shaped, with chromosome breaks, deletions, and translocations. The mitotic rate is usually increased. In the cytoplasm, intracellular structures appear disorganized and change in size and shape. Changes in microtubules, which support the cell and are essential for controlling nearly all cellular functions, have a major impact. Mitochondria also become disorganized and irregularly shaped[18].

P53 causes accumulation of various mutations. P53 functions to detect DNA damage and activate repair mechanisms.

Genomic instability also causes increased epigenetic silencing or modulation of gene function. Increased DNA methylation in the promoter region of tumor suppressor genes is also found in many types of cancer. Increased DNA methylation is also associated with changes in histone modifications in chromatin and often correlates with DNA methylation. Changes in the promoter region of genes cause gene deletion or changes in gene expression [19].

In this study, the biomarker studied was Tumor suppressor p53. This biomarker analysis was conducted to understand the expression of tumor suppressor p53 and its relationship to the development of hepatocellular carcinoma (HCC). The results of immunohistochemistry tests in this study showed the accumulation of tumor suppressor p53 in the cytoplasm, not entering the nucleus of liver tissue in mice that received rosemary leaf ethanol extract. These results indicate that tumor suppressor p53 is inactive. The results of other liver tissue immunohistochemistry tests showed the presence of active tumor suppressor p53 (brown), where the location of this active biomarker was at the edge of the tumor cell diameter of the liver tissue sample. Liver tissue in hepatocellular carcinoma model mice in this study showed abnormal cell shapes. During development, normal cells will differentiate. Differentiation means that a cell becomes specialized in its structure and function, and gathers with similarly differentiated cells. The more specialized a cell is, the higher the differentiation, the less often the cell enters the cell cycle and reproduces and divides. Cells that do not reproduce are highly differentiated cells.

The results of the immunohistochemical test of this study showed that the expression of tumor suppressor p53 differed between the normal group and the group treated with DMBA and rosemary ethanol extract. This study showed that Rosemary ethanol extract could inhibit the decrease in tumor suppressor p53 expression significantly in the group treated with DMBA, with an increase seen at doses of 400 mg/kg BW and 800 mg/kg BW. Rosemary leaf ethanol extract can inhibit the decrease in tumor suppressor p53 expression

because it has strong overall antioxidant activity.

4. CONCLUSION

The effect of 800 mg/kg BW of rosemary leaf ethanol extract could not inhibit the increase in arginase-1 expression of hepatocellular carcinoma in male Wistar rats induced by p-Dimethylamino benzaldehyde (DMBA). Ethanol extract of rosemary leaves 400 mg/kg BW and 800 mg/kg BW inhibited the decrease in the expression of the tumor suppressor p53 from hepatocellular carcinoma in male Wistar rats induced by p-Dimethylamino benzaldehyde (DMBA).

DISCLAIMER(ARTIFICIALINTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

This study is a true experimental laboratories in vivo. This research protocol has met the ethical principles of research, the research protocol was reviewed and approved by the Udayana University ethics committee with a letter or Ethical Clearance document Number: B / 259 / UN14.2.9 / PT.01.04 / 2023. This research protocol has received approval for the implementation of research from the Integrated Biomedical Laboratory with a letter or document Number: 1568 / UN14.2.2.VII.6 / LT / 2023.

REFERENCES

- [1] P. R. Galle *et al.*, "EASL Clinical Practice Guidelines: Management of hepatocellular carcinoma," *J. Hepatol.*, vol. 69, no. 1, pp. 182–236, 2018, doi: 10.1016/j.jhep.2018.03.019.
- [2] K. Oura, A. Morishita, S. Hamaya, K. Fujita, and T. Masaki, "The Roles of Epigenetic Regulation and the Tumor Microenvironment in the Mechanism of Resistance to Systemic Therapy in Hepatocellular Carcinoma," *Int. J. Mol. Sci.*, vol. 24, no. 3, 2023, doi: 10.3390/ijms24032805.

- [3] J. W. Park, 2022 *KLCA-NCC Korea practice guidelines for the management of hepatocellular carcinoma*, vol. 28, no. 4. 2022. doi: 10.3350/cmh.2022.0294.
- [4] M. S. Torbenson, "Hepatocellular carcinoma: making sense of morphological heterogeneity, growth patterns, and subtypes," *Hum. Pathol.*, vol. 112, pp. 86–101, 2021, doi: 10.1016/j.humpath.2020.12.009.
- [5] F. Piñero, M. Dirchwolf, and M. G. Pessôa, "Biomarkers in Hepatocellular Carcinoma: Diagnosis, Prognosis and Treatment Response Assessment," *Cells*, vol. 9, no. 6, pp. 1–27, 2020, doi: 10.3390/cells9061370.
- [6] E. Shahini, G. Pasculli, A. G. Solimando, C. Tiribelli, R. Cozzolongo, and G. Giannelli, "Updating the Clinical Application of Blood Biomarkers and Their Algorithms in the Diagnosis and Surveillance of Hepatocellular Carcinoma: A Critical Review," *Int. J. Mol. Sci.*, vol. 24, no. 5, pp. 1–39, 2023, doi: 10.3390/ijms24054286.
- [7] et al Stroth, "Telomeres and Telomerase in the Development of Liver Cancer," *Cancers (Basel)*, 2020.
- [8] T. M. Grzywa, A. Sosnowska, P. Matryba, and Z. Rydzynska, "Myeloid Cell-Derived Arginase in Cancer Immune Response," *Front. Immunol.*, vol. 11, no. May, pp. 1–24, 2020, doi: 10.3389/fimmu.2020.00938.
- [9] F. Niu *et al.*, "Biomedicine & Pharmacotherapy Arginase : An emerging and promising therapeutic target for cancer treatment," *Biomed. Pharmacother.*, vol. 149, no. March, 2022, doi: 10.1016/j.biopha.2022.112840.
- [10] W. Chen, J. Chen, Q. Zheng, J. Dong, and Y. Zhu, "The Oncogenic Role of ARG1 in Progression and Metastasis of Hepatocellular Carcinoma," *Biomed Res. Int.*, vol. 2018, 2018, doi: 10.1155/2018/2109865.
- [11] J. Hu *et al.*, "Targeting mutant p53 for cancer therapy: direct and indirect strategies," *J. Hematol. Oncol.*, vol. 14, no. 1, pp. 1–19, 2021, doi: 10.1186/s13045-021-01169-0.
- [12] Y. Zhu, B. Zheng, H. Wang, and L. Chen, "New knowledge of the mechanisms of sorafenib resistance in liver cancer," *Acta Pharmacol. Sin.*, pp. 614–622, 2017, doi: 10.1038/aps.2017.5.
- [13] C. Gonçalves, D. Fernandes, I. Silva, and V. Mateus, "Potential Anti-Inflammatory Effect of Rosmarinus officinalis in Preclinical In Vivo Models of Inflammation," *Molecules*, vol. 27, no. 3, 2022, doi: 10.3390/molecules27030609.
- [14] S. M. Hussain *et al.*, "Cognition enhancing effect of rosemary (Rosmarinus officinalis L.) in lab animal studies: a systematic review and meta-analysis," *Brazilian J. Med. Biol. Res.*, vol. 55, pp. 1–14, 2022, doi: 10.1590/1414-431X2021e11593.
- [15] J. Moore, M. Yousef, and E. Tsiani, *Anticancer effects of rosemary (Rosmarinus officinalis L.) extract and rosemary extract polyphenols*, vol. 8, no. 11. 2016. doi: 10.3390/nu8110731.
- [16] R. E. Menjivar *et al.*, "Arginase 1 is a key driver of immune suppression in pancreatic cancer," pp. 1–25, 2023.
- [17] M. Lacroix, R. Riscal, G. Arena, L. K. Linares, and L. Le Cam, "Metabolic functions of the tumor suppressor p53: Implications in normal physiology, metabolic disorders, and cancer," *Mol. Metab.*, vol. 33, no. October 2019, pp. 2–22, 2020, doi: 10.1016/j.molmet.2019.10.002.
- [18] El. J. Corwin, "Buku Saku Patofisiologi." pp. 541–2, 2009.
19. Huether, Sue E., McCance, Kathryn L., Brashers, Valentina L., Rote, Neal S., 2019. Buku Ajar Patofisiologi. Edisi Ke-Enam. 216-248. Singapore. Elsevier.