

Residue Retention in a Maize-Chickpea Cropping System under Conservation Agriculture Influence Soil biochemical attributes in Vertisol of Central India

ABSTRACT

This study examined the biochemical properties of vertisol under residue retention (RR) practices of 0%, 30%, and 90% in a maize-chickpea cropping system under conventional tillage (CT). The parameters assessed were total organic carbon (TOC), β -glucosidase activity, dehydrogenase activity (DHA), fluorescein diacetate hydrolysis activity (FDA), and stratification ratio. Soil samples were collected from 0–15 cm and 15–30 cm depths at the end of the cropping cycle to evaluate the effects of different residue management strategies. At the surface soil (0–15 cm), TOC was significantly higher under 90% RR (15.24 g kg⁻¹) and 30% RR (12.16 g kg⁻¹) compared to CT (9.39 g kg⁻¹). Enzymatic activities also showed significant improvements with increased residue retention. DHA at 0–15 cm was highest under 90% RR (103.57 μ g TPF g⁻¹ day⁻¹), followed by 30% RR (84.63 μ g TPF g⁻¹ day⁻¹) and CT (70.75 μ g TPF g⁻¹ day⁻¹). A similar trend was observed for FDA, where 90% RR recorded 26.13 μ g fluorescein g⁻¹ h⁻¹, exceeding CT (22.91 μ g fluorescein g⁻¹ h⁻¹). β -glucosidase activity was also highest under 90% RR (169.60 μ g PNG g⁻¹ soil h⁻¹), with reduced values at greater soil depths. TOC and enzymatic activities demonstrated strong positive correlations, confirming their sensitivity to management practices. Stratification ratios did not vary significantly across residue retention levels, likely due to the high clay content protecting TOC and enzymes. These findings highlight the potential of residue retention to enhance soil health and serve as reliable indicators of soil quality in sustainable cropping systems.

Keywords: Residue retention, Tillage, cropping system, Total organic carbon, soil enzymatic activity,

INTRODUCTION

Crop residue retention is a key element of conservation agriculture (CA), alongside (a) minimal mechanical soil disturbance, (b) permanent soil cover, and (c) crop rotation, as highlighted by Ojeda *et al.* (2015) and Ye *et al.* (2019). Crop residues help mitigate the negative impacts of conventional farming practices and enhance soil quality, facilitating better adaptation to climate-related risks (Das *et al.*, 2020; Thierfelder *et al.*, 2018). Land degradation continues to be a significant global issue, particularly in India, where approximately 44% of the total land area is affected (Bhattacharyya *et al.*, 2015; Mythili & Goedecke, 2016). The incorporation of crop residues enhances soil organic matter (OM), conserves soil moisture, and fosters biological activity (Huang, Xu, & Chen, 2008). Residues, especially from staple crops like maize and chickpea, provide a vital source of organic carbon that can stimulate microbial populations involved in soil carbon cycling. Maize (*Zea mays L.*), an emerging versatile crop with wider adaptability and photo-insensitivity under the different ecological scenario. It has the potential to address issues such as, water scarcity and climate change (Parihar *et al.* (2018).

Similarly, chickpea (*Cicer arietinum*) is a protein-rich and best among all legume proteins, with the most production centered in India. Maize-based rotations with improved soil management practices enhanced soil properties Aulakh *et. al* (2008).

Concerns regarding soil deterioration and quality losses have recently increased the significance of soil quality indices, which enable the evaluation of patterns and modifications in various soil management techniques. The agroecosystem's productivity and sustainability are determined by the quality of the soil. To get empirical data on how conservation techniques impact soil qualities, a residue retention management experiment is consequently required. To ascertain the impact of various soil management strategies, a number of indices that combine the physical, chemical, and biological characteristics of soil have been employed (Doran *et al.* 1994). Analyzing soil biological processes is crucial for assessing soil quality. A good short-term predictor of soil biological and biochemical fertility is microbial activity (Melero *et al.*, 2008; Nannipieri *et al.*, 1990). Soil enzymes play a crucial role in driving numerous reactions related to the breakdown of soil organic matter, nutrient cycling, and the formation of soil structure. They are also considered indicators of soil health, as they respond quickly to changes in soil management practices (Gianfreda *et al.*, 1996). The stratification ratio serves as a useful measure for assessing the variation of soil nutrients with depth. The distribution of soil organic carbon across different depths has gained significance due to its role in nutrient retention, boosting biological activity, preventing erosion, and supporting agricultural productivity (Franzuebbers *et al.*, 2007).

We hypothesize that by improving soil microbial activity and soil fertility, residue retention would be beneficial. Total organic carbon (TOC), dehydrogenase (DHA), fluorescein diacetate (FDA), and β -glucosidase were evaluated in soils with varying levels of residue retention in comparison to soils managed using conventional tillage.

MATERIALS AND METHODS

Experimental Site

The study was conducted at the research farm of the Indian Institute of Soil Science (IISS) in Bhopal, India. The experimental site is geographically situated at coordinates 23°18'28.26"N and 77°24'26.00"E, with an elevation of 500 meters above sea level. The area receives an average annual rainfall of 1,146 mm, with more than 80% of it occurring between June and September. The region experiences an average annual air temperature of 25°C and has a generally humid subtropical climate. The summer season starts in the latter half of March and lasts until mid-June, while winter peaks in January, with temperatures occasionally dropping close to freezing at night. The soil at the experimental site is classified as Vertisols (black soils) from the montmorillonite isohyperthermic family of typic haplustert, characterized by its alkaline nature and distinct swell-shrink properties (Aher *et al.*, 2018).

The experiment was conducted using a Randomized Block Design (RBD), with each treatment replicated five times and randomized. Each plot measured 7m x 6m. The treatments included residue retention (RR) of (0%), 30%, and 90% in a maize and chickpea cropping system under zero tillage, and conventional tillage management. After harvesting both maize and chickpea, the residues were collected, chopped, and left on the soil surface prior to planting the next crop in each cycle. Nutrient supply was primarily through urea (46% N), Di-ammonium Phosphate (DAP: 18% N and 46% P), and Muriate of Potash (MOP: 60% K). The recommended fertilizer doses were 120:24:33 kg of N:P:K per hectare for maize and 40:26:24 kg of N:P:K per hectare for chickpea, applied during each cropping season.

Soil Sampling and Chemical Analysis

At the end of second year experiment, soil samples were collected from two different depths: 0-15 cm and 15-30 cm, after the harvest of each maize and chickpea crop in 2022-2024. The freshly collected

soil samples were passed through a 2 mm sieve and immediately stored in plastic bags, loosely tied to allow proper aeration and prevent moisture loss, at 4°C until microbiological and enzyme activity assessments. The remaining soil was air-dried for chemical analysis, which was conducted within two weeks.

Soil pH and electrical conductivity (EC) were assessed using the 1:2.5 soil-to-water ratio method. Total organic carbon content in the soil was analysed through dry combustion with a TOC analyser.

Soil Enzymatic Analysis

β -glucosidase activity was evaluated following the method of Eivazi and Tabatabai (1988), by incubating soil with p-nitrophenyl β -D-glucopyranoside and measuring p-nitrophenol absorbance at 400 nm. Dehydrogenase activity was assessed using the procedure described by Thalmann (1968), through soil incubation with 2,3,5-triphenyl-tetrazolium chloride (TTC) and measuring the absorbance of triphenyl formazan (TPF) at 546 nm. Fluorescein diacetate (FDA) hydrolysis was determined based on the method of Adams and Duncan (2001), by incubating soil with fluorescein diacetate and recording fluorescein absorbance at 490 nm.

The stratification ratio for total organic carbon and enzymatic activities was determined as the ratio of their values in the surface soil layer (0–15 cm) to those in the deeper layer (15–30 cm), following the method outlined by Franzluebbers (2002). All microbial analyses were performed in triplicate for each sample, and the results were expressed on an oven-dry weight basis.

Statistical Analysis

Statistical analysis was conducted using MS-Excel, and the results were reported as mean values. Significant differences between residue retention management practices were determined using the student's t-test at ($p = 0.05$). Analysis of covariance (ANOVA) was performed to evaluate the variability of all parameters for each treatment across different soil depths. A correlation matrix for various properties was constructed based on Pearson correlation coefficients ($p = 0.05$).

RESULT

Soil Chemical Parameters

At the end of the 2022–2024 cropping season, soil pH, EC, and TOC in residue retention practices were measured at both depths (0–15 cm and 15–30 cm). Across all treatments in both cropping years, the pH and EC of the soil were found to be significant at the surface and non-significant in the subsurface (Table 1). The mean soil pH range at the surface (0–15 cm) ranged from 7.5 to 7.9, with a lower value (7.5) seen under 90% crop residue retention treatment. Although there was a slight rise in pH in subsurface soil, residue retention treatments did not noticeably affect this parameter. The average soil EC range was 0.13 dS/m to 0.23 dS/m at the surface (0–15 cm), with the 90% crop residue retention treatment having the lowest value (0.13 dS/m). However, the mean values of pH and EC at both depths do not significantly differ in the case of no residue and CT.

The distribution of total organic carbon (TOC) varied under different residue retention (RR) levels and conventional tillage (CT) (Table 1). Residue retention significantly increased TOC at both 0–15 cm and 15–30 cm soil depths in the maize-chickpea cropping system. The increase in TOC due to residue retention management ranged from 38.4% to 10.3% at the 0–15 cm depth and from 18.0% to 4.2% at the 15–30 cm depth. After two years of conservation agriculture, TOC levels under 90%, 30%, and 0% RR were 38.4%, 22.8%, and 10.3% higher, respectively, than those under CT at the 0–15 cm depth. TOC values ranged from 15.24 g kg⁻¹ (0–15 cm) to 8.66 g kg⁻¹ (15–30 cm) under 90% RR, 12.16 g kg⁻¹ (0–15 cm) to 7.88 g kg⁻¹ (15–30 cm) under 30% RR, 10.47 g kg⁻¹ (0–15 cm) to 7.41 g kg⁻¹ (15–30 cm) under 0% RR, and 9.39 g kg⁻¹ (0–15 cm) to 7.10 g kg⁻¹ (15–30 cm) under CT. The mean TOC values showed a decreasing trend with increasing soil depth, with the highest TOC concentration recorded at the 0–15 cm depth under 90% RR.

Dehydrogenase (DHA), Fluorescein Diacetate (FDA) and B-Glucosidase Activity

Significant variations in dehydrogenase activity (DHA) was observed in both surface and subsurface soils under different levels of residue retention (RR) compared to conventional tillage (CT) (Fig 1). At the 0–15 cm depth, DHA activity ranged from 103.57 to 70.55 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$, while at the 15–30 cm depth, it varied from 64.19 to 44.77 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$. The increase in DHA ranged from 31.88% to 4.14% at 0–15 cm and from 30.3% to 2.9% at 15–30 cm. DHA levels under 90%, 30%, and 0% RR were 31.8%, 19.07%, and 4.14% higher, respectively, than those under CT at the 0–15 cm depth. DHA values ranged from 103.57 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (0–15 cm) to 64.19 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (15–30 cm) under 90% RR, 87.18 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (0–15 cm) to 53.06 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (15–30 cm) under 30% RR, 73.60 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (0–15 cm) to 46.12 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (15–30 cm) under 0% RR, and 70.55 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (0–15 cm) to 44.77 $\mu\text{g TPF g}^{-1} \text{ soil } 24 \text{ h}^{-1}$ (15–30 cm) under CT. The mean DHA values showed a decreasing trend with increasing soil depth, with the highest DHA concentration recorded under 90% RR at the 0–15 cm depth. In the subsurface, DHA activity was significantly higher under 90% RR, showing a 59.6% increase compared to CT.

Statistical analysis revealed a significant effect of different residue retention (RR) on fluorescein diacetate (FDA) activity (Fig. 1). FDA activity ranged from 26.13 to 16.27 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ at the 0–15 cm depth and from 11.97 to 8.57 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ at the 15–30 cm depth. The increase in FDA activity due to RR management ranged from 37.8% to 17.2% at 0–15 cm and from 28.4% to -4.4% at 15–30 cm. FDA activity under 90%, 30%, and 0% RR was 37.8%, 12.6%, and 17.2% higher, respectively, than under CT at the 0–15 cm depth. FDA ranged from 26.13 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 11.97 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 90% RR, 18.62 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 8.81 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 30% RR, 19.64 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 8.21 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 0% RR, and 16.27 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 8.57 $\mu\text{g fluorescein g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under CT. The mean FDA values exhibited a decreasing trend with increasing soil depth, with the highest FDA concentration recorded under 90% RR at the 0–15 cm depth. In the subsurface (15–30 cm), FDA activity was significantly higher under 90% RR, showing a 28.4% increase compared to CT.

β -glucosidase, a key enzyme in the soil carbon cycle, showed activity levels ranging from 169.6 to 83.7 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ in surface and subsurface soils. The increase in β -glucosidase activity due to residue retention (RR) management ranged from 16.7% to 12.6% at the 0–15 cm depth and from 19.9% to -4.7% at the 15–30 cm depth. At the 0–15 cm depth, β -glucosidase activity was 16.7%, 12.6%, and 7.9% higher under 90%, 30%, and 0% RR, respectively, compared to conventional tillage (CT). β -glucosidase activity ranged from 169.60 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 104.62 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 90% RR, 161.58 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 89.47 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 30% RR, 153.36 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 79.99 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under 0% RR, and 141.29 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (0–15 cm) to 83.75 $\mu\text{g PNG g}^{-1} \text{ soil h}^{-1}$ (15–30 cm) under CT. The mean β -glucosidase activity showed a declining trend with increasing soil depth. The highest activity was recorded at the 0–15 cm depth under 90% RR. At the subsurface level (15–30 cm), β -glucosidase activity was significantly higher under 90% RR, with a 19.9% increase compared to CT.

No significant differences were observed in the stratification ratios of total organic carbon and enzymatic activities between the different residue retention levels and conventional tillage (CT) treatments (Fig 2). Generally, enzymatic activities (β -glucosidase, DHA, and FDA) exhibited a strong correlation ($p < 0.01$) with total organic carbon content and were also strongly correlated (Table 3).

Table 1. Mean value of pH, electrical conductivity (EC) and total organic carbon (TOC) in soil under different Residue Retention (RR) level and Conventional tillage (CT) at the different depths.

Soil parameters	Soil depths (cm)	Residue Retention			CT	C.D value ($p=0.05$)
		RR-0%	RR-30%	RR - 90 %		

pH	0-15	7.81	7.64	7.54	7.92	0.19
	15-30	7.74	7.64	7.54	7.87	N/S
EC (dS m ⁻¹)	0-15	0.23	0.19	0.13	0.23	0.02
	15-30	0.15	0.14	0.15	0.15	N/S
TOC (g kg ⁻¹)	0-15	10.47	12.16	15.24	9.39	2.74
	15-30	7.41	7.88	8.66	7.10	1.05

RR- residue retention, CT- conventional tillage




Figure 1. Total organic carbon (TOC) in soil samples collected from fields under different residue retention and tillage practices. Soil enzymatic activities (b) DHA, (c) FDA, (d) β -glucosidase were estimated under different residue retention (RR) and conventional tillage at the different depth 0-15 cm and 15-30 cm. The residue retentions are no residue retention (0%), 30% residue retention and 90% residue retention. Each data point represents average with error bar as standard error of three replicated observations. Significant difference between treatments was indicated with $p < 0.05$.




Figure 2. Stratification ratio (0-15 cm/15-30cm) for total organic carbon (TOC) and soil enzymatic activities (DHA, dehydrogenase activity; β -glu, β -glucosidase activity; FDA, Fluorescein diacetate activity) under residue retention (RR); 0%, 30% and 90% and conventional tillage (CT). Vertical bar represents a standard error. Significant difference between treatment is indicated with $p < 0.05$.

Table 2. correlation coefficient between biochemical and chemical properties (TOC) in soil samples (n=40)

	TOC	DHA	FDA	β -glucosidase
TOC	-			
DHA	0.84**	-		
FDA	0.78*	0.94**	-	
β -glucosidase	0.78*	0.95**	0.92**	-

Correlation is significant at * $p = 0.05$ level, ** $p = 0.01$ level

DISCUSSION

Climate-smart agricultural practices influence soil enzyme activities to varying degrees. Significant variations in enzyme activities were observed under different levels of crop residue retention (90%, 30%, 0%) compared to conventional tillage. Notable differences in SOC and enzyme activities were recorded in both surface and subsurface soils among the residue retention treatments. Our findings showed that SOC was significantly higher under 90% crop residue retention in the surface soil compared to CT. This is consistent with studies by Kumar *et al.*, (2017), Hati *et al.*, (2015), and McCarty *et al.*, (1997), who reported that conservation tillage, particularly no-till (NT), results in higher SOC concentrations in the topsoil and alters its distribution throughout the soil profile. The greatest differences in SOC concentration between tillage treatments were observed in the uppermost soil layer, with the following order: RT > NT > CT. Several studies have shown that long-term conservation tillage systems (NT and RT) maintain higher SOC levels in the surface soil compared to CT (Conant *et al.*, 2007; Lopez-Fando *et al.*, 2009). Increased SOC concentrations are typically attributed to a variety of interacting factors, including minimal soil disturbance, enhanced residue retention and addition, reduced surface soil temperatures, improved soil moisture, and lower erosion risks (Ismail *et al.*, 1994). Crop residues contribute to the SOC pool, and returning more crop residues to the soil is linked to higher SOC concentrations (Dolan *et al.*, 2006). SOC is one of the soil biological properties most influenced by tillage practices (Somasundaram *et al.*, 2014).

The DHA activity in 90%, 30%, and conventional tillage varied from 70.55 to 103.57 $\mu\text{g TPF g}^{-1} \text{ day}^{-1}$ in the surface layer. Residue retention in no-tillage systems had a significant effect on DHA activity. A similar finding was reported by Parihar *et al.*, (2016), where surface soil under no-tillage (NT) practices showed significantly higher DHA (122.35 $\mu\text{g TPF g}^{-1} \text{ day}^{-1}$) compared to CT (77.07 $\mu\text{g TPF g}^{-1} \text{ day}^{-1}$). The decomposition of retained maize residues releases carbon, which can be available to soil microbes, leading to higher DHA activity in the surface soils under a maize-chickpea cropping system. Kumar *et al.*, (2017) reported significantly higher DHA in soybean + pigeon pea rotations, followed by maize-gram systems.

Dehydrogenase activity is a well-established indicator of biological activity in soils, as the enzyme exists as an integral part of microbial cells but does not accumulate extracellularly. The oxidation of soil organic matter by dehydrogenase involves the transfer of protons and electrons from substrates to acceptors, and is considered linked to the respiration pathways of microorganisms (Das *et al.*, 2011). DHA activity was significantly influenced by the availability of organic matter, soil temperature, and soil moisture. Madejonet *et al.*, (2007) and Tao *et al.*, (2009) observed higher DHA activity under conservation agriculture systems with legume rotations compared to CT.

The mean FDA activity values for 90%, 30%, 0%, and conventional tillage ranged from 22.91 to 30.85 $\mu\text{g fluorescein g}^{-1} \text{ h}^{-1}$ in both the surface and subsurface layers. Residue retention management practices had a significant effect on the surface layer during the crop cycle. Similar findings were reported by Perez-Brandan *et al.*, (2012) and Gajda *et al.*, (2013), who observed higher soil microbial enzymatic activities under conservation agriculture with legume rotations compared to conventional tillage. A significant influence of soil organic matter (SOM) on various biological properties of soil has been documented (Askari *et al.*, 2014; Sinha *et al.*, 2014; Sinha NK, 2014; Marinari *et al.*, 2006). In this study, SOM concentration significantly influenced total organic carbon, DHA, FDA, and β -glucosidase activities, as supported by the strong correlations between soil biological properties and soil organic carbon content.

Higher β -glucosidase activity was observed in surface soils compared to subsurface soils in the maize-chickpea cropping system, likely due to increased carbon input from the fibrous root mass of maize in the previous year. This finding was consistent with the higher soil carbon concentration under 90% residue retention compared to conventional tillage, and the significant positive correlations observed between β -glucosidase activity, residue load, and total organic carbon. Martin-Lammerding *et al.*, (2015) and Acar *et al.*, (2018) noted that β -glucosidase activity was highest under no-tillage (NT), followed by reduced tillage (RT), with CT showing the lowest levels of β -glucosidase activity. Jat *et al.*, (2021) and Acar *et al.*, (2018) also reported that β -glucosidase activity was significantly higher in rhizospheric soils compared to bulk soils.

CONCLUSION

The soil biological parameters in the Vertisol of central India under the maize-chickpea cropping system were significantly impacted by varying levels of crop residue retention management. The results clearly demonstrated that higher residue retention, particularly 90% and 30%, supported greater biological activity compared to conventional tillage. Residue retention, as a key component of conservation tillage, triggered a rapid response in soil microbial activity. Biological activities in the soil were predominantly concentrated in the upper layer (0–15 cm), where TOC, DHA, FDA, and β -glucosidase activities were notably higher. This increase could be attributed to the accumulation of organic matter from crop residues, litterfall, root biomass, and root and soil biota exudates, as well as the enhanced interaction between the soil surface and atmospheric conditions, fostering soil biodiversity. The strong correlations observed between total organic carbon and enzymatic activities (DHA, FDA, and β -glucosidase) in the surface layer highlight the critical role of organic carbon in promoting microbial activity and sustaining soil biodiversity.

References

1. Franzluebbers, A. J. "Soil organic matter stratification ratio as an indicator of soil quality." *Soil and Tillage Research* 66, no. 2 (2002): 95-106.
2. Adam, Gillian, and Harry Duncan. "Development of a sensitive and rapid method for the measurement of total microbial activity using fluorescein diacetate (FDA) in a range of soils." *Soil biology and biochemistry* 33, no. 7-8 (2001): 943-951.
3. Aher, Satish Bhagwatrao, Brij Lal Lakaria, Swami Kaleshananda Swami Kaleshananda, Amar Bahadur Singh, Sivakoti Ramana Sivakoti Ramana, Jyoti Kumar Thakur, Ashish Kumar Biswas, Pramod Jha, Madhab Chandra Manna, and Dharmendra Singh Yashona. "Soil microbial population and enzyme activities under organic, biodynamic and conventional agriculture in semi-arid tropical conditions of central India." (2018): 763-773.
4. Askari, Mohammad Sadegh, and Nicholas M. Holden. "Quantitative soil quality indexing of temperate arable management systems." *Soil and Tillage Research* 150 (2015): 57-67.
5. Aulakh, Milkha S., and Cynthia A. Grant. "Integrated nutrient management for sustainable crop production." (*No Title*) (2008).
6. Kumar, Awanish, Anusuiya Panda, L. K. Srivastava, and V. N. Mishra. "Effect of conservation tillage on biological activity in soil and crop productivity under rainfed Vertisols of central India." *Int. J. Chem. Stud* 5, no. 6 (2017): 1939-1946.
7. Bhattacharyya, Ranjan, Birendra Nath Ghosh, Prasanta Kumar Mishra, Biswapati Mandal, Cherukumalli Srinivasa Rao, Dibyendu Sarkar, Krishnendu Das et al. "Soil degradation in India: Challenges and potential solutions." *Sustainability* 7, no. 4 (2015): 3528-3570.
8. Bielińska, Elżbieta, and Agnieszka Mocek-Plóćiniak. "Impact of the tillage system on the soil enzymatic activity." *Archives of Environmental Protection* (2012)..
9. Perez-Brandán, Carolina, José L. Arzeno, Jorgelina Huidobro, Betiana Grümberg, Cinthia Conforto, Sally Hilton, Gary D. Bending, José M. Meriles, and Silvana Vargas-Gil. "Long-term effect of tillage systems on soil microbiological, chemical and physical parameters and the incidence of charcoal rot by *Macrophomina phaseolina* (Tassi) Goid in soybean." *Crop Protection* 40 (2012): 73-82.
10. Stewart, Catherine E., Keith Paustian, Richard T. Conant, Alain F. Plante, and Johan Six. "Soil carbon saturation: concept, evidence and evaluation." *Biogeochemistry* 86 (2007): 19-31.
11. Das, Shonkor Kumar, and Ajit Varma. "Role of enzymes in maintaining soil health." *Soil enzymology* (2011): 25-42.
12. Das, Anup, Jayanta Layek, Ramkrushna Gandhiji Idapuganti, Savita Basavaraj, Rattan Lal, Krishnappa Rangappa, Gulab Singh Yadav, Subhash Babu, and Shishomvanoo Ngachan. "Conservation tillage and residue management improves soil properties under a upland rice-rapeseed system in the subtropical eastern Himalayas." *Land Degradation & Development* 31, no. 14 (2020): 1775-1791.
13. Martín-Lammerding, Diana, Mariela Navas, María del Mar Albarrán, José Luis Tenorio, and Ingrid Walter. "LONG term management systems under semiarid conditions: Influence on labile organic matter, β -glucosidase activity and microbial efficiency." *Applied Soil Ecology* 96 (2015): 296-305.
14. Dolan, M. S., C. E. Clapp, R. R. Allmaras, J. M. Baker, and J. A. E. Molina. "Soil organic carbon and nitrogen in a Minnesota soil as related to tillage, residue and nitrogen management." *Soil and Tillage Research* 89, no. 2 (2006): 221-231.
15. Eivazi, F., and M. A. Tabatabai. "Glucosidases and galactosidases in soils." *Soil Biology and Biochemistry* 20, no. 5 (1988): 601-606.
16. Gajda, A. M., B. Przewloka, and K. Gawryjolek. "Changes in soil quality associated with tillage system applied." *International Agrophysics* 27, no. 2 (2013).
17. Jat, Hanuman Sahay, Ashim Datta, Madhu Choudhary, Parbodh Chander Sharma, Bharti Dixit, and Mangi Lal Jat. "Soil enzymes activity: Effect of climate smart agriculture on rhizosphere and bulk soil under cereal based systems of north-west India." *European Journal of Soil Biology* 103 (2021): 103292.
18. Harasim, Elżbieta, Jacek Antonkiewicz, and Cezary A. Kwiatkowski. "The effects of catch crops and tillage systems on selected physical properties and enzymatic activity of loess soil in a spring wheat monoculture." *Agronomy* 10, no. 3 (2020): 334.
19. Hati, K. M., R. S. Chaudhary, K. G. Mandal, K. K. Bandyopadhyay, R. K. Singh, Nishant K. Sinha, M. Mohanty, J. Somasundaram, and R. Saha. "Effects of tillage, residue and fertilizer nitrogen on crop yields, and soil physical properties under soybean-wheat rotation in vertisols of Central India." *Agricultural Research* 4 (2015): 48-56.
20. Huang, Zhiqun, Zhihong Xu, and Chengrong Chen. "Effect of mulching on labile soil organic matter pools, microbial community functional diversity and nitrogen transformations in two hardwood plantations of subtropical Australia." *Applied Soil Ecology* 40, no. 2 (2008): 229-239.
21. Ismail I, Ismail, Isro, R. L. Blevins, and W. W. Frye. "Long-term no-tillage effects on soil properties and continuous corn yields." *Soil Science Society of America Journal* 58, no. 1 (1994): 193-198.
22. Doran, John W., and Timothy B. Parkin. "Defining and assessing soil quality." *Defining soil quality for a sustainable environment* 35 (1994): 1-21.
23. Gianfreda, L., and J. M. Bollag. "Influence of natural and anthropogenic factors on enzyme activity in soil." (1996): 123-193.
24. Liu, Chang, Meng Lu, Jun Cui, Bo Li, and Changming Fang. "Effects of straw carbon input on carbon dynamics in agricultural soils: a meta-analysis." *Global change biology* 20, no. 5 (2014): 1366-1381.
25. López-Fando, Cristina, and M. T. Pardo. "Changes in soil chemical characteristics with different tillage practices in a semi-arid environment." *Soil and Tillage Research* 104, no. 2 (2009): 278-284.
26. Acar, Mert, Ismail Celik, Hikmet GÜNAL, Nurullah Acir, Zeliha Barut Bereket, and Mesut Budak. "Tillage effects on soil organic carbon, microbial biomass carbon and beta-glucosidase enzyme activity in a typical haploxerert soil." *Scientific Papers. Series A. Agronomy* 61 (2018).
27. Madejón, Engracia, F. Moreno, J. M. Murillo, and Francisco Pelegrín. "Soil biochemical response to long-term conservation tillage under semi-arid Mediterranean conditions." *Soil and Tillage Research* 94, no. 2 (2007): 346-352..
28. Marinari, Sara, Roberto Mancinelli, Enio Campiglia, and Stefano Grego. "Chemical and biological indicators of soil quality in organic and conventional farming systems in Central Italy." *Ecological Indicators* 6, no. 4 (2006): 701-711.
29. McCarty, G. W., and J. J. Meisinger. "Effects of N fertilizer treatments on biologically active N pools in soils under plow and no tillage." *Biology and Fertility of Soils* 24 (1997): 406-412.

29. Mythili, Gurumurthy, and Jann Goedecke. "Economics of land degradation in India." *Economics of land degradation and improvement—a global assessment for sustainable development* (2016): 431-469
30. Dw, Nelson. "Total carbon, organic carbon and organic matter." *Chemical and Microbial Properties* 9 (1982): 530-594.
31. Ojeda, Gerardo, Stefania Mattana, Anna Àvila, Josep Maria Alcañiz, Martin Volkman, and Jörg Bachmann. "Are soil–water functions affected by biochar application?." *Geoderma* 249 (2015): 1-11.
32. Nannipieri, Paolo, STEFANO GRECO, and Brunello Ceccanti. "Ecological significance of the biological activity in soil." *Soil biochemistry* (2017): 293-356.
33. Parihar, Chiter M., Malu R. Yadav, Aditya K. SINGH, Bhupender KUMAR, Vijay POONIYA, Sanatan PRADHAN, Rakesh K. VERMA, Muli D. PARIHAR, Hari S. NAYAK, and Yashpal S. SAHARAWAT. "Long-term conservation agriculture and intensified cropping systems: Effects on growth, yield, water, and energy-use efficiency of maize in northwestern India." *Pedosphere* 28, no. 6 (2018): 952-963.
34. Roldán, A., J. R. Salinas-García, M. M. Alguacil, and F. Caravaca. "Changes in soil enzyme activity, fertility, aggregation and C sequestration mediated by conservation tillage practices and water regime in a maize field." *Applied Soil Ecology* 30, no. 1 (2005): 11-20.
35. Melero, Sebastiana, EngraciaMadejón, Juan Francisco Herencia, and Juan Carlos Ruiz. "Effect of implementing organic farming on chemical and biochemical properties of an irrigated loam soil." *Agronomy Journal* 100, no. 1 (2008): 136-144.
36. Sinha, Nishant K., Usha K. Chopra, Anil K. Singh, M. Mohanty, J. Somasundaram, and R. S. Chaudhary. "Soil physical quality as affected by management practices under maize–wheat system." *National Academy Science Letters* 37 (2014): 13-18.
37. Sinha, Nishant K., M. Mohanty, Bharat P. Meena, Hiranmoy Das, Usha K. Chopra, and Anil K. Singh. "Soil quality indicators under continuous cropping systems in the arid ecosystem of India." *African Journal of Agricultural Research* 9, no. 2 (2014): 285-293.
38. Somasundaram, J., B. L. Lakaria, R. Saha, N. K. Sinha, P. Jha, R. S. Chaudhary, R. K. Singh et al. "Management of stressed soils of dryland agriculture in semi-arid tropics-areview." *Indian Journal of Soil Conservation* 42, no. 2 (2014): 178-187.
39. Tao, Jun, Bryan Griffiths, Shujie Zhang, Xiaoyun Chen, Manqiang Liu, Feng Hu, and Huixin Li. "Effects of earthworms on soil enzyme activity in an organic residue amended rice–wheat rotation agro-ecosystem." *Applied Soil Ecology* 42, no. 3 (2009): 221-226.
40. Thalmann, A. "ZurMethodik der Bestimmung der Dehydrogenaseaktivitätim Boden mittelstriphenyltetrazoliumchlorid (TTC)." *Landwirtsch. Forsch.* 21 (1968): 249-258..
41. Thierfelder, Christian, Frédéric Baudron, Peter Setimela, Isaiah Nyagumbo, Walter Mupangwa, Blessing Mhlanga, Nicole Lee, and Bruno Gérard. "Complementary practices supporting conservation agriculture in southern Africa. A review." *Agronomy for Sustainable Development* 38 (2018): 1-22.
42. Walkley, Aldous, and I. Armstrong Black. "An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method." *Soil science* 37, no. 1 (1934): 29-38.
43. Ye, Xinxin, Yin Ye, Rushan Chai, Junli Li, Chao Ma, Hongying Li, QizhongXiong, and Hongjian Gao. "The influence of a year-round tillage and residue management model on soil N fractions in a wheat-maize cropping system in central China." *Scientific Reports* 9, no. 1 (2019): 4767.