

Isolation and Characterization of Naga king chilli rhizobacteria

Abstract

The Northeast region of India is recognized for its chilli diversity with number of variants noted from this region. Amongst the many landraces of chilli that are cultivated in this region, the Naga King Chilli (*Capsicum chinense* Jacq.) is one of the best known worldwide. Rhizosphere is the soil around the root system of a plant considered to be rich in nutrients due to the accumulation of various plants exudates, providing a rich source of nutrients for soil microbes and is reflected on the number of microbes being higher in this region when compared with the bulk soil. The bacteria colonizing this region of the soil are called rhizobacteria and are believed to play a vital role in plant growth and development. An investigation was carried out in order to know the indigenous rhizobacterial population of Naga King Chilli rhizosphere collected from the farmers' field Nagaland, where the crop is popularly grown. 27 rhizobacteria were isolated on nutrient agar and king's B agar medium and characterized based on their colony characters and some biochemical test including gram reaction, KOH test, gelatine hydrolysis and starch hydrolysis test. The results suggested that the rhizosphere of Naga king chilli is dominated by gram negative bacteria with 21 isolates being gram negative. All the isolated shows positive catalyst and gelatine hydrolysis activity with 18 isolates being able to hydrolyse starch.

Keywords: Characterization, Naga King chilli, Rhizobacteria

Introduction

The Northeast region of India is recognized as hot-spot for chilli diversity (Mathur *et al.*, 2000) and amongst the many landraces of chilli that are cultivated in this region, the Naga King Chilli (*Capsicum chinense* Jacq.) is the best known worldwide. It is a self-pollinated plant; however, considerable cross pollination (upto 10%) may occur when insect population is high. It behaves as a semi perennial herb if grown under favourable condition (Borgohain and Devi, 2007). Chilli fruits constitute large amounts of beneficial compounds including antioxidants,

carbohydrates, minerals, phytochemicals, proteins, amino acids, and vitamins (Olatunji and Afolayan, 2018). The narrow zone of soil directly surrounding the root system is referred to as rhizosphere (Walker *et al.*, 2003). This zone is rich in nutrients when compared with the bulk soil due to the accumulation of a variety of plant exudates, such as amino acids and sugars, providing a rich source of energy and nutrients for bacteria (Gray and Smith, 2005). The rhizosphere is populated by a diverse range of microorganisms and the bacteria colonizing this habitat are called rhizobacteria (Schroth and Hancock, 1982) which increases the absorption capacity of nutrients and protection against phytopathogens (Pérez-García *et al.*, 2023). Knowledge of the native bacterial population, their characterization, and identification is required for understanding the distribution and diversity of indigenous bacteria in the rhizosphere of specific crops (Chahboune *et al.*, 2011).

Materials and method

A field survey was undertaken for the collection of rhizobacteria from the farmer's field of Peren district, Nagaland. Where, rhizospheric bacteria were isolated from the rhizosphere of healthy plant at flowering stage.

Isolation and maintenance of bacteria

Soil dilution plate technique described by Waksman, 1927 was followed for the isolation of rhizobacteria and the suspension was serially diluted to 10^{-8} . 0.1 ml of the last 3 serial dilution were spread with a glass spreader on nutrient agar and King's B medium plates and incubated for 48h at $26 \pm 1^\circ\text{C}$. Colonies with different morphological appearances were selected from the countable plates and re-streaked on a new plate containing the same media to obtain pure colonies (Patel *et al.*, 2014).

Characterization of rhizobacteria

Morphology of bacterial colonies were characterized based on various traits such as size, shape, elevation, surface, optical properties, margin, and pigmentation on bacterial cultures previously grown for 48 h on NA medium and incubated at 28°C .

A series of biochemical tests were conducted using the criteria of Bergey's Manual of Systemic Bacteriology (1994).

Catalase test

To detect the production of catalase a loop full of 24-48 hours old test bacterium was smeared on the slide and then covered it with a few drops of 3% hydrogen peroxide. The smear was observed for bubble production. (Koche and Gade, 2013)

KOH test

A loopful of bacteria was mixed in a drop 3% aqueous KOH solution for not more than 10 seconds. The inoculating loop was raised a few centimetres from the microslide and the formation of a mucoid thread was recorded. Gram positive bacterium do not produce strands even on repeated strokes of the inoculating loop while gram negative bacterium do.

Starch hydrolysis

Starch hydrolysis was evaluated using nutrient agar amended with 0.2% starch. After incubating the test bacterium on the medium for 7 days, the agar plates were then flooded with Lugol's iodine and allowed to act for few minutes. The cultures showing a clear zone was considered to be positive reaction.

Gelatine liquefaction test

Stab method described by Koche and Gade. (2013) was followed for gelatine liquefaction test (Peptone-10g; Beef extract-5g; Gelatin-20g and Distilled water - 1000ml). Inoculation was done by stabbing a straight inoculating needle charged with 48 hours old growth of the test bacterium. The tubes were incubated at 20°C and observation were recorded for liquefaction of gel column.

Result and discussions

Morphological characters of the rhizobacterial isolates are presented in Table 1. The isolates were observed to be predominantly smooth, round, either orange or milky white in colour, smooth surface, convex with entire margin and translucent Result on biochemical characterization are shown in Table2. It was Observed that out of the 27 isolated rhizobacteria eight were found to be Gram positive and 19 Gram negative, five isolates were recorded to utilize lactose whereas, only six isolates were found to

be negative for KOH test. All the isolates were recorded to liquify gelatine. However, eight isolates could not hydrolyse starch.

The findings of the present investigation were found to be in contrary to the findings of Banerjee *et al.* (2011) who reported that the bacterial population in chilli rhizosphere was dominated by gram positive bacteria with white, irregular, opaque colonies as the Naga king chilli rhizosphere was found to be dominated by Gram negative, orange coloured, opaque colonies. This difference in observations may be due to the fact that different ecological and environmental conditions favour the growth, development and establishment of different bacterial population and hence reflected in the population of the rhizobacteria. The results on catalase productions were found to be in conformity with the findings of Patel and Desai (2015) who observed that all rhizobacterial isolates were positive for catalyst production and hence, are aerobic in nature.

Table 1: Morphological characterization of rhizobacteria

Isolates	Size	Shape	Colour	Surface	Elevation	Margi n	Opacit y
T ₁	Small	Round	White	Smooth	Flat	Erode	Translu cent
T ₂	Small	Round	White	Smooth	Convex	Entire	Translu cent
T ₃	Small	Round	Creamy white	Smooth	Convex	Entire	Opaque
T ₄	Small	Irregula r	Light yellow	Smooth	Raised	Undula ted	Translu cent
T ₅	Small	Round	Orange	Smooth	Convex	Entire	Opaque
T ₆	Small	Round	Light orange	Smooth	Convex	Entire	Opaque
T ₇	Small	Irregula r	Orange	Corrugate d	Raised	Undula ted	Opaque
T ₈	Small	Round	Reddish	Smooth	Convex	Entire	Opaque
T ₉	Small	Round	Yellow	Smooth	Pulvinate	Entire	Opaque
T ₁₀	Small	Round	Yellow	Smooth	Convex	Entire	Translu

	l						cent
T ₁₁	Small	Round	Pale yellow	Smooth	Convex	Entire	Opaque
T ₁₂	Small	Round	Greenish yellow	Smooth	Convex	Entire	Opaque
T ₁₃	Medium	Round	White	Smooth	Convex	Entire	Opaque
T ₁₄	Small	round	Yellow	Smooth	Raised	Entire	Opaque
T ₁₅	Small	Round	Milky white	Smooth	Convex	Entire	Translucent
T ₁₆	Medium	Round	Orange	Smooth	Convex	Entire	Opaque
T ₁₇	Medium	Round	Dull white	Smooth	Convex	Entire	Translucent
T ₁₈	Small	Round	Orange	Corrugated	Raised	Undulated	Opaque
T ₁₉	Small	Round	Light orange	Smooth	Flat	Entire	Translucent
T ₂₀	Small	Round	Pale white	Smooth	Convex	Entire	Translucent
T ₂₁	Small	Round	Orange	Smooth	Convex	Entire	Opaque
T ₂₂	Small	Round	Yellow	Smooth	Flat	Entire	Opaque
T ₂₃	Small	Round	Orange	Smooth	Convex	Entire	Opaque
T ₂₄	Small	round	Pale white	Smooth	Flat	Entire	Translucent
T ₂₅	Small	Round	Creamy white	Smooth	Convex	Entire	Opaque
T ₂₆	Medium	Round	Dull white	Smooth	Convex	Entire	Opaque
T ₂₇	Small	Round	Pale yellow	Smooth	Convex	Entire	Opaque

Table 2. Biochemical characterization of the rhizobacteria

Isolates	Gram staining	Starch Hydrolysis	KOH test	Gelatine liquification	Catalyst test
T ₁	-	-	+	+	+
T ₂	-	-	+	+	+

T ₃	-	+	+	+	+
T ₄	-	+	+	+	+
T ₅	+	-	-	+	+
T ₆	-	+	+	+	+
T ₇	-	+	+	+	+
T ₈	-	+	+	+	+
T ₉	-	+	+	+	+
T ₁₀	-	+	+	+	+
T ₁₁	-	-	+	+	+
T ₁₂	-	-	+	+	+
T ₁₃	-	+	+	+	+
T ₁₄	-	+	+	+	+
T ₁₅	+	+	-	+	+
T ₁₆	-	-	+	+	+
T ₁₇	+	+	-	+	+
T ₁₈	-	+	+	+	+
T ₁₉	-	-	+	+	+
T ₂₀	+	+	-	+	+
T ₂₁	-	-	+	+	+
T ₂₂	+	+	-	+	+
T ₂₃	-	+	+	+	+
T ₂₄	-	+	+	+	+
T ₂₅	+	+	-	+	+
T ₂₆	-	+	+	+	+
T ₂₇	-	+	+	+	+

References

- Banerjee S, Datta M., Palit R., Sengupta C and Pandit MK. 2011. Plant growth promoting rhizobacteria enhance growth and yield of chilli (*Capsicum annuum* L.) under field conditions. *Australian Journal of Crop Science*. **5**(5):531-536.
- Borgohain R and Devi J. 2007. The hottest chilli: A new horizon in agri-entrepreneurship. *Science Tech Entrepreneur*. <http://www.assamagribusiness.nic.in>.
- Chahboune R, Barrijal S, Moreno S, and Bedmar EJ. 2011. Characterization of *Bradyrhizobium* species isolated from root nodules of *Cytisus villosus* grown in Morocco. *Systematic and Applied Microbiology*. **34**: 440–445.
- Gray EJ and Smith DL. 2005. Intracellular and extracellular PGPR: commonalities and distinctions in the plant–bacterium signalling processes. *Soil Biology and Biochemistry*. **37**: 395–412.
- Koche MD and Gade RM. 2013. Effect of sodium chloride, streptomycin sulphate and different agro-chemical on growth of *Pseudomonas fluorescens*. *Journal of Mycopathological Research*. **51**(2): 295-299.
- Mathur R, Dangi RS, Dass SC. and Malhotra RC. 2000. The hottest chilli variety in India. *Current Science*. **79**: 287–288.
- Olatunji, T. L. and Afolayan, A. J. 2018. The suitability of chili pepper (*Capsicum annuum* L.) for alleviating human micronutrient dietary deficiencies: A review. *Food Science & Nutrition*. **6**:2239-2251.
- Patel, P. V. and Desai, P. B. 2015. Isolation of Rhizobacteria from Paddy Field and Their Traits for Plant Growth Promotion. *Research Journal of Recent Sciences*. **4**: 34-41.
- Patel M, Sheth U, Hajela P, Joshi B and Animasaun DA. 2014. Isolation and diversity analysis of rhizobacteria from sugarcane and its biocontrol potential against *Rhizoctonia solani* a common plant pathogen. *International Journal of Current Microbiological Applied Science*. **3**(12): 69-76.
- Pérez-García, L.A., Sáenz-Mata, J., Fortis-Hernández, M., Navarro-Muñoz, C. E., Palacio-Rodríguez, R. and Preciado-Rangel, P. 2023. Plant-Growth-Promoting Rhizobacteria Improve Germination and Bioactive Compounds in Cucumber Seedlings. *Agronomy*. **13**(315):1-10.
- Schroth MN. and Hancock JG. 1982. Disease-suppressive soil and root-colonizing bacteria. *Science*. **216**:1376-81.
- Walker TS, Bais HP, Grotewold E. and Vivanco JM. 2003. Root exudation and rhizosphere biology. *Plant Physiology*. **132**: 44–51.




Plate 1. Biochemical test

- a. Gram reaction
- b. Lactose utilization test
- c. KOH test
- d. Gelatine liquification test
- e. Starch Hydrolysis test
- f. Catalase test