

# DETERMINATION OF THE FUNCTIONAL NUTRITIONAL AND SENSORY PROPERTIES OF SEASONING POWDERS PRODUCED FROM FERMENTED AFRICAN LOCUST BEAN CHICKEN AND SHIITAKE MUSHROOMS

## Abstract

Rewrite again due to

Background .....

Aims .....

Methodology .....

Results .....

Conclusion .....

*Food seasoning powders were produced from a combination of fermented African locust bean (iru), chicken and shiitake mushroom powders. Four combinations were made containing different proportions of iru, chicken and mushroom (I/C/M) as follows 1 (16.6/16.6/66.7), 2(16.6/66.7/16.6), 3(66.7/16.6/16.6), 4(33.3/33.3/33.3).the functional, nutritional and sensory properties of the seasoning powders were evaluated using standard analytical procedures. There was no significant difference at  $P>0.05$  in the bulk densities of all the four blends. The blend containing 66.7 % chicken powder had the highest protein content  $50.32\pm1.18\%$ .all the seasoning powder blends had less than 1% monosodium glutamate while free glutamic acid was found to most abundant  $443.39\pm 1.44\text{mg}/100\text{g}$  in the blend containing more iru: 3(66.7/16.6/16.6)(I/C/M). This same blend was considered to be the most acceptable product as it had the highest general acceptability score  $8.22 \pm 0.17$ .*

**Key words:** Seasoning powder, Iru, chicken, Shiitake Mushroom.

## Introduction

According to European Spice Association (ESA), seasoning is a blend of permitted food ingredients added as necessary to achieve the purpose for which it is designed (e.g. to improve the taste, eating quality and/or functionality of a food).Seasoning powders can be defined as mixtures of different compounds whose main role is to enhance taste and flavor of ready to eat food to which they are added.Generally commercial powdered seasonings are added before or during cooking.Food spices, whether natural or artificial, are in high demand in daily meals because they enhance the flavor of common dishes and make them more attractive when added. There are different types of food seasonings sold on the market, but what they have in common is that they are all produced in the laboratory using different ingredient ratios and techniques. (Nguyen Minh Thuy *et al.*, 2024).Besides their taste, aroma and composition, seasoning powders also need to have certain physical and functional properties, which are very important during processing, packaging, handle, manipulation, storage, and application in the kitchen.The most important physical properties of food seasoning powders are bulk density, granulation, wettability, dispersibility and flowability. Bulk density is a ratio of weight of powdered material and volume which that certain powder weight occupies. It depends on granulation, particle size and shape. Particles with irregular shape have lower bulk density. Bulk density is important for packaging and storage space. Dispersibility is important for uniform mixing after addition to liquid food medium. Flowability is important for transport operations during processing and good flowability without caking is required.

.Commercial spice blends or seasonings consist of spices, vegetables and other ingredients, including flavorings, salt, sugar, dextrose, corn syrup solids, starches, maltodextrin, yeast extracts or hydrolysates, hydrolysed proteins, monosodium glutamate, or nucleotides. In the production of commercial seasoning blends, spices and other ingredients are first plated on some sort of carrier (e.g. salt, sugar, dextrose, or maltodextrin), before being mixed with other spices or ingredients. Anti-caking agents are usually added in required amounts below 2% [2]. According to FAO standards, powdered vegetable based food seasonings must satisfy following requirements: moisture  $\leq 5\%$ , NaCl  $\leq 60\%$ , monosodium glutamate  $\leq 33\%$  and dried vegetables  $\geq 8\%$  [3].

African Locust Bean (*Parkia biglobosa*) is among the leguminous plants used by man particularly in some African countries for the production of local condiment. African locust bean seeds are rich in protein and usually fermented to a tasty food condiment called dawadawa which is used as a flavour intensifier for soups and stews and also adds protein to a protein- poor diet (Yahaya Mode *et al.*, 2018). Dawadawa or iru is one of the most fully exploited traditional additives, used almost every day; in most food preparations, for its aroma and flavouring enhancing effects. Apart from serving as a food additive that enhances the organoleptic properties of the food, dawadawa may also serve as a source of proteins (Murtala *et al.*, 2016).

Chicken meat is a substantial food for consumers all over the world. It is rich in essential amino acids and involves large amount of proteins (Cui *et al.*, 2003). Chicken breast meat (a type of white meat) was selected because it has a lower fat content and is normally perceived as a healthier meat option by consumers. (Charlton *et al.*, 2002). The chicken breast can be dehydrated and then is made into a famous natural seasoning (chicken powders) at home and abroad. Chicken powders can be healthy, delicious, and rich in nutritional ingredient (Ran *et al.*, 2003).

Mushrooms are excellent sources of proteins, minerals and vitamins which are important role for nutritional purpose. Shiitake mushroom is the medicinal and edible mushroom known for its potential health benefits. It is scientifically known as (*Lentinula edodes*). It is the main essential medicinal mushroom which second ranks in terms of total mushroom production in the world only next to *Agaricus* species. Shiitake is a high-quality delicious taste and texture with mushroom and many types of bioactive compounds are found in this mushroom which is as follows Lentinan  $\beta$ -(1glucan/polysaccharide), Lentinus edodes mycelium (LEM), Eritadenine, Ergothioneine (an amino acid that can act as an antioxidant. These compounds protect to various types of human diseases. These are important bioactive compounds that have lipid-lowering, antimicrobial, and antioxidant properties. Shiitake mushrooms have a rich, earthy flavor and a distinctive taste best described as meaty (Prince *et al.*, 2023).

The research problem

Objective

Literature Review

Conceptual Framework

## Materials and Methods

### Production of Fermented African Locust Bean Powder


This was done according to the method described by (Adejumo *et al.*, 2013 and Zannou *et al.* 2018) (Fig 1). Dried *P. biglobosa* seeds were washed and sorted in order to eliminate spoiled grains and foreigner particles. The seeds underwent the first cooking which is the first important operation that seeds undergo during condiments production and it consists of boiling seed in water for 6 to 24 hours. This operation allows not only a loss of astringency or bitterness of the seed but also the softening of the seminal integument. After this step, the cooked seeds were let to cool and washed in order to take off any remaining impurity and astringency. Seeds were then ready for dehulling. Dehulling is the second major operation involving in the production of the fermented condiments and consists of removing seed coats. Seed coats were taken off by pressing between palms of hands. The second cooking is a short 1 to 2 hours boiling of cotyledons in water. After the cooking, cotyledons will be well drained. The fermentation represents the last most important operation involved in the production of seeds condiments. Although above-cited operations prepare cotyledons to render the fermentation most successful, this step is the most sensitive amongst all operations. It gives to cotyledons all their nutritional capacities, microbiologic properties, and organoleptic characteristics. Cotyledons were spread in basket trays, calabash trays or containers, wrapped with heavy cloths, jutes backs and/ or polyethylene bags and then left for fermentation which lasts for 48 hr. After the fermentation, the condiment was oven- dried ground to a fine powder and packaged.

### Production of Chicken Powder

Chicken flavouring powder was prepared as shown in Fig 2. Muscle from slaughtered homebred chicken was used. It was washed, sliced into small pieces, salt herbs spices and water were added to it and it was boiled for one hour. It was then ground to a paste using a kitchen blender. The paste was spread in a laboratory drying oven and dried at 60°C for 12hr. It was then ground to a fine powder, sieved and packaged in polythene bags.

### Production of Shiitake Mushroom Powder

Mushroom powder was prepared as shown in Figure 3 according to the method described by Balaji *et al.*, 2009 (Fig 3). Dried shiitake mushrooms were soaked in water in a ratio 3:1(w/v) to rehydrate them. According to Wu and Wang (2000), the aroma of dried shiitake mushrooms is more flavourful and meaty than that of fresh mushrooms. They were washed with clean water to remove impurities and soil particles. They were then boiled in water for one hour together with herbs spices and salt after which they were blended into a paste using a kitchen blender. The pastes was oven dried at 90°C for 6 hr. after which it was ground to a fine powder, sieved and packaged polythene bags and stored at room temperature.






Figure 1: Flow Chart for the Production of *ITA* Powder. Source: (Adejumo *et al.*, 2013, Zannou *et al.*, 2018,)

Figure 2: Flow Chart for the Production of Chicken Powder




Figure 3: Flow Chart for the Production of Mushroom Powder  
Source: Balaji *et al.*, 2009



Plate 1: Iru (I), Chicken (C) And Shiitake Mushroom(M) Seasoning Powders

### Blend Formulation

Four different blends were prepared of combinations of iru, chicken and mushroom powders as shown in Table 1.

Table 1: Blend Formulation

Experimental Blend	Iru	Chicken Powder	Shiitake Powder %	Mushroom
1	16.6	16.6	66.7	
2	16.6	66.7	16.6	
3	33.3	33.3	33.4	
4	66.7	16.6	16.6	

## Analytical Procedures

### Determination of Bulk Density

The bulk density of the flour samples were determined by the method described by Onwuka (2005). The sample (50g) was weighed into 100 mL graduated cylinder, then, this was gently tapped at the bottom several times on a laboratory bench, until no further diminution of the sample level. The bulk density was calculated as:

$$\text{Bulk density } \left( \frac{\text{g}}{\text{mL}} \right) = \frac{\text{Weight of sample (g)}}{\text{Volume of sample (mL)}} \quad (1)$$

### 3.8.2 Determination of Swelling Index

The swelling index was determined as described by AOAC (2016). The sample was transferred into clean, graduated (50 mL) cylinder up to the 10 mL mark and the volume noted. Distilled water was added to the flour sample up to the 50 mL mark. The cylinder was swirled and allowed to stand for 2 min after which it was swirled again. It was allowed to stand for another 8 min after which the volume of the swollen sample was taken. The swelling index or power of the sample was calculated as:

$$\text{Swelling Index} = \frac{\text{Sample volume after swelling (mL)}}{\text{Sample volume before swelling (mL)}} \quad (2)$$

### 3.8.3 Determination of Water Absorption Capacity

The AOAC (2016) method was used. One gram of flour was weighed into a graduated conical flask and 10 mL of water was added to the weighed sample and mixed well. The sample was allowed to stand at room temperature for 30 min and then centrifuged at 5000 rpm for 30 min. The supernatant was carefully decanted and measured to know the volume of the free water or oil. The absorption capacity was expressed as grams of water absorbed per gram of sample or in percentage. Calculation: Water absorption capacity of the sample was calculated as:

$$WAC \left( \frac{\text{g}}{\text{g}} \right) = (V_1 - V_2) \times \text{density of water (g/mL)} \quad (3)$$

Where  $V_1$  is initial volume of distilled water added

$V_2$  is volume of supernatant decanted

### 3.8.4 Determination of Oil Absorption Capacity

The AOAC (2016) method was used. One gram of flour was weighed into a graduated conical flask and 10 mL of oil was added to the weighed sample and mixed well. The sample was allowed to stand at room temperature for 30 min and then centrifuged at 5000 rpm for 30 min. The supernatant was carefully decanted and measured to know the volume of the free oil. The absorption capacity was expressed as grams of oil absorbed per gram of sample or in percentage. Calculation: Water absorption capacity of the sample was calculated as:

$$OAC \left( \frac{\text{g}}{\text{g}} \right) = (V_1 - V_2) \times \text{density of oil (g/mL)} \quad (4)$$

Where  $V_1$  is initial volume of oil added

$V_2$  is volume of supernatant decanted

## Determination of Proximate Composition of Seasoning Powders

Moisture, ash, crude fat and crude fibre were determined in accordance with the official methods of the Association of Official Analytical Chemists [5]. Moisture content was determined by oven drying of a weighed sample to a constant weight at 105°C. Crude protein content was determined by Kjeldahl method using 6.25 as the conversion constant after the determination of each sample's nitrogen. Crude fat content was determined by Soxhlet method using n-hexane as solvent. The ash content was determined gravimetrically after ignition at 550°C. Carbohydrate content was calculated by difference. All analyses were carried out in triplicates.

#### **Determination of Taste Compounds**

MSG content in the seasoning powders was determined by HPLC using the method described by Verni *et al.*, 2010. Free glutamic acid content was determined using an amino acid analyser. IMP and GMP contents were determined by HPLC using the method described by Qiu *et al.*, 2016.

#### **Sensory Evaluation**

Sensory evaluation was carried out on aqueous solutions (10g in 100ml distilled water ) of the seasoning powders prepared using a 20 member panel of semi trained panellists composed of both male and female students from the university of Bamenda Cameroon. A nine point hedonic scale was used to grade the seasoning powders with 1 being extremely dislike and 9 being extreme like. The parameters evaluated were appearance, taste and general acceptability.

#### **Statistical Analysis**

All analysis was done in triplicates. Results were expressed as the mean  $\pm$  standard deviation. The data obtained was subjected to statistical analysis by ANOVA at 5% level of significance. Separation of means was carried out using Duncan's Multiple Range test.

#### **Results and Discussion**

##### **Functional Properties of Seasoning Powder Blends**

Functional properties are intrinsic physicochemical characteristics which affect the behaviour of properties in food system during processing, manufacturing, storage and preparations. These properties include emulsion capacity and stability. Functional properties are the fundamental physico- chemical properties that reflect the complex interaction between the composition, structure, molecular conformation and physico-chemical properties of food components together with the nature of environment in which these are associated and measured (Kaur and Singh , 2006). Functional characteristics are required to evaluate and possibly help to predict how new proteins, fat, fibre and carbohydrates may behave in specific systems as well as demonstrate whether or not such protein can be used to stimulate or replace conventional protein (Siddiq *et al.*, 2009). The food property is characterized of the structure, quality, nutritional value and /or acceptability of a food product. A functional property of food is determined by physical, chemical, and/or organoleptic properties of a food. Prominent properties are water absorption capacity and oil absorption capacity, bulk density, viscosity and swelling index (Aremu *et al.*, 2007). Table 2 shows the functional properties of the blends of seasoning powders made from ALB, chicken and mushroom.

There was no significant difference in the Bulk densities of the seasoning powders at 5% significance level. The blend with equal quantities of the three powders (I, C, M) had a bulk density of  $0.59 \pm 0.04$ g/ml. Bulk density is a vital parameter that determines the suitability of flours for the ease of packaging and transportation of particulate foods as well as for infant formulations (Shittu *et al.*, 2005). Bulk density depends on the particle size and initial moisture content of flours. The compressibility of the powder determines the appearance of a container's contents upon reaching the consumer, and may serve as an index of cohesion (see below). Bulk density control, therefore, is a prime objective of many food processes, especially spray drying and grinding. bulk density is one of its most important characteristics.



It determines the choice of the container size and strength of the reconstituted food, if prepared from a given volume (e.g., tablespoon, cup, etc.). (Micha, 2003) Differences in the bulk density is generally affected by the particle size and the density of the flour and it is very important in determining the packaging requirement, material handling and application in wet processing in the food industry. (Adebowale *et al.*, 2008). The fact that all the seasoning powders had statistically the same bulk densities means that their packaging requirements should be the same and their Hausner ratios and Carr's indices would also be similar. Hausner ratio and Carr index of compressibility are indicators of flowability and cohesiveness. Lower values of these two properties indicate better flowability and lower cohesiveness. (Geldart *et al.*, 2006)

Swelling index values ranged from  $1.60 \pm 0.04$  in sample 1 (16.6/16.6/66.7) (I/C/M) to  $1.90 \pm 0.01$  in sample 3 (66.7/16.6/16.6) (I/C/M). The blend with equal parts of the three ingredients had a swelling index of  $1.73 \pm 0.21$ . It was observed that the blend with more mushroom (16.7/16.7/66.6 I/C/M) had a swelling index of  $1.60 \pm 0.04$ , while the blend with more iru (66.7/16.6/16.8 I/C/M) had a swelling index of  $1.90 \pm 0.01$ . Finally the blend with more chicken powder (16.6/66.7/16.6 I,C,M) had a swelling index of  $1.72 \pm 0.02$  which was not significantly different from the blend with equal quantities of the three ingredients and 95 % confidence level.

Water absorption capacity or characteristics represent the ability of a product to associate with water under conditions where water is limited (Singh, 2001). Water absorption capacity is a critical function of protein in various food products like soups, dough and baked products. The absorption capacity of water by flour is a useful functionality for protein utility in aqueous food formulations, especially those involving high handling (Osungbaro *et al.*, 2010).

Water absorption capacity of the powders in this study was observed to increase with increasing quantities of shiitake mushroom powder. Therefore the blend containing more mushroom powder 1 (16.6/16.6/66.7) (I/C/M) had a water absorption capacity of  $40.16 \pm 0.04$  %. The matrix of protein with water is advantageous to properties, such as hydration, swelling power, solubility and gelation (Etudaiye *et al.*, 2009).

The OAC also makes flour suitable in facilitating enhancement in flavor and mouth feel when used in food preparation. Due to these properties, the protein probably could be used as functional ingredient in foods such as whipped toppings, sausages, chiffon dessert, angel and sponge cakes etc. The ability of the proteins of these flours to bind with oil makes it useful in food system where optimum oil absorption is desired. This makes flour to have potential functional uses in foods such as sausage production and seasoning cube production. Oil absorption capacity values ranged from  $15.48^a \pm 0.37\%$  in the blend with (16.6/16.6/66.7) (I/C/M) to  $18.31^c \pm 0.02\%$  in the seasoning powder with more chicken powder (16.6/66.7/16.6) (I/C/M). The blend with equal quantities of the three ingredients had an oil absorption capacity of  $17.82 \pm 0.10\%$ . The high oil absorption capacity observed in the flour samples suggests the presence of good lipophilic constituents and therefore may be suitable as recipes for sausage, soups, and cakes (Aremu *et al.*, 2006). High oil absorption capacity observed in these blends suggests the presence of good lipophilic constituents and therefore may be suitable as recipes for sausage, soups, and cakes (Aremu *et al.*, 2006).

Peak viscosity is often correlated with the final product quality. It also provides an indication of the viscous load likely to be encountered during mixing. Higher swelling index is indicative of higher peak viscosity (Maziya-Dixon *et al.*, 2004). The highest viscosity  $375.86 \pm 0.55$  RVU was registered in the sample containing 16.6% Iru, 66.7% chicken powder and 16.6% mushroom powder. The lowest viscosity  $311.06 \pm 0.02$  RVU was observed in the sample (66.7/16.6/16.6) (I/C/M).

#### **Proximate Composition of Blends of Seasoning Powders**



Table 3 shows the proximate composition of the seasoning powder blends produced from iru, chicken and shiitake mushrooms. There was no significant difference in the moisture contents of the seasoning powder blends. Moreover all the seasoning powders had moisture contents lower than 6%.. Generally, foods high in moisture are susceptible to microbial attack. Therefore, the low moisture exhibited in the flavouring powders may make them stable to microbial growth and thus confer on them extended shelf life. Lower moisture content too will prevent caking of the powder during storage and avoid the use of chemical anticaking agents.

It was observed that protein content generally increased with increasing concentrations of iru and chicken. The highest protein content  $50.32 \pm 1.18\%$  was observed in the blend with high concentration of chicken powder (16.6/66.7/16.6) (I/C/M). The blend with equal quantities of all three ingredients had a protein content of  $35.28 \pm 0.41\%$ . Proteins are essential for body building.

Lipid or crude fat content in the seasoning powders ranged from  $6.11 \pm 0.01\%$  in the powder that contained more mushroom (16.6/16.6/66.7)(I/C/M) to  $15.41 \pm 0.37\%$  in the blend that contained more chicken (16.6/66.7/16.6) (I/C/M). Lipid content generally increased with increasing concentration of chicken and ALB. Lipids contribute to flavour in food products. Crude fibre on the other hand increased with increasing concentration of mushroom powder. Thus the highest crude fibre content  $9.47 \pm 0.28\%$  was observed in 1(16.6/16.6/66.7) (I/C/M) the sample with more mushroom(16.6/16.6/66.7)(I/C/M) .Fibre in foods generally offers a variety of health benefits and it is essential in reducing the risk of chronic diseases (Food Science Avenue, 2008)..

Ash content ranged from  $6.25^a \pm 0.12\%$  in the powderblend with equal quantities of all three ingredients (33.3/33.3/33.3)(I/C/M) to  $7.12 \pm 0.01\%$  in the blend with more chicken powder(16.6/66.7/16.6) (I/C/M). High ash content implies high mineral contents which helps retard the growth of certain micro-organisms and some minerals are necessary in diets for health benefits (McClements, 2003).

The blend with equal quantities of the three ingredients had a carbohydrate content of  $41.72 \pm 0.55\%$ . This same blend had a total energy content of  $407.01 \pm 0.96 \text{ Kcal/100g}$ . the lowest carbohydrate content  $24.04^c \pm 1.47\%$  was observed in the blend with more chicken powder (16.6/66.7/16.6) (I/C/M) and the highest carbohydrate content  $60.41^g \pm 0.29\%$  was seen in the blend with more mushroom (16.6/16.6/66.7).(I/C/M).

Total energy contents in the seasoning powders ranged from  $365.59 \pm 0.41 \text{ Kcal/100g}$  in the blend with more mushroom powder (16.6/16.6/66.7).(I/C/M) to  $437.05 \pm 1.51 \text{ Kcal/100g}$  in the blend with more iru (66.7/16.6/16.6).

**Table 2: Functional Properties of Blends of Seasoning Powders**

Sample I/C/M %	Bulk Density (g/ml)	Swelling Index	WAO (%)	OAC (%)	VISCOSITY(RVU)
1(16.6/16.6/66.7)	0.62 <sup>a</sup> ±0.01	1.60 <sup>a</sup> ± 0.04	40.16 <sup>d</sup> ± 0.04	15.48 <sup>a</sup> ± 0.37	342.95 <sup>c</sup> ± 1.47
2(16.6/66.7/16.6)	0.63 <sup>a</sup> ±0.00	1.72 <sup>b</sup> ±0,02	34.63 <sup>b</sup> ± 0.04	18.31 <sup>c</sup> ± 0.02	375.86 <sup>d</sup> ± 0.55
3(66.7/16.6/16.6)	0.63 <sup>a</sup> ±0.04	1.90 <sup>c</sup> ± 0.01	32.67 <sup>a</sup> ± 0.40	18.24 <sup>c</sup> ± 0.18	311.06 <sup>a</sup> ± 0.02
4(33.3/33.3/33.3)	0.59 <sup>a</sup> ±0.04	1.73 <sup>b</sup> ± 0.21	36.23 <sup>c</sup> ± 0,05	17.82 <sup>b</sup> ±0.10	328.84 <sup>b</sup> ± 2.50

Results are reported as mean ± standard deviation of triplicate determinations. Means with the same superscript are not significantly different at p<0.05  
 Key: I is Iru, C is chicken and M is shiitake Mushroom

**Table 3: Proximate Composition of Seasoning Powders Blends**

Sample I/C/M %	Moisture %	Crude Protein %	Crude Fats %	Crude Fibre %	Ash %	Carbohydrates %	Total Energy Kcal/100g
1(16.6/16.6/66.7)	5.66 <sup>a</sup> ±0.04	17.24 <sup>a</sup> ± 0.18	6.11 <sup>a</sup> ± 0.01	9.47 <sup>d</sup> ±0.28	6.77 <sup>b</sup> ± 0.17	60.41 <sup>c</sup> ±0.29	365.59 <sup>a</sup> ± 0.41

2(16.6/66.7/16.6)	5.12 <sup>a</sup> ± 0.01	50.32 <sup>c</sup> ±1.18	15.41 <sup>d</sup> ± 0.37	3.11 <sup>b</sup> ± 0.02	7.12 <sup>b</sup> ± 0.01	24.04 <sup>a</sup> ±1.47	436.13 <sup>c</sup> ±1.90
3(66.7/16.6/16.6)	5.21 <sup>a</sup> ± 0.01	35.11 <sup>b</sup> ±10.11	13.57 <sup>c</sup> ±0.25	1.34 <sup>a</sup> ± 0.08	6.36 <sup>a</sup> ±0.22	43.62 <sup>b</sup> ± 0.13	437.05 <sup>c</sup> ± 1.51
4(33.3/33.3/33.3)	5.01 <sup>a</sup> ±0.01	35.28 <sup>b</sup> ± 0.41	11.01 <sup>b</sup> ±0.03	5.77 <sup>c</sup> ± 0.07	6.25 <sup>a</sup> ± 0.12	41.72 <sup>e</sup> ± 0.55	407.01 <sup>b</sup> ±0.96

Results are reported as mean ± standard deviation of triplicate determinations. Means with the same superscript are not significantly different at p<0.05  
Key: I is Iru, C is chicken and M is shiitake Mushroom

### Umami Compounds in Seasoning Powders

The compounds that elicit umami taste present in the different blend formulations are presented in table 4. All the seasoning powder blends had negligible quantities of MSG with values ranging from  $0.18 \pm 0.04\%$  in the blend 16.6/66.7/16.6 (I/C/M)  $0.24^c \pm 0.02\%$  in the powder with equal quantities of the three ingredients F(33.3/33.3/33.3). There was no significant difference in the MSG content of this sample with that of the sample with more chicken (66.7/16.6/16.6). Glutamic acid content was highest  $443.39^i \pm 1.44\text{mg}/100\text{g}$  in the powder with more iru (66.7/16.6/16.6) and  $290.61^a \pm 0.33$  in the powder with equal quantities of the three ingredients. This same blend with equal quantities of the three ingredients also had the lowest IMP content  $56.18 \pm 0.16$  while the highest IMP content  $70.96 \pm 0.17$  was observed in the blend that contained more of iru (66.7/16.6/16.6 I/C/M). Guanosine monophosphate (GMP) however was found to be most prominent  $209.10 \pm 0.24$  in the blend with more chicken 16.6/66.7/16.6 (I/C/M). The lowest value  $99.30 \pm 0.16$  of GMP was found in the blend that contained equal quantities of the three ingredients.

**Table 4: Umami Taste Compounds in Seasoning Powders**

Sample I/C/M %	MSG %	Glutamic acid (mg/100g)	IMP (mg/100g)	GMP (mg/100g)
1(16.6/16.6/66.7)	$0.22^b \pm 0.08$	$363.44^c \pm 1.17$	$62.79^b \pm 0.46$	$192.95^c \pm 0.65$
2(16.6/66.7/16.6)	$0.18^a \pm 0.04$	$352.40^b \pm 0.32$	$63.34^b \pm 0.18$	$209.10^c \pm 0.24$
3(66.7/16.6/16.6)	$0.24^c \pm 0.01$	$443.39^d \pm 1.44$	$70.96^c \pm 0.17$	$110.62^b \pm 0.33$
4(33.3/33.3/33.3)	$0.24^c \pm 0.02$	$290.61^a \pm 0.33$	$56.18^a \pm 0.16$	$99.30^a \pm 0.16$

Results are reported as mean  $\pm$  standard deviation of triplicate determinations. Means with the same superscript are not significantly different at  $p < 0.05$

### Sensory Properties of the Seasoning Powders

Table 5 shows the sensory properties of the seasoning powders prepared by blending Iru, chicken and shiitake mushroom powders. The highest score for appearance was observed in the blend containing 16.6% iru, 66.7% chicken and 16.6% mushroom. On the other hand the blend containing 66.7% iru 16.6% chicken and 16.6% mushroom had the highest score for taste and appearance. This could be as a result of its high glutamic acid content. This particular blend was therefore considered to be the most acceptable product.

**Table 5: Sensory Properties of the Seasoning Powders**

Sample I/C/M %	Appearance	Taste	General Acceptability
1(16.6/16.6/66.7)	$7.11^a \pm 0.08$	$6.31^a \pm 1.17$	$6.27^a \pm 0.46$
2(16.6/66.7/16.6)	$8.18^c \pm 0.24$	$7.22^b \pm 0.32$	$6.95^b \pm 0.18$

3(66.7/16.6/16.6)	7.52 <sup>a</sup> ± 0.21	8.76 <sup>d</sup> ± 1.44	8.22 <sup>d</sup> ± 0.17
4(33.3/33.3/33.3)	7.98 <sup>b</sup> ± 0.02	7.67 <sup>c</sup> ± 0.43	7.55 <sup>c</sup> ± 0.16

### Conclusion

Food seasoning powder can be produced from a combination of fermented African locust bean, chicken and mushroom powders. The most acceptable product was that which contained 66.7% locust bean, 16.6% chicken and 16.6% mushroom powders. This seasoning powder blend has a high protein content, and when used as a spice, it can provide good nutrition for human health, been selected as the best formula with a good combination of other ingredients. This product has higher protein content than seasoning powder made from other sources, so it is very suitable for people with protein needs and can be used to replace part of seasoning powder from meat/fish/shrimp. This seasoning powder also contains very little monosodium glutamate unlike other seasoning powders commonly sold in the markets. This provides a healthier solution for people sceptical about consuming the controversial MSG which is said to be the cause of many health issues. However, there also needs to be technological improvements applied and tested on a larger production scale to develop this new product for the food technology industry with the goal of utilizing locally available products for food seasoning.

### Knowledge contribution

.....  
.....  
.....

### Recommendation

.....  
.....  
.....  
.....  
.....

## References

- Akabanda Fortune, Charles Parkouda, Jennifer Suurbaar, Addai-Mensah Donkor and 1James Owusu-Kwarteng. (2018). EFFECTS OF MECHANICAL DEHULLING ON MICROBIOLOGICAL CHARACTERISTICS AND CHEMICAL CHANGES DURING PROCESSING OF *Parkia biglobosa* SEEDS INTO dawadawa, A WEST AFRICAN ALKALINE FERMENTED CONDIMENT. *Journal of Ghana Science Association*, 17(2), 14-19.
- Aswal, D.S. Bhakuni, A.K. Goel, K. Kar, B.N. Mehrotra, K.C. Mukherjee. ((2004)). *Screening of Indian plants for biological activity*.
- Conway and Byrne . (2013). Determination of the Total Volatile Basic Nitrogen (TVBN) in Fish according to Conway and Byrne method . *VELP Scientifica*.
- Elman I, Ana P, Cristiane Kr, Maria E. (2013). Characterization of umami taste sensitivity in children with and without cancer;. *Journal of Human Growth and Development*, 23(2), 136-143.
- Food.gov.uk. (2010). "Current EU approved additives and their E Numbers." (<http://www.food.gov.uk/safereating/chemsafe/additivesbranch/numberlist>) .
- Geha R S, Beiser A, Ren C, Patterson R, Greenberger PA, et al. . (2000). ) Review of Alleged Reaction to Monosodium Glutamate and Outcome of a Multicenter Doubl. *Journal of Nutrition* , 10580-10625.
- Jan, S., Rafiq, S.I., Saxena, D.C.: . (2015). Effect of physical properties on flow ability of commercial rice flour/powder for effective bulk handling. (In: *International Conference on Advancements in Engineering and Technology* (pp. 0975–8887). Int. J. Comput. Appl.
- Mouritsen, O. G. & Styrbaek, K. Umami: . ( 2014). Unlocking the Secrets Behind the Fifth Taste . (Columbia University Press, New York).
- Ninomya K, Rozin E. . (2004). *The fifth taste of human being umami*: . london : Umami Internacional Center.the world. Londres: Cross Media Limited;.
- Phongpaichit, J. Nikom, N. Rungjindamai, J. Sakayaroj, N. Hutadilok-Towatana, V. Rukachaisirikul and K. Kirtikara. (2007). Antioxidant levels . *FEMS Immunol. Med. Microbiol*, 51(3), 517-525.
- Saxena, J. Saxena, R. Nema, D. Singh, A. Gupta.. (2013). Phytochemistry of Medicinal Plants. . *Journal of Pharmacognosy and Phytochemistry*, 168-182.
- Taghvaei and S. M. Jafari,. (2015). Plant antioxidants. *International Journal of Food Science and Technology*, 52(3), 1272-1282.
- The United States Pharmacopeia. . (2002.). . *USP25/NF20* . Rockville:: United States Pharmacopeial Convention, Inc.
- ABDEL-KADER, M. R.1, A. A. ASRAN2, A. A. R. AL-GENDY1 and K. E. KALEAL3. (2014). FOOD PREFERENCE FOR ALBINO RATS AND ALBINO MICE UNDER LABORATORY CONDITIONS. *Egypt. Journal of Agricultural . Research* , 92 ( 4 ),, 92(4), 1279-1289.
- Aburto, N. J., Ziolkovska, A., Hooper, L., Elliott, P., Cappuccio, F. P., & Meerpohl, J. K. (2013 ). Effect of lower sodium intake on health: Systematic review and meta- analyses. . *BMJ*, 346, , 6.
- Acharya, R., Mishra, M., Patro, J. and Panda, M. K. . (2008). Effect of Vitamins C and E on Spermatogenesis in Mice Exposed to Cadmium. *Reproductive Toxicology*, 25,, 2(4), ): 89 – 95.
- Achi, O. K. (2005). Reviewon on Traditional fermented protein condiments in Nigeria . *African Journal of Biotechnology*, 4(13), 1612-1621,.
- Adebowale, A.A., Sanni, L.O., and Awonorin, S.O. ., (2008). . Effect of texture modifiers on the physicochemical and sensory properties of dried fufu. *Food Science and Technology International*, 11(5), 373-382.
- ADEJUMO, A.A; AZEEZ, I.O; GEPLY, J.J. OBOITE; F.O. . (2013). PROCESSING, UTILIZATION AND CHALLENGES OF AFRICAN LOCUST BEAN (*Parkia biglobosa*, Jacque Benth) IN ARIGIDI AKOKO, ONDO STATE, NIGERIA. *Journal of Agriculture and Social Research*, 13(1).
- Afolayan M., Bawa G. S., Sekoni A. A., Abeke F. O., Inekwe V. O., Odegbile E. O. (2014). Phytochemical and nutritional evaluation of locust bean fruit puld. *Journal of Emerging Trends in Engineering and Applied Sciences*, 5, 44–47.



- Agbaire PO, Oyewole EO. . (2012). Nutritional and anti- nutritional levels of some local vegetables from Delta State, Nigeria. *African Journal Food Sciences*, 6, 8-11.
- Ajayi,O.A.,Akinrinde,O.O.AkinwuniO.O. (2015). Towards the development of shelf stable 'iruu' (Parkia biglobosa) condiment bouillon cubes using corn,cassava and potato starch extracts as binders. *Nigerian Food Journal*, 38(1), 63-72.
- Akçay, C., Ceylan, F., Arslan, R. ( 2023.). Production of oyster mushroom (Pleurotus ostreatus) from some waste lignocellulosic materials and FTIR characterization of structural changes. . *Science Report*, 13, 12897.
- Alli. (2009). Determination of the chemical composition of Senna-siamea (cassia leaves). *Pakistan Journal of Nutrition*, 8, 119-121.
- Andrea, P. (2005). The Cultural History of Plants. . *Routledge. Prance, Ghilleen; Nesbitt, Mark (eds.)*., 30. ISBN 0415927463.
- Ansari,S.A.,Rustamani,M.A.,Kakar,A.W.,Khooharo,A.A.,Fozia, D. and Baloch, H.B. (2005). Grain losses caused by house rat, Rattus rattus L. and its control. *Pakistan Journal of Zoology*, 37(1), 33-38.
- Antonella Dalle Zotte 1,\* , Elizabeth Gleeson 1, Daniel Franco 2 , Marco Cullere 1 and José Manuel Lorenzo 2. (2020). Proximate Composition, Amino Acid Profile, and Oxidative Stability of Slow-Growing Indigenous Chickens Compared with Commercial Broiler Chickens. *Foods*, 9, 546.
- AOAC. ((2000)). Official Methods of Analysis,,. *Association of Official Analytical Chemists*, (p. 17th edn. ). Washington DC .
- AOAC. (2016). . *Official methods of analysis, association of official analytical chemists. 20th ed.* Washington D.C.: AOAC.
- Arampath Palitha and Silva G. (2019). Development of Spice cubes for chicken and fish curries. *Annual Symposium of export Agricultural crops* , (p. 8). Sri Lanka.
- Aremu1, E. Y. Awala2, O. D. Opaluwa2, R. Odoh1 and T. O. Bamidele3. (2015). Effect of Processing on Nutritional Composition of African Locust Bean. *Communications in Applied Sciences*, 3(1), 22-41.
- Augustine I Airaodion1\*, Emmanuel O Ogbuagu2, Etinosa U Osemwowa3, Uloaku Ogbuagu1, Chimdi E Esonu1, Aanu P Agunbiade1, Davidson Okereke1 and Abiodun P Oloruntoba1. (2019). Toxicological Effect of Monosodium Glutamate in Seasonings on Human Health. *Global Journal of Nutrition & Food Science*, 1(5), 1-9.
- Australian. (2010). *standard code food standards new zealand australia*.
- Bak, W.C., Park, J.H., Park, Y.A., Ka, K.H., . (2014). . Determination of glucan contents in the fruiting bodies and mycelia of Lentinula edodes cultivars. *Mycobiology* , 42,, 301–304.
- Bakare RI, Magbagbeola OA, Akinwande AI, Okunowo OW.. ( 2010). Nutritional and chemical evaluation of Momordica charantia. . *Journal of Medicinal Plants Research* 4, , 2189-2193.
- Barnett SM, Diako C, Ross CF . . (2019). Identification of a salt blend: Application of 443 the electronic tongue, consumer evaluation, and mixture design 444 methodology. *J Food Science* 2019, 84 445, 2, 327-338.
- Bellisle, F. (1999). Glutamate and the Umami taste: sensory, metabolic, nutritional and behavioural considerations a review of the literature published in the last 10 years. :*Neurosc Biobehav Rev*, 23(3), 423-38.
- Beyreuther K, Biesalksi HK, Fernstrom . (2007). Consensus meeting: monosodium glutamate – an update. . *European Journal of Clinical Nutrtrion*, 61, 304-313.
- Bich Thuy Thi, N., Ve Van, L., Huyen Trang Thi, N., Luyen Thi, N., Thuy Trang Thi, T., Nghien Xuan, N., . ( 2021). . Nutritional requirements for the enhanced mycelial growth and yield performance of Trametes versicolor. *Journal of Applied Biology and Biotechnology*.
- Black. (2003). Zinc deficiency, infectious disease and mortality in the developing world. *Journal of Nutrition*, 133, 1485S-9S.

- Boardman, A., Greenberg, D., Vining, A., & Weimer, D. . (2011). *Cost-Benefit Analysis: Concepts and Practice*. 4th edition.
- Burdge, G. C. and Calder, P. C. (2005). Alpha-linolenic acid metabolism in adult humans: the effects of gender and age on conversion to longer-chain polyunsaturated fatty acids. *Universal Journal of lipid Science and Technology*, 107,, 426-439.
- Buzayehu Desisaa,c, Diriba Muletaa,\* , Mulissa Jidab, Tatek Dejenec, Abayneh Goshub, Tadesse Negid, Pablo Martin-Pintoe. (2024). Improvement of nutritional composition of shiitake mushroom (*Lentinula edodes*) using formulated substrates of plant and animal origins. *Future Foods*, 9, 2666.
- Canada Gov Inspection. (2014). *protein, elements in nutrition* . canadia gov inspection .
- Carver J, Walker W. (1995). . The role of nucleotides in human nutrition. *Journal of Nutritional Biochemistry*, 58-72.
- Cheung, P. (2013). Mini-review on edible mushrooms as source of dietary fiber: preparation and health benefits. *Food Science and Human Wellness* , 2, 162–166.
- Cobos, J.D.V., Viej'o, F.G., Cobo, J.C., Sulca, R.L., Verdugo, D.N., Moncayo, M.F.G., Endara, A.G.,. (2021). Chemical and productivity characterization of parental and hybrid strains of *Lentinula edodes* cultivated in different agricultural residues. *Emirate Journal of Food Agriculture* , 260, 263.
- Cristiane Vieira Helm, F.B., Charles Windsnon Isidoro Haminiuk, E.A.D.L., Marcelo Barba Bellettini, . (2019). Influence of cultivation methods on the chemical and nutritional characteristics of *Lentinula edodes*. . *Emiater. Journal of Food Agriculture.*, 1006.
- Crowe. (2009). *UMAMI in Foods:* . : EAL Workgroup.
- CuiL,LiDJ,LiuCQ.. ( (2012)). Effectoffermentationonthenutritive valueofmaize. . *International Journal of Food Science and Technology*, 47:, 755–60.
- Dasagupta S,Bussen N,Probst K. (2011). The economics of small scale soft shell crayfish production. *Aquacultural Economics and Management*, 15(3), 166-192.
- Dermiki, M., Phanphensophon, N., Mottram, D. S., & Methven, L.. (2013). Contributions of non-volatile and volatile compounds to the umami taste and overall flavour of shiitake mushroom extracts and their application as flavour enhancers in cooked minced meat. *Food Chemistry*,, 141, 77-83.
- Diawara, B. and Diasso, K. (2004). Bilan de mati`eres dans le proc´ed´e traditionnel de production du Soumbala. Valorisa- tion technologique et nutritionnelle du n´er´e ou Parkia biglo- bosa (Jacq.) Benth: une esp`ece agro foresti`ere, Diawara B. and Jakobsen M.(eds.), . *DANIDA-KVLCNRST/IRSAT*,, 132-133.
- Dimopoulou, M., Kolonas, A., Mourtakos, S., Androutsos, O., Gortzi, O. (2022). Nutritional composition and biological properties of sixteen edible mushroom species. *Applied. Scence*. , 12, 8074.
- Divrikli, U., Horzum, N., Soylak, M.,. (2006). Trace heavy metal contents of some spices and herbal plants from western Anatolia. *Turkey. International Journal of Food SciencesandTechnique*, 41, 712-716.
- Domínguez, R.; Pateiro, M.; Gagaoua, M.; Barba, F.J.; Zhang, W.; Lorenzo, J.M. . (2019). A comprehensive review on lipid oxidation in meat and meat products. . *Antioxidants*, 8,, 429.
- Dyta, V. P. (2018). Analysis of disodium 5'-guanylate and disodium 5'-inosinate as flavor enhancer in food spices by thin-layer chromatography-densitometry,.. *International Journal of Applied Pharmaceutics*, 10(Special Issue).
- Edema M.O. and Fawole O. y,. (2006). . Evaluation and Optimization of Critical Control Points in the Production of Iru. . *Research Journal of Microbiolog*, 1(16), 503-511.
- Ejigui J, Savoie L, Marin J, Desrosiers T. (2005). Beneficial changes and drawback of a traditional fermentation process on chemical composition and anti- nutritional factors of yellow maize (*Zea mays*). *Journal of Biological Science*, 5, 590– 6.

- Elkanah, F.A., Oke, M.A., Adebayo, E.A... (, 2022). Substrate composition effect on the nutritional quality of *Pleurotus ostreatus* (MK751847) fruiting body. *Heliyon*, 8.
- Esenwah CN, Ikenebomeh MJ. (2008). Processing effects on the nutritional and anti-nutritional contents of African locust bean (*Parkia-biglobosa* (Benth) Seed. *Pakistan Journal of Nutrition*, 7(2), 214-217.
- Etudaiye, H.A., Nwabueze, T.U. and Sanni, L.O). . (2009). Quality of fufu processed from cassava mosaic disease (CMD) resistant varieties. *African Journal of Food Science*,, 3(3), 61-67.
- Ezugwu EN, Okoye JI, Ene GI. (2018). Effects of processing on the nutrient and anti- nutrient contents of African locust bean (*Parkia biglobosa*) flour. . *International Journal of Food Science and Nutrition*, 3(4), 107-111.
- FAO,. (2004., September , 21–30). Vitamin and mineral requirements in human nutrition. (I. 9. 4, Ed.) *Second edition: report of a joint FAO/WHO expert consultation*,, p. .154612 3.
- Feng, X.M., A.R.B. Eriksson and J. Schnure. (2005.). Growth of lactic acid bacteria and *Rhizopus oligosporus* during barley tempeh fermentation. *International . Journal of Food Microbiology*, 104, 249-256.
- FLAVIA POP, CRISTINA ANAMARIA SEMENIUC. (2002). Oxidative Stability of Avocado and Peanut Oils Under Different Heating Temperatures. *Revista de Chimie* <https://revistadechimie.ro> [https://doi.org/10.37358/Rev.Chim.1949, 73\(1\), 1-8](https://doi.org/10.37358/Rev.Chim.1949, 73(1), 1-8).
- Food Science Avenue . (2008). . *Crude fibre*. Retrieved from [www.fooscience-avenue.com/2008/04/crude-fibre.html?m=1](http://www.fooscience-avenue.com/2008/04/crude-fibre.html?m=1) Assessed February 2014.
- Friedrich-Ebert-Allee. (2017). *Poultry Production in Cameroon:How the import restriction affects the Cameroonian poultry sector*. Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) GmbH.
- Gao, S., Huang, Z., Feng, X., Bian, Y., Huang, W., Liu, Y. ( 2020). . Bioconversion of rice straw agro-residues by *Lentinula edodes* and evaluation of non-volatile taste compounds in mushrooms. *Science . Report , 10*, 1814.
- Garcia, E.J., Alencar, S.M.D., Reis, A., Loguercio, A.D., Helena, R. and Grande, M. (2012). Antioxidant Activity by DPPH Assay of Potential Solutions to be Applied on Bleached Teeth. *Brazilian Dental Journal*,, 23, 22–27. <https://doi.org/10.1590/S0103- 6440>.
- Geldart, D., Abdullah, E.C., Hassanpour, A., Nwoke, L.C., Wouters, I.: . (2006). Characterisation of powder flowability using measurement of angle of repose. . *China Particuol.*, 4(3-4), 104–107.
- Gernah D. I., Atolagbe M. O., Echeowo C. C. . (2007). Nutritional composition of the African locust bean (*Parkia biglobosa*) fruit pulp. . *Nigerian Food Journal*, 25, 190–196.
- Gernah, D.I., Inyang, C.U., & Ezeora, N.L. . (2007). Incubation and fermentation of African locust beans (*Parkia biglobosa*) in production of "dawadawa". J. *Journal of Food Processing and Preservation*,, 31(2), 227-239.
- Gernah,D.I.,Inyang,C.,Ezeora,N. (2005). Effect of incubation materials on fermentation of African locust beans(*Parkia biglobosa*) in the production of dawadawa. *Nigerian Food Journal*.
- Gilbert, P. A., & Heiser, G. . . (2005). Salt and health: The CASH and BPA perspective. , 30(1), 62–69.
- Golden. (2009). Nutrient requirements of moderately malnourished populations of children., . *Food Nutrition Bulletin*, 30, 267-342.
- Guedes M.D.V., Marques M.S., Guedes P.C., Contri R.V., Kulkamp Guerreiro I.C. (2021). The use of electronic tongue and sensory panel on taste evaluation of pediatric medicines: A systematic review. *Pharmaceutical Development and Technology*, 26(2), :119–137.
- Harith, N., Abdullah, N., Sabaratnam, V. (2014). . Cultivation of *Flammulina velutipes* mushroom using various agro-residues as a fruiting substrate. *Pesqui. Agropecu. Bras.*, 49, 181–188.
- Harmita S. (2006). *Textbook of Physicochemical*. Depok. Jakarta: Faculty of Pharmacy, Universitas Indonesia; 2006.
- Harmita, S. (2004). *Instructions on how to validate the method and how to calculate it*. Jakarta: .Majalah Ilmu Kefarmasian.

- He, F. J., Tan, M., Ma, Y., & MacGregor, G. A., . (2020). *Journal of the American College of CardiologyJACC* , 75(6), 632–647.
- Heise H., Crisan A., Theuvsen L. (2015). The poultry market in Nigeria: Market Structure and potential for investment in the market. *International Food and Agribusiness Management Review* , 18( special issue A:), 197–222.
- Hough G, Wakeling I, Mucci A, Chambers E, Gallardo IM, Alves LR: (6):. (2006). Number of 358 consumers necessary for sensory acceptability tests. *Food Qual Pref* 359 2006, 6, 522-526. [http://www.fao.org/docrep/005/y4640e/y4640e02.htm#P919\\_1309276](http://www.fao.org/docrep/005/y4640e/y4640e02.htm#P919_1309276). (n.d.).
- Ikeda. (1908). *Patent No. 14804*. Japanese.
- Ikeda, K. (2022). *Umami Information Center* .
- Ilana E., Ana Paula G., Cristiane K., Maria Elisabeth M. (2013). CHARACTERIZATION OF UMAMI TASTE SENSITIVITY IN CHILDREN WITH AND WITHOUT CANCER. *Journal of Human Growth and Development*, 23(2), 136-143.
- International Glutamate Information Service.. (2017). *Monosodium Glutamate & Umami"*. . Retrieved 3 June 2015 Archived from the original on 25 June.
- Irdasari. (2009). Analysis of Consumer Attitudes Toward the Performance of the Masako to Enrich the Flavor. *Thesis. Bogor.* : Faculty of Agriculture, Intitut Pertanian Bogor.
- Islam, S., Karatzas, C., Rodenhiser, A., Alwattari, A., Huang, Y., & Turcotte, C. . (2006). Methods and apparatus for spinning spider silk protein:. *Google Patents*.
- Janick, J. (2008). *Parkia biglobosa* African Locust Bean. *The encyclopedia of fruit & nuts*, 395- 400. Wallingford, CABI North American Office., U.K.
- Jump, D. B. (2002). The biochemistry of n-3 polyunsaturated fatty acids. *Journal of Biology and Chemistry* , 277, 8755-8758.
- Karou, S., Tchacondo, T., Tchibozo, M. D., Abdoul-Rahaman, S., Anani, K., Koudouvo, K.,. (2011). Ethnobotanical study of medicinal plants used in the management of diabetes mellitus and hypertension in the Central Region of Togo..I. *PharmBio*, 49(12), 1286-1297.
- Katz, S. E. ( (2012).). *The Art of Fermentation: An In-Depth Exploration of Essential Concepts and Processes from Around the World.* . (I. 978-1603582865., Ed.) .Chelsea: Green Publishing. .
- Kaur M, Singh . (2006). Relationships between selected properties of seeds, flours, and starches from different chickpea cultivars. . *Int. J. Food Prop*, 9, :597-608.
- Kiers, J L, Van Laeken, A E A, Rombouts, F M, and Nout, M J R . . (2000). In vitro digestibility of Bacillus fermented soya bean. *Intenationsl Journal of Food Microbiology*, 160, 163–169.
- Kim Y, F. Lian, K.-J. Yeum, N. Chongviriyaphan, S.-W. Choi, R. M. Russell and X.-D. Wangr. (2007). Antioxidants. *International Journal of Cancer*, 120(9), 1847-1854.
- Kondoh T, Mallick H, Torii K. (2009). Activation of the gut-brain axis by dietary gluatamate and physiologic significance in energy homeostasis. *American journal of Clinical nutrition.*, 90, :832S-837S.
- Kong, Y., Zhang, L. L., Zhao, J., Zhang, Y. Y., Sun, B. G., & Chen, H. T. .. (2019). Isolation and identification of the umami peptides from shiitake mushroom by consecutive chromatography and LC-Q-TOF-MS. *Food Research International*, 121, 463–470.
- Kourouma, K; Jean C.G; Achille, E.A and Clement,A. . . (2011). Ethnic differences in use values and use patterns of *Parkia biglobosa* in Northern Benin. *Journal of Ethnobiology and Ethnomedicine*, 3.
- Krasnow M, Bunch T, Shoemaker C, Loss C. . (2012). Effects of cooking temperatures on the physicochemical properties and consumer acceptance of chicken stock. . *Journalof Food Science*, 71, :S19-S23.
- Kris-Etherton PM, Hecker KD, Bonanome A, Coval SM, Binkoski AE, Hilpert KF, Griel AE, Etherton TD.. (2002). Bioactive compounds in foods: Their role in the prevention of cardiovascular disease and cancer. . *American Journal of Medicine*, 113, 71-83.

- Krishna VN, Karthika D, Surya DM, Rubini M, Vishalini M, Pradeepa Y. (2010). Analysis of monosodium l-glutamate in food products by high-performance thin layer chromatography. *Journal of Young Pharmacists*, 2, 297-300.
- Kurihara. (2009). Glutamate from discovery as a food Flavorto role as a basic taste (umami). *The American Journal of Clinical Nutrition*, 90(3), 719-722.
- Labuza, T. S. (1970). Water content and stability of low moisture and intermediate moisture foods. *Food Technology*, 24, :543-550.
- Ledesma-Amaro R, Jiménez A, Santos M, Revuelta J. (2013). Biotechnological production of feed nucleotides by microbial strain improvement. . *Process Biochemistry*, 48, 1263-70.
- Li, S., Wang, A., Liu, L., Tian, G., Wei, S., Xu, F.. (2018). Evaluation of nutritional values of shiitake mushroom (*Lentinus edodes*) stipes. . *Journal of Food Measures and characteristics* , 12, 2012–2019.
- Luscombe-Marsh, N. D., Smeets, A. J. & Westerterp-Plantenga, M. S. . (2008). Taste sensitivity for monosodium glutamate and an increased liking of dietary protein. *Br. J. Nutr.*, 99,, 904–908.
- Massey LK, Palmer RG, Horner HT. . (2001). Oxalate content of soybean seeds (Glycine max: Leguminosae), Soya foods and other edible legumes. *Journal of Agricultural and Food Chemistry*, 49, 4262-4266.
- Mateljan, G. (2007.). *The world's healthiest foods: Vitamin C*. . <http://www.whfoods.org/vitaminC> .
- Mau J.L., Tseng J., Wu C.R., Chen C.H., Lin S.D. (2021). Chemical, nutritional, and bioactive compositions of fresh, washed, and blanched shiitake. *Czech Journal of Food Science*, 39(6), 00.
- Maziya-Dixon B, Dixon A.G.O., and Adebowale, A.A. . (2004). Targeting different end uses of cassava: genotypic variations for cyanogenic potentials and pasting properties. *A paper presented at ISTRC-AB Symposium*,, 5-12.
- McClements, D. (2003). . *Analysis of Food Products*. . Retrieved from [www.people.umass.edu/~mcclemen/581Ash&Minerals.html](http://www.people.umass.edu/~mcclemen/581Ash&Minerals.html) Assessed February .
- Media, Umami International Center and International Glutamate Information Service. . (2004). *The fifth taste of human being umami*:. london: the world. London: Cros.
- Mejia E.G., Aguilera-Gutiérrez Y., Martín-Cabrejas M.A. and Mejia L . (2015). . Industrial processing of condiments and seasonings and its implications for micronutrient fortification. *Annals of the New York Academy of Sciences*, 1357(1), 8-28.
- Meydani SN, Barnett JB, Dallal GE, Fine BC, Jacques PF, Leka LS, et al. (2007). Serum zinc and pneumonia in nursing home elderly. *Annual Journal of Clinical Nutrition*, 86, :1167-73.
- Micha, P. (2003). Physical Properties of Food Powders -. *FOOD ENGINEERING*, 1, 1-3.
- Micheal, B. (2020). The Strange Case of Dr Ho Man Kwok. *Colgate Magazine* .
- Ministry of Health, Republic of Indonesia. (2013). . *Regulation of the Head of the Drug and Food Supervisory Agency No 23 n 2013 About the Maximum Limit of the use of Taste Strengthening Food Additives*. . jakarta: Ministry of Health, Republic of Indonesia.
- Ministry of Health, Republic of Indonesia. (2012). *Regulation of Ministry of Health Regarding Food Additives*. . Jakarta. : Ministry of Health, Republic of Indonesia No 033 i.
- Mojet, J., Heidema, J., & Christ-Hazelhof, E. (2004). . Effect of concentration on taste-taste interactions in foods for elderly and young subjects. . *Chemical Senses*,, 29(8), 671–681.
- Mouritsen, O. G. & Khandelia, H. . (2012). Molecular mechanism of the allosteric enhancement of the umami taste sensation. . *FEBS J*. 279,, 3112–3120 .
- Murray R, Granner D, Rodwell V. (2006). *Harper's Illustrated Biochemistry*. (27 ed.). New York: Lange Medical Books/McGraw-Hill;.
- Murtala Y, 1Babandi A, 1Mashi JA, 1Ubayi HM, 2Ibrahim S, 1Ibrahim A, 3Shehu D, 1Alhassan AJ. (2016). Proximate, Vitamins and Mineral Compositions of Locust and Soya Beans-Based Daddawa sold in Kano Metropolis, Kano, Nigeria. *Journal of Biomedical Science*, 1(1), 1-7.
- NATALIA D.K., HARMITA\*, TAUFIQ INDRA RUKMANA. (2018). DEVELOPMENT OF A HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY METHOD FOR ANALYZING DISODIUM 5'-

- GUANYLATE AND DISODIUM 5'-INOSINATE LEVELS IN FLAVOR ENHANCERS. *international Journal of applied Pharmaceutics*, 10(1).
- Nathan, J. (2024). Wet and Dry basis Sumary. In J. Nathan, *LearnChemE*.
- Naylor A.,Morrow A. (2001). *Developmentalreadinessof normal full term infants to progress from exclusive breastfeeding to the introduction of complementary foods*. WashingtonED.C: linkages /Wellstart international.
- Nestle. (2011). *Annual Report of Nestle Nigeria*, . Retrieved from [https://www.nestle-cwa.com/sites/g/files/pyd-noa346/files/asset-library/documents/nestle\\_2011\\_annual\\_report\\_.pdf](https://www.nestle-cwa.com/sites/g/files/pyd-noa346/files/asset-library/documents/nestle_2011_annual_report_.pdf).
- Nguyen Minh Thuy1\*, Nguyen Ngoc Huong Anh1, Nguyen Thi Kim Xuyen1, Nguyen Hoang Yen Vi1, Nguyen Hoang Thuy Quyen1, Tran Ngoc Giao1, Hong Van Hao1, Ngo Van Tai2, Nguyen Van Hoa3. (2024). Development of seasoning powder from foam-mat dried *Artemia franciscana* biomass. *Journal of Applied Biology & Biotechnology*, 12(5), 198-203,.
- Ninomya K, Rozin E. (2004). The fifth taste of human being umami:.
- Ninomya K., Rozin, E. (2007). The fifth taste of human being umami:.. (U. I. Center., Ed.) 80.
- Ninomya K., Rozin, E. (2007). The fifth taste of human being umami:.. (U. I. Center., Ed.) 80.
- Ninomya K., Rozin, E. (2007). *The fifth taste of human being umami:..* (U. I. Center., Ed.) london: Cross Media Limited.
- Nnam, N. (2000). Chemical Evaluation of Multimixes formulated from some Local Staples for Use as Complementary Foods in Nigeria. *Plant Food for Human Nutrition*, 55, 255-263.
- Nnam, N. M., & Obiakor, P. N. (2003). Effect of fermentation on the nutrient and anti nutrient composition of baobab (*Adansonia digitata*) seeds and rice( *Oryza sativa*) grains. *Ecology and Food and Nutrition*, 42(4-5), 265-277.
- Non-Wood. (2009). *News An Information Bulletin on Non-Wood Forest Products*. ItalyRome: published by FAO,.
- Ntui, V. O., Uyoh, E. A., Urua, I. S., Ogbu, U., & Okpako, E. C. . (2012). Regeneration of *Parkia biglobosa* Benth.: An important tree species of Africa. *Journal of Microbiology and Biotechnology Research*, 2(1), 169-177.
- Nwosu, O. M. (2008). Vitamins, phytochemicals and toxic elements in the pulp and seed of raphia palm fruit (*Raphia hookeri*). *Fruits*, 63, 297–302.
- Odunfa, S.A. and O.B. Oyewole. (1986. ). Identification of *Bacillus* species from iru, a fermented African locust bean product. . *Journal of Basic Microbiology*, 26, 101-108.
- Ojewumi, M.E., Omoleye, J.A., Ajayi, A.A. and Obanla, O.M. (2022). "Molecular Compositions and Morphological Structures of Fermented African Locust Bean Seed ( *Parkia biglobosa* )", . *Lett. Appl. NanoBioscience*, ., 22(1), . 3111–3119.
- Okafor UI, Omemu AM, Obadina AO, Bankole MO, Adeyeye SAO.. (2018). Nutritional composition and anti-nutritional properties of maize ogi co-fermented with pigeon pea. *Food Science and Nutrition*, 6, 1–15.
- Olalusi, A.P., Omosebi, M.O. and Agbola, O.Y.,. (2019,). Influence of Drying Methods on the Drying Characteristics and Nutritional Quality of Fermented Locust Beans. *International Journal of Environmental and . Agricultural. Biotechnology* , 4(19), 1695–1703.
- Onnyi O., Odediran O. F., Ajuebor N. 22:.. (2004). Federal Institute of Industrial Research, I. . *Nigeria Food Journa*, 22, 203–208.
- Onyeka, E.U., Okehie, M.A. (2018). Effect of substrate media on growth, yield and nutritional composition of domestically grown oyster mushroom (*Pleurotus ostreatus*). *African Journal of Plant Science*, 12, 141–147.
- Oodee Blessing B.1, Ojoye N. Briggs1\*, Adline E. Ben-Chioma1, Edna O. Nwachuku1. (2023). Effects of Administration of Bouillon Cubes on Insulin Resistance, Lipid Profile and Renal Function Parameters in Female Albino Rats. *Annals of Innovation in Medicine (AIM) ISSN:*, 2(1), 1-9.



- Osungbaro, T.O., Jimoh, D. and Osundeyi, E. . (2010). . Functional and pasting properties of composite cassava-sorgium flour meals. *Agriculture and Biology Journal of North America*, 1(4), 715-720.
- Ozcelik, E., Peksen, A. . (2007). Hazelnut husk as a substrate for the cultivation of shiitake mushroom (*Lentinula edodes*). . *Bioresour. Technol*, 98,, 2652–2658.
- Paul, M. V., Abhilash, M., Varghese, M. V., Alex, M., & Nair, R. H.). (2012). Protective effects of alpha-tocopherol against oxidative stress related to nephrotoxicity by monosodium glutamate in rats. *Toxicology Mechanisms and Methods*, 22(8), 625-630.
- Petracci, M.; Bianchi, M.; Mudalal, S.; Cavani, C. . (2013). . Functional ingredients for poultry meat products. *Trends in Food Science and Technology*, 33,, 27–39.
- Pil Nam Seong<sup>1</sup>, Soo Hyun Cho<sup>1</sup>, Kuyng Mi Park<sup>1</sup>, Geun Ho Kang<sup>1</sup>, Beom Young Park<sup>1</sup>, Sung Sil Moon<sup>2</sup>, and Hoa Van Ba<sup>1</sup>, \*. (2015). Characterization of Chicken By-products by Mean of Proximate and Nutritional Compositions. *Korean Journal of Food Science*, 35(2), 179–188.
- Pillai SL, Nair . (2013). Proximate composition, mineral elements and antinutritional factors in *Cleome viscosa* L. and *Cleome burmanni* W. & A. (Cleomaceae). . *International Journal of Pharmacy and Pharmaceutical Sciences* , 384-387.
- Pop, F. (2018). *Studia UBB Chem*, 63(2), 43-52.
- Populin T, Moret S, Truant S, Conte L. (2007). A survey on the presence of free glutamic acid in foodstuffs, with and without added monosodium glutamate. *Food Chemistry* , 104, .1712-1717.
- Prince., Singh, G., Verma, P. K., Rahmani, U. and Nandi,. (2023). Shiitake Mushrooms: A Medicinal Mushroom for Human Health. *Vigyan Varta*, 4(8), 214-217.
- Qiu W, Chen S, Xie J, Qu Y, Song X. (2016). Analysis of 10 nucleotides and related compounds in *Litopenaeus vannamei* during chilled storage by HPLC-DAD. *LWT Food Sci Tech*, 67, :187-93.
- Rahman. (2007). Aging. *Clinical. Intervention*, 2(2), 219-236.
- Ray S., Partha S. S., Rooh U. A., Anil K. N., Saroj K. S., Amiya K. R., Akash M.:. (2013). Prolapse in laying hens: Its pathophysiology and management: A review. *Indian Journal of Animal Production and Management*, 29, 17–24.
- Rohman, A., Riyanto, S., Mistriyani, Shuhaira and Nugroho, A.E. (2017). I: Study from Two Cultivars Antiradical Activities of Rambutan Peel. . *Research Journal of Phytochemistry*, 11(1), 42–47. [https:// doi.org/10.3923/rjphyto.2017.42.47](https://doi.org/10.3923/rjphyto.2017.42.47).
- Roper, S. & Chaudhari, N. . (2017). Taste buds: Cells, signals and synapses. . *Nat. Rev. Neurosci.* , 18, 485–497.
- Sadiku, O. (2010). Processing Methods Influence the Quality of Fermented African Locust Bean (*Iru/ogiri/dawadawa*) *Parkia biglobosa*. *Journal of Applied Sciences Research*, 6, 1656–1661.
- Sahni, S., Shere, D.M.:I., . (2017). Comparative evaluation of physico-chemical and functional properties of apple, carrot and beetroot pomace powders. *Int. J. Food Ferment Techno*, 7(2), 317–323.
- Salim A. S., Simeons A. J., Waruhiu A., Orwa C., Anyanjo C. . (2002). *Agroforest tree database: A tree species reference and selection guide*. Online at [http:// www.icraf.org](http://www.icraf.org) .
- Sanodiya B., Thakur G., Baghel R., Prasad G. and Bisen P. (2009). *Ganoderma lucidum*: A Potent Pharmacological Macrofungus. *Current Pharmaceutical Biotechnology* , 10(8), 717-742.
- Sawadogo-Lingani, H., Diawara, B., Ganou, L., Gouyahali, S., Halm, M., Amoa-Awua, W., and Jakobsen, M. (2003). Effet du d'écorticage mécanique sur la fermentation des graines de n'ér e *Parkia biglobosa*, en soumbala. e *Parkia biglobosa*, en soumbala. *Annales des Sciences Agronomiques du Benin*, 5, 67-84.
- SCHMIDT CHARLOTTE VINTHER, OLE G. MOURITSEN . (2022). Umami taste as a driver for sustainable eating. *international journal of food design*, 7(2).
- Schmidt, Charlotte Vinther Karsten Olsen & Ole G. Mouritsen. (2020). *Umami synergy as the scientific principle behind taste-pairing champagne and oysters* . [www.nature.com/scientificreports](http://www.nature.com/scientificreports).

- Seah, R. Siriporogutikorn, Sand Usawakesumem W. (2010). Anti-oxidant and antibacterial properties of Keung-kleoung paste and its ingredients. *Assian Journal of Food and Agricultural Industries*, 3, 210-213.
- Senga, P., Opota, D., Tamba, A., Tona, G., Kambu, O., Covaci, A., Apers, S., Pieters, L., & Cimanga, K. (2013). Chemical composition and nutritive value study of the seed oil of *Adenanthera pavonina* L. (Fabaceae) growing in Democratic Republic of Congo. *International Journal of PharmTechResearch*, 5(1), 205-216.
- Shantanu, T. (2019). The Shiitake mushroom- A history in magic and folklore. *The Mycophile*, 59(1), 1-4.
- Shittu, T.A., Sanni, L.O., Awonorin, S.O., Maziya-Dixon, B. and Dixon, A. (2005). Use of multivariate techniques in studying flour making characteristics of some cassava mosaic disease resistant cassava clones. *African Crop Science Conference Proceeding* 7, (pp. 621-630.).
- Siddiq M, Nasir M, Ravi R, Dolan KD, Butt MS . (2009). Effect of defatted maize germ addition on the functional and textural properties of wheat flour. *Int. J. Food Prop*, 12, :860-870.
- Silpa Sasi, Sathu T, Sunanda C and Pavan M . (2020). Proximate composition and sensory properties of poultry meat cocktail nuggets . *The Pharma Innovation Journal*, 9(7), 198-201.
- Sina S. and Traoré S.A. (2002). *Parkia biglobosa* (Jacq.) R.Br. ex G.Don. Record from Protabase. . . *ROTA Ressources végétales de l'Afrique tropicale*) *Plant Resources of Tropical Africa*, Wageningen, the Netherland.
- Sobowale SS, Olatidoye OP, Olorode OO, Akinlotan JV. (2011). . Nutritional potentials and chemical values of some tropical leafy vegetables consumed in South West Nigeria. *Journal of Sciences and Multidisciplinary Research*, 3, 55-65.
- Sosa M, G. (2006). Sensory acceptability of menus and sweet snacks among children and adults from low- and medium-income households in Argentina. *Food Quality and Preference*, 17(7-8), 590-597.
- Steve and Julie Rockcastle . (2013). *Best Management Practices for Log-Based Shiitake Cultivation in the northeastern united states*. Northeast SARE Research and Education Cornell University .
- Sunil Kochhar, Barbara Mouratou, and Philipp Christen. (n.d.). Amino Acid Analysis by High-Performance Liquid Chromatography after Derivatization with 1-Fluoro-2,4-dinitrophenyl-5-L-alanine Amide (Marfey's Reagent). In N. P. C. Cooper, *Methods in molecular biology :Amino Acid Analysis Protocols* (Vol. 159, pp. 49-54). New Jersey : Humana Press .
- Tabata, Kazuoko, Hiisako. (2006). Comparison of Chemical Compositions of Shiitake (*Lentinus edodes* (Berk.) Sing) Cultivated. *Food Science and Technology Research*, 12(4), 252-255.
- Tester, R. F., Karkalas, J., Qi, Xi. (2004). Starch-composition, fine structure and architecture. *Journal of Cereal Sciences*, 39, 151– 165.
- The United States Pharmacopeia. (2002.). . USP25/NF20 .
- Thiombiano, D. N., Lamien, N., Dibong, D. S., Boussim, I. J., & Belem, B. . (2012). The role of woody species in managing food shortage in Burkina Faso. *Sécheresse*, 23(2), 86-93.
- Tian HX, Zhang YJ, Qin L, Chen C, Liu Y, Yu HY:. (2018). Evaluating taste contribution of brown sugar in chicken seasoning using taste compounds, sensory evaluation, and electronic tongue. *Int J Food Properties* 2018, 21 515 , 1, 471-483.
- Torrico DD, Jirangrat W, Wang J, Chompreeda P, Sriwattana S, Prinyawiwatkul W:. (2018). Novel modelling approaches to characterize and quantify carryover effects on sensory acceptability . *Foods* 2018, 7 364, 11, 186-197.
- Umami Information Center. . (2012). Umami Rich Foods. Retrieved from <http://www.umamiinfo.com/> 3. Yamaguchi S, Ninomiya K. Umami and Food Palatability. *The Journal of Nutrition.*, 921S-926S.
- Umedum, N. L., Udeozo, I. P., Muoneme, O., Okoye, N., & Iloamaka, I. (2013). Proximate analysis and mineral content of three commonly used seasonings in Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, 5, 11-14.
- USDA. (2013.). *International egg and poultry*.

- Valverde, M.E., Hernández-Pérez, T., Paredes-López, O., . (2015). . Edible mushrooms: improving human health and promoting quality life. *International Journal of . Microbiology*, 15, 1–14.
- Veni N Krishna, D Karthika, [...], and YJ Pradeepa. (2010). Analysis of Monosodium L-Glutamate in Food Products by High-Performance Thin Layer Chromatography. *journal of young pharmacists*, 2(3), 297–300.
- Vinay TN, Park CS, Kim HY, Jung SJ. . (2013:). Toxicity and dose determination of quillaja saponin, aluminum hydroxide and squalene in olive flounder (*paralichthys olivaceus*). . *Vetrinary Immunology Immunopathology*, 158(1-2), 73-85.
- Vlaardingen, G. S.-3. (2009). *Patent No. EP 2 217 097 B1*. EUROPEAN PATENT SPECIFICATION.
- WHO. (2014). *Global Status Report On Non-Communicable Diseases*. Geneva: World Health Organization.
- Wintergerst ES, Maggini S, Hornig DH. (2007). .Contribution of selected vitamins and trace elements to immune function. *Ann NutrMetab.*, 200, :301-323.
- Wright LT, Nancarrow C, Brace I. (2000). Researching taste: layers of analysis. *British Food Journal*., 102(5-6), 429-440.
- Wu, C.-M., & Wang, Z.). (2000). Volatile compounds in fresh and processed shiitake mushrooms (*Lentinus edodes* sing.). *Food Science and Technology Research*, , 6(3), 166–170.
- Yahaya Mode, \*1Adamu, Shahidah A., 2Salau, I. A. and 2Sambo, S. (2018). Processing Effect of Nutritional and Anti- Nutritional Content of African Locust Bean Seeds (*Parkia biglobosa* Benth). *Greener Journal of Agricultural Sciences*., 8(12), 370-375,.
- Yamaguchi S, Ninomiya K.. . (2000). Umami and food palatability. *Journal of nutrition*, 130:, 921S-926S.
- Yamaguchi, Shizuko; Ninomiya, Kumiko. (1889). "What is umami?". *Food Reviews International*., 14(3), 123–3.
- Yasutada, I.T. (2001). A method for estimating food preference of the commensal Rat. *Medical entomology and Zoology*., 52(3), 231-239.
- Zhang J., Sun-Waterhouse D., Su G., Zhao M.. (2019). New insight into umami receptor, umami/umami-enhancing peptides and their derivatives: A review. *Trends in Food Science & Technology*, 88, 429–438.
- Zhang, F. B. et al. . (2008). Molecular mechanism for the umami taste synergism. *Proc. Natl. Acad. Sci. USA* 105,, 20930–20934.
- Zhang, N., Chen, H., Zhang, Y., Ma, L., & Xu, X. (2013). Comparative studies on chemical parameters and antioxidant properties of stipes and caps of shiitake mushroom as affected by different drying methods... *Journal of the Science of Food and Agriculture*, 93(12), 3107–3113.
- Zhao, H., Wang, L., Brennan, M., Brennan, C. (2022). . How does the addition of mushrooms and their dietary fibre affect starchy foods. *J. Future Foods*, 2, 18–24.