

**ISOLATION AND SCREENING OF PLANT GROWTH PROMOTING
RHIZOBACTERIA (PGPR) ISOLATS DURING DIFFERENT SEASONS FROM
SOUTH GUJARAT**

UNDER PEER REVIEW

Abstract

Plant Growth Promoting Rhizobacteria (PGPR) a major constituent of Rhizobacteria, encourage the plant growth through their diverse mechanisms. In this investigation, 33 different Isolates isolated from the rhizosphere soils of various research stations in south Gujarat were screened for their plant growth promoting activity. All the 33 tested isolates in pre monsoon and post monsoon were screened and recorded different PGPR activities. Thirteen isolates of pre-monsoon season and twenty isolates of post monsoon season selected for further study. Based on colonial characteristics and morphology of isolates, 33 elected for Siderophore, HCN, IAA production phosphate solubilizing and antagonistic activities. Among 33 isolates, 10 isolates showed phosphate solubilizing properties, 10 isolates had Siderophore production, 6 isolates had IAA activity and no one organism produce HCN. All the 33 isolates have found with nitrogen fixing potential. Among these, those isolated before the monsoon season have more number of Gram positive bacteria while those isolated after monsoon were Gram positive as well as Gram negative.

Comment [o1]: Add Space

Keywords: Bacillus, Siderophore, Nitrogen fixation, IAA

Comment [o2]: Where is bacillus in abstract

INTRODUCTION

Microorganisms play vital role in agriculture in order to promote the transfer of plant nutrients and reduce application of chemical fertilizers. Among different microbes, Plant Growth-Promoting Rhizobacteria (PGPR) have potential to exert a positive effect on plant growth[3]. Beneficial plant microbe interactions in the rhizosphere can influence plant growth and soil fertility. These beneficial effects of PGPR have direct or indirect performance on plants. Direct promotion of growth by PGPR including production of metabolites that enhances plant growth such as auxins [1], cytokinins, gibberellins and through the solubilization of phosphate minerals [5]. Indirect growth promotion occurs via the removal of pathogens by the production of secondary metabolites such as hydrogen cyanide and siderophores [6]. Plants are regularly concerned in interactions with a broad range of bacteria that colonize the rhizosphere (rhizobacteria), the phyllosphere (epiphytes), and the inside of plant tissues (endophytes) [4]. The application of PGPR as crop inoculants for biofertilization, phytostimulation, and biocontrol would be an attractive alternative to decrease the use of chemical fertilizers which also effect

environmental pollution [1]. *Bacillus* sp. is widespread bacteria in agricultural soils and has many traits that make them well-matched as PGPR. Their plant growth promoting activities include production of HCN, siderophores, protease, antimicrobials and phosphate solubilizing enzymes [7]. In the present study, we investigated the production of HCN, siderophores, nitrogen fixation, antimicrobials and phosphate solubilization by 17 isolates of PGPR isolated from soils of different localities in Gujarat. The objectives of study were, (1) Isolation of different free living nitrogen fixing bacteria with respect to different seasons (2) Biochemical Characterization of isolates and (3) Evaluation of PGP and antimicrobial activities.

Comment [o3]: Need space between Bacillus and sp

MATERIALS AND METHODS

Site description of soil sampling

Soil samples from different sites were collected in sterile bag. Sampling was carried out before monsoons and post monsoons seasons, Soil samples were collected from different depths (0-15 cm) of soil.

Enumeration of total bacterial population and free living nitrogen fixing bacteria

All collected soil samples were brought to laboratory and processed immediately for isolating PGPR organisms using Ashby's Mannitol agar medium.

Comment [o4]: Start with next paragraph

Total viable count was carried out by serial dilution of 1 gm soil and resulting two dilution were spread over the Nutrient agar plate. For the counting of free living nitrogen fixing bacteria serially dilute sample was spread over Ashby's mannitol agar plate. Their results are depicted in Table 1 and 2 respectively.

Comment [o5]: Correct spelling of Ashby's

Comment [o6]: Space between Ashby's and mannitol

Comment [o7]: Correct spelling of Ashby's

Comment [o8]: Space between Ashby's and mannitol

PGPR activities of different isolates

HCN activity was checked by method given by Bakker and Schipper (1987) (2). Phosphate solubilizing activity was checked on Pikovaskya's agar medium having insoluble phosphate. Siderophore production was checked on Chrome Azurol-S (CAS) agar plates. IAA production was checked by Salkowsky method. Antagonistic activity of isolates were checked against *Fusarium* and *Sclerotium* and they were tested by dual culture technique. Bacterial isolates were streaked at one side of Petri dish (one cm away from the edge) containing PDA culture medium. 9 mm mycelial disc from seven days old PDA culture of *Fusarium* and *Sclerotium* were placed at the opposite side of Petri dishes perpendicular to the bacterial streak

Comment [o9]: Add space before siderophore

Comment [o10]: Space before IAA

Comment [o11]: Add space before and

Comment [o12]: Space before were

respectively and incubated at $27\pm 2^{\circ}\text{C}$ for 5-7 days. Petri dishes inoculated with fungal discs alone served as control.

Nitrogen estimation was carried out by Kjeldahl method in which nitrogen free medium was inoculated with active culture isolates (uninoculated flask also kept as blank) and incubated on rotary shaker at room temperature for 48 hrs. After incubation, broth is analyzed for estimating total nitrogen.

RESULTS AND DISCUSSION

Results showed that total number of viable count was high during the monsoon season compared to summer season (Table-1). Number of free living nitrogen fixing bacteria were also higher during post monsoon season (Table-2). Isolation of free living nitrogen fixing bacteria was carried out on Ashbay's Mannitol agar medium before and after monsoon seasons. Thirteen isolates of pre-monsoon season and twenty isolates of post monsoon season were selected for further study. Based on colonial characteristics and morphology of isolates, 33 selected for Siderophore, HCN, IAA production, phosphate solubilizing and antagonistic activities (Table-3). Among 33 Screened, 10 isolates showed phosphate solubilizing properties, 10 showed Siderophore production, 6 Isolates had IAA activity and no one organism produce HCN (Figure 1, 2 and 3). It was found that the isolates of A19, A24 and A28 Isolates have multiple characteristics to produce Siderophore, phosphate solubilization and nitrogen fixing ability. A12 isolate has nitrogen fixing, Phosphate solubilising and IAA production potential while A21 isolate has Nitrogen fixing, siderophore and IAA production potential. Zone of solubilization of isolates were measured and result showed that isolates of A12 and A16 have highest Zone of utilization/colony ratio (Table-4). IAA quantitative analysis indicates that isolates A15 and A26 produce highest amount of IAA is 38.6ppm and 39.0ppm respectively (Table-5). Out of 33 nitrogen isolates, 13 organisms isolated before monsoon season and 20 organisms isolated after monsoon seasons. Among these, those isolates isolated before the monsoon season have more number of Gram positive bacteria while those isolated after monsoon season Gram positive as well as Gram negative. No antagonistic effect was by isolates against test organisms. Findings of biochemical characterization of selected isolates is shown in Table-6. Result of nitrogen estimation indicates that isolates of A15, A20 and A30 fix nitrogen in higher amount as compared to other isolates. However, all the isolates have potential to fix atmospheric nitrogen (Table-7).

Comment [o13]: Check spelling and add space before Mannitol

Comment [o14]: Space after for

Comment [o15]: Space before Among

Comment [o16]: Space before 2

Table-1: Total Microbial count in different soil samples collected during various seasons

Sr. No.	Location of soil sample	cfu/gm (before Monsoon season)	cfu/gm (after Monsoon season)
1.	Regional Cotton Research Station, Maktampur, Bharuch	1.5x 10 ⁸	1.85x 10 ⁸
2.	NARP, Bharuch	1.6x 10 ⁸	2.1x 10 ⁸
3.	Agricultural Research Centre, Tanchha	1.2x 10 ⁸	4.9x 10 ⁸
4.	Agricultural Research Station, Achhalia	1.2x 10 ⁸	3.6x 10 ⁸

Table-2: Total viable count of free living nitrogen fixing In soil samples collected during different seasons

Comment [o17]: I small

Sr. No.	Location of soil sample	cfu/gm (before monsoon)	cfu/gm (after monsoon)
1.	Regional Cotton Research Station, Maktampur, Bharuch	2.6x 10 ⁴	5.8x 10 ⁵
2.	NARP, Bharuch	1.2x 10 ⁴	2.1x 10 ⁵
3.	Agricultural Research Centre, Tanchha	0.8x 10 ⁴	2.9x 10 ⁵
4.	Agricultural Research Station, Achhalia	1.9x 10 ³	1.6x 10 ⁴

Table-3: Different properties of free living nitrogen fixing bacteria

Properties	Nitrogen Fixation	Phosphate solubilisation	HCN production	Siderophore production	IAA production
Isolates Isolated monsoon season					
A1	Positive	-	-	-	-
A2	Positive	-	-	-	Positive
A3	Positive	Positive	-	-	-
A4	Positive	-	-	Positive	-
A5	Positive	-	-	-	-
A6	Positive	Positive	-	-	-
A7	Positive	Positive	-	-	-
A8	Positive	-	-	Positive	-

A9	Positive	-	-	-	-
A10	Positive	-	-	Positive	-
A11	Positive	-	-	Positive	-
A12	Positive	Positive	-	-	Positive
A13	Positive	-	-	-	-
Isolates isolated monsoon season					
A14	Positive	-	-	Positive	-
A15	Positive	-	-	-	Positive
A16	Positive	Positive	-	-	-
A17	Positive	-	-	-	Positive
A18	Positive	-	-	-	-
A19	Positive	Positive	-	Positive	-
A20	Positive	-	-	-	-
A21	Positive	-	-	Positive	Positive
A22	Positive	Positive	-	-	-
A23	Positive	-	-	-	-
A24	Positive	Positive	-	Positive	-
A25	Positive	-	-	-	-
A26	Positive	-	-	-	Positive
A27	Positive	-	-	-	-
A28	Positive	Positive	-	Positive	-
A29	Positive	-	-	-	-
A30	Positive	-	-	-	-
A31	Positive	-	-	Positive	-
A32	Positive	-	-	-	-
A33	Positive	Positive	-	-	-

Figure: 1.Siderophore productionby different isolates

Comment [o18]: Space before by




Figure: 2. Phosphate solubilization by different isolates




Figure: 3. IAA Production by different isolates




Table-4: Phosphate solubilizing activity on pikovasky medium (Zone of solubilization)

Isolates	Zone ratio	Isolates	Zone ratio
A3	4.00	A19	1.33
A6	1.25	A22	2.00
A7	2.50	A24	1.25
A12	1.20	A28	1.10
A16	1.25	A33	1.06

Table-5 : IAA production by different isolates

Isolates	IAA (ppm)
A2	23.5
A12	12.8
A15	38.6
A17	10.0
A21	18.5
A26	39.0

Table-6. Biochemical Characterization of different isolates

Biochemical tests were carried out for selected isolates with different colonial characteristics and PGP activities.

Sr. No.	Test	A12	A19	A21	A24	A28
1	Lactose	Positive	-	-	-	-
2	Xylose	Positive	-	-	-	Positive
3	Maltose	Positive	-	-	-	Positive
4	Fructose	Positive	Positive	Positive	Positive	Positive
5	Dextrose	Positive	Positive	Positive	Positive	Positive
6	Galactose	Positive	-	-	-	-
7	Raffinose	Positive	-	-	-	Positive
8	Trehalose	Positive	Positive	Positive	-	Positive
9	Melibiose	Positive	-	-	-	Positive
10	Sucrose	Positive	Positive	Positive	Positive	Positive
11	L- arabinose	Positive	Positive	Positive	Positive	Positive
12	Mannose	Positive	-	Positive	-	-
13	Inulin	Positive	-	Positive	Positive	Positive
14	Na- Gluconate	-	-	Positive	-	Positive
15	Glycerol	Positive	Positive	Positive	Positive	Positive
16	Salicin	Positive	Positive	Positive	Positive	-
17	Dulcitol	-	-	-	Positive	-
18	Inositol	-	-	Positive	-	-
19	Sorbitol	-	-	Positive	Positive	-

20	Mannitol	Positive	-	Positive	Positive	Positive
21	Adonitol	Positive	-	-	-	-
22	Arabitol	Positive	-	-	-	-
23	Erythritol	Positive	-	-	-	--
24	α - methyl $-D$ -glucoside	-	-	-	-	-
25	Rhamnose	-	-	-	Positive	-
26	Cellobiose	Positive	-	-	Positive	-
27	Melezitose	-	-	-	Positive	Positive
28	α - methyl $-D$ -Manoside	-	-	-	Positive	-
29	Xylitol	-	-	-	Positive	-
30	ONPG	-	-	-	-	Positive
31	Esculin hydrolysis	Positive	Positive	Positive	Positive	Positive
32	D- arabinose	Positive	-	Positive	Positive	-
33	Citrate utilization	-	-	-	-	-
34	Malonate utilization	-	-	-	-	Positive
35	Sorbose	-	-	-	Positive	Positive
36	ONPG	-	Positive	-	-	Positive
37	Lysine Utilization	-	-	-	-	-
38	Ornithine Utilization	-	-	Positive	-	-
39	Urease	Positive	Positive	Positive	Positive	Positive
40	Phenylalanine deamination	-	-	-	-	-

41	Nitrate reduction	Positive	-	Positive	Positive	Positive
42	H ₂ S production	-	-	-	-	-
43	Citrate Utilization	-	-	-	-	-
44	Voges-Proskauer	Positive	-	Positive	Positive	Positive
45	Methyl red	-	-	-	-	-
46	Indole	-	-	Positive	-	Positive

Table:7 Nitrogen estimation of isolates by kjeldahl

Isolates	% N	Isolates	% N
A1	0.00168	A18	0.00128
A2	0.00114	A19	0.00147
A3	0.00228	A20	0.00321
A4	0.00270	A21	0.00178
A5	0.00290	A22	0.00254
A6	0.00200	A23	0.00198
A7	0.00228	A24	0.00226
A8	0.00319	A25	0.00178
A9	0.00159	A26	0.00245
A10	0.00228	A27	0.00178
A11	0.00250	A28	0.00125
A12	0.00410	A29	0.00245
A13	0.00228	A30	0.00345
A14	0.00200	A31	0.00149
A15	0.00364	A32	0.00267
A16	0.00224	A33	0.00267
A17	0.00219		

CONCLUSION:

Microbial population along with free living nitrogen microbes is increased in post monsoon season. Beforemonsoon season, number of Gram positive bacterial population is higher than the Gram negative. Isolates A19(*Streptomyces coelicolor*), A24(*Enterobactersp.*), and A28(*Bacillus altitudinis*) have potential characteristics to produce Siderophore as well as nitrogen fixing and phosphate solubilizing properties. The isolate A12 has characteristic to produce IAA along with nitrogen fixation and phosphate solubilization Properties.

Comment [o19]: Space between before and monsoon

Comment [o20]: Space before have

Comment [o21]: Add references related to isolates A19, A24, and A28.

REFERENCES

1. Ali B, Sabri AN, Hasnain S (2010). Rhizobacterial potential to alter auxin content and growth of *Vigna radiata* (L.). *World J Microbiol Biotechnol* 26: 1379-1384.

Comment [o22]: Add and before last author

Comment [o23]: Add full name of journal

2. Bakker, A.W. and Schipper, B. (1987). Microbial cyanide production in the rhizosphere in relation to potato yield reduction and *Pseudomonas* spp. mediated plant growth stimulation. *Soil Biol. Biochem.*, 19: 451-457.

Comment [o24]: Add full name of journal

Comment [o25]: Add and before pandey

3. Dastager SG, Deepa CK, Pandey A (2011). Potential plant growth-promoting activity of *Serratia nematodiphia* NII-0928 on black pepper (*Piper nigrum* L.). *World J Microbiol Biotechnol* 27: 259-265.

Comment [o26]: Space between serratia and nematodiphia

Comment [o27]: Add full name of journal

4. Deepa CK, Dastager SG, Pandey A (2010). Isolation and characterization of plant growth promoting bacteria from non-rhizospheric soil and their effect on cowpea (*Vigna unguiculata* (L.) Walp.) seedling growth. *World J Microbiol Biotechnol* 26: 1233-1240.

Comment [o28]: Add and before pandey

Comment [o29]: Space between vigna and unguiculata

Comment [o30]: Add full name of journal

5. El-Hadad ME, Mustafa MI, Selim SM, Mahgoob AEA, El-Tayeb TS, et al. (2010). In vitro evaluation of some bacterial isolates as biofertilizers and Biocontrol agents against the second stage juveniles of *Meloidogyne incognita*. *World J Microbiol Biotechnol* 26: 2249-2256.

Comment [o31]: Add all author name

Comment [o32]: Add space after meloidogyne

Comment [o33]: Add full name of journal

6. Idris HA, Labuschagne N, Korsten L (2010). Suppression of *Pythium ultimum* root rot of sorghum by rhizobacterial isolates from Ethiopia and South Africa. *Biological Control* 45: 72-84.

Comment [o34]: Add and before korsten

Comment [o35]: Space after pythium

7. Suresh A, Pallavi P, Srinivas P, Kumar VP, Chandra SJ, et al. (2010). Plant growth promoting activities of fluorescent pseudomonads associated with some crop plants. *Afr J MicrobiolRes* 4: 1491-1494.

Comment [o36]: Add all author name

Comment [o37]: Add full name of journal

UNDER PEER REVIEW