

Estimation of Ursolic acid in *Bauhinia racemosa* Lam. by HPTLC method

Abstract

Plants have been utilized since ancient times to cure and manage a variety of diseases. Their capacity to heal or prevent ailments is emanates from the presence of various phytochemicals such as alkaloids, flavonoids, glycosides, terpenoids, steroids, tannins, phenols, etc. Pentacyclic triterpenoids have attracted the interest of researchers due to their diverse pharmacological activities. One of the Pentacyclic triterpenoids is Ursolic acid (UA) with significant biological effects, including anti-inflammatory, anticancer, antidiabetic, antioxidant, and antibacterial properties. *Bauhinia racemosa* Lam. is used in folk medicines for the treatment of various disorders. In the present work, a subtle HPTLC method has been developed for determination of Ursolic acid in *Bauhinia racemosa* Lam. The demonstrated method is simple, rapid, accurate and can be used for routine qualitative testing.

Keywords: Secondary metabolite, Pentacyclic triterpenoids, Ursolic acid, *Bauhinia racemosa* Lam, HPTLC

1. Introduction

Medicinal plants and their secondary metabolites are being utilized more often as supplemental medicine to cure diseases. A range of potent phytoconstituents may be found in the therapeutic plants [1]. The importance of plant secondary metabolites in industry, agriculture, and medicine has shed light on several studies concerning their biological activity, synthesis, and biosynthesis. It has been estimated that these active natural commodities are the source of at least 40% of medications [2]. Terpenes and their derivatives are an important class of natural compounds. Recently, pentacyclic triterpenoids have gained popularity as a class of terpenes. Ursolic acid (UA) (Figure 1) belongs to the C₃₀ class of isoprenoid compounds. It is a pentacyclic triterpenoid with a molecular weight of 456.68 g/mol and a melting point of 283–285 °C [3].

Commented [HM1]: Assertion needs to be cited




Figure 1. Structure of Ursolic acid

Different pharmacological activities of ursolic acid are established, including anti-inflammatory [4], hepatoprotective [5], hair growth stimulant, and anti-tumor activity [6]. Pharmacological studies have shown that UA can interfere with a variety of signaling pathways involved in the survival and development of cancer and treat inflammatory or oxidative stress-related disorders, including cancer, cardiovascular disease and neurodegeneration [7].

Bauhinia racemosa Lam belongs to the Fabaceae family. It occurs in India, Sri Lanka, and China. Many applications of plant parts in folk medicine have been described, including flowers as a diuretic. The treatment for dysentery includes the use of flowers, buds, and dried leaves. The root bark is utilized in inflammation of the liver [8,9]. The seeds are tonic and aphrodisiac. Leaves are anti-diabetic in nature [10]. General phyto-pharmacological screening of the plant has showed that the ethanol extract of *B. racemosa* leaves has analgesic, antipyretic, and antiplasmodic properties [11] as well as antimicrobial activity and antihistaminic effect [10,12]. The plant's fresh flower buds demonstrated antiulcer action [13], as well as hypotensive and hypothermic activity [14,15].

Chromatographic technology is essential for both qualitative and quantitative research. HPTLC is an advanced method to estimate biomarker levels. Synthetic medications increase the chance of developing illnesses such as cancer, diabetes, and neurological disorders, endangering global health. Natural herb-based treatments are urgently needed to address this issue. Indigenous remedies can mitigate the negative effects of manufactured pharmaceuticals, providing a viable solution to the global health issue.

Commented [HM2]: cite

Commented [HM3]: cite

Commented [HM4]: cite

The aim of the study is to develop a simple and rapid method for ursolic acid estimation since it has many pharmacological properties and wide application in therapeutics.

Commented [HM5]: independent assertions in this paragraph need to be cited adequately

2. Material and Methods

2.1 Collection and Authentication of plant material

B.racemosa Lam. healthy leaves were obtained from Rani Baug in Mumbai, India. The species was identified and verified at St. Xavier's College Blatter Herbarium (Accession no. PD-431) Mumbai. Leaves were dried in the shade, then chopped into small pieces and roughly powdered.

2.2 Preparation of extract

The leaf powder was extracted using Soxhlet apparatus using hydroalcoholic solvent (60:40). The extraction method comprised 12 siphon cycles lasting 8 hours, producing a dark green solvent that separated phytoconstituents from the plant. The extract was collected, and the solvent was removed using simple evaporation at ambient temperature. The concentrated extract was used for further analysis.

2.3 Preparation of standard stock solution

To make stock solutions of Ursolic acid, dissolve 1 mg in 1 mL of methanol (final concentration $\mu\text{g/mL}$). A standard graph was prepared using a series of dilutions ranging from 100 to 500 $\mu\text{g/ml}$.

2.4 Estimation of Ursolic acid [16]

A 200 μl methanolic solution was placed into a test tube and evaporated in a boiling water bath. The tube was filled with 0.3 ml of 5% vanillin glacial acetic acid and 1 ml of perchloric acid solution. The sample solutions were heat at 60°C for 45 minutes and cooled under tap water. To the cooled solution, 5 ml of glacial acetic acid was added. The reaction mixture was stirred evenly using a vortex. The OD was measured at 544 nm on UV Vis Spectrophotometer.

3. HPTLC method development

3.1 Preparation of stock

Stock solution - Suspend 10 mg of Ursolic Acid in 10 ml of methanol in a volumetric flask. Shake and sonicate the solution for 5 to 10 minutes. Further dilute the solution to 10 ml of methanol to prepare working solution.

3.2 Sample preparation

Accurately weighed 0.2 gm of sample in a 5 ml volumetric flask. Add methanol, shake and sonicate the solution for 5 to 10 minutes. Dilute the solution to 5 ml with methanol.

3.3 Optimization of chromatographic conditions

Merck cat. No. 5548 HPTLC silica gel 60 F254 was used as a stationary phase. 10 μ l of sample and 1 μ l of standard were loaded with a band width of 8mm.

3.4 Chromatography Determination and Derivatization

The pre-derivatized plates were developed with a mixture of Toluene: Ethyl acetate : Formic Acid (5: 1.5: 0.5) (v/v/v) as a mobile phase. The plate was sprayed with freshly prepared Vanillin Sulphuric Acid Reagent. Heat the plate at 110°C for 5-10 minutes and detection was done at 540 nm wavelength.




Figure 1a. Typical Chromatogram of standard ursolic acid




Figure 1b. Chromatogram of ursolic acid in *B. racemosa*




Figure 2. 3D spectra of Ursolic acid + *B.racemosa* Figure 3. Spectral comparison of Standard Ursolic acid and *B.racemosa*




Figure 4. Graphical representation of the calibration curve of standard Ursolic acid

Commented [HM6]: I believe that these figures will be more useful if moved to the results section and positioned appropriate to illustrate the results discussed

4. Results and Discussion

In the current study, the quantification of the contents of Ursolic acid and the HPTLC method was developed for *B. racemosa*. The chromatograms (Figure 1a,1b,2,3) were obtained upon scanning at 540 nm. The content of ursolic acid in the present plant sample was found to be 0.09 ng/µg. The calibration plots (Figure 4) were linear in the range of 100-500 (µg/ml), and the correlation coefficient (r) was 0.9729 which indicates good linear dependence of peak area on concentration.

Chemical fingerprinting has been shown to be a thorough and realistic way of assessing the quality and identifying the species of traditional medications. It employs chromatographic methods to create distinct patterns of identification for herbal drugs,

which may then be utilized to assess not only the presence or absence of certain markers, but also the proportion of all detected analytes [17]. HPTLC uses much less solvents, making it more cost-effective and environmentally friendly. Furthermore, it is a rapid method of analysis that allows for the simultaneous processing of a high number of data while minimizing memory effects. Moreover, it enables the identification of all molecules, including those that are substantially maintained at baseline [18].

Commented [HM7]: This text is not necessary under results and discussion section. You can move it to the background/ introduction

A novel HPTLC approach has been developed for detecting and quantifying ursolic acid in apple peel extract [19]. UA has been discovered in the cuticular wax of numerous edible fruits, including cherry, loquat, pear, peach, quince, apple, and bilberry. The cuticular wax of five Rosaceae fruits contained high levels of this triterpene [20]. Several HPTLC investigations have been published for the quantitative measurement of ursolic acid in numerous plant extracts, such as the aerial portions of *Nepeta deflersiana* [21] and *Wattakaka volubilis* [22]. An HPTLC densitometric approach was devised to quantify ursolic acid in *U. dioica* leaves [23]. In *Alstonia scholaris*, a quantitative method was developed and quantified for monitoring ursolic acid [24].

Moreover, UA is gaining popularity because of its therapeutic benefits, which include anti-inflammatory, antioxidant, anti-apoptotic, and anti-carcinogenic effects [25]. This study attempted to establish a method which was found to be simple, accurate, specific and robust for the analysis of ursolic acid in crude drug samples.

Commented [HM8]: This is material for the introduction

5. Conclusion

In summary, leaves of *B. racemosa* Lam. can be considered as a good source of Ursolic acid. Since the diverse biological activities of ursolic acid reported in folk medicine and recent studies (such as antibacterial, antifungal, antiviral, anti-inflammatory, antioxidant, anti-wrinkle, anticancer, anti-hepatotoxic, antipyretic, and antidiabetic), and the fair content of that compound in the leaves was observed. We could presuppose that this plant may be appreciated in domains of pharmacology, pharmacognosy, cosmetology, and oncology.

Abbreviations

UA: Ursolic acid

HPTLC: High-performance thin layer chromatography

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

References

1. Darbhe J, Neve V, Bhalchim V, Tonpe A, Khandave P, Mane S, Yewale R. Estimation of Ursolic Acid in Methanolic Extract of *Momordica dioica* by HPTLC Method. *Pharmacognosy Research.* 2023;15(4). <http://dx.doi.org/10.5530/pres.15.4.077>
2. Gershenzon J, Kreis W. Biochemistry of terpenoids: monoterpenes, sesquiterpenes, diterpenes, sterols, cardiac glycosides and steroid saponins. *Biochemistry of plant secondary metabolism.* 1999 Sep 21;2:222-99.
3. Shaikh n, sawant g, dixit n. A review on ursolic acid: A naturally obtained pentacyclic triterpene. *Int J Pharm Pharm Sci.* 2021 Jun. 1 [cited 2024 Sep. 13];13(6):1-5. [10.22159/ijpps.2021v13i6.41106](https://doi.org/10.22159/ijpps.2021v13i6.41106)
4. Hirota M, Mori T, Yoshida M, Iriye R. Suppression of tumor promoterinduced inflammation of mouse ear by ursolic acid and 4, 4-dimethylcholestane derivatives. *Agricultural and Biological Chemistry.* 1990;54(4):1073-5. <https://doi.org/10.1080/00021369.1990.10870043>
5. Shukla B, Visen PK, Patnaik GK, Tripathi SC, Srimal RC, Dayal R, Dobhal PC. Hepatoprotective activity in the rat of ursolic acid isolated from *Eucalyptus hybrid*. *Phytotherapy Research.* 1992 Mar;6(2):74-9. <https://doi.org/10.1002/ptr.2650060205>
6. Lee KH, Lin YM, Wu TS, Zhang DC, Yamagishi T, Hayashi T, Hall IH, Chang JJ, Wu RY, Yang TH. The cytotoxic principles of *Prunella vulgaris*, *Psychotria serpens*, and *Hyptis capitata*: ursolic acid and related derivatives I. *Planta Medica.* 1988 Aug;54(04):308-11. <https://doi.org/10.1055/s-2006-962441>
7. Kashyap D, Tuli HS, Sharma AK. Ursolic acid (UA): A metabolite with promising therapeutic potential. *Life sciences.* 2016 Feb 1;146:201-13. <https://doi.org/10.1016/j.lfs.2016.01.017>

8. Davis SN. Insulin, oral hypoglycemic agents and the pharmacology of the endocrine pancreas. Goodman and Gilman's the Pharmacological Basis of Therapeutics. Brunton, LL (Ed.). McGraw-Hill, New York. 2006:1613-45.
9. Kumar RS, Gupta M, Mazumdar UK, Rajeshwar Y, Kumar TS, Gomathi P, Roy R. Effects of methanol extracts of *Caesalpinia bonducella* and *Bauhinia racemosa* on hematology and hepatorenal function in mice. *The Journal of toxicological sciences*. 2005;30(4):265-74. <https://doi.org/10.2131/jts.30.265>
10. Nirmal SA, Dhasade VV, Laware RB, Rathi RA, Kuchekar BS. Antihistaminic effect of *Bauhinia racemosa* leaves. *Journal of Young Pharmacists*. 2011 Apr 1;3(2):129-31. <https://doi.org/10.4103/0975-1483.80301>
11. Gupta M, Mazumdar UK, Kumar RS, Gomathi P, Rajeshwar Y, Kakoti BB, Selven VT. Anti-inflammatory, analgesic and antipyretic effects of methanol extract from *Bauhinia racemosa* stem bark in animal models. *Journal of ethnopharmacology*. 2005 Apr 26;98(3):267-73. <https://doi.org/10.1016/j.jep.2005.01.018>
12. Ali MS, Azhar I, Amtul Z, Ahmad VU, Usmanghani K. Antimicrobial screening of some *Caesalpinaceae*. *Fitoterapia*. 1999 Jun 1;70(3):299-304. [https://doi.org/10.1016/S0367-326X\(99\)00015-5](https://doi.org/10.1016/S0367-326X(99)00015-5)
13. Akhtar AH, Ahmad KU. Anti-ulcerogenic evaluation of the methanolic extracts of some indigenous medicinal plants of Pakistan in aspirin-ulcerated rats. *Journal of Ethnopharmacology*. 1995 Apr 1;46(1):1-6. [https://doi.org/10.1016/0378-8741\(94\)01220-t](https://doi.org/10.1016/0378-8741(94)01220-t)
14. Dhawan BN, Dubey MP, Mehrotra BN, Tandon JS. Screening of Indian plants for biological activity: Part IX. *Indian Journal of Experimental Biology*. 1980;18:594-602.
15. Kumar RS, Sivakumar T, Sunderam RS, Gupta M, Mazumdar UK, Gomathi P, Rajeshwar Y, Saravanan S, Kumar MS, Muruges K, Kumar KA. Antioxidant and antimicrobial activities of *Bauhinia racemosa* L. stem bark. *Brazilian journal of medical and biological research*. 2005;38:1015-24. <https://doi.org/10.1590/s0100-879x2005000700004>
16. Gupta A, Maheta P, Chauhan R, Pandey S, Yadav JS, Shah S. Simultaneous quantification of bioactive triterpene acids (ursolic acid and oleanolic acid) in different extracts of *eucalyptus globulus* (L) by HPTLC method. *Pharmacognosy Journal*. 2018;10(1). <http://dx.doi.org/10.5530/pj.2018.1.30>

17. Saxena HO, Parihar S, Pawar G, Rao GR. Simultaneous densitometric determination of β -sitosterol and lupeol through validated HPTLC method in different plant parts of *Uraria picta* (Jacq.) Desv. ex DC.–A dashmool species. *Acta Chromatographica*. 2023 Jan 20;35(1):99-105. <https://doi.org/10.1556/1326.2022.01018>
18. Jamal M, Amir M, Ali Z, Mujeeb M. A comparative study for the extraction methods and solvent selection for isolation, quantitative estimation and validation of ursolic acid in the leaves of *Lantana camara* by HPTLC method. *Future Journal of Pharmaceutical Sciences*. 2018 Dec 1;4(2):229-33. <https://doi.org/10.1016/j.fjps.2018.07.002>
19. Nikam PH, Kareparamban JA, Jadhav AP, Kadam VJ. Development and validation of high-performance thin layer chromatographic method for ursolic acid in *Malus domestica* peel. *Indian Journal of Pharmaceutical Sciences*. 2013 Jul;75(4):486. <https://doi.org/10.4103/0250-474x.119815>
20. Ludeña-Huaman MA, Ramos-Inquiltupa DA. Determination of the content of ursolic and oleanolic acid in the cuticular wax of fruits of different species of Rosaceae. *Revista Colombiana de Química*. 2019 Aug;48(2):15-20. [10.15446/rev.colomb.quim.v48n2.77046](https://doi.org/10.15446/rev.colomb.quim.v48n2.77046)
21. Orfali R, Siddiqui NA, Alam P, Alhowiriny TA, Al-Taweel AM, Al-Yahya S, Majrashi NM, Mehmood R, Khan SI, Perveen S. Biological Evaluation of Different Extracts of Aerial Parts of *Nepeta deflersiana* and Standardization of Active Extracts Using 8-Epi-7-Deoxyloganic Acid and Ursolic Acid by Validated HPTLC Method. *Evidence-Based Complementary and Alternative Medicine*. 2018;2018(1):8790769.
22. Gopal V, Mandal V, Mandal SC. HPTLC evaluation of oleanolic acid and ursolic acid from the methanol extract of *Wattakaka volubilis*. *Journal of Acute Disease*. 2014 Jan 1;3(1):59-61. [https://doi.org/10.1016/S2221-6189\(14\)60013-5](https://doi.org/10.1016/S2221-6189(14)60013-5)
23. Shailajan S, Hande H, Singh D, Tiwari B. Estimation of ursolic acid from *Urtica dioica* L. using validated HPTLC method. *Journal of Applied Pharmaceutical Science*. 2014 May 27;4(5):092-5. <https://dx.doi.org/10.7324/JAPS.2014.40517>
24. Shetty P, Mangaonkar K, Sane RT. HPTLC determination of ursolic acid in *Alstonia scholaris* R. Br. *JPC–Journal of Planar Chromatography–Modern TLC*. 2007 Feb;20:65-8. <https://doi.org/10.1556/jpc.20.2007.1.11>

25. Seo DY, Lee SR, Heo JW, No MH, Rhee BD, Ko KS, Kwak HB, Han J. Ursolic acid in health and disease. *The Korean journal of physiology & pharmacology: official journal of the Korean Physiological Society and the Korean Society of Pharmacology*. 2018 May;22(3):235. <https://doi.org/10.4196/kjpp.2018.22.3.235>

UNDER PEER REVIEW