

~~Haploinsufficiency~~Haploinsufficiency in human diseases: A mini-review

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Abstract

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, (Asian) Chinese (Simplified, PRC)

Human disease studies have identified several hundred haploinsufficient genes affecting a range of phenotypes and cause of several diseases. ~~Haploinsufficiency~~Haploinsufficiency (HI) results due to single functional copy of a gene. In this mini-review we review the topic of HI with reference to causes, study models and detection methods. Further, we highlight the genetic, clinical, and pathological implications of the mechanism. Finally, the potential clinical applications are discussed. The study of HI therefore assumes significance since its spectrum extends to the realm of biophysics, gene expression, chromosome biology, quantitative traits, and evolutionary biology.

Key words: ~~Haploinsufficiency~~Haploinsufficiency, Dosage effects, heterozygous mutations, Loss of ~~functions~~function, Fitness consequences.

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold

1. Introduction

~~Haploinsufficiency~~Haploinsufficiency (HI) is defined as insufficient function to maintain a wild-type phenotype in the presence of one wild-type allele and one mutant allele (Adam F Johnson et al., 2019). (HI) can arise through mutations that lead to loss of function, or partial or complete loss of expression of one allele. Occasionally ~~haploinsufficiency~~haploinsufficiency can be additive or unlinked. Such genetic interactions are interesting to biologists for as they provide ~~evidencesevidence that~~ two gene products physically interact or cooperate in a network. In case of sex-linked linked to mutation of one gene copy may help identify pathways that are limiting for cellular function. Whereas, in the case of autosomal dominance the HI interfere with wild-type function from mutations that do not interfere with function of the remaining wild-type allele(Fuller et al., 2019). Heterozygosity may be deleterious in diploid organisms because it may disrupt a proper balance between various components within a complex or network. Many critical cellular processes must be exquisitely sensitive to the dosage of their effectors.

Historical studies of dosage preceding HI-The link between gene deletions and disease was not recognized until advancements in human karyotyping made reliable analysis possible (Lejeune et al., 1963). Most mutant alleles are recessive, meaning a single normal allele (haplosufficient, HS) maintains the typical phenotype. The dominance of wild-type alleles has been debated for nearly a century. Fisher (1928) suggested that modifier genes evolved to counteract mutant effects, while Wright (1934) argued that recessiveness arises from biochemical properties, as enzyme activity plateaus beyond a certain concentration. Later studies favored Wright's simpler model (Kondrashov & Koonin, 2004). Kacser & Burns (1981) suggested metabolic control analysis, showing that a 50% enzyme reduction often has minimal impact, especially in long metabolic pathways. Additional evidences using *S. cerevisiae* genome-wide knockout library (Deutschbauer et al., 2005) found that ~3% of yeast genes exhibit HI. Notably, essential and nonessential HI strains show similar fitness, and only 98 of 1102 essential genes are HI suggesting a complex link between gene essentiality and HI.

All branches of life, from bacteria to humans, carry HI which they accumulate throughout development and life term. As a result their genome may contain amplifications and deletions, inactivated genes, or gains and loss of whole chromosomes. At the allelic level they include spectrum of mutations involve gene inactivation-point mutations, single nucleotide insertions and deletions

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold

(<https://www.nature.com/scitable/>). The mutational event depends on the cellular process affected and location in the genome. For example, sub-chromosomal rearrangements are particularly likely to occur in repetitive areas of the genome, whereas whole chromosomal gains and losses are **only more** likely to occur during cell division. Finally, HI can occur due to errors in mitosis or meiosis can result from perturbation to the mitotic machinery, or the chromosome surveillance mechanisms. (Rendy Hosea et al., 2024). Whole genome duplication (WGD)-throughout eukaryotic evolution, genomes undergoes increase in ploidy (Anthony Levasseur and Pierre Pontarotti 2011 14). Model organisms such as *Saccharomyces cerevisiae*, *Caenorhabditis elegans* and *Drosophila melanogaster* have been instrumental in the cellular and biological studies of HI.

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

2. Genetic and cellular causes

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, (Asian) Chinese (Simplified, PRC)

A phenotype may manifest when the reduced function at two separate loci present in combination (termed unlinked non-complementation) exhibits the additive nature of HI. According to several researchers, mechanistic and evolutionary forces drive HI. An example is the imbalance due to *Arid1b* protein. HI produces abnormal gamma-aminobutyric acid (GABA) neurotransmission Moffat et al. (2019). The gene has been linked to autism spectrum disorder (ASD) and intellectual disability (ID) (Santen et al., 2012; Santen et al., 2014). Altered epigenetic modifications are associated with HI. The *ARID1B* gene encodes 1B (ARID1B), an AT-rich-interactive-domain-containing protein that is a subunit of the chromatin remodeling complex, BRG1/BRM-associated factor (BAF), which regulates expression via nucleosome remodeling (Singhal et al., 2010; Alver et al., 2017). An example of cellular manifestation is the survival motor neuron gene (*SMN2*). *SMN2* is a paralogue of the survival motor neuron 1 (*SMN1*) gene present in humans, but not other primates (Levasseur and Pontarotti 2020). *SMN1* mutations cause spinal muscular atrophy disease associated with progressive loss of motor neurons. Because of a noncoding point mutation, the *SMN2* mRNA splicing pattern typically lacks exon 7 and produces a truncated protein unable to complement the lack of SMN protein in patients.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Major theories aimed to explain the mechanisms of HI include the dosage balance hypothesis and the insufficient amounts hypothesis (Johnson et al., 2019; Morrill and Amon, 2019). The latter attributes HI to a stoichiometric imbalance of members of a protein complex, whereas the former claims that the reduced quantity of protein that is produced is insufficient for keeping its

normal function. Explanation by several researchers on the insufficient amounts hypothesis suggests that transcriptional regulators work close to a threshold level, thus making the system extremely dosage sensitive (Veitia, 2002). The hypothesis is based on transcription factor (TF) cooperativity and positive feedback. Figure 2 is a pictorial depiction of the mechanisms. Using a systems-biology perspective, Roman Zug's (2022) explain cell fate decisions as bi-stable switches, requiring both positive feedback and cooperativity. The author proposes that HI of transcriptional regulators, as well as associated (developmental disorders (DDs), result from disruption of bi-stability (i.e., disruption of positive feedback or cooperativity) in the gene regulatory network (GRN) underlying the cell fate decision in several cells involved in hematopoiesis (Palii et al., 2019), neurogenesis (Faure et al., 2020), gonad development (Stévant et al., 2019), and other cells.

Several TFs (in their activation domains) and cofactors possess substantial intrinsically disordered regions (IDRs), which serve as the source of cooperativity (Liu et al., 2006; Dyson and Wright, 2016). Few examples include the PAX6 and SOX2 during eye development (Kamachi et al., 2001 30), PAX3 and SOX10 during neural crest development (Bondurand et al., 2000), and FLI1 and GATA2 during hematopoiesis (Shi et al., 2014), and several others are crucial to cell fate determination (and whose HI results in DDs). The positive feedback loops between master TFs and their super enhancers (SEs) become particularly sensitive to changes in TF/cofactor concentration owing to the high level of cooperativity found at (SEs), through several binding sites of TF and transcriptional regulators (Hnisz et al. (2017). When cells shift towards one fate, the dose sensitivity becomes evident. Furthermore, decline in TF levels causes de-activation. Finally, ~~e oo operativity cooperativity~~ cooperativity may be mediated by a number of indirect processes, such as cofactors, or by direct TF–TF interactions (Spitz and Furlong, 2012; Reiter et al., 2017).

3. Models to study ~~haploinsufficiency~~ haploinsufficiency (cellular and animal)

3.1. Cellular advances in modeling human diseases using patient-cultured cells—fibroblasts, HeLa cells, and pluripotent stem cells (iPSCs)—have revolutionized biomedical research, paving the way for human disease research. These methods have been complemented with gene editing methods such as clustered regularly interspaced short palindromic repeats (CRISPR). Several

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

researchers have successfully demonstrated the use of these methods to study human mutations) in wide spectrum of neuropsychiatric diseases. The preceding paragraphs enlist a representative list.

a. human embryonic stem cells (hESCs). Arel-Gallily et al., 2022 using human embryonic stem cells (hESCs) and the CRISPR method, established a heterozygous genome-wide loss-of-function (HI) library. The study highlights over 600 essential (HI) genes enriched in several pathways, with the WNT and TGF- β pathways being the major affected. In another example, Niggle et al., 2023 report that in the heterogeneous nuclear ribonucleoprotein (HNRNPC) gene (HI) affects alternative splicing of intellectual disability using induced pluripotent stem cells (iPSCs) due defective mRNA processing. With several advances and modifications in the methods, several neuropsychiatric diseases are now amenable to iPSC and several researchers have used them to study cellular and molecular mechanisms of disease.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

b. Patient cultured cells (fibroblast) Schuster et al., 2022, report the generation of dermal fibroblasts from a patient with Dravet syndrome carrying a deletion on chromosome 2 encompassing SCN1A and flanking genes. Dravet syndrome is an early-onset, devastating epilepsy syndrome caused by heterozygous mutations in SCN1A. In another example, James et al., 2024 demonstrates altered mitochondrial function in fibroblast cell lines of carriers of SMA, an childhood-onset neuromuscular disease. Major findings include depolarized mitochondrial membrane potential, increased levels of reactive oxygen species, and reduced citrate synthase activity.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

c. HeLa cell lines. Salviati et al., 2012 report HI of genes involved in CoQ10 biosynthesis. (COQ4)-The coenzyme Q4 gene encodes a protein that organizes the multi-enzyme complex for the synthesis of coenzyme Q10 (CoQ10). HeLa cells demonstrated reduction of CoQ10. Neuromuscular symptoms include mental retardation, encephalomyopathy, and dysmorphic features. In another example Alghusen et al., 2023 report O-GlcNAc misregulation in the mitochondrial integrated stress response by regulating ATF4 in Alzheimer's disease (AD). Several neuropsychiatric diseases are now cultured in cellulo to gain insights into the cellular and molecular mechanisms of diseases.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

d. Yeast. Yeasts have been instrumental in gaining insights into the molecular basis of many human disorders, particularly those resulting from impaired cellular metabolism owing to the conservation of several cellular and metabolic proteins. Ficociello et al., 2018 used yeasts as a model system to study the molecular basis of Hailey-Hailey disease (HHD), a skin disorder caused by HI of the gene ATPase secretory pathway Ca^{2+} transporting 1 (ATP2C1) gene. The study report defects in PMR1 yeast ortholog of the gene with similarities with HHD-derived keratinocytes. Another example is of Parkinson's disease (PD) in which Fanning et al., 2019 demonstrate α -synuclein (αS) pathologically impacting the brain, a highly lipid-rich organ. Lipidomic investigations in human αS -expressing yeast show alterations in αS or lipid/fatty acid homeostasis.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

3.2. Laboratory methods

At the clinical laboratory level, HI detection methods rely on CNV and exome sequencing. Variations in the number of copies of specific DNA segments can be detected using fluorescent in situ hybridization (FISH), comparative genomic hybridization (CGH), and high-resolution array-based tests based on array comparative genomic hybridization (or aCGH), SNP array technologies, and high-resolution microarrays. Exome sequencing identifies genetic variations in the protein-coding regions (exomes) of DNA. The method relies on the size advantage of the human exome and cost-effectiveness. A few examples of CNV and sequencing are listed below.

a. CNV (copy number variation array): Xi et al. 2024 report 1q22 microdeletion of 936.3 Kb in ASH1-like histone lysine methyltransferase gene using CNVs array in developmental delay and intellectual disability (DD/ID) in patients. The microdeletion involves 24 genes, including the ASH1L (also known as KMT2H and encoding a histone lysine methyltransferase) gene. The protein has a role in transcription and in excision repair of chemical induced mutations. In another example Houwer et al., 2024 report two novel variants in the GRN (progranulin) gene in frontotemporal lobar degeneration (FTLD). The pathogenicity of variants was supported by typical neuropathological features and reduced PGRN levels.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

b. Exome sequencing: Wu and Li. 2022 report a total of 488 (33.5%) pathogenic variations among 1,457 global developmental/intellectual disabilities (GDD/ID) children cohorts, including cases of monogenic mutations and chromosomal microdeletions or micro-duplications. Rat

hippocampal neurons overexpressing the variant showed an increase in NMDAR-induced currents; additionally, immunocytochemistry revealed reduced localization between the GluN2C subunit and 14-3-3 proteins during membrane trafficking. In the ionotropic receptor N-methyl-D-aspartate receptor (NMDA) type subunit-2C (GluN2C) gene (GRIN2C), Rubino et al., 2025 report a rare damaging variant in familial late-onset AD. The study enabled genetic discernment of late-onset of disease.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

3.3. Animal models. Several animal models have been used to study HI; these include mice, Drosophila, zebrafish, and Caenorhabditis elegans. The preceding paragraph only lists a few examples to describe the range and applications.

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

a. Mice: Longakit et al., 2022 demonstrate Loss of Neurofibromatosis type 1(NF1) gene accelerated uveal and intra-dermal melanoma tumorigenesis. NF1 encodes the multifunctional tumor suppressor protein neurofibromin which regulates MAPK signaling. Transgenic mice demonstrate accelerated melanoma formation and/or growth in melanocytes and Schwann cells of peripheral nerves. RNA-seq analysis reveals cAMP signaling and myogenesis. In another example, Kaijie Ma et al., 2024, demonstrate autism-like social deficits in Ash1l HI mouse in the absent, small, or homeotic-like 1 a histone methyltransferase (ASH1L) gene. Mice displayed social deficits, increased self-grooming, and cognitive impairments. The neurons demonstrated the excitability of p neurons in the prefrontal cortex (PFC) with enhanced glutamatergic synaptic transmission, and diminished GABAergic synaptic inhibition.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

b. Drosophila: Matsuoka et al., 2024 report HI of syntaxin-binding protein 1 (STXBP1) gene in encephalopathy with α -synucleinopathies. The protein plays an important role in synaptic vesicle docking and fusion. Pathology in Drosophila melanogaster overexpression analysis demonstrates partial loss-of-function mechanisms leading to reduction in brain size. Also exacerbation of eye, degeneration, locomotor dysfunction, and dopaminergic neurodegeneration with a significant increase in detergent-insoluble α -Syn levels. In another example Huang et al., 2024, report loss-of-function in retinoblastoma-binding protein (5RBBP5) results in a syndromic neurodevelopmental disorder associated with microcephaly. RBBP5 encodes a core member of the protein complex that methylate's the histone-3 lysine-4. Epigenetic dysregulation has been

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

associated with the disease. Variants affect evolutionarily conserved amino acids located at the interface between RBBP5 and the nucleosome.

c. Fish: Wu et al.,⁵² report impaired development of neural-crest cell-derived organs and (ID) caused by the mediator complex subunit 13Lp (MED13L) gene HI in zebrafish. Knockout of med13b ortholog showed early defective migration of cranial neural crest cells (NCCs) contributing to cartilage deformities. Abnormal distribution of developing neurons in different brain regions was observed. Finally, transcriptome analysis revealed differential expression of components of Wnt and FGF signaling pathways. In another example Sadler et al., 2020, report rare and de novo coding variants in chromo domain genes in Chiari I malformation (CM1), characterized by displacement of the cerebellum, a pediatric neurological condition. Genetic screening demonstrated significant enrichment of rare and de novo non-synonymous variants in the chromo domain. Finally, HI in zebrafish demonstrated macrocephaly and posterior hindbrain displacement reminiscent of CM1.

d. Caenorhabditis elegans: Robert et al., 2023 report HI in SIN3 Transcription Regulator Family Member A (SIN3) gene as the underlying cause of Witteveen-Kolk syndrome and related (ID) and autism syndromes. SIN-3 regulates the germline transcriptional program in C. elegans and gene expression through multiple interactions, such as histone deacetylases. In another example Lazar et al., 2015, 55 report HI mutations in the human progranulin gene resulting in causing frontotemporal lobar degeneration with TAR DNA-binding protein-43 (TDP-43) inclusions. The progranulin cleavage products and granulins, exacerbate TDP-43 toxicity. C. elegans model demonstrated complete loss of the progranulin gene; however, it did not worsen TDP-43 toxicity, whereas the progranulin heterozygosity did.

4. ~~Haploinsufficiency~~Haploinsufficiency in human diseases

The spectrum of human diseases affected by HI includes organs and organ systems and affects diverse cellular and metabolic pathways. In the following paragraphs, representative examples are discussed to highlight the aforementioned spectrum. For a detailed and updated list, the reader is referred to OMIM and PubMed. Table-1 is a brief list of human diseases associated with ~~haploinsufficiency~~haploinsufficiency.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

A novel condition with cytopenia and susceptibility to hematological malignancies is defined by the ETS-related gene (ERG) gene HI. The transcription factor ERG is essential for stem cell activity and hematopoiesis. ERG localizes to promoters or enhancers of genes that control hematopoiesis. Variants of the gene exhibit loss-of-function (LOF) traits, which interfere with nuclear localization, DNA binding, and/or transcriptional transactivation. In mouse model fetal liver cells are unable to induce myeloid differentiation as well as cytokine-independent proliferation. Zerella et al., 2024. Another example is the Mitochondrial Dynamin-Like GTPase (OPA1) gene. Mutations in the gene induce adult-onset and progressive auditory neuropathy in mice. OPA1, which codes for a major GTPase associated with mitochondrial dynamin. Ultrastructural examination reveals selective loss of sensory inner hair cells and degeneration of the axons and myelin sheaths. Also a higher level of Opa1 mRNA expression supports HI as a disease mechanism. Affortit et al., 20. HI in the complement dysregulation is a key component in the pathogenesis of age-related macular degeneration (AMD) and early-onset macular drusen (EOMD). iPSCs of variants with loss of complement factor (CFH) and factor H-like protein 1 (FHL-1) display reduced mRNA expression and protein expression. Under inflammatory or oxidative stress conditions, CFH and FHL-1 expression paralleled that of controls, Lim RR 2024. In the genes expressed in kidney heterozygosity, inverted formin-2 (INF2) gene mutations cause focal segmental glomerulosclerosis (FSGS) with or without Charcot-Marie-Tooth disease. Knockout mice are susceptible to glomerular disease, in contrast to INF2 knockout mice after response to puromycin amino nucleoside (PAN)-induced kidney injury. Cellular measurements showed that mutation confers a gain-of-function effect. RNA expression analysis showed altered adhesion and mitochondria-related pathways. Podocytes in mice and human kidney organoids recapitulate adhesion and mitochondrial phenotypes. In skeletal disorders Subramanian et al., 2024 report pathogenic variants in the desmoplakin (DSP) gene encoding DSP, leading to heterogeneous skin, adnexa and heart-related phenotypes. DSP is a desmosomal component expressed in skin and heart, essential for desmosome stability and intermediate filament connection. Keratinocytes showed fragile cell-cell connections and perinuclear retracted intermediate filaments. EGFR upregulation was observed, implicating the use of EGFR inhibitors as a therapeutic alternative. Epidermal growth factor receptor (EGFR) is a transmembrane protein expressed in the basal epidermal layer involved in proliferation and differentiation processes (Daniela et al., 2024).

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, No underline, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Several genes confer HI in diabetes, such as the heterozygous mutations in the genes encoding transcriptional regulators, hepatocyte nuclear factor (HNF)-1alpha and HNF-4alpha, which cause maturity-onset diabetes of the young (MODY). MODY is a result of defective pancreatic islet cell functions. HNF-1alpha functions through genetic network that is activated during differentiation, allowing cells to maintain critical specialized functions. At the core lies a cross-regulatory feedback circuit between HNF-1alpha and HNF-4alpha. The loss of either gene alleles can increase the feedback circuit resulting in decreased expression of all four alleles selectively in beta-cells (Ferrer et al., 2002). Of the several genes expressed in the liver, the FERM Domain Containing Kindlin2 (FERMT2) gene codes for the Kindlin-2 protein. HI protects against fatty liver and glucose intolerance without affecting energy metabolism in mice by targeting. Kindlin-2 binds and stabilizes Foxo1 by inhibiting its ubiquitination and degradation through the Skp2 E3 ligase. Kindlin-2 overexpression in the liver exacerbates lipid metabolism disorder in hepatocytes through Foxo1 phosphorylation (Huanqing et al., 2022). Several genes are implicated in human intellectual disability and other associated diseases. AT-hook DNA binding motif containing 1 (AHDC1) HI causes Xia-Gibbs syndrome (XGS), a rare syndromic disorder characterized by developmental delay with intellectual disability, muscular hypotonic, brain anomalies, and nonspecific dysmorphic features. The interstitial deletion (1p36.11p35.3) encompasses the coding region of the AHDC1 gene (Bertrand et al., 2024). In the gastrointestinal (GI) Lynch syndrome (LS), the tract is susceptible to carcinogenesis. According to Knudson's two-hit hypothesis, tumors of (LS) patients exhibit high levels of microsatellite instability (MSI), which is caused by failure of DNA mismatch repair (MMR). Colon of knock-out mice indicate variable MLH1 levels are leading to tissue-specific MSI in advanced stages of neoplasia. Shrestha et al., 2021.

HI in several genes and networks involved in several human disorders, such as asthma, diabetes, cancer, and several other disorders, has been reported by several investigators. In the foregoing paragraph we describe a few representative genes and associated pathways. The reader is referred to PubMed for a detailed literature review. Few genes belonging to this category are listed in Table-2. Transcription factor 3 (TCF3) is a transcription factor contributing to early lymphocyte differentiation. Germline mono-allelic dominant negative and bi-allelic loss-of-function (LOF) null TCF3 mutations cause a fully penetrant severe immunodeficiency with

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

reduced transcription or translation resulting in reduced wild-type TCF3 protein expression. and in B-cell defects. Murine TCF3 HI resulted in a reduction of circulating B cells RNA sequencing analysis implies gene-dosage effect (Boast et al., 2023). Yan Li et al., 2021 report HI in the Hedgehog interacting protein (HHIP) gene. HHIP protein represses airway remodeling and metabolic reprogramming in chronic obstructive pulmonary disease (COPD)-derived airway smooth muscle cells.

Several genes are implicated in cancer, which include oncogenes, tumor suppressor genes, and genes coding for several cellular pathways. AaRS-interacting multifunctional protein 3 (AIMP3), previously known as p18, a tumor suppressor gene, is shown to up-regulate p53 in response to DNA damage. AIMP3 HI disrupts oncogene p53-induced activation and genomic stability. Growth factor- or Raps-dependent induction of p53 was blocked by loss of AIMP3. Heterozygous cells were susceptible to cell transformation induced by oncogenes Ras or Myc alone and displayed cell division and chromosomal structure abnormalities (Bum-Joon Park et al., 2006). Another example is the serine-/arginine-rich splicing factor 2 gene (SRSF2). Recurrent mutations in the gene are common in neoplastic diseases. The gene plays pivotal roles in pre-mRNA processing and gene transcription and regulates bi-directional transcription of DNA replication and repair genes. HI induces DNA damage without halting the cell cycle. In mice, deletion or homozygous mutation both cause extensive DNA damage. Frost, Erin L. 2024. Numerous tumor suppressor genes (TSGs), including p27Kip1, p53, DMP1, NF1, and PTEN, have been found to exhibit HI, resulting in the cell's incapacity to carry out regular cellular tasks, aiding the growth of tumors. AML1, EGR1, TGFβR1/2, and SMAD4 are among the recently identified TSGs in which (HI) is demonstrated. Both human samples and gene knockout mice models have validated these results. Multiple (HI) TSGs may collaborate in 5q-, 7q-, and 8q- disorders, according to recent research (Inoue and Fry 2017). Network analyses of genes in cancer are also reported. In ovarian cancer (OV) HI network (KEGG) pathways indicate autophagy and compensatory proteolysis pathways as significant (Delaney et al., 2017).

Glycogen synthase kinase 3 (GSK-3) gene, which encodes a serine-threonine kinase with two isoforms (α - and β), is implicated in type 2 diabetes mellitus (T2D). Animal models of HI suggest an isoform-specific role of GSK-3 in high-fat diet (HFD)-induced obesity and glucose

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, Font color: Auto, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted ... [1]

Formatted ... [2]

Formatted ... [3]

intolerance. Tamoxifen-inducible global GSK-3 α/β gain implicates the dose-response effect of GSK-3 isoforms in heterozygote mice. Post-HFD, GSK-3 α heterozygous and controls displayed a comparable phenotype; however they were significantly protected against glucose intolerance (Gupte et al., 2024). Another gene is the regulatory factor X 6 (RFX6), indicated in pancreatic endocrine development and differentiation. RFX6 HI predisposes to diabetes through impaired beta cell function. Heterozygous RFX6 $+/-$ SC-islets exhibited HI associated with reduced beta cell maturation. Allelic series suggest defective pancreatic beta cells, mirroring the phenotype observed in Mitchell-Riley syndrome (Ibrahim et al., 2024).

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

HI affects several genes in the cardiac system. An example is the TGF-beta-activated kinase 1/MAP3K7 binding protein 2 gene (MAP3K7) causing congenital heart defects (CHDs). The role of HI of this gene is further supported by its conserved expression in the developing zebrafish heart and dosage sensitivity (Thienpont et al., 2010). Mutations in the lamin A/C (LMNA) gene which encodes nuclear membrane protein cause dilated cardiomyopathy (DCM) and apoptosis-triggered cardiac conduction system disease. HI caused early-onset programmed cell death of atrioventricular (AV) nodal myocytes and progressive electrophysiological disease. Conduction system investigations indicate (AV) nodal myocytes had abnormally shaped nuclei and active apoptosis. Telemetric and in vivo electrophysiological studies showed arrhythmias and impaired contractility (Wolf et al., 2008).

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Finally, HI is demonstrated in several human multi-organ disorders; an example is reported by (Del Viso et al., 2025), where HI of SF3B2 is associated with anomalies such as skeletal extra craniofacial, cardiac, renal, and abnormalities of the central nervous system (CNS). De novo or inherited variants have been associated with either or both inter- and interfamilial variability. In summary, HI affects several genes involved in several organs, organ, and multi-organ systems, reflecting their abrogating roles in the spectrum of cellular pathways and cellular mechanisms (functions, DNA replication, and recombination) and surveillance (DNA repair). Also, they affect metabolic and cardiac, blood, renal, and skeletal disorders.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

5. Applications of HI and conclusion

The findings on model organisms have clarified few aspects regarding the molecular basis of HI. In the budding yeast, *Saccharomyces cerevisiae*, dosage-sensitive genes are believed to be either that interact widely with protein partners. In flowering plants, interaction networks (mathematical) or part of an essential metabolic pathway crucial elements shaping HI (Navarro-Quiles et al., 2024). Hence the phenomenon of HI extends to other eukaryotes. Hence, it can be used to test several gene products and hypothesis.

From a human genetics perspective recent advancements in high-throughput sequencing and computational biology have enabled an exhaustive genome-wide study of HI loci in the human genome. The observation of dosage and balance hypothesis which explains why mutations affecting HI genes are often associated with dominantly inherited conditions, from congenital malformations to neoplasias. This will enable genetic-testing in the clinical setup (prenatal). Large-scale sequencing efforts to identify human genes that are intolerant to or under strong selection against heterozygous loss-of-function are sparking tremendous interest in HI and its role in disease etiology (Kaplanis et al., 2020; Karczewski et al., 2020). With increased recognition of age-related health problems globally, the role of insufficient DNA methylation and its effects on healthy aging is another area of intense research (Liu et al., 2011). In the area of basic research, such as the pluripotent reprogramming factors, iPSC quality and efficiency studies the role of HI in pluripotent cells and affect developmental potential and their cell derivatives given significance (Benchetrit et al., 2019; Sebban and Buganim 2015). A complete knockout of a single key pluripotency gene may drastically affect embryonic stem cell function and epigenetic reprogramming. Additionally, methylation can complicate the program. Defective telomerase activity is a key cause of various diseases due to telomerase RNA component (TERC) (Calado and Young 2009b). HI cause telomere shortening and is implicated in several diseases.

In summary, high-throughput genomic data, computational modeling, and comparative analyses from different biological systems converge into an integrated understanding of the molecular and evolutionary basis for HI in humans. Such integration signifies that gene dosage, molecular interactions, and selective pressures are intimately interconnected, and thus provide vital insights into the pathogenesis of dosage-sensitive diseases and into the design of appropriate therapeutic strategies. The

Formatted: Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold, Not Italic

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold, Not Italic

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold, Not Italic

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Font: +Body, 11 pt, Not Bold, Not Italic

Formatted: Font: +Body, 11 pt, Not Bold

Formatted: Default Paragraph Font, Font: +Body, 11 pt

Formatted: Font: +Body, 11 pt, Not Italic

Formatted: Font: +Body, 11 pt, Not Italic

Formatted: Font: +Body, 11 pt, Not Italic, Font color: Auto

Formatted: Font: +Body, 11 pt, Not Italic

study of HI has important implications for human biomedical research. The author acknowledges that several areas of relevance to human biology, genetics have not been included in the review.

UNDER PEER REVIEW

References

1. Adam F Johnson, Ha T Nguyen 1, Reiner A Veitia. Causes and effects of haploinsufficiency. *Biol Rev Camb Philos Soc*. 2019 Oct;94(5):1774-1785. doi: 10.1111/brv.12527. Epub 2019 May 31. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
2. Fuller ZL, Berg JJ, Mostafavi H, Sella G, Przeworski M. Measuring intolerance to mutation in human genetics. *Nat Genet*. 2019 May;51(5):772-776. doi: 10.1038/s41588-019-0383-1. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
3. LEJEUNE J. AUTOSOMAL DISORDERS. *Pediatrics*. 1963 Sep;32:326-37. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
4. Fisher RA. Correspondence. *Eugen Rev*. 1928 Apr;20(1):72. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
5. S Wright. An Analysis of Variability in Number of Digits in an Inbred Strain of Guinea Pigs. *Genetics*. 1934 Nov;19(6):506-36. doi: 10.1093/genetics/19.6.506. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
6. Kondrashov FA, Koonin EV. A common framework for understanding the origin of genetic dominance and evolutionary fates of gene duplications. *Trends Genet*. 2004 Jul;20(7):287-90. doi: 10.1016/j.tig.2004.05.001. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
7. H Kacser, J A Burns. The molecular basis of dominance. *Genetics*. 1981 Mar-Apr;97(3-4):639-66. doi: 10.1093/genetics/97.3-4.639. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
8. Deutschbauer AM, Jaramillo DF, Proctor M, Kumm J, Hillenmeyer ME, Davis RW, Nislow C, Giaever G. Mechanisms of haploinsufficiency revealed by genome-wide profiling in yeast. *Genetics*. 2005 Apr;169(4):1915-25. doi: 10.1534/genetics.104.036871. Epub 2005 Feb 16. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
9. www.nature.com/scitable/ Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
11. Hosea R, Hillary S, Naqvi S, Wu S, Kasim V. The two sides of chromosomal instability: drivers and brakes in cancer. *Signal Transduct Target Ther*. 2024 Mar 29;9(1):75. doi: 10.1038/s41392-024-01767-7. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
12. Anthony Levasseur, Pierre Pontarotti. The role of duplications in the evolution of genomes highlights the need for evolutionary-based approaches in comparative genomics. *Biol Direct*. 2011 Feb 18;6:11. doi: 10.1186/1745-6150-6-11. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
13. Jeffrey J. Moffat, Eui-Man Jung, Minhan Ka, Amanda L. Smith, Byeong Tak Jeon, Gijis W.E. Santen, Woo-Yang Kim. The role of ARID1B, a BAF chromatin remodeling complex subunit, in neural development and behavior. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. Volume 89, 8 March 2019, Pages 30-38. <https://doi.org/10.1016/j.pnpbp.2018.08.021>. Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic
14. Santen GW, Aten E, Sun Y, Almomani R, Gilissen C, Nielsen M et al. Mutations in SWI/SNF Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

chromatin remodeling complex gene ARID1B cause Coffin-Siris syndrome. Nat Genet 2012; 44(4): 379–380.

15. Santen GW, Clayton-Smith J, consortium ABC. The ARID1B phenotype: what we have learned so far. Am J Med Genet C Semin Med Genet 2014; 166C(3): 276–289.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

16. Nishant Singhal, Johannes Graumann, Guangming Wu, Marcos J. Araújo-Bravo, Dong Wook Han, Boris Greber, Luca Gentile, Matthias Mann, Hans R. Schöler, Chromatin-Remodeling Components of the BAF Complex Facilitate Reprogramming, Cell, Volume 141, Issue 6, 2010, Pages 943-955.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

17. Alver BH, Kim KH, Lu P, Wang X, Manchester HE, Wang W, Haswell JR, Park PJ, Roberts CW. The SWI/SNF chromatin remodelling complex is required for maintenance of lineage specific enhancers. Nat Commun. 2017 Mar 6;8:14648. doi: 10.1038/ncomms14648. PMID: 28262751; PMCID: PMC5343482

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

18. Anthony Levasseur, Pierre Pontarotti The role of duplications in the evolution of genomes highlights the need for evolutionary-based approaches in comparative genomics. Biol Direct. 2011 Feb 18;6:11. doi: 10.1186/1745-6150-6-11.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

19. Johnson, A. F., Nguyen, H. T. and Veitia, R. A. (2019). Causes and effects of haploinsufficiency. Biol. Rev. 94, 1774-1785. doi:10.1111/brev.12527

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

20. Morrill, S. A. and Amon, A. (2019). Why haploinsufficiency persists. Proc. Natl. Acad. Sci. USA 116, 11866-11871

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

21. Veitia, R. A. (2002). Exploring the etiology of haploinsufficiency. BioEssays 24, 175-184. doi:10.1002/bies.10023

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

22. Roman Zug. Developmental disorders caused by haploinsufficiency of transcriptional regulators: a perspective based on cell fate determination. 2022. The Company of Biologists. doi:10.1242/bio.058896

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

23. Palii, C. G., Cheng, Q., Gillespie, M. A., Shannon, P., Mazurczyk, M., Napolitani, G., Price, N. D., Ranish, J. A., Morrissey, E., Higgs, D. R. et al. (2019). Single cell proteomics reveal that quantitative changes in co-expressed lineage-specific transcription factors determine cell fate. Cell Stem Cell 24, 812-820. doi:10.1016/j.stem.2019.02.006.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

24. Ferrer, J. (2002). A genetic switch in pancreatic β -cells: implications for differentiation and haploinsufficiency. Diabetes 51, 2355-2362. doi:10.2337/diabetes.51.8.2355

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

25. Stévant, I., Kühne, F., Greenfield, A., Chaboissier, M.-C., Dermizakis, E. T. and Nef, S. (2019). Dissecting cell lineage specification and sex fate determination in gonadal somatic cells using single-cell transcriptomics. Cell Rep. 26, 3272-3283. doi:10.1016/j.celrep.2019.02.069

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

26. ▲ Liu, J., Perumal, N. B., Oldfield, C. J., Su, E. W., Uversky, V. N. and Dunker, A. K. (2006). Intrinsic disorder in transcription factors. *Biochemistry* 45, 6873-6888. doi:10.1021/bi0602718
27. ▲ Dyson, H. J. and Wright, P. E. (2016). Role of intrinsic protein disorder in the function and interactions of the transcriptional coactivators CREB-binding protein (CBP) and p300. *J. Biol. Chem.* 291, 6714-6722. doi:10.1074/jbc.R115.692020
28. ▲ Kamachi, Y., Uchikawa, M., Tanouchi, A., Sekido, R. and Kondoh, H. (2001). Pax6 and SOX2 form a co-DNA-binding partner complex that regulates initiation of lens development. *Genes Dev.* 15, 1272-1286. doi:10.1101/gad.887101
29. ▲ Bondurand, N., Pingault, V., Goerich, D. E., Lemort, N., Sock, E., Le Caignec, C., Wegner, M. and Goossens, M. (2000). Interaction among SOX10, PAX3, and MITF, three genes altered in Waardenburg syndrome. *Hum. Mol. Genet.* 9, 1907-1917. doi:10.1093/hmg/9.13.1907
30. ▲ Shi, X., Richard, J., Zirbes, K. M., Gong, W., Lin, G., Kyba, M., Thomson, J. A., Koyano-Nakagawa, N. and Garry, D. J. (2014). Cooperative interaction of Etv2 and Gata2 regulates the development of endothelial and hematopoietic lineages. *Dev. Biol.* 389, 208-218. doi:10.1016/j.ydbio.2014.02.018
31. ▲ Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. and Sharp, P. A. (2017). A phase separation model for transcriptional control. *Cell* 169, 13-23. doi:10.1016/j.cell.2017.02.007
32. ▲ Spitz, F. and Furlong, E. E. M. (2012). Transcription factors: from enhancer binding to developmental control. *Nat. Rev. Genet.* 13, 613-626. doi:10.1038/nrg3207
33. ▲ Reiter, F., Wienerroither, S. and Stark, A. (2017). Combinatorial function of transcription factors and cofactors. *Curr. Opin. Genet. Dev.* 43, 73-81. doi:10.1016/j.gde.2016.12.007
34. ▲ Roni Sarel-Gallily, Tamar Golan-Lev, Atilgan Yilmaz, Ido Sagi, Nissim Benvenisty. Genome-wide analysis of haploinsufficiency in human embryonic stem cells. *Cell Rep.* 2022 Mar 29;38(13):110573. doi: 10.1016/j.celrep.2022.110573.
35. ▲ Eva Niggel, Arjan Bouman, Lauren C Briere 3, Remco M Hoogenboezem, Ilse Wallaard, Joohyun Park, Genomics England Research Consortium; Undiagnosed Diseases Network. HNRNPC haploinsufficiency affects alternative splicing of intellectual disability-associated genes and causes a neurodevelopmental disorder. *Am J Hum Genet.* 2023 Aug 3;110(8):1414-1435. doi: 10.1016/j.ajhg.2023.07.005.
36. ▲ Jens Schuster, Claudia de Guidi, Rekha Tripathi, Joakim Klar, Niklas Dahl. Generation of a human iPSC line (UIGPi015-A) from a patient with Dravet syndrome and a 2.9 Mb deletion spanning SCN1A on chromosome 2. *Stem Cell Res.* 2022 Apr;60:102712. doi: 10.1016/j.scr.2022.102712. Epub 2022 Feb 16.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font color: Auto, Pattern: Clear

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

37. Rachel James, Kiterie M. E. Faller, Ewout J. N. Groen, Brunhilde Wirth & Thomas H. Gillngwater. Altered mitochondrial function in fibroblast cell lines derived from disease carriers of spinal muscular atrophy. *Communications Medicine* volume 4, Article number: 86 (2024).
38. Leonardo Salviati 1, Eva Trevisson, Maria Angeles Rodriguez Hernandez, Alberto Casarin, Vanessa Pertegato et al., Haploinsufficiency of COQ4 causes coenzyme Q10 deficiency. *J Med Genet.* 2012 Mar;49(3):187-91. doi: 10.1136/jmedgenet-2011-100394.
39. Ibtihal M Alghusen , Marisa S Carman, Heather M Wilkins, Taylor A Strope , Caleb Gimore, Halyna Fedosyuk , Jad Shawa , Sophiya John Ephrame , Aspin R Denson, Xiaowan Wang , Russell H Swerdlow, Chad Slawson . O-GlcNAc impacts mitophagy via the PINK1-dependent pathway. *Front Aging Neurosci.* 2024 Aug 8;16:1387931. doi: 10.3389/fnagi.2024.1387931.
40. Graziella Ficociello, Azzurra Zonfrilli , Samantha Cialfi, Claudio Talora, Daniela Uccelletti. Yeast-Based Screen to Identify Natural Compounds with a Potential Therapeutic Effect in Hailey-Hailey Disease. *Int J Mol Sci.* 2018 Jun 20;19(6):1814. doi: 10.3390/ijms19061814.
41. Saranna Fanning, Aftabul Haque, Thibaut Imberdis, Valeriya Baru, M Inmaculada Barrasa, Silke Nuber, et al., Lipidomic Analysis of α -Synuclein Neurotoxicity Identifies Stearoyl CoA Desaturase as a Target for Parkinson Treatment. *Mol Cell.* 2019 Mar 7;73(5):1001-1014.e8. doi: 10.1016/j.molcel.2018.11.028. Epub 2018 Dec 4.
42. Hui Xi, Ying Peng, Wanqin Xie, Jialun Pang , Na Ma, Shuting Yang, Jinping Peng, Hua Wang. A chromosome 1q22 microdeletion including ASH1L is associated with intellectual disability in a Chinese family. *Mol Cytogenet.* 2020 Jun 4;13:20. doi: 10.1186/s13039-020-00483-5. eCollection 2020.
43. Julie F H De Houwer, Elise G P Dopfer, Ana Rajcic, Renee van Buuren, Marina Arcaro, Daniela Galimberti, Guido J Breedveld, Two novel variants in GRN: the relevance of CNV analysis and genetic screening in FTLN patients with a negative family history. *J Neurol.* 2024 Dec 16;272(1):64. doi: 10.1007/s00415-024-12758-7.
44. Dandan Wu and Rong Li. Genetic analysis of neurodevelopmental disorders in children. *Front Child Adolesc Psychiatry.* 2022 Dec 7;1:987339. doi: 10.3389/frcha.2022.987339. eCollection 2022.
45. Elisa Rubino, Maria Italia, Elisa Giorgio , Silvia Boschi , Paola Dimartino, Tommaso Pippucci, Fausto Roveta. Exome sequencing reveals a rare damaging variant in GRIN2C in familial late-onset Alzheimer's disease. *Alzheimers Res Ther.* 2025 Jan 14;17(1):21. doi: 10.1186/s13195-024-01661-y.
46. Anne Nathalie Longakit, Oscar Urtatiz , Amy Luty, Christina Zhang, Chloe Hess, Alyssa Yoo, Hannah Bourget, Catherine D Van Raamsdonk. Loss of NF1 accelerates uveal and intra-dermal melanoma tumorigenesis and oncogenic GNAQ transforms Schwann cells. *Cancer Res Commun.* 2025 Jan 13. doi: 10.1158/2767-9764.CRC-24-0386. Online ahead of print.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

47. ▲ Kaijie Ma, Kylee McDanie, Daoqi Zhang, Maria Webb, Luye Qin. Chemogenetic Inhibition of Prefrontal Cortex Ameliorates Autism-Like Social Deficits and Absence-Like Seizures in a Gene-Trap Ash11 Haploinsufficiency Mouse Model. *Genes* (Basel). 2024 Dec 18;15(12):1619. doi: 10.3390/genes15121619.
48. ▲ Taro Matsuoka, Hideki Yoshida, Takashi Kasai, Takenori Tozawa, Tomoko Iehara, Tomohiro Chiyonobu. α -Synuclein pathology in *Drosophila melanogaster* is exacerbated by haploinsufficiency of Rop: connecting STXBP1 encephalopathy with α -synucleinopathies. *Human Molecular Genetics*, Volume 33, Issue 15, 1 August 2024, Pages 1328–1338.
49. ▲ Yue Huang, Kristy L Jay, Alden Yen-Wen Huang, Jijun Wan, Sharayu V Jangam, Odelia Chorin, Annick Rothschild, Ortal Barel, et al., Loss-of-function in RBBP5 results in a syndromic neurodevelopmental disorder associated with microcephaly. *Genet Med.* 2024 Nov;26(11):101218. doi: 10.1016/j.gim.2024.101218. Epub 2024 Jul 19.
50. ▲ Kagistia Hana Utami, Cecilia Lanny Winata, Axel M Hillmer, Irene Aksoy, Hoang Truong Long, Herty Liany, et al., Impaired development of neural-crest cell-derived organs and intellectual disability caused by MED13L haploinsufficiency. *Hum Mutat.* 2014 Nov;35(11):1311–20. doi: 10.1002/humu.22636. Epub 2014 Sep 23.
51. ▲ Brooke Sadler 1, Jackson Wilborn 2, Lilian Antunes 3, Timothy Kuensting 2, Andrew T Hale 4, Stephen R Gannon 5, Kevin McCall 3, Carlos Cruchaga 6, Rare and de novo coding variants in chromodomain genes in Chiari I malformation. *Am J Hum Genet.* 2021 Jan 7;108(1):100–114. doi: 10.1016/j.ajhg.2020.12.001. Epub 2020 Dec 21.
52. ▲ Valerie J Robert, Matthieu Caron, Loic Gely, Annie Adrait, Victoria Pakulska, Yohann Couté, Manon Chevalier, Christian G Riedel. SIN-3 acts in distinct complexes to regulate the germline transcriptional program in *Caenorhabditis elegans*. *Development.* 2023 Oct 17;150(21):dev201755. doi: 10.1242/dev.201755
53. ▲ Dominique A Salazar, Victoria J Butler, Andrea R Argouarch, Tsung-Yuan Hsu, Amanda Mason, Ayumi Nakamura, Helen McCurdy, David Cox. The Progranulin Cleavage Products, Granulins, Exacerbate TDP-43 Toxicity and Increase TDP-43 Levels. *J Neurosci.* 2015 Jun 24;35(25):9315–28. doi: 10.1523/JNEUROSCI.4808-14.2015.
54. ▲ Jiarna R Zerella, Claire C Homan, Peer Arts, Xuzhu Lin, Sam J Spinelli, Parvathy Venugopal, et al., Germ line ERG haploinsufficiency defines a new syndrome with cytopenia and hematological malignancy predisposition. 2024 Oct 24;144(17):1765–1780.
55. ▲ Corentin Affortit, Carolanne Coyat, Anissa Rym Saidia, Jean-Charles Ceccato, Majida Charif, et al., The human OPA1 delTTAG mutation induces adult onset and progressive auditory neuropathy in mice. *Cell Mol Life Sci.* 2024 Feb 9; 81(1):80.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

56. ▲ Rayne R Lim, Sharlene Shiralí, Jessica Rowlan, Abbi L Engel, Marcos Nazario Jr , Kelie Gonzalez , Aspen Tong , Jay Neitz, Maureen Neitz , Jennifer R Chao .CFH Haploinsufficiency and Complement Alterations in Early-Onset Macular Degeneration. Invest Ophthalmol Vis Sci. 2024 Apr 1;65(4):43.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

57. ▲ Balajikarthick Subramanian , Sarah Williams , Sophie Karp , Marie-Flore Hennino , Sonako Jacas , Miriam Lee , Cristian V Riella, Seth L Alper , Henry N Higgs , Martin R Pollak.INF2 mutations cause kidney disease through a gain-of-function mechanism. Sci Adv. 2024 Nov 15;10(46):eadr1017.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

58. ▲ Daniela Andrei , Jeroen Bremer , Duco Kramer , Albertine M Nijenhuis , Marije van der Molen , Gilles F H Diercks , Peter C van den Akker , Mathilde C S C Vermeer , Peter van der Meer , Maria C Bolling 1.Epidermal growth factor receptor inhibition leads to cellular phenotype correction of DSP-mutated keratinocytes.Exp Dermatol. 2024 Mar; 33(3):e15046.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

59. ▲ Jorge Ferrer A genetic switch in pancreatic beta-cells: implications for differentiation and haploinsufficiency. Diabetes. 2002 Aug; 51(8):2355-62.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

60. ▲ Huangqing Gao , Liang Zhou , Yiming Zhong , Zhen Ding, Sixiong Lin , Xiaoting Hou , Xiaoqian Zhou , Jie Shao , Fan Yang , Xuenong Zou , Huiling Cao, Guozhi Xiao .Kindlin-2 haploinsufficiency protects against fatty liver by targeting Foxo1 in mice.Nat Commun. 2022 Feb 23; 13(1):1025.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

61. ▲ Miriam Bertrand , Gulalai Shah , Brent S Pedersen , Alexander Schulz , Anja Weise , Thomas Liehr , Peter Huppke , Stephanie DiTroia , Aaron R Quinlan , Tobias B Haack , Ralf A Husain .De novo AHDC1 Deletions Identified by Genome Sequencing in Two Individuals with Xia-Gibbs Syndrome. Mol Syndromol. 2024 Oct; 15(5):389-397.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

62. ▲ Kul S Shrestha , Minna M Tuominen , Liisa Kauppi .Mlh1 heterozygosity and promoter methylation associates with microsatellite instability in mouse sperm. Mutagenesis. 2021 Jul 7; 36(3):237-244.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

63. ▲ Brigitte Boast , Shubham Goel , Luis I González-Granado , Julie Niemela , Jennifer Stoddard , Emily S J Edwards , Sandali Seneviratne. et al., TCF3 haploinsufficiency defined by immune, clinical, gene-dosage, and murine studies. J Allergy Clin Immunol. 2023 Sep; 152(3):736-747.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

64. ▲ Yan Li , Li Zhang, Francesca Polverino , Feng Guo , Yuan Hao , Taotao Lao, Shuang Xu , Lijia Li , Betty Pham , Caroline A Owen , Xiaobo Zhou .Hedgehog interacting protein (HHIP) represses airway remodeling and metabolic reprogramming in COPD-derived airway smooth muscle cells. Sci Rep. 2021 Apr 27; 11(1):9074.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

65. ▲ Bum-Joon Park , Young Sun Oh, Seung Yong Park, So Jung Choi, Cornelia Rudolph, Brigitte Schlegelberger, Sunghoon Kim. AIMP3 haploinsufficiency disrupts oncogene-induced p53 activation and genomic stability. Cancer Res. 2006 Jul 15;66(14):6913-8.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

66. [Mmm](#)
67. [Kazushi Inoue , Elizabeth A Fry 1.Haploinsufficient tumor suppressor genes. Adv Med Biol. 2017;118:83-122.](#)
68. [Joe Ryan Delaney , Chandni B Patel , Katelyn McCabe Willis , Mina Haghighiabyaneh , Joshua Axelrod , Isabelle Tancioni , Dan Lu , Jaidev Bapat , Shanique Young et al.,Haploinsufficiency networks identify targetable patterns of allelic deficiency in low mutation ovarian cancer. Nat Commun. 2017 Feb 15;8:14423.](#)
69. [Manisha Gupte, Prachi Umbarkar 2, Jacob Lemon, Sultan Tousif, Hind Lal.Animal models of haploinsufficiency revealed the isoform-specific role of GSK-3 in HFD-induced obesity and glucose intolerance.Am J Physiol Cell Physiol. 2024 Dec 1;327\(6\):C1349-C1358. doi: 10.1152/ajpcell.00552.2024. Epub 2024 Sep 30.](#)
70. [Hazem Ibrahim , Diego Balboa , Jonna Saarimäki-Vire , Hossam Montaser , Oleg Dyachok , Per-Eric Lund , Muhmmad Omar-Hmeadi 3, Jouni Kvist RFX6 haploinsufficiency predisposes to diabetes through impaired beta cell function](#)
[Diabetologia. 2024 Aug;67\(8\):1642-1662. doi: 10.1007/s00125-024-06163-y. Epub 2024 May 14.](#)
71. [Bernard Thienpont 1, Litu Zhang, Alex V Postma, Jeroen Breckpot, Léon-Charles Tranchevent, Peter Van Loo, Kjeld Møllgård et aal.,Haploinsufficiency of TAB2 causes congenital heart defects in humans. Am J Hum Genet. 2010 Jun 11; 86\(6\):839-49.](#)
72. [Cordula M Wolf , Michael Arad, Ferhaan Ahmad, Atsushi Sanbe, Scott A Bernstein, Okan Toka, Tetsuo Konno, Gregory Morley, Jeffrey Robbins, J G Seidman, Christine E Seidman, Charles I Berul.Reversibility of PRKAG2 glycogen-storage cardiomyopathy and electrophysiological manifestations.Circulation. 2008 Jan 15; 117\(2\):144-54.](#)
73. [Florescia Del Viso , Dihong Zhou , Susan Starling , Emily Fleming , Carol Saunders .SF3B2 Haploinsufficiency Associated With Hirschprung Disease and Complex Cardiac Defect Without Craniofacial Microsomia. Case Reports Am J Med Genet A. 2025 Feb; 197\(2\):e63886.](#)
74. [Carla Navarro-Quiles, Samuel Daniel Lup , Tamara Muñoz-Nortes, Héctor Candela , José Luis Micol.The genetic and molecular basis of haploinsufficiency in flowering plantsTrends Plant Sci. 2024 Jan;29\(1\):72-85. doi: 10.1016/j.tplants.2023.07.009. Epub 2023 Aug 24.](#)
75. [Joanna Kaplanis, Kaitlin E Samocha, Laurens Wiel, Zhancheng Zhang, Kevin J Arvai, Ruth Y Eberhardt, Giuseppe Gallone 1, Stefan H Lelieveld, Hilary C Martin, et al., Deciphering Developmental Disorders Study.Evidence for 28 genetic disorders discovered by combining healthcare and research data.Nature. 2020 Oct;586\(7831\):757-762. doi: 10.1038/s41586-020-2832-5.](#)
76. [Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alfoldi J, Wang Q, Collins RL, Laricchia](#)

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

KM, Ganna A, Birnbaum DP, Gauthier LD, Brand H, et al., Genome Aggregation Database Consortium; Neale BM, Daly MJ, MacArthur DG. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*. 2020 May;581(7809):434-443. doi: 10.1038/s41586-020-2308-7. Epub 2020 May 27.

77. Hana Benchetrit, Mohammad Jaber, Valery Zayat, Shulamit Sebban, Avital Pushett, Kirill Makedonski, Direct Induction of the Three Pre-implantation Blastocyst Cell Types from Fibroblasts. *Cell Stem Cell*. 2019.

78. Shulamit Sebban, Yosef Buganim. Nuclear Reprogramming by Defined Factors: Quantity Versus Quality. *Trends Cell Biol*. 2016 Jan;26(1):65-75. doi: 10.1016/j.tcb.2015.08.006. Epub 2015 Oct 1.

79. Rodrigo T Calado. Telomere diseases. *Neal S Young N Engl J Med*. 2009 Dec 10;361(24):2353-65. doi: 10.1056/NEJMra0903373.

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic

Figures and tables

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Not Italic, (Asian) Chinese (Simplified, PRC)




Figure-1 . Possible genetic and cellular consequences of Haploinsufficiency(HI).Adapted from Roman Zug 2022.




Figure-2. Molecular consequences due to gene dosage alterations. Adapted from M. Felicia Basilicata¹D, Claudia Isabelle Keller Valsecchi¹ 2021.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Table1. Brief list of human diseases associated with Haploinsufficiency

Formatted: Font: +Body, 11 pt, Not Italic

Formatted: Font: +Body, 11 pt, Not Italic

Sl. no	Disease	Gene involved	Organ involved/Major anomalies	Associated pathology	Reference
1	Xia-Gibbs Syndrome	AT-hook DNA binding motif containing 1AHDC1)	Intellectual disability, muscular hypotonia, brain anomalies.	Foot deformity, skin and connective tissue abnormalities.	Bertrand M <i>et al.</i> , 2024
2	Nedlaad disorder	cell cycle associated protein 1(CAPRIN1)	Dysmorphic facial and digital, hand anomalies feature. Respiratory difficulties. Central nervous system.	Delay in walking, language delay, anomalies mainly of the corpus callosum. Mild hearing loss.	Pavinato L <i>et al.</i> , 2024
3	Von recklinghausen disease	NF1 (Neuro fibromin gene)	Skin Lesions and café-au-lait spots on skin. Lisch Nodules on Iris.	Fibromatous tumors on skin. Benign and malignant tumor of skin.	William <i>et al</i> 2009
4	Autoimmune	cytotoxic T-	Abnormal	Inflammatory lung	Lopez

	ne lymphoproliferative disease	lymphocyte associated protein 4(CTLA4)	lymphocytic infiltration of non lymphoid organs include brain,lung and gastrointestinal tract.	lesions. Lymphocytic enteropathy.Seizures and Headache.	nevado <i>et al.</i> , 2021
5	Leri-weill dyschondrosteosis	SHOX (homeobox SHOX)	Skeletal dysplasia of bone and cartilage,medulung wrist deformity includes proximal carpal bones.	Tibiofibular disproportion. Problems on limbs.	Ross <i>et al.</i> , 2005
6	Niemann-pick disease	NPC intracellular cholesterol transporter 1(NPC1)	Neurological abnormalities include ataxia, seizures. Loss of speech and myoclonic jerks.	Amentia. Extension of spine and legs. Retinal degeneration. Myoclonus or Bulbar involvement. Hepatosplenomegaly.	Vance 2006
7	Darier-white disease	ATPase sarcoplasmic/endoplasmic reticulum Ca2+ transporting 2ATP2A2	Crusted plaques, painfull erosions,warty papules and plaques in seborrheic areas and distinctive nail abnormalities.	Bullous lesions.Multiple bone cysts resulting in non-traumatic fractures of the long bones.	Castori <i>et al.</i> , 2009
8	Mowat-wilson syndrome	zinc finger E-box binding homeobox 2(ZEB2)	Impaired intellectual development. Delayed motor development.	Microcephaly. Mental retardation. Epilepsy.	Mowat <i>et al.</i> , 2003
9	Charcot-marie-tooth disease, neuronal, type 2d	glycyl-tRNA synthetase 1(GARS1)	Muscle weakness.Distal sensory Impairment. Induced hand cramps.	Severe muscle weakness and atrophy of the hands and mild to moderate weakness of the feet. Scoliosis.	Yalcouye, A <i>et al.</i> , 2019
10	William-beuren syndrome	membrane associated guanylate kinase, WW and PDZ domain containing	Dental anomalies and peripheral pulmonary artery stenosis.	Abnormal scarring, wrinkles, soft skin, difference in skin elasticity.	Kozel BA <i>et al.</i> , 2014

		2(MAGI2)			
--	--	----------	--	--	--

Table-2.Human clinical syndromes associated with ~~haploinsufficiency~~haploinsufficiency

Formatted: Font: +Body, 11 pt, Not Italic

Sl.no	Gene involved	Associated condition/brief description	Reference
1	transcription factor 3(TCF3)	Immunodeficiency/B-cell Defects and reduced serum immunoglobulin	Boast B <i>et al.</i> , 2023
2	Hedgehog interacting protein (HHIP)	Chronic obstructive pulmonary disease (COPD) Inhibited aerobic glycolysis and repressed cell proliferation.	Li Y <i>et al.</i> , 2021
3	The serine-/arginine-rich splicing factor 2 (SRSF2)	Squamous cell carcinoma. DNA damage with cell cycle alterations. clonal expansion.	Frost, Erin L <i>et al.</i> , 2024
4	The arginine methyltransferase (PRMT5)	Inflammatory bowel disease and colorectal cancer. reduced mucosal defense	Hernandez JE <i>et al.</i> , 2023.
5	Filamin C (FLNC)	Cardiomyopathies,myofibrillar myopathy,cardiac arrhythmias.	Job A J Verdonshot <i>et al.</i> , 2020
6	Paired like homeodomain 1(Pitx1)	Clubfoot, muscle hypoplasia	David M Alvarado <i>et al.</i> , 2010.

7	OTU deubiquitinase with linear linkage specificity (OTULIN)	Fasciitis and skin necrosis, auto-inflammation	Rob J W Arts <i>et al.</i> , 2023
8	Melanocortin 4 receptor (Mc4r)	Monogenic severe obesity, hyperphagia, hyperinsulinemia, and hyperglycemia.	Navneet Matharu <i>et al.</i> , 2019
9	Protein associated with LIN7 1, MAGUK p55 family member (PALS1)	Defects in renal functions, proteinuria and cyst formation.	Thomas Weide <i>et al.</i> , 2017
10	Nuclear receptor coactivator 5 (NCOA5)	Nonalcoholic fatty liver disease (NAFLD), insulin resistance. Non-alcoholic Steatohepatitis.	Yueqi Zhang <i>et al.</i> , 2024.

UNDER PEER REVIEW

Page 11: [1] Formatted	SDI 1084	2/5/2025 3:57:00 PM
------------------------	----------	---------------------

Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)

Page 11: [2] Formatted	SDI 1084	2/5/2025 3:57:00 PM
------------------------	----------	---------------------

Font: (Default) Times New Roman, 12 pt, Not Italic, (Asian) Chinese (Simplified, PRC)

Page 11: [3] Formatted	SDI 1084	2/5/2025 3:57:00 PM
------------------------	----------	---------------------

Default Paragraph Font, Font: (Default) Times New Roman, 12 pt, (Asian) Chinese (Simplified, PRC)