

A Review of the Botanical, Ethnomedicinal Uses, and Phytochemistry of *Dalbergia* (Fabaceae) Species

ABSTRACT

Secondary metabolites found in plants are recognized to have medicinal qualities and are utilized in both mainstream and traditional medicine. More than 80% of people on the planet use medicinal plants to treat a variety of illnesses, including fever, malaria, stomachaches, and ulcers. One significant family of flowering plants with therapeutic qualities is the Fabaceae, sometimes called the *Leguminoceae*. The *Fabaceae* subfamily *Faboideae* of the pea family includes the large genus *Dalbergia*, which has over 300 species. Its tiny to medium-sized trees, bushes, and lianas are found in tropical parts of Central and South America, Africa, Madagascar, and Southern Asia. This work is focused on the general review of the Biology/botanical aspect, ethnomedicinal properties and the phytochemistry of *Dalbergia Fabaceae*. Many species of *Dalbergia* are commonly utilized in diverse communities around the world to treat a variety of diseases. Phytoconstituents commonly found in *Dalbergia* include isoflavonoids, neoflavonoids, glycosides, cinnamylphenols, quinones, and furans.

Keywords: *Dalbergia*, *Fabaceae*, phytochemistry, Botanical, medicinal, Africa

INTRODUCTION

Plant-based secondary metabolites are used as food additives, bio-pesticides, flavorings, fragrances, colors, agrochemicals, and pharmaceuticals. Higher plants are the source of two-thirds of all new chemicals found annually (Al-Snafi, 2017). Plants have always been a part of medical research, from the beginning of human civilization to the current era of synthetic drugs. Around the world, herbal medicines are used in conventional disease treatment practices (Chaachouay and Zidane, 2024).

Nature contains a wide variety of medicinal plants that can help treat a range of illnesses (Saurabh et al., 2012). Despite the availability of several effective synthetic drugs, herbal medicine is a triumph of popular therapeutic diversity. The use of plant-based health products has increased dramatically in both developed and developing nations in recent years, which has resulted in a global explosion of herbal goods (Srivastav et al., 2011). Numerous products with high activity profiles have been produced as a result of the extraction and characterization of different active phyto-compounds from these green factories (Bijauliya et al., 2020). Natural

Commented [Ma1]: • Mentioning specific challenges or gaps (e.g., limited pharmacological studies) in the abstract could strengthen its relevance.

• Clarify if all *Dalbergia* species were studied or only a subset

Commented [Ma2]: Add – as medicinal qualities and

Commented [Ma3]: Emphasize the novelty of this review. What makes this compilation unique compared to previous studies on *Dalbergia* species?

Grammatical Issues:

• Long sentences reduce clarity. For example: "Around the world, herbal medicines are used in conventional disease treatment practices." This can be rephrased for conciseness.

• Avoid phrases like "despite the availability of several effective synthetic drugs..." without substantiating with references.

Commented [Ma4]: Verify claims like "two-thirds of all new chemicals annually are derived from higher plants." Are recent statistics available?

Commented [Ma5]: Italic

prescription drugs made from plant sources are becoming more and more well-liked worldwide due to their accessibility, affordability, non-toxicity, and lack of synthetic ingredients. Flavonoids and phenolic chemicals are found in many plants with antioxidant properties (Lakshmi *et al.*, 2014).

The genus *Dalbergia* contains 300 species, 37 of which are found in Africa and just a small number in Nigeria. According to Vasudeva *et al.* (2009), several *Dalbergia* species are important timber trees that are valued for their beautiful and often fragrant wood that is rich in aromatic oils. *Dalbergia* is a genus name honoring the 18th-century Swedish brothers Nils and Carl Dalberg (Sudhakar *et al.*, 2013). Plants in the genus *Dalbergia Fabaceae* are rich in isoflavones and neoflavones. Several *Dalbergia* species have been reported to possess medicinal properties (Radha *et al.*, 2015). They are used in the treatment of diarrhea, leucoderma, dyspepsia, dysentery, syphilis, gonorrhea, stomach ache, leprosy, eye illnesses, scabies, pain, and ringworm in traditional systems of medicine all throughout the world (Kazembe *et al.*, 2012; Saha *et al.*, 2013).

Of the 300 species in the genus *Dalbergia*, 37 are found in Africa, and just a few are found in Nigeria. ~~A number of~~ *Dalbergia* species are significant timber trees that are prized for their exquisite, frequently scented wood that is abundant in aromatic oils (Vasudeva *et al.*, 2009). The Swedish brothers Nils and Carl Dalberg, who lived in the 18th century, are honored by the genus name *Dalbergia* (Sudhakar *et al.*, 2013). Neoflavones and isoflavones are abundant in plants belonging to the genus *Dalbergia Fabaceae*. It has been claimed that a number of *Dalbergia* species have therapeutic qualities (Radha *et al.*, 2015).

Botanical Aspect of *Dalbergia Fabaceae*

Small to medium-sized trees, shrubs, and lianas in the broad genus *Dalbergia* are members of the pea family's *Fabaceae* subfamily *Faboideae*. The genus is widely distributed and indigenous to tropical parts of Madagascar, Africa, southern Asia, and Central and South America (Rahaingoson *et al.*, 2022). In the genus *Dalbergia*, the leaves of small to medium-sized trees, shrubs, climbers, and lianas are leathery, alternate, and compound. According to Chopra *et al.*, (1980), the leaflets are glabrous, alternating, and sometimes sub-opposite. Kidney-shaped seeds have a light brown color and are thin and flat. It is rather typical for the roots to nodulate. According to Saha *et al.*, (2013), a number of species are cultivated for their colored wood, which is used in the furniture business.

Commented [Ma6]: Grammatical Issues:

Ensure scientific names are italicized throughout this section, e.g., *Dalbergia sissoo*. Sentences like "With 730 genera and more than 19,400 species, the Leguminosae family also referred to as the Fabaceae is one of the largest families of flowering plants" could be streamlined.

Suggestions:

Add figures or tables summarizing species distribution or characteristics. The redundancy in describing the wood's aromatic properties should be addressed.

With 730 genera and more than 19,400 species, the *Leguminosae* family also referred to as the *Fabaceae* is one of the largest families of flowering plants (Stevens, 2008). The *Faboideae* subfamily, which has 274 ILDIS-accepted species worldwide, especially in tropical and subtropical regions, includes the genus *Dalbergia* (Mamta et al., 2014). Due to their beautiful and aromatic wood, most *Dalbergia* species make good timber trees (Chopra et al., 1980).

Dalbergia sissoo

This tree, which may grow to a height of around 25 meters, is medium to large, with a grey-yellow trunk, longitudinal crack, and downcast twig. Approximately five alternate leaflets on leathery, pinnately complex leaves. The petiole of the leaf stalk is about 15 cm long, and each leaflet is about 6 cm long with a finely pointed tip. Pale to pink, fragrant, nearly sessile, and up to 1.5 cm long, the flowers are grouped in dense clusters that are 5–10 cm long. Pods are light brown, oblong, flat, thin, and strap-like; they are 4–8 cm long and 1–8 cm wide. According to Sudhakar et al., (2013), they have one to five flat, bean-shaped seeds that are 8 to 10 mm long.

They have several suckers on the surface roots and a long taproot. Large upper branches support a spreading crown, while juvenile shoots are downy and drooping. Established stems have light brown to dark grey bark that is up to 2.5 cm (0.98 inches) thick and sheds in narrow strips. The plant fixes nitrogen from the atmosphere using bacteria that are present in nodules in the root system. The leaf litter that builds up and breaks down contributes to soil fertility by providing the soil with more nitrogen, potassium, iron, manganese, and organic carbon. Both seeds and root suckers are employed in the process of proliferation. Several parts of the plant are used to make medicines (Sudhakar et al., 2013).

Dalbergia saxatilis

It can grow as little as 2 to 3 meters in height, but it can also be a robust climbing shrub with stems up to 20 meters long (Burkill, 1985). The stem's diameter around the base can reach up to 5 cm, but it can potentially reach up to 15 cm. Some of the branches are very convoluted since they have been transformed into woody spine hooks. The wood and local medicinal uses are the main reasons for collecting the plant in the wild. It is sometimes used as a leafy vegetable and is occasionally sold in local markets (Tropical, 2022).

Spinosa Dalbergia

Tiny trees or shrubs with spines at the tips of their branches, they can grow up to 3–6 meters tall. Small, purplish-white, bisexual, and zygomorphic flowers with a campanulate calyx, a free ovary, an incurved style, a capitate stigma, a papilionaceous corolla, papilionaceous petals, monadelphous stamens, and minute anthers. One-seeded fruits with flat, falcate, brownish pods (PROTA, 2017).

Dalbergia latifolia

The tree features clusters of tiny white flowers, pinnately complex leaves, and grey bark that peels into long fibers. Young seedlings of *Dalbergia latifolia* have a strong taproot and hardly any secondary roots. At first, seedlings develop slowly. Nodules created by symbiosis with nitrogen-fixing bacteria have already been seen on seedling roots. Although growth rates are reported to vary widely, young trees also grow slowly. *Dalbergia latifolia* trees are often surrounded by many root suckers. Seedlings of *Dalbergia latifolia* may withstand some shade, but the plant needs moderate light. In open spaces, it has a tendency to develop twisted and branchy (PROTA, 2017).

Dalbergia melanoxylon

The bark surface is white to pale grey or greyish brown, thin, smooth, but can become rough and fissured or flaking, and the inner leaves are arranged in a spiral pattern. It is a deciduous, spiny shrub or small tree that can grow up to 12(–20) m tall, frequently multiple-stemmed, and heavily branched. The bole is typically short, branchless for up to 2(–3.5) m, often gnarled and fluted, reaching up to 50(–100) cm in diameter. A terminal or axillary panicle that is 2–12 cm long, has many flowers, is short-hairy to almost glabrous, and branches laxly. The calyx is campanulate, 2–3(–4) mm long, with lobes shorter than tube, bottom lobe longest, and upper lobe fused; the corolla is white, with obovate standard and clawed wings and a keel; the flowers are bisexual, papilionaceous, and 4–6 mm long, nearly sessile; and the stamens are typically 9, fused into a tube, but free in upper part; ovary superior (PROTA, 2017).

Ethnomedicinal Properties of *Dalbergia Fabaceae*

The ethnomedicinal usage of several *Dalbergia* genus members have been recognized and published. Numerous species are frequently used to cure a range of illnesses in varied populations throughout (Ninh, 2017). *Dalbergia sissoo* Roxb. Fabaceae is used as an antipyretic, expectorant, anthelmintic, aphrodisiac, and abortifacient. Additionally, it is recommended for syphilis, leucoderma, ulcers, and dyspepsia (Lakshmi et al., 2014).

Commented [Ma7]: Grammatical Issues:

- "The ethnomedicinal usage of several *Dalbergia* genus members have been recognized and published." This should be rephrased to "The ethnomedicinal uses of several *Dalbergia* species have been well-documented."
- Avoid unnecessary repetition. For instance, the uses of *Dalbergia sissoo* are described multiple times in similar terms.

Sissoo oil, which is made from the seeds, is used to treat scabies, skin burning, and blue irritation. To ease heavy menstruation, a finely ground paste of the leaves mixed with palm candy is administered in the morning. The leaves can be brewed three times a day to cure boils, acne, and painful micturition. The leaf juice is used to clear the urine of pus and treat jaundice. Breast edema can be lessened by using the leaf decoction. Powdered bark or leaf decoction is used to cure leprosy and gonorrhoea. When combined with milk and water, sissoo nectar can be used to treat fevers of any kind (Sudhakar et al., 2013).

In India, particularly in rural regions, traditional medicine practitioners use the leaves of *Dalbergia* species to treat diarrhoea (Mulla et al., 2011; Mohanty et al., 1998). The leaves of *Dalbergia saxatilis* are powdered and burned to repel insects (Namadina et al., 2019; Okwute et al., 2009). In traditional medicine, the leaves, bark, and roots of *Dalbergia saxatilis* are utilized to treat various ailments, including cough, smallpox, skin lesions, bronchial disorders, and toothaches (Saha et al., 2013; Hassan et al., 2016). *Dalbergia spinosa* Roxb is traditionally used to treat a range of conditions, such as fever, pain, skin infections, and urinary tract infections (Senthamarai et al., 2003). Additionally, to counteract the effects of alcohol, a teaspoon of *Dalbergia spinosa* Roxb root powder is dissolved in water (Senthamarai et al., 2003). The fruits of *Dalbergia* species are traditionally used as an antipyretic and tonic. The leaves and stem bark serve as febrifuges and anthelmintics (Naskar, 2004; Saha et al., 2013).

The stem bark of *Dalbergia latifolia* contains tannins, which are used in the treatment of leprosy, obesity, and intestinal worms (Kirtikar and Basu, 2005; Khalid et al., 2011). Additionally, the bark is utilized to alleviate body pain, while the whole plant is used to treat diarrhoea, dyspepsia, stomach ailments, as an anthelmintic, and as a bitter tonic (Ghani, 1999; Khare, 2007; Saha et al., 2013). In Senegal, the stem and root bark of *Dalbergia melanoxydon* are combined with baobab or tamarind fruits to relieve diarrhoea.

The smoke from burning its roots is inhaled to treat colds, bronchitis, and headaches. In Sudan, inhaling the smoke from burned stems helps ease rheumatism (Kamanula et al., 2018). An infusion of the roots is also used as an anthelmintic and aphrodisiac to treat gonorrhoea, stomachaches, and abdominal pain. The bark decoction of *Dalbergia spinosa* is applied to treat wounds, while leaf decoctions are used to alleviate joint pain, and leaf sap is used to treat inflammation of the mouth and throat (Bolza et al., 1972; Neuwinger et al., 2000; Saha et al., 2013).

Commented [Ma8]: Suggestions:

Include a summary table linking species, traditional uses, and cited regions.

Some medicinal claims (e.g., *Dalbergia saxatilis* for smallpox) require stronger supporting evidence.

Phytochemistry of *Dalbergia* **Fabaceae**

Several phytoconstituents have been identified from *Dalbergia* species, including isoflavonoids, neoflavonoids, glycosides, cinnamylphenols, quinones, and furans (Saha *et al.*, 2013). The separation of betulinic acid (**1**) from the root wood of *Dalbergia saxatilis* was reported by Okwute and Isyaka (2014). Fatty acid esters, sterols, and phenols have been found in the bark and leaves of *Dalbergia saxatilis* (Okwute *et al.*, 2009), 4-phenyl coumarin (**2**) like dalbergin (**3**) (Ahluwalia and Seshadri, 1957), methyl dalbergin (**4**), allyl quinone (**5**) like dalbergenone (**6**) (Ahluwalia and Seshadri, 1963), 4-phenylchromone (**7**) like dalbergichromene (**8**) (Mukerjee *et al.*, 1971; Lakshmi *et al.*, 2014), isodalbergin (**9**) (Mukerjee *et al.*, 1971; Saha *et al.*, 2013). Bijauliya *et al.* (2020) isolated quercetin (**10**) from *Dalbergia sissoo* ethanolic leaf extracts. Genstein (**11**), biochanin A (**12**), biochanin A 7-O-glucoside (**13**), pratensein (**14**), Caviunin 7-O- $[\beta$ -D-apiofuranosyl-(1 \rightarrow 6)- β -D-glucopyranoside] (**15**) was identified in the leaves of *Dalbergia sissoo* by Dixit *et al.*, (2012).

From the leaves of *Dalbergia sissoo*, Farag *et al.*, (2001) isolated Biochanin A 7-O- $[\beta$ -D-apiofuranosyl-(1 \rightarrow 5)- β -D-apiofuranosyl-(1 \rightarrow 6)- β -D-glucopyranoside] (**16**), biochanin A 7-O-apiosyl-(1 \rightarrow 6)-glucoside (**17**), 7-Methyltectorigenin 4'-O- $[\beta$ -D-apiofuranosyl-(1 \rightarrow 6)- β -D-glucopyranoside] (**18**), dalbergichromene (**8**), and nordalbergin (**19**) were isolated from the stem bark/heartwood of *Dalbergia sissoo* (Sharma *et al.*, 1980). Radha *et al.*, (2015) isolated 5-hydroxy-6-methoxy-3',4-methylenedioxy-7- $[(6$ -O-D-apiofuranosyl-D-glucopyranosyl)oxy] isoflavone (**20**), as well as dalspinin (**21**), dalspinosin (**22**), and caviunin (**23**), from the ethanol extract of *Dalbergia spinosa* roots. The leaves and stem bark of *Dalbergia spinosa* were found to contain 7-O-Methyltectorigenin 4'-O-galactoside (**24**), prunasin (**25**), and Prunetin 4'-O-galactoside (**26**) according to Narayanan *et al.*, (1988).

From the heartwood of *Dalbergia tonkinensis* Prain's methanol extract, Son *et al.*, (2017) identified carboxyethyl flavanones [(2S)-8-Carboxyethylnaringenin] (**27**) and [(2S)-6,8-Dicarboxyethylpinocembrin] (**28**). Buteaspermanol (**29**), homoferreirin (**30**), dalbergin (**3**), and medicarpin (**31**), Linalool (**32**), β -damascenone (**33**), β -ionone (**34**), geranyl acetone (**35**), and α -ionone (**36**) were discovered in the volatile oil of *Dalbergia frutescens* leaves (Mendes *et al.*, 2014), as well as formononetin (**37**) from the bark (Mendes *et al.*, 2014). From *Dalbergia latifolia* seeds, Shyam and Chibber (1978) extracted 12a-hydroxyrotenoid, dalbinol (**38**), and sisafolin coumarin (**39**). Rastogi and Mehrotra (1993), found dalbergichromene (**8**), lupeol (**40**),

Commented [Ma9]: Grammatical Issues:


Sentences listing phytochemicals are lengthy and difficult to read. Example: "The separation of betulinic acid (1) from the root wood of *Dalbergia saxatilis* was reported by Okwute and Isyaka (2014)." Consider breaking these into simpler phrases.

Repeated use of authors' names (e.g., Mukerjee *et al.*, 1971) in consecutive sentences can be consolidated.

Commented [Ma10]: Remove


latifolin (**41**), and dalbergin (**3**) in the tree's bark, as well as latinone (**42**), a substituted phenanthrene-1, 4-quinone (**43**), neoflavonoid (**44**) in the tree's heartwood (Thurlough *et al.*, 1981; Khalid *et al.*, 2011). The heartwood of *Dalbergia melanoxylon* was found to contain 2,3-Dihydrobenzofuran (**45**) and melanoxin (**46**), according to (Chalo *et al.*, 2023).

Structures




Commented [Ma11]: A detailed diagram or table of chemical structures would enhance this section.


Expand on the pharmacological significance of the identified phytochemical with references in table.




8




9




10




11




12




13




14




15




16




17




18




19




20




21




22




23















24




25



26





CONCLUSION

Several species of *Dalbergia* are known for their medicinal properties and are used in traditional medicine systems across the globe to treat a wide range of conditions, including diarrhea, leucoderma, dyspepsia, dysentery, syphilis, gonorrhoea, stomach aches, leprosy, eye disorders, scabies, pain, and ringworm. However, only a limited number of *Dalbergia* species have been subjected to pharmacological research to validate their traditional uses. Additionally, only a few compounds have been identified based on their phytochemical constituents. Despite the existence of over 300 *Dalbergia* species worldwide, only a small fraction has been studied in detail.

Commented [Ma12]: Grammatical Issues:

Avoid redundancy: "Despite the existence of over 300 *Dalbergia* species worldwide, only a small fraction has been studied in detail." The phrase "only a small fraction" already implies scarcity.

Suggestions:

Highlight future research opportunities, such as pharmacological validation of traditional uses or exploring under-researched species.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

REFERENCES

- Al-Snafi, A. E. (2017). Chemical constituents and pharmacological effects of *Dalbergia sissoo*-A review. *IOSR Journal of Pharmacy*, 7(2), 59-71.
- Ahluwalia V. K. & Seshadri T. R. (1957); Constituion of Dalbergin. Part II. *Journal of Chemical Society*, 970.
- Ahluwalia V. K. & Seshadri T. R. (1963); Dalbergenone from heartwood of *Dalbergia sissoo*. *Current science*, 32, 455.
- Sudhakar, K., Bharath, M., Tulasi, E. L. R., , & Eswaraiah, M. C. (2013). *Dalbergia sissoo* DC– An important medicinal plant. *Int J Res Pharm Chem*, 3(2), 384-388.
- Bijauliya, R. K. K., Kannoja, P., Mishra, P., & Pathak, G. K. (2020). Isolation and Structure Elucidation of Quercetin like Structure from *Dalbergia sissoo* (Fabaceae). *Journal of Drug Delivery and Therapeutics*, 10(3-s), 6-11.
- Burkill, H. M., (1985). Entry for *Desmodium scorpiurus* (Sw.) Desv. [family LEGUMINOSAE-PAPILIONOIDEAE]. In: *Useful Plants of West Tropical Africa*, Vol 3, Royal Botanic Gardens, Kew, UK
- Bolza E., Keating W. G. (1972). African timbers: the properties, uses and characteristics of 700 species. Division of Building Research, CSIRO, Melbourne, Australia, pp 710.
- Chaachouay, N., & Zidane, L. (2024). Plant-Derived Natural Products: A Source for Drug Discovery and Development. *Drugs and Drug Candidates*, 3(1), 184-207. <https://doi.org/10.3390/ddc3010011>
- Chalo DM, Franke K, Nchiozem-Ngnitedem VA, Kakudidi E, Origa-Oryem H, Namukobe J, Kloss F, Yenesew A, Wessjohann LA. (2023). Prenylated Isoflavanones with Antimicrobial Potential from the Root Bark of *Dalbergia melanoxylon*. *Metabolites*. 13(6):678. doi: 10.3390/metabo13060678. PMID: 37367838; PMCID: PMC10304121.
- Chibber S. S., Khera U. (1978). Dalbinol - a new 12a-hydroxyrotenoid from *Dalbergia latifolia* seeds. *Phytochemistry* 17, 1442-1443.

Commented [Ma13]:

Grammatical Issues:

Ensure proper formatting of references. For instance, italicize journal titles like *Journal of Chemical Society* and scientific names in citations.

Uniformly format author names (e.g., include initials after surnames).

Suggestions:

Cross-check all references for relevance and accuracy. Some references (e.g., *Kazembe et al., 2012*) seem overused

Commented [Ma14]: Italic

- Chopra, R. N., Nyer S. L. and Chopra I. C. (1980). Supplement to the glossary of Indian medicinal plants, CSIR, New Delhi. pp: 90.
- Dalbergia latifolia (PROTA). (2017). *Plant Use English*. [https://uses.plantnet-project.org/e/index.php?title=Dalbergia_latifolia_\(PROTA\)&oldid=325964](https://uses.plantnet-project.org/e/index.php?title=Dalbergia_latifolia_(PROTA)&oldid=325964).
- Dixit P., Chillara R., Khedgikar V., Gautam J., Kushwaha P., Kumar A., Singh D., Trivedi R., Maurya R. (2012). Constituents of *Dalbergia sissoo* Roxb. Leaves with osteogenic activity. *Bioorganic & Medicinal Chemistry Letters* 22, 890-897.
- Farag S. F., Ahmed A. S., Terashima K., Takaya Y., Niwa M. (2001). Isoflavonoid glycosides from *Dalbergiasissoo*. *Phytochemistry* 57, 1263-1268.
- Ghani A. (1998). Medicinal plants of Bangladesh: chemical constituents and uses. 1st ed. Asiatic Society of Bangladesh, pp. 155-156.
- Hassan F. I., Zezi A. U., Danmalam U. H., Yaro A. H. (2016); Effect of methanol leaf extract of *Dalbergia saxatilis* Hook.f (fabaceae) on renal function. *Avicenna J Phytomed.* 6 (6): 592-596.
- Kazembe T., Munyarari E., Charumbira I. (2012). Use of traditional herbal medicines to cure malaria. *Bulletin of Environment, Pharmacology and Life Sciences* 1, 63-85.
- Kamanula M, Munthali CR, Dziwapo A, Kamanula JF. (2018). Mineral and phytochemical composition of baobab (*Adansonia digitata L.*) root tubers from selected natural populations of Malawi. *Malawi Med J.* 30(4):250-255.
doi: 10.4314/mmj.v30i4.7. PMID: 31798803; PMCID: PMC6863415.
- Khalid M., Siddiqui H. H., Freed S. (2011). *In-vitro* assessment of antioxidant activity of *Dalbergia latifolia* barks extract against free radicals. *American-Eurasian Journal of Scientific Research*, 6, 172-177.
- Khare C. P. (2007). *Indian Medicinal Plants: an Illustrated Dictionary*. Springer-Verlag, New York, USA, pp. 199-201.
- Kirtikar K. R. and Basu B. D. (2005). *Indian medicinal plants*. International Book Distributor, Dehradun, India. pp: 824.
- Lakshmi, T. M., Radha, R., & Jayshree, N. (2014). In vitro antioxidant activity, total phenolic and total flavonoid content in extracts from the bark of *Dalbergia sissoo roxb*. *International Journal of Pharma Science Research*, 5(5), 226-231.

- Mamta B. , Archana S., Chhaya R., (2014). Dalbergia sissoo - An Important Medical Plant. *Journal of Medicinal Plants Studies*. 2(2): 76-82.
- Mohanty R. B., Dash S. K., Padhy S. N. (1998); Traditional phytotherapy for diarrhoeal diseases in India- a review, *Ethnobotany*. 10, 103-111
- Mendes, C. E., Flach, A., da Costa, L. A., Denardin, R. B., & Moura, N. F. D. (2014). Chemical composition and multivariate analysis of the volatile oil of *Dalbergia frutescens* (Vell.) Britton (Fabaceae). *Journal of the Brazilian Chemical Society*, 25, 1326-1330.
- Mukerjee S. K., Saroja T., Seshadri T. R. (1971). Dalbergichromene: A new neoflavonoid from stem bark and heartwood of *Dalbergia sissoo*. *Tetrahedron* 27, 799-803.
- Mulla WA, Chopade AR, Bhise SB, Burade KB, Khanwelkar CC. (2011). Evaluation of anti-diarrheal and in vitro antiprotozoal activities of extracts of leaves of *Alocasia indica*. *Pharm Biol*. 49(4):354-61. doi: 10.3109/13880209.2010.517211. PMID: 21428863.
- Namadina, M. M Aliyu, B.S, Hayatu, M., Aliko, A.A., Bako, A.T., Aminu, M.A., Hamisu, H., Kamal, R.M., Sunusi, U., Idris, A.M., Mukhtar, A.U., Ahmad, F.U., Bashir, A.A., Muttaka, A., Sanusi, H., Hafiz, S.S., Umar, M.K., Adamu, M.M., Sale, A.I., Karaye, S.I . (2019). Pharmacognostic and Acute Toxicity Study of *Dalbergia saxatilis* Stem Bark. *Dutse Journal of Pure and Applied Sciences (DUJOPAS)*, 5(2b):91-103.
- Narayanan V., Nacarajan N. S. (1988). Two isoflavone galactosides from *Dalbergia spinosa*. *Phytochemistry*, 27, 2364-2365.
- Ninh The S. (2017). A Review on the Medicinal Plant *Dalbergia odorifera* Species: Phytochemistry and Biological Activity. *Evid Based Complement Alternat Med*. ;2017:7142370. doi: 10.1155/2017/7142370.
- Naskar K. (2004). *Manual of Indian mangroves*, 1st ed. Daya Publishing House, Delhi, India.
- Neuwinger H. D. (2000). *African Traditional Medicine: a Dictionary of Plant Use and Applications*. Medpharm Scientific, Stuttgart, Germany, pp 589.
- Okwute S. K., Onyia R., Anene C., Amodu O. P. (2009). Protectant, insecticidal and antimicrobial potentials of *Dalbergia saxatilis* Hook f. (Fabaceae). *African Journal of Biotechnology* 8, 6556-6560.
- Okwute, S. K., & Isyaka, M. S. (2014). Betulinic acid from antimicrobial root wood extract of *Dalbergia saxatilis* Hook F. (Fabaceae). *European Journal of Medicinal Plants*, 4(6), 687-694.

- Radha, R., Vasantha, V. S., & Pitchumani, K. (2015). A new isoflavone apioglucoside from the roots of *Dalbergia spinosa*. *Natural Product Communications*, 10(11), 1934578X1501001138.
- Rahaingoson, F. R., Oyebanji, O., Stull, G. W., Zhang, R., & Yi, T.-S. (2022). A Dated Phylogeny of the Pantropical Genus *Dalbergia* L.f. (Leguminosae: Papilionoideae) and Its Implications for Historical Biogeography. *Agronomy*, 12(7), 1612. <https://doi.org/10.3390/agronomy12071612>
- Rastogi RP, Mehrotra BN. (1993). Compendium of Indian Medicinal Plants. Vol III, Publication and Information Directorate, CSIR, New Delhi. 351.
- Saha S., Anisuzzman M., Islam M. K., Mondal H., Talukder C. (2013). Antibacterial and cytotoxic potential of *Dalbergia spinosa* Roxb. Leaves. *International Journal of Pharmaceutical Sciences and Research* 4, 512-515.
- Senthamarai R., Umadevi G., Jai Ganesh K. P. (2003). Antimicrobial activity of root extracts of *Dalbergia spinosa* Roxb. *Ancient Science of Life* 22, 84-87.
- Son, N. T., Harada, K., Cuong, N. M., & Fukuyama, Y. (2017). Two new carboxyethylflavanones from the heartwood of *Dalbergia tonkinensis* and their antimicrobial activities. *Natural Product Communications*, 12(11), 1934578X1701201115.
- Stevens P. F. (2008). Angiosperm Phylogeny Website Version 9, June 2008.
- Sharma A., Chibber S. S., Chawala H. M. (1980). Isocaviunin-7-gentioisobioside, a new isoflavone glycoside from *Dalbergia sissoo*. *Phytochemistry* 19, 715.
- Saurabh A., Mishra A. S., Gupta S. (2012); A review on medicinal plant which may effective in the treatment of ulcer or which show antiulcer activities; *International journal of biopharmaceutical & toxicological research*. 2(1): 266-276.
- Srivastav S., Singh P. (2011); *Achyranthes aspera* - An important medicinal plant: A review; *J. Nat. Prod. Plant Resources*. 1 (1): 1-14.
- Shyam, S. and K. U. Chibber, (1978). Dalbinol-a new 12a-hydroxyrotenoid from *Dalbergia latifolia* seeds. *Phytochem*. 17: 1442-1443.
- Thurlough, O., Criodain M. O. S., Mary J. M. and Dervilla M. X. D., (1981). Latinone, a phenanthrene-1,4- quinone from *Dalbergia latifolia*. *Phytochem*. 20: 1089-1092.

Tropical Plants Database, Ken Fern. (2022). tropical.theferns.info.
<tropical.theferns.info/viewtropical.php?id=Dalbergia+saxatilis>

Vasudeva N., Vats M., Sharma S. K., Sardana S. (2009); Chemistry and biological activities of the genus *Dalbergia* - A review, Pharmacognosy Reviews. 3(6): 307-319.

General Recommendations:

- **Consistency:** Use a uniform style for headings, tables, and scientific names.
- **Figures & Tables:** Add graphical representations to summarize key findings.
- **Clarity:** Rewrite lengthy sentences and avoid redundancy across sections.

UNDER PEER REVIEW