

# A Survey to assess Banded Leaf and Sheath Blight disease severity in major maize growing areas of Andhra Pradesh

## ABSTRACT

*Rhizoctonia solani* f.sp. *sasakii*, the causal agent of maize banded leaf and sheath blight (BLSB) disease, is the most destructive soil borne pathogen capable of infecting maize at all growth stages starting from seedling to maturity. To estimate the severity of BLSB an extensive roving survey was conducted during *rabi*, 2018 in major maize growing areas of Andhra Pradesh i.e., West Godavari, East Godavari, Vizianagaram, Guntur and Kurnool to record the severity index of disease. The disease occurs all over the state with maximum disease severity was recorded in West Godavari district (61.35%) followed by East Godavari (58.05%), Vizianagaram (46.87%) and Guntur (25.85%) while least PDI was recorded in Kurnool district (23.02 %). Disease symptoms were noticed during the survey under field conditions.

**Key words:** Banded leaf and sheath blight (BLSB); Maize; Disease severity; Survey

## 1. INTRODUCTION

Maize (*Zea mays* L.) is one of the most important cereal crops in the world originated from Mexico, raises agricultural economy, serves as food, feed and raw material for industry. Maize contains approximately 72% starch, 10% protein, and 4% fat, supplying an energy density of 365 Kcal/ 100 gm (Nuss and Tanumihardjo, 2010). In India, maize is the third most important cereal crop after rice and wheat, grown in a wide range of environments extending from extreme semi-arid to sub-humid and humid regions. Since its introduction in India, maize has become an important crop as well as an integral component of the cropping systems of Indian agriculture. Despite very high yield potential of maize, one of the major deterrents to high grain yield is its sensitivity to several diseases. Nearly 112 diseases have been reported, of which, 65 are known to occur in India (Saxena, 2002). Seed rot, seedling blight, leaf spots, blights, downy mildews, stalk rots, banded leaf and sheath blight and smut are the most important diseases of maize crop (Hafiz, 1986). BLSB was caused by most destructive pathogen *Rhizoctonia solani* f.sp. *sasakii*, causes nearly 60% yield loss. A high positive correlation between disease index and premature death of plants that resulted in drastic reduction on grain yield as high as 97 per cent (Butchaiah, 1977).

BLSB was first reported from Sri Lanka in 1927 as Sclerotial disease (Bertus, 1927). In India, it was first reported in 1960 from Tarai region of Uttar Pradesh. The causal organism was *Hypochochilus sasakii* (Payak and Renfro, 1966) and it was appeared in epidemic form in 1972 in Mandi district of Himachal Pradesh (Thakur *et al.*, 1973). The pathogen characterized by the formation of sclerotial bodies in advanced stages of infection. Due to the shortage of water for rice cultivation in *rabi* season, farmers in most of the districts of Andhra Pradesh opting for maize. Rice fallow maize cultivation (zero tillage) has become popular in Krishna, Guntur, Prakasam, East and West Godavari districts due to better yields, higher productivity and monetary returns. Keeping in view the present study was conducted to assess the severity of this disease in major maize growing areas of Andhra Pradesh.

## 2. MATERIALS AND METHODS

A roving survey was conducted during *rabi* 2018-2019 to assess the disease severity of banded leaf and sheath blight in major maize growing areas of Andhra Pradesh based on maize crop statistics, five predominant maize growing districts i.e., Vizianagaram, West Godavari, East Godavari, Guntur and Kurnool were chosen for survey. Two mandals from each district, four villages from each mandal and two fields from each village were surveyed.

**Commented [A1]:** Replace it with- occurred in all the districts and

**Commented [A2]:** Include the conclusion of the study in brief.

**Commented [A3]:** Delete comma

**Commented [A4]:** Use is (present tense)

**Commented [A5]:** Add and before causes

**Commented [A6]:** Add was reported before that

**Commented [A7]:** Add full stop before starting the new line.

**Commented [A8]:** The scientific name should be italicized

**Commented [A9]:** Delete was

**Commented [A10]:** Add is after pathogen

**Commented [A11]:** Add are after Pradesh

**Commented [A12]:** Add these factors after keeping

**Commented [A13]:** Start a new line from based

In each field, disease severity was recorded in one square meter areas at five different plants, one each from four corners (excluding the border rows) and one from the centre. The severity score was assessed using the 1-5 scale given by Shekhar and Kumar (2012) (Table 1).

**Table 1** Disease severity scale for banded leaf and sheath blight in maize (Shekhar and Kumar, 2012)

Scale	Percentage of infection
1	Infection is on one leaf sheath, lesions are one or few, non-coalescent
2	Infection is on two to three leaf sheaths, lesions are few and non-coalescent on third leaf sheath from ground level.
3	Infection is not up to the ear shoot but on more than two leaf-sheaths
4	Infection is on all leaf sheaths up to the ear shoot but shank is not infected
5	Infection presents beyond the ear shoot; reduced ear size, husk leaves bleached and caked with or without sclerotial development and kernel formation absent or rudimentary.

Banded leaf and sheath blight infected plants having typical disease symptoms were used for isolation of the pathogen. Data on isolates collected, area of the crop, cropping system, hybrid/cultivar grown was collected during survey.

### 3. RESULTS AND DISCUSSION

#### 3.1 SURVEY

The severity of maize banded leaf and sheath blight was assessed by roving survey method during *rabi*, 2018-19 in major maize growing districts *i.e.*, West Godavari, East Godavari, Vizianagaram, Guntur and Kurnool of Andhra Pradesh were surveyed for BLSB disease severity (Plate 1).

Banded leaf and sheath blight was predominant in all the five districts of Andhra Pradesh surveyed. Per cent disease Index (PDI) ranged from 23.02 to 61.35% is a clear indication for its significance in causing potential yield losses. Among the districts surveyed the highest PDI was recorded in West Godavari district (61.35%) followed by East Godavari (58.05%), Vizianagaram (46.87%) and Guntur (25.85%) while least PDI was recorded in Kurnool district (23.02 %) (Table 2 and Fig. 1).


Per cent disease Index (PDI) varied widely in mandals of all the five districts. The highest PDI was recorded in Koyyalagudem mandal of West Godavari district (61.80%) followed by Dwaraka Tirumala mandal (60.90%) of West Godavari and the lowest PDI was recorded in Peddakadubur mandal (15.65%) of Kurnool district.

In West Godavari district, among the four villages surveyed the highest PDI was recorded in the Dwaraka Tirumala village (63.40%) while the lowest was recorded in the Narayanapuram village (58.40%). Similarly, from East Godavari district maximum PDI was recorded from Peddapuram village (61.30%) and minimum was recorded in Kattamuru village (56.00%). Manyapuripeta village of Vizianagaram district recorded higher PDI (48.80%) while the lowest (45.40%) was recorded in Jarajapuripeta village. In Guntur district maximum PDI was recorded from Budampadu village (36.20%)

**Commented [A14]:** Delete this part of the sentence

**Commented [A15]:** Add the standard formula for calculating PDI in Materials and Methods.

and no disease was recorded in Appikatla village. Highest PDI of 31.30 % was recorded in Peddakadubur village from Kurnool district and no disease was recorded in Chinnakadubur village.



**Plate 1. Pictorial representation of disease severity scale for maize BLSB**

**Table 2. Prevalence of maize banded leaf and sheath blight in major maize growing districts of Andhra Pradesh during *rabi* 2018-19.**

District name	Mandal name	Village name	Field code	PDI (%) Sample mean	PDI (%) Village mean	PDI (%) Mandal mean	PDI (%) District mean & Spacing	Crop area (acres)	Variety /Hybrid	Cropping system							
West Godavari	Koyyalagudem	Bayyanagudem	RWGB-1	57.60	63.20	61.80	61.35 (60x15 cm)	2	P-3396	Rice-Maize-Black gram							
			RWGB-2	68.80				1.2	Advanta PAC 751	Rice-Maize-Black gram							
		Koyyalagudem	RWGK-1	58.40	60.40			2	P-3344	Rice-Maize-Black gram							
			RWGK-2	62.40				1	P-3396	Rice-Maize-Black gram							
	Dwaraka Tirumala	Narayanapuram	RWGN-1	59.20	58.40	60.90		2.3	Kaveri gold	Rice-Maize-Black gram							
			RWGN-2	57.60				2	Advanta PAC 741	Rice-Maize-Green gram							
		Dwaraka Tirumala	RWGD-1	60.60	63.40			2	P-3396	Rice-Maize-Black gram							
			RWGD-2	66.20				1	Kaveri 50	Rice-Maize-Green gram							
			East Godavari	Peddapuram				Peddapuram	REGP-1	67.40	61.30	59.35	58.05 (60x15 cm)	2	Advanta PAC 741	Rice-Maize-Green gram	
									REGP-2	55.20				1.5	Kaveri 50	Rice-Maize-Green gram	
Katravulapalli	REGK-1	55.20			57.40	2	Advanta PAC 741	Rice-Maize-Green gram									
	REGK-2	59.60				1	P-3344	Rice-Maize-Black gram									
Jaggampeta	Jaggampeta	REGJ-1		63.20	57.50	57.00	2	Kaveri gold	Rice-Maize-Black gram								
		REGJ-2		51.80			1.5	P-3396	Rice-Maize-Black gram								
	Kattamuru	REGK-1		58.40	56.00		2	Kaveri gold	Rice-Maize-Green gram								
		REGK-2		53.60			2	P-3396	Rice-Maize-Black gram								
		Vizianagaram		Gurla			SR Peta	RVZS-1	44.80	45.40	47.10			46.87 (60x15 cm)	0.5	Advanta PAC 751	Maize-Maize
								RVZS-2	46.00						1	Advanta PAC 751	Maize-Maize
Manyapuripeta	RVZM-1		48.00		48.80	1.5	Kaveri 50	Maize-Maize									
	RVZM-2		49.60			1	Kaveri 50	Maize-Maize									
Nellimarla	Buradupeta		RVZB-1	44.80	46.00	46.65	0.5	P -3355	Maize-Maize								
			RVZB-2	47.20			0.5	Kaveri 50	Maize-Maize								
	Jarajapupeta		RVZJ-1	54.40	47.30		1	P -3355	Maize-Maize								
			RVZJ-2	40.20			1.5	P -3355	Maize-Maize								

District name	Mandal name	Village name	Field code	PDI (%) Sample mean	PDI (%) Village mean	PDI (%) Mandal mean	PDI (%) District mean & Spacing	Crop area (acres)	Variety /Hybrid	Cropping system		
Guntur rural	Bapatla	Bapatla	RGUB-1	66.40	33.20	16.60	25.85 (70x25 cm)	0.25	P-3344	Rice-Maize-Black gram		
			RGUB-2	0.00				0.5	P-3546	Rice-Maize-Black gram		
		Appikatla	RGUA-1	0.00	0.00			0.5	Syngenta NK 30	Rice-Maize-Black gram		
			RGUA-2	0.00				1	Syngenta NK 30	Rice-Maize-Black gram		
	Gunturrural	Guntur	RGUG-1	68.20	34.10	35.15		1.5	P-3344	Rice-Maize-Black gram		
			RGUG-2	0.00				0.5	Syngenta NK 7720	Rice-Maize-Black gram		
		Budampadu	RGUB-1	72.40	36.20			1.5	P-3344	Rice-Maize-Black gram		
			RGUB-2	0.00				1	P-3546	Rice-Maize-Black gram		
	Kurnool	Peddakadubur	Peddakadubur	RKLP-1	62.60	31.30		15.65	23.02 (70x25 cm)	0.5	P-3396	Maize-Maize-Groundnut
				RKLP-2	0.00					1	P-3546	Maize-Maize-Chick pea
Chinnakadubur			RKLC-1	0.00	0.00	1	P-3546			Maize-Maize-Groundnut		
			RKLC-2	0.00		0.5	P-3546			Maize-Maize-Groundnut		
C. Belagal		Rangapuram	RKLR-1	58.20	29.10	30.40	1.5	P-3396		Maize-Maize-Chick pea		
			RKLR-2	0.00			0.5	P-3546		Maize-Maize-Groundnut		
		C. Belagal	RKLC-1	63.40	31.70		1.5	P-3396		Maize-Maize-Groundnut		
			RKLC-2	0.00			1	P-3546		Maize-Maize-Chick pea		




Fig. 1. Prevalence of banded leaf and sheath blight in major maize growing districts of Andhra Pradesh

### 3.2 SYMPTOMATOLOGY

Banded leaf and sheath blight disease was initiated on 40 to 45 days old plants. Symptoms appeared on lower leaf sheaths as small water soaked, irregularly globular white lesions that extended to upper leaf sheaths as enlarged spots with irregular margins. Initially, spot colour varied from purple to brown tinge. They turned in to straw coloured necrotic patches giving banded appearance, which quickly spread to leaf blades and developed rapidly resulting in thin, papery leaf that resembled like snake skin casting. The pathogen caused damage to all the aerial parts except tassel (Plate 2 and 3).



Plate 2. Symptoms of BLSB on maize under field conditions: a-Rind spotting; b-Sclerotial bodies on rind of stalk; c-BLSB symptoms on leaves; d-Different degrees of lesion development on rind of stalk





**Plate 3. Symptoms of BLSB on maize under field conditions: e-BLSB symptoms on outermost husk leaves; f-Sclerotial bodies on cob; g-Shredding of husk leaves; h- Shredding of husk leaves; i- Stalk breakage; j-Severely infected plants with BLSB**

#### 4. DISCUSSION

The results are in accordance with the finding of Madhavi (2012), who reported banded leaf and sheath blight disease severity to range from 28.33 to 66.67% in Andhra Pradesh, Devi *et al.* (2018) reported average disease severity index of 41.30% to 66.50% maize BLSB in five districts of Himachal Pradesh similarly Rajput and Harlapur (2014) reported BLSB incidence was low (10.75%) to severe form (52.45%) among the four surveyed districts of northern Karnataka and further they reported narrow genetic makeup of the commercial hybrids, intensive cultivation of maize, season after season, year after year in addition to limited disease management practices and congenial weather conditions were suggested to be the reasons for higher disease incidence in Northern Karnataka.

Patra (2007) reported that in West Bengal, disease intensity was moderate to severe in male inbred line CML-163 and female inbred line CML-193-1 of hybrid maize (HQPM-1) at pre-flowering stage. Joye *et al.* (1990) reported that aerial blight of soybean caused by *R. solani* decreased disease with increased row to row spacing which might have not favoured secondary spread of the disease. Surveys conducted by Nishat (2009) from July to October 2006 revealed the prevalence of maize disease in Bahraiah, Uttar Pradesh, India and stated Banded leaf and sheath blight (BLSB) as predominant disease

**Commented [A16]:** It should be full stop

**Commented [A17]:** Start a new line from here

**Commented [A18]:** Delete the comma

**Commented [A19]:** Delete it



of maize. The incidence of BLSB was higher in crops sown in late *kharif*. Lee *et al.* (1989) revealed moderate occurrence of banded leaf and sheath blight of maize caused by *R. solani* during July and to the end of August as severe. Dense planting was reported to increase disease levels. Akhtar *et al.* (2009) reported maximum disease severity of 80.46 per cent was reported from Hisri Chauli block followed by Jirabar with severity of 50.30 per cent in Ranchi district of Jharkhand.

Commented [A20]: Delete it

The symptoms observed were in accordance to the description of Ahuja and Payak (1982) reported initially, disease initiated as irregularly globular to elongated lesions appear as water-soaked areas due to loss of chlorophyll later the affected areas bleached, become straw colored, necrotic and it spreads rapidly from the lower to upper sheaths rapid under favorable conditions. In an advanced stage of infection, husk leaves also show shredding, followed by the development of sclerotia. Lu *et al.* (2012) reported that crop damage caused due to loss of photosynthetic leaf area upon foliar infection and stalk rot lead to crop lodging.

Commented [A21]: Correct the sentence

## 5. CONCLUSION

The results obtained in the present study concluded that the maximum disease severity in Godavari districts (61.35% and 58.05%) and Vizianagaram district (46.87%) might be due to genotype and spacing (60 x 15 cm<sup>2</sup>) used. In regions where, wider spacing (70 x 25 cm<sup>2</sup>) was followed disease severity was low as seen in Guntur (25.85%) and Kurnool (23.02%). Genotypes growing in these regions were also different from the remaining three districts surveyed, which might have also contributed for limited disease severity. Besides, inoculum load in soil might have also resulted in higher severity.

Commented [A22]: Write revealed in place of concluded

## REFERENCE

- Ahuja, S.C and Payak, M.M. (1982). Symptoms and signs of banded leaf and sheathblight of maize. *Phyto parasitica*, **10**: 41-49.
- Akhtar, J., Jha, V.K., Kumar, A and Lal, H.C. (2009). Occurrence of banded leaf and sheath blight of maize in Jharkhand with reference to diversity in *Rhizoctonia solani*. *Asian Journal of Agricultural Sciences*. **1(2)**: 32-35.
- Bertus, L.S. (1927). A sclerotial disease of maize (*Zea mays* L.) due to *Rhizoctonia solani* Kuhn. *Year book*, Department of Agriculture, Ceylon. 44-46.
- Butchaiah, K. (1977). Studies on Banded Sclerotial diseases of maize. *M.Sc. Thesis*. G.B. Pant University of Agriculture and Technology, Pantnagar, India.
- Devi, B., Thakur, B.R and Singh, G. (2018). A Survey on Banded Leaf and Sheath Blight of Maize Incited by *Rhizoctonia solani* f. sp. *Sasakii* in Himachal Pradesh, *International Journal of Pure and Applied Biosciences*. **6 (5)**: 960-964.
- Hafiz, A. (1986). Plant Disease. *Pakistan Agricultural Research Council*, Islamabad. 93-102.
- Joye, G.F., Berggren, G.T and Berner, D.K. (1990). Effects of row spacing and within-row plant population on *Rhizoctonia* aerial blight of soybean and soybean yield. *Plant Disease*. **74**: 158-160.
- Lee, S.B., Kim, J.G., Han, M.S and Han, H.J. (1989). Studies on the occurrence of corn sheath blight caused by *Rhizoctonia solani*-influence of growth season and cultural environment on the



- occurrence of sheath blight disease. *Journal of the Korean Society of Grassland Science*. **9 (3)**: 174-178.
- Lu, Y.L., Xu, J., Yuan, Z.M., Hao, Z.F., Xie, C.X., Li, X.H., Shah, T., Lan, H., Zhang, S.H., Rong, T.Z and Xu, Y.B. (2012). Comparative LD mapping using single SNPs and haplo types identifies QTL for plant height and biomass as secondary traits of drought tolerance in maize. *Molecular Breeding*. **30**: 407-418.
- Madhavi, M. (2012). Variability in *Rhizoctonia solani* f. sp. *Sasakii* (kuhn) exner the incitant of banded leaf and sheath blight of maize (*Zea mays* L.). *Ph.D. Thesis*. Acharya N. G. Ranga Agricultural University, Rajendranagar, Hyderabad.
- Nishat, A.M.T.P. (2009). Prevalence of maize diseases in Bahraich (U.P). *Annals of Plant Protection Sciences*. **17 (2)**: 512-513.
- Nuss, E.T and Tanumihardjo, S.A. (2010). Maize: a paramount stable crop in the context of global nutrition. *Comprehensive Reviews in Food Science Food Safety*. **9**: 417-436.
- Patra, D. K. (2007). Occurrence of banded leaf and sheath blight diseases of maize in West Bengal. *Journal of Mycopathology Research*. **45 (1)**: 137-138.
- Payak, M.M and Renfro, B.L. (1966). Diseases of maize new to India. *Indian Phytopathology*. **3**: 14-18.
- Rajput, L.S and Harlapur, S.I. (2014). Studies on banded leaf and sheath blight of maize caused by *Rhizoctonia solani* f.sp. *sasakii* Exner. *Karnataka Journal of Agricultural Sciences*. **27(1)**:82-84.
- Saxena, S.C. (2002). Bio-Intensive Integrated Disease Management of Banded Leaf & Sheath Blight of Maize. In *Proceedings of the 8th Asian Regional Maize Workshop*, Bangkok, Thailand. 380-390.
- Shekhar, M and Kumar, S. (2012). Inoculation methods and disease rating scales for maize diseases. *Directorate of Maize Research*. Indian Council of Agricultural Research. New Delhi. 8-11.
- Thakur, M.S., Sharma, S.L and Munjai, R.L. (1973). Correlation studies between incidence of banded sclerotial disease and ear yield in maize. *Indian Journal of Mycology and Plant Pathology*. **3**: 180-181.