

A Statistical Approach for Analysis of Trend Pattern of Pigeon Pea in India

ABSTRACT

The present paper deals with the analysis of trend pattern in area, production, and yield of pigeon pea in India. The analysis is carried out using secondary time series data on area, production, and yield of pigeon pea pertaining to the period 2001-2023. The trend values have been computed on fitting well-known statistical models viz., linear model, exponential model, quadratic model, and cubic model. Moreover, the accuracy of the concerned fitted models have been evaluated using various statistical measures viz., coefficient of determination (R^2), root mean square error (RMSE) and relative mean absolute percentage error (RMAPE). The results of the investigation reveal that the concerned models fitted well for exploring the trend in area, production, and yield of pigeon pea. Moreover, the cubic model is slightly more precise as compared to the other fitted models, and hence it could be utilized for forecasting the scenario of pigeon pea in India. The findings of the study provide useful insights towards policy formulation regarding enhancement of pigeon pea production for meeting global food demand and nutritional security.

Comment [AG1]: Rewrite the abstract according to the journal format...

Keywords: *Linear model; exponential model; quadratic model; cubic model; coefficient of determination; root mean square error; relative mean absolute percentage error.*

1. INTRODUCTION

Pulses are essential pillars of the human diet, serving as a vital source of protein, particularly for vegetarians. In cooked form, pulses offer a rich blend of essential amino acids, dietary fibers, vitamins, and minerals.

India is the largest producer and consumer of pulses in the world (Source: Directorate of Pulses Development, Govt. of India [1]). In India during 2023-2024, the area under pulses was 27.51 million hectares with a production of 24.25 million tons, reporting a yield of 881 kg/hectare (Source: Directorate of Economics & Statistics, Govt. of India [2]). The most commonly grown pulses in India include chickpea, pigeon pea, urdbean, mungbean, and lentil.

Comment [AG2]: Add either number or reference for citation..

Pigeon pea (*Cajanus cajan* L.), natively known as Arhar in India, belongs to Fabaceae family, and is a short-lived perennial legume. It holds a significant place in the Indian diet. Primarily grown in semi-arid and tropical regions, pigeon pea is highly valued for its adaptability to diverse climatic conditions, drought tolerance, and capacity to improve soil fertility through nitrogen fixation. As a staple in Indian cuisine, particularly in the form of split grains (*dal*), it serves as an affordable and high-quality protein source, which is essential for the predominantly vegetarian population. India is the world's leading producer of pigeon pea, making it a centrepiece for the nation's food security and agricultural sustainability (Source: Directorate of Pulses Development, Govt. of India [1]). The major pigeon pea growing states in India include Maharashtra, Karnataka, Uttar Pradesh, Gujarat, Jharkhand and Telangana. During the year 2023-24, India reported production of 3.42 million tons of pigeon pea under 4.13 million hectares of area, witnessing a yield of 827 kg/hectare (Source: Directorate of Economics & Statistics, Govt. of India, [2]).

In recent years, several scientists and researchers have conducted studies on pulses. Kim *et al.* [3] conducted a systematic review and meta-analysis of randomized controlled trials to summarize and quantify the effects of dietary pulse consumption on body weight, waist circumference, and body fat. Gurusamy *et al.* [4] demonstrated the aspects of pulse grains for healthy foods that can tackle protein energy malnutrition and the management of diseases. Kumar *et al.* [5] investigated anti-nutritional compounds present in different pulses including their fractions, significance, and beneficial and adverse effects on human health. Yousif *et al.* [6] quantified and compared the level of mechanization across selected crops, including pulses, within the Gezira scheme, Sudan. The data about production practices, were collected from the farmers and engineers. Moreover, the level of mechanization was calculated for two scenarios viz., traditional and improved farming systems.

In the present scenario, the time series analysis of pulses is of utmost importance for policy formulation regarding food security and meeting the global nutritional demand. Considering the given fact, several scientists and researchers conducted statistical analysis for pulses in various geographical regions. Sharma *et al.* [7] examined the growth and trend of pulse production in India using time series data on area, production, yield, and trade pertaining to the period 1980-81 to 2008-09. Sharma *et al.* [8] conducted detailed investigation of various regions of Uttar Pradesh for analysis of growth rates and instability of pulse crops for two different periods viz., before and after the launch of technological missions on pulse production in the country. The selected pulse crops were arhar, pea, gram, and lentil. Vishwajith *et al.* [9] forecasted the production scenario of pulses in major pulses growing states of India using Autoregressive Integrated Moving Average (ARIMA) and Generalised Autoregressive Conditional Heteroscedastic (GARCH) models. Vani and Mishra [10] investigated the impact of irrigation on pulses production in India using regression and linear decomposition analysis. Ray and Bhattacharyya [11] investigated the trend behavior of pulses production, productivity, and net availability of India. The time series models viz., linear, quadratic, exponential, logarithmic, power, ARIMA, and ARIMAX models were used in the analysis. Akah *et al.* [12] evaluated pulse production, consumption and utilization in Nigeria, and investigated trend and growth rates in area, production and yield of cowpea in Nigeria and other major regions during the period 1996-98 to 2016-2018. Mishra *et al.* [13] analyzed the trend in production of total pulses in India using autoregressive integrated moving average (ARIMA) model on the basis of time series data pertaining to the period 1961 to 2019. Biam *et al.* [14] conducted trend and decomposition analysis of pulses production in North-East India across four periods viz., Phase I (1978-79 to 1991-92); Phase II (1992-93 to 2005-06), Phase III (2006-07 to 2019-20) and Pool (1978- 79 to 2019-2020). Balai *et al.* [15] reported the growth, decomposition and instability in area, production and productivity of rabi pulse crops viz., gram and lentil in Madhya Pradesh. Kumar *et al.* [16] carried out the comparative assessment of instability and growth rate in area, production, and yield of major pulses (viz., chickpea, pigeon pea, urdbean, and mungbean) in India using secondary time series data pertaining to the period 2000-2021. Prioty *et al.* [17] examined growth and instability in area, production, and yield of minor pulses in Bangladesh using secondary time series data pertaining to the period 1981-2020. Some other noteworthy contributions towards statistical modeling and time series analysis of crops, other than pulses, have been made by Kumar and Menon [18], Rana and Kumar [19], Kumar *et al.* [20], and Prakash *et al.* [21].

The objective of the present paper is to analyze the trend pattern in area, production, and yield of pigeon pea in India. The analysis is carried out by fitting well-known statistical models (viz., linear, exponential, quadratic, and cubic models) to the time series data on pigeon pea pertaining to the period 2001-2023, and estimating the trend values. The accuracy of the concerned fitted models have been evaluated using coefficient of determination (R^2), root mean square error (RMSE) and relative mean absolute percentage error (RMAPE).

2. MATERIALS AND METHODS

In the present paper, the secondary time series data on pigeon pea pertaining to the period 2001-2023 is utilized for the analysis of trend pattern in area, production, and yield of pigeon pea in India. The time series data have been obtained from the records of Directorate of Economics & Statistics, DAC&FW, Govt. of India.

In order to analyze the trend pattern in area, production, and yield of pigeon pea, the trend values are obtained by fitting linear, exponential, quadratic, and cubic models to the time series data as follows:

(a) Linear Model:

$$y_t = a + bt \dots (1)$$

where y_t denotes the observed time series value of area, production, or yield (as the case may be) of pigeon pea at time t .

The values of constants ' a ' and ' b ' are obtained by using the principle of least squares on solving the following normal equations:

$$\sum y_t = na + b \sum t \dots (2)$$

$$\sum ty_t = a \sum t + b \sum t^2 \dots (3)$$

where ' n ' represents the number of observed values.

(b) Exponential Model:

$$y_t = ae^{bt} \dots (4)$$

Taking natural log on both sides of above equation, we have

$$\log_e y_t = \log_e a + bt \log_e e$$

$$\text{i.e., } Y_t = A + bt \dots (5)$$

$$\text{where } Y_t = \log_e y_t, A = \log_e a, \text{ and } \log_e e = 1$$

The normal equations for estimating the values of 'A' and 'b' are as follows:

$$\sum Y_t = nA + b \sum t \dots (6)$$

$$\sum tY_t = A \sum t + b \sum t^2 \dots (7)$$

Finally, the value of 'a' is obtained on using

$$a = \text{antilog}(A)$$

(c) Quadratic Model:

$$y_t = a + bt + ct^2 \dots (8)$$

The values of constants 'a', 'b', and 'c' are obtained on using the principles of least squares as follows:

$$\sum y_t = na + b \sum t + c \sum t^2 \dots (9)$$

$$\sum ty_t = a \sum t + b \sum t^2 + c \sum t^3 \dots (10)$$

$$\sum t^2 y_t = a \sum t^2 + b \sum t^3 + c \sum t^4 \dots (11)$$

(d) Cubic Model:

$$y_t = a + bt + ct^2 + dt^3 \dots (12)$$

The values of constants 'a', 'b', 'c' and 'd' are obtained on solving the following normal equations.

$$\sum y_t = na + b \sum t + c \sum t^2 + d \sum t^3 \dots (13)$$

$$\sum ty_t = a \sum t + b \sum t^2 + c \sum t^3 + d \sum t^4 \dots (14)$$

$$\sum t^2 y_t = a \sum t^2 + b \sum t^3 + c \sum t^4 + d \sum t^5 \dots (15)$$

$$\sum t^3 y_t = a \sum t^3 + b \sum t^4 + c \sum t^5 + d \sum t^6 \dots (16)$$

3. RESULTS AND DISCUSSION

The secondary time series data on area, production, and yield of pigeon pea in India is presented in Table 1. The trend values are obtained on fitting linear, exponential, quadratic and cubic models to the data on area, production, and yield, and the findings are depicted in Tables 2, 3 and 4, respectively. Moreover, the fitted model equations for area, production, and yield of pigeon pea are presented in Table 5.

Table 1. Time series data on area, production, and yield of pigeon pea in India

Year	Area (in Million Hectares)	Production (in Million Tons)	Yield (in Tons / Hectare)
2001	3.33	2.26	0.68
2002	3.36	2.19	0.65
2003	3.52	2.36	0.67
2004	3.52	2.35	0.67
2005	3.58	2.74	0.77
2006	3.56	2.31	0.65
2007	3.73	3.08	0.83
2008	3.38	2.27	0.67
2009	3.47	2.46	0.71
2010	4.37	2.86	0.65
2011	4.01	2.65	0.66
2012	3.89	3.02	0.78
2013	3.90	3.17	0.81
2014	3.85	2.81	0.73
2015	3.96	2.56	0.65
2016	5.34	4.87	0.91
2017	4.44	4.29	0.97
2018	4.55	3.32	0.73
2019	4.53	3.89	0.86
2020	4.72	4.32	0.92
2021	4.90	4.22	0.86
2022	4.07	3.31	0.81
2023	4.13	3.42	0.83

(Source: Directorate of Economics & Statistics, DAC&FW, Govt. of India)

Table 2. Trends values for area of pigeon pea

Year (t)	y_t	Trend Values			
		Linear	Exponential	Quadratic	Cubic

		Model (L_t)	Model (E_t)	Model (Q_t)	Model (C_t)
2001	3.33	3.33	3.35	3.21	3.45
2002	3.36	3.39	3.40	3.30	3.40
2003	3.52	3.45	3.46	3.39	3.38
2004	3.52	3.51	3.51	3.48	3.38
2005	3.58	3.57	3.57	3.57	3.42
2006	3.56	3.64	3.62	3.65	3.48
2007	3.73	3.70	3.68	3.73	3.55
2008	3.38	3.76	3.73	3.80	3.64
2009	3.47	3.82	3.79	3.88	3.74
2010	4.37	3.88	3.85	3.95	3.85
2011	4.01	3.94	3.91	4.01	3.96
2012	3.89	4.00	3.97	4.08	4.08
2013	3.90	4.07	4.03	4.14	4.19
2014	3.85	4.13	4.10	4.19	4.29
2015	3.96	4.19	4.16	4.25	4.38
2016	5.34	4.25	4.22	4.30	4.46
2017	4.44	4.31	4.29	4.34	4.52
2018	4.55	4.37	4.36	4.39	4.56
2019	4.53	4.44	4.42	4.43	4.58
2020	4.72	4.50	4.49	4.47	4.56
2021	4.90	4.56	4.56	4.50	4.52
2022	4.07	4.62	4.63	4.53	4.44
2023	4.13	4.68	4.70	4.56	4.32

Table 3. Trends values for production of pigeon pea

Year (t)	y_t	Trend Values
--------------	-------	--------------

		Linear Model (L_t)	Exponential Model (E_t)	Quadratic Model (Q_t)	Cubic Model (C_t)
2001	2.26	2.12	2.19	2.09	2.45
2002	2.19	2.21	2.26	2.19	2.34
2003	2.36	2.29	2.32	2.28	2.28
2004	2.35	2.38	2.39	2.38	2.27
2005	2.74	2.47	2.46	2.47	2.29
2006	2.31	2.56	2.53	2.56	2.34
2007	3.08	2.64	2.60	2.65	2.43
2008	2.27	2.73	2.67	2.74	2.53
2009	2.46	2.82	2.75	2.83	2.66
2010	2.86	2.90	2.83	2.92	2.80
2011	2.65	2.99	2.91	3.01	2.94
2012	3.02	3.08	2.99	3.09	3.09
2013	3.17	3.16	3.07	3.18	3.24
2014	2.81	3.25	3.16	3.27	3.39
2015	2.56	3.34	3.25	3.35	3.52
2016	4.87	3.42	3.35	3.43	3.64
2017	4.29	3.51	3.44	3.52	3.74
2018	3.32	3.60	3.54	3.60	3.82
2019	3.89	3.68	3.64	3.68	3.86
2020	4.32	3.77	3.74	3.76	3.87
2021	4.22	3.86	3.85	3.84	3.84
2022	3.31	3.94	3.96	3.92	3.77
2023	3.42	4.03	4.07	4.00	3.65

Table 4. Trends values for yield of pigeon pea

Year (t)	y_t	Trend Values			
		Linear Model (L_t)	Exponential Model (E_t)	Quadratic Model (Q_t)	Cubic Model (C_t)
2001	0.68	0.65	0.65	0.66	0.68
2002	0.65	0.66	0.66	0.67	0.68
2003	0.67	0.67	0.67	0.67	0.67
2004	0.67	0.68	0.68	0.68	0.67
2005	0.77	0.69	0.69	0.69	0.68
2006	0.65	0.70	0.70	0.70	0.68
2007	0.83	0.71	0.71	0.71	0.69
2008	0.67	0.72	0.72	0.72	0.70
2009	0.71	0.73	0.72	0.72	0.71
2010	0.65	0.74	0.73	0.73	0.72
2011	0.66	0.75	0.74	0.74	0.74
2012	0.78	0.76	0.75	0.75	0.75
2013	0.81	0.77	0.76	0.76	0.77
2014	0.73	0.78	0.77	0.77	0.78
2015	0.65	0.79	0.78	0.78	0.80
2016	0.91	0.80	0.79	0.79	0.81
2017	0.97	0.81	0.80	0.80	0.82
2018	0.73	0.82	0.81	0.81	0.83
2019	0.86	0.83	0.82	0.83	0.84
2020	0.92	0.84	0.83	0.84	0.85
2021	0.86	0.85	0.84	0.85	0.85
2022	0.81	0.86	0.85	0.86	0.85
2023	0.83	0.87	0.87	0.87	0.85

In Tables 2, 3 and 4, the term ' y_t ' denotes the observed value of area (in million hectares), production (in million tons), and yield (in tons / hectare) of pigeon pea for the year ' t ' ($t = 2001, 2002, \dots, 2023$). Moreover, ' L_t ' denotes the linear trend value of area, production, and yield of pigeon pea for the year ' t '. In a similar manner, ' E_t ' denotes the exponential trend value, ' Q_t ' denotes the quadratic trend value, and ' C_t ' denotes the cubic trend value.

The relative influence of the trend values on the observed values of area, production, and yield of pigeon pea are graphically demonstrated in Figs. 1 to 12.

Table 5. Models equations for area, production, and yield of pigeon pea in India

	Linear Model	Exponential Model	Quadratic Model	Cubic Model
Area	$y_{t'} = 4 + 0.06 t'$	$y_{t'} = 3.97e^{0.015 t'}$	$y_{t'} = 4.08 + 0.06 t' - 0.0016t'^2$	$y_{t'} = 4.08 + 0.11t' - 0.0016t'^2 - 0.0006t'^3$
Production	$y_{t'} = 3.08 + 0.09 t'$	$y_{t'} = 2.99e^{0.028 t'}$	$y_{t'} = 3.09 + 0.09 t' - 0.0004 t'^2$	$y_{t'} = 3.09 + 0.15 t' - 0.0004t'^2 - 0.0008t'^3$
Yield	$y_{t'} = 0.76 + 0.01 t'$	$y_{t'} = 0.75e^{0.013t'}$	$y_{t'} = 0.75 + 0.01 t' + 0.0001t'^2$	$y_{t'} = 0.75 + 0.01 t' + 0.0001t'^2 - (6 \times 10^{-5})t'^3$

(Note: $t' = t - 2012$)




Fig. 1 Linear trend values for area of pigeon pea




Fig. 2 Linear trend values for production of pigeon pea




Fig. 3 Linear trend values for yield of pigeon pea




Fig. 4 Exponential trend values for area of pigeon pea




Fig. 5 Exponential trend values for production of pigeon pea




Fig. 6 Exponential trend values for yield of pigeon pea




Fig. 7 Quadratic trend values for area of pigeon pea




Fig. 8 Quadratic trend values for production of pigeon pea




Fig. 9 Quadratic trend values for yield of pigeon pea




Fig. 10 Cubic trend values for area of pigeon pea




Fig. 11 Cubic trend values for production of pigeon pea




Fig. 12 Cubic trend values for yield of pigeon pea

The accuracy of the concerned fitted models have been evaluated using various statistical measures viz., coefficient of determination (R^2), root mean square error (RMSE) and relative mean absolute percentage error (RMAPE) using the following formulae:

$$R^2 = 1 - \frac{\sum_{t=1}^n (y_t - \hat{y}_t)^2}{\sum_{t=1}^n (y_t - \bar{y})^2}$$

$$RMSE = \sqrt{\frac{1}{n} \sum_{t=1}^n (y_t - \hat{y}_t)^2}$$

and

$$RMAPE = \frac{1}{n} \sum_{t=1}^n \left| \frac{y_t - \hat{y}_t}{y_t} \right| \times 100$$

Comment [AG3]: Give the equations and explain this in methodology part

where \bar{y} denotes the mean value of variable Y , i.e., area, production, or yield (as the case may be) of pigeon pea. Also, \hat{y}_t represents the trend value of variable Y , which is obtained on fitting the concerned models (such as linear model, exponential model, quadratic model, or cubic model) to the variable Y .

The values of R^2 , RMSE and RMAPE for the concerned fitted models on area, production, and yield of pigeon pea are computed and the findings are summarized in Table 6.

Table 6. Model evaluation for area, production, and yield of pigeon pea in India

	Model	R^2	RMSE	RMAPE
Area	Linear	0.586	0.343	5.583
	Exponential	0.625	0.347	5.608
	Quadratic	0.600	0.337	5.953
	Cubic	0.674	0.305	5.702
Production	Linear	0.574	0.495	11.415
	Exponential	0.633	0.501	11.017
	Quadratic	0.575	0.495	11.584
	Cubic	0.635	0.458	10.893
Yield	Linear	0.441	0.073	7.678
	Exponential	0.449	0.073	7.594
	Quadratic	0.444	0.072	7.627
	Cubic	0.465	0.071	7.088

The following results are obtained from Table 6:

- (i) The values of R^2 for the fitted models are greater than 0.5 for area and production of pigeon pea, whereas for the yield $R^2 < 0.5$ for the concerned fitted model.
- (ii) The cubic model reported highest R^2 in area, production, and yield of pigeon pea, as compared to the other models.
- (iii) The cubic model reported least RMSE in area, production, and yield of pigeon pea, as compared to the other models.
- (iv) In terms of production and yield, the cubic model exhibited least RMAPE as compared to the other fitted models. However, in terms of area, all the fitted models have nearly the same RMAPE values.

Hence, on considering the above mentioned points, it can be revealed that the concerned fitted models are appropriate for analysis of trend pattern in area, production, and yield of pigeon pea in India. Moreover, the cubic model seems to be more precise, as compared to the other models, for exploring the scenario of trend pattern in pigeon pea.

4. CONCLUSION

In the present paper, trend pattern analysis is carried out for area, production, and yield of pigeon pea in India using time series data pertaining to the period 2001-2023. The trend values have been estimated by fitting linear, exponential, quadratic, and cubic models to the concerned time series data on pigeon pea. Moreover, the accuracy of the fitted models have been measured using coefficient of determination (R^2), root mean square error (RMSE) and relative mean absolute percentage error (RMAPE).

The results of the analysis reveal that the concerned fitted models are appropriate for estimating the trend in area, production, and yield of pigeon pea. Moreover, on the basis of values of R^2 , RMSE, and RMAPE, it can be concluded that the cubic model is slightly more reliable as compared to the other fitted models for forecasting the scenario of pigeon pea in India.

The findings of the investigation provide useful insights towards policy formulation regarding enhancement of pigeon pea production for meeting global food demand and nutritional security. In order to boost the pigeon pea production, the potential farmers could be encouraged for its cultivation.

Disclaimer (Artificial Intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

REFERENCES

1. Anonymous (2024). Annual Report of Ministry of Agricultural & Farmers Welfare. *Directorate of Pulses Development, Government of India.*
2. Anonymous (2024). Agricultural Statistics at a Glance. *Directorate of Economics & Statistics, DAC&FW, Government of India.*
3. Kim, S. J., De Souza, R. J., Choo, V. L., Ha, V., Cozma, A. I., Chiavaroli, L. Mirrahimi, A., Mejia, S.B., Buono, M.D., Bernstein, A.M., Leiter, L.A., Kris-Etherton, P.M., Vuksan, V., Beyene, J., Kendall, C.W.C., Jenkins, D.J.A. and Sievenpiper, J.L.(2016). Effects of dietary pulse consumption on body weight: a

systematic review and meta-analysis of randomized controlled trials. *The American Journal Of Clinical Nutrition*, 103(5): 1213-1223.

4. Gurusamy, S., Vidhya, C.S., Khasherao, B.Y. and Shanmugam, A. (2022). Pulses for health and their varied ways of processing and consumption in India- A review. *Applied Food Research*, 2: 1-9.
5. Kumar, Y., Basu, S., Goswami, D., Devi, M., Shivhare, U.S. and Vishwakarma, R.K. (2022). Anti-nutritional compounds in pulses: Implications and alleviation methods. *Legume Science*, 4:1-13.
6. Yousif, L.A., Dahab, M. and Ahmed, S.B. (2025). Mechanization levels of some crops in the Gezira scheme, Sudan. *Asian Journal of Research in Agriculture and Forestry*, 11(1): 1-9.
7. Sharma, M.K., Sisodia, B.V.S. and Lal, K. (2013). Growth and trends of pulse production in India. *Journal of Food Legumes*, 26(1&2): 86-92.
8. Sharma, M.K., Sisodia, B.V.S. and Kumar, V. (2014). Pattern of growth and scenario of pulse production in Uttar Pradesh: A regional study. *International Journal of Innovative Social Science & Humanities Research*, 2 (1): 1-16.
9. Vishwajith, K.P., Dhekale, B.S., Sahu, P.K., Mishra, P. and Noman, M.D. (2014). Time series modeling and forecasting of pulses production in India. *Journal of Crop and Weed*, 10(2): 147-154.
10. Vani, G.K. and Mishra, P. (2019). Impact of irrigation on pulses production in India: A time-series study. *Legume Research*, 42(6): 806-811.
11. Ray, S. and Bhattacharyya, B. (2020). Time series modeling and forecasting on pulses production behavior of India. *Indian Journal of Ecology*, 47(4): 1140-1149.
12. Akah, N.P., Kunyanga, C.N., Okoth, M.W. and Njue, L.G. (2021). Pulse production, consumption and utilization in Nigeria within the regional and global context. *Sustainable Agriculture Research*, 10(2): 48-64.
13. Mishra, P., Yonar, A., Yonar, H., Kumari, B., Abotaleb, M., Das, S.S. and Patil, S.G. (2021). State of the art in total pulse production in major states of India using ARIMA techniques. *Current Research in Food Science*, 4: 800-806.

14. Biam, K.P., Singh, N.U., Gowda, C., Tripathi, A.K., Singh, S.B., Pual, P., Nivetina L., Amrutha T. and Dkhar, H. (2022). Pulses production in North-East India: trend and decomposition analysis. *Indian Journal of Hill Farming*, 35: 134-141.
15. Balai, H.K., Meena, P.C., Meena, M.L., Bairwa, K.C., Kumar, S., Meena, R.P. and Bairwa, S.K. (2024). Growth, decomposition and instability analysis of major rabi pulse crops in Madhya Pradesh, India. *Asian Journal of Agricultural Extension, Economics & Sociology*, 42(7): 79-88.
16. Kumar, M., Singh, G., Singh, S. and Mishra, A. (2024). Performance of the major pulses crop in India: growth and instability. *Asian Journal of Research in Crop Science*, 9(4): 348-357.
17. Prioty, J.K., Rahman, K.S. and Miah, M.A.M. (2024). Growth and instability analysis of minor pulses in Bangladesh. *Food Science and Engineering*, 5(1): 200-207.
18. Kumar, M. and Menon, S.V. (2022). Statistical modeling and trend analysis of jackfruit production in the districts of Kerala in India. *International Journal of Agriculture, Environment and Biotechnology*, 15(03): 745-752.
19. Rana, S.K. and Kumar, M. (2022). Growth rate and instability analysis of sugarcane in selected states of India. *International Journal of Agriculture, Environment and Biotechnology*, 15(04): 837-843.
20. Kumar, M., Prakash, G. and Rana, S.K. (2024). Statistical modeling for analysis of growth and trend pattern of wheat production in selected states of India. *Asian Journal of Research in Crop Science*, 9(1): 66-75.
21. Prakash, G., Kumar, M., Rana, S.K. and Gowda K.E., S. (2025). A statistical approach for assessment of growth rate and instability of wheat in selected states of India. *Journal of Modern Applied Statistical Methods*, 24(1): 76-89.