

## **Effect of Biochar as Substitute in Nursery Potting Mixture on Growth of Coffee Seedlings and Soil Properties**

### **ABSTRACT**

Biochar is a carbon rich product derived from agricultural waste biomasses through pyrolysis and its application in agriculture reported to enhance the soil fertility. Cherry husk is the waste biomass generated during the process of hulling the cherry coffee at the coffee curing factory. Published reports regarding biochar as a substitute mostly pertains to the replacement of peat in the growth medium. Reports regarding use of biochar as substitute in the nursery potting medium to grow coffee seedlings are not available. Therefore, the current study was aimed to assess the effect of biochar obtained from cherry husk as substitute in the nursery potting medium and its effect on growth of coffee seedlings and soil properties. The biochar from cherry husk was produced using an electric biomass pyrolizer set at 500<sup>0</sup>C and pyrolyzed for one hour. The biochar so produced was used as substitute in nursery potting medium to assess the growth of coffee seedlings and soil properties. The experiment consisted of four treatments: T<sub>1</sub>-jungle soil: farm yard manure: sand (control); T<sub>2</sub>-biochar: - farm yard manure: sand; T<sub>3</sub>-jungle soil: biochar: sand; T<sub>4</sub>-jungle soil: farm yard manure: biochar. The results indicated that growth of coffee seedlings grown in T<sub>1</sub> and T<sub>4</sub> treatments were statistically similar. The growth of coffee seedlings grown in T<sub>2</sub> and T<sub>3</sub> treatments were comparatively poor. The findings of the study clearly indicated that biochar can be a potential alternate to sand in the nursery growth medium to grow coffee seedling.

### **Revised Abstract:**

Biochar is a carbon-rich product derived from agricultural waste biomasses through pyrolysis, widely reported to enhance soil fertility. Coffee cherry husk, a byproduct of the hulling process in coffee curing factories, presents an opportunity for biochar production. While previous studies have explored biochar as a peat substitute in growth media, research on its role in nursery potting mixtures for coffee seedlings is lacking. This study aimed to

evaluate the effects of biochar derived from cherry husk on the growth of coffee seedlings and soil properties. Biochar was produced using an electric biomass pyrolyzer at 500°C for one hour and incorporated into the nursery potting medium. The experiment included four treatments: T1 (jungle soil: farmyard manure: sand, control), T2 (biochar: farmyard manure: sand), T3 (jungle soil: biochar: sand), and T4 (jungle soil: farmyard manure: biochar). Results showed that seedling growth in T1 and T4 was statistically similar, while growth in T2 and T3 was comparatively lower. These findings suggest that biochar can serve as a viable alternative to sand in nursery potting mixtures, potentially improving soil properties and promoting sustainable seedling production.

**Key words:** Coffee; Cherry husk; Biochar; Coffee seedlings.

---

## 1. INTRODUCTION

Coffee is believed to be originated in Ethiopia. Currently, coffee is grown over eighty countries in the world with Brazil, Vietnam and Colombia being the largest coffee producers accounting for over half of the world's coffee production. Although there are more than eighty coffee species under the genus *Coffea*, arabica (*Coffea arabica* L.) and robusta (*Coffea canephora* Pierre ex Froehner) are commonly grown *Coffea* species worldwide on commercial scale. In India, coffee is grown to an extent of 4.70 lakh hectares (Arabica-2.42 lakh hectares; Robusta-2.28 lakh hectares) with the total production of 3.43 lakh metric tonne of green coffee bean in 2022-23 crop year (Anonymous, 2024).

Coffee being the second largest traded commodity in the world market, it generates copious amount of solid waste biomasses. According to International Coffee Organization (ICO), global coffee industry generates about 39 million tons of biological waste biomasses each year (Anonymous, 2023). While in India, approximately 8 lakh tonnes of waste biomass are produced yearly from coffee processing (Anonymous 2024). Coffee processing generates various solid waste biomasses viz., pulp (fruit skin), parchment husk, cherry husk, coffee silver skin and spent coffee ground. Coffee pulp is a good starter material for the production of compost or vermi-compost. The other waste biomasses are transformed into fuel briquettes. In our earlier work, biochar was produced from cherry husk, parchment husk, uprooted diseased coffee stem and pepper stem waste using biomass pyrolyzer unit besides the physicochemical characterization of biochar obtained from these coffee farm waste biomasses (Gopinandhan et al., 2022).

In recent years, biochar is drawing attention worldwide due to its potential to improve soil fertility. Biochar is produced from the combustion of various agri-waste biomasses under limited or nil oxygen condition and considered as soil ameliorant. Biochar is rich in carbon content that are more stable in the soil (Effa and Otie, 2023). Due to the versatile physicochemical characteristics of biochar, its application in soil is reported to influence several attributes of soil and the changes brought out by the biochar in the edaphic environment reported to improve agricultural productivity (Sheng and Zhu, 2018; Dai et al., 2020). Further, owing to its unique specific surface area and cation exchange capacity, biochar exhibits remarkable efficiency in removing pollutants like antibiotics, dyes, heavy metals, herbicides and pesticides (Oliveira, 2017). Biochar has also been used as a management tool to control various plant diseases (Navdeep and Adesh, 2020) and plant parasitic nematode (Dwi and Niken, 2017).

The successful establishment coffee husbandry starts with the production of healthy and vigorous seedlings. Under Indian condition, raising of coffee seedlings involves two stages *viz.*, sowing of coffee seeds in the germination bed and transplanting of sprouted seedlings into polybag containing nursery growth medium and this stage is known as polybag (basket) nursery. The well-developed seedlings which are normally four to six to months old are planted in the field during August and September months. The composition of growth medium for the successful establishment of coffee seedlings normally consists of jungle soil, compost and sand in a judicious proportion. However, due to slow decline in the availability of jungle soil caused by gradual shrinking of forest cover, non-availability of good quality farm yard manure (or) compost and escalating of cost of sand are compelling to look for alternate growth medium.

Though several studies on biochar have been focussed on its application in the soil system of various crops, only limited reports are available about its application in coffee farming (Samuel et al., 2019; Herviyanti et al., 2020; Jessica et al., 2022; Leta et al., 2022; Sanchez-Reinoso et al., 2022; Sanchez-Reinoso et al., 2023<sup>ab</sup>). Furthermore, published reports regarding biochar as a substitute mostly pertains to replacement of peat substrate in the growth medium (Ostos et al., 2008; Li et al., 2009; Tian et al., 2012; Vaughn et al., 2013; Zhang et al., 2014; Dispenza et al., 2016; Nieto et al., 2016; Picca et al., 2023). There are no reports pertaining to use of biochar as substitute in the nursery potting mixture to grow coffee seedlings. Biochar being a low-cost and carbon rich natural product, the present study aimed

to assess the effect of biochar obtained from cherry husk, as substitute in the growth medium and its effect on growth of coffee seedlings and soil properties.

## 2. MATERIALS AND METHODS

The study was carried out in a poly house nursery available at the research farm in central coffee research institute (CCRI) located at Chikkamagaluru district in Karnataka state during 2023-24 season. The institute is situated in the southern hill zone of Karnataka state at 13°34 north latitude and 75°48 east longitudes with an elevation of 825 meter above mean sea level. In general, CCRI receives an annual average rainfall of 2,500 mm distributed over 4 to 5 months from June to September with a total annual rainy day ranging from 113 to 133 days. The temperature and relative humidity ranges from 18°C to 34°C and 60% to 90%, respectively.

The coffee cherry husk (CCH) was collected from a coffee curing factory at Chikkamagaluru district in Karnataka State. The biochar from CCH was prepared using a biomass pyrolyzer unit. The temperature of the furnace was ramped at a rate of 10°C in 60 minutes to the reach pyrolysis holding temperature (500°C). The pyrolysis holding temperature was then sustained for 120 minutes and the furnace was allowed to cool at an average rate of approximately 30°C in 60 minutes. Later, the biochar was grounded to pass through 2 mm test sieve (M/s. Techno Instruments Company, Bangalore, India) and the fine biochar powder was used for the nursery trails.

The preparation of nursery mixture/growth medium started with the sieving of sun-dried jungle soil and well-rotted farm yard manure (FYM). The fine biochar powder was mixed to the sieved jungle soil and FYM according to the treatment levels. The treatments included in the current study were control T<sub>1</sub> (growth medium consisting of jungle soil: farmyard manure: sand in 6:2:1 proportion), T<sub>2</sub> (growth medium with biochar: farmyard manure: sand in 6:2:1 proportion), T<sub>3</sub> (growth medium with jungle soil: biochar: sand in 6:2:1 proportion) and T<sub>4</sub> (growth medium with jungle soil: farm yard manure: biochar in 6:2:1 proportion). The biochar was mixed with other constituents of nursery mixture based on the bulk density of biochar (213 kg/m<sup>3</sup>). Accordingly, the percentage of biochar in T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> treatments was 14%, 10% and 1.8%, respectively.

The nursery mixture was filled up into the nursery bags (one liter capacity polybag of 22.5 x 15 cm size with 150-gauge thickness and eight to ten holes of 3 mm size at the bottom half of the polybag to facilitate drainage of excess water). The nursery mixture was

moistened with water and filled into nursery bags firmly. The bags were arranged within the rectangular frames of bamboo reapers (ten to twelve bags per row). These frames are held in position with bamboo (or) wooden props driven firmly into the ground at suitable distances. The experiment was carried out with five replications per treatment and fifty seedlings per replication.

The sprouted seedlings of *Coffea arabica* (variety no.13) and *Coffea canephora* (variety no.3) were transplanted from germination bed to polybags. The sprouted seedlings were gently lifted from the primary bed using a stick to avoid the roots from breaking. Then, the root portion of seedlings dipped in Carbendazim 50 WP (one gram/liter) to prevent fungal infestation followed by dipping the root portion in cow dung slurry to avoid desiccation of root tips. Prior to transplanting, the polybags were watered and a vertical hole about 5 cm deep was made at the center of the basket. Seedlings were planted into the hole after nipping off the tip of the tap root and then lifted slightly to open out the roots. Care was exercised to keep the shoot system of the seedlings just above soil surface. After transplanting, the seedlings were maintained following the standard nursery management practices.

The percent mortality of seedlings in each treatment was recorded by visual assessment. Three seedlings from each replication were sampled for analysis purpose on every two months up to six months. The shoot length was measured as the distance from the collar region to the apex of the plant. The root length was measured as the distance from the collar region to the end of the longest root. Then, the stem and root portions were separated for biomass estimation. The fresh weight of stem and root portions were determined. Then, the shoot and root portions were dried separately at 70<sup>0</sup>C until constant weight was achieved. A representative soil sample was collected from all the four treatments after four and six months of seedlings growth in the polybag nursery. The air-dried soil samples were analysed for pH, EC and percent organic carbon following standard protocols (Jackson, 1973; Walkley and Black, 1934).

The data were subjected to analysis of variance (ANOVA) according to LSD test to indicate statistically significant differences between variables following the Agress package (version 3.01 data entry module and version 7.01 ANOVA package for researchers).

### **3. RESULTS AND DISCUSSION**

The data on growth parameters viz., shoot length, root length, fresh shoot weight, fresh root weight, dry shoot biomass weight and dry root biomass weight recorded in the present investigation showed that the growth of coffee seedlings steadily increased as the period of growth extended regardless of treatments (growth mediums) and coffee varieties (Tables 1 a, b, c).

The data on above-ground biomass (shoot length, fresh shoot biomass and dry shoot biomass weights) indicated that in case of arabica, among the four growth mediums evaluated, the T<sub>4</sub> growth medium recorded statistically highest shoot length, fresh shoot weight and dry shoot weight in six months old seedling followed by T<sub>1</sub> growth medium. While in Robusta, the T<sub>4</sub> growth medium recorded the highest shoot length, fresh shoot weight and dry shoot biomass weight in six months old seedling followed by T<sub>1</sub> growth medium and these two treatments were statistically on par to each other. Regardless of coffee varieties, the lowest above-ground biomasses were witnessed in T<sub>2</sub> growth medium in six months old seedling. In the current study, the reduction of above ground biomass observed in T<sub>2</sub> and T<sub>3</sub> growth mediums was in good agreement with the report of Dispenza et al., (2016) who observed that increased level of biochar resulted in reduction of shoots production in ornamental *Euphorbia* plant.

The data on below ground biomass (root length, fresh root biomass and dry root biomass weights) showed that in case of arabica, of the four growth medium studied, the T<sub>4</sub> growth medium recorded the highest root length and fresh root weight in six months old seedling followed by T<sub>1</sub> growth medium and these two treatments were statistically similar. While, the T<sub>4</sub> growth medium recorded statistically highest dry root weight followed by T<sub>1</sub> growth medium. In Robusta, the T<sub>4</sub> growth medium recorded the highest root length and dry root weight in six months old seedling followed by T<sub>1</sub> growth medium and these two treatments were statistically similar. Whilst, the T<sub>4</sub> growth medium recorded statistically highest fresh root weight followed by T<sub>1</sub> growth medium. Irrespective of coffee varieties, the lowest below ground biomasses were seen in T<sub>2</sub> growth medium in six months old seedling. The reduction of below ground biomass as seen in T<sub>2</sub> and T<sub>3</sub> growth mediums was similar to those of Samuel et al., (2019) who demonstrated that excess level of biochar increases the pH of growth medium which tends to cause copper and boron deficiencies. Boron deficiency results in stunted root growth and premature death of thin root tips in coffee.

**Table 1a:** Influence of biochar on growth parameters of arabica and robusta seedlings (2 months old seedlings)

Treatments	Arabica						Robusta					
	SL	RL	FSW	FRW	DSW	DRW	SL	RL	FSW	FRW	DSW	DRW
T <sub>1</sub>	14.12 <sup>a</sup> (±2.4)	8.67 <sup>b</sup> (±1.5)	2.32 <sup>a</sup> (±0.7)	0.21 <sup>b</sup> (±0.08)	0.59 <sup>a</sup> (±0.1)	0.06 (±0.01)	12.25 <sup>a</sup> (±1.8)	12.25 <sup>a</sup> (±2.3)	2.83 <sup>a</sup> (±0.8)	0.65 <sup>a</sup> (±0.1)	0.72 <sup>a</sup> (±0.13)	0.16 <sup>a</sup> (±0.06)
T <sub>2</sub>	5.09 <sup>b</sup> (±1.1)	4.23 <sup>c</sup> (±0.9)	0.48 <sup>c</sup> (±0.1)	0.16 <sup>b</sup> (±0.04)	0.16 <sup>c</sup> (±0.04)	0.05 (±0.02)	8.52 <sup>c</sup> (±1.7)	7.65 <sup>b</sup> (±1.2)	0.31 <sup>c</sup> (±0.07)	0.12 <sup>c</sup> (±0.04)	0.22 <sup>c</sup> (±0.08)	0.10 <sup>b</sup> (±0.03)
T <sub>3</sub>	4.48 <sup>b</sup> (±0.7)	4.51 <sup>c</sup> (±0.8)	0.47 <sup>c</sup> (±0.1)	0.10 <sup>bc</sup> (±0.03)	0.13 <sup>c</sup> (±0.02)	0.10 (±0.01)	7.83 <sup>c</sup> (±0.5)	6.41 <sup>b</sup> (±1.1)	0.35 <sup>c</sup> (±0.06)	0.13 <sup>c</sup> (±0.04)	0.23 <sup>c</sup> (±0.04)	0.09 <sup>b</sup> (±0.02)
T <sub>4</sub>	13.18 <sup>a</sup> (±1.8)	11.03 <sup>a</sup> (±2.2)	1.69 <sup>b</sup> (±0.8)	0.33 <sup>a</sup> (±0.09)	0.40 <sup>b</sup> (±0.07)	0.09 (±0.02)	9.88 <sup>b</sup> (±1.7)	12.44 <sup>a</sup> (±2.4)	1.31 <sup>b</sup> (±0.4)	0.24 <sup>b</sup> (±0.06)	0.57 <sup>b</sup> (±0.15)	0.14 <sup>a</sup> (±0.04)
CD (5%)	2.01	1.86	0.49	0.07	0.09	NS	2.40	2.01	0.52	0.12	0.12	0.03

**Table 1b:** Influence of biochar on growth parameters of arabica and robusta seedlings (4 months old seedlings)

Treatments	Arabica						Robusta					
	SL	RL	FSW	FRW	DSW	DRW	SL	RL	FSW	FRW	DSW	DRW
T <sub>1</sub>	20.91 <sup>a</sup> (±2.8)	16.20 <sup>a</sup> (±2.6)	5.15 <sup>a</sup> (±1.7)	1.03 <sup>a</sup> (±0.3)	1.43 <sup>a</sup> (±0.4)	0.39 <sup>a</sup> (±0.12)	19.41 <sup>a</sup> (±2.8)	17.69 <sup>a</sup> (±2.61)	9.02 <sup>a</sup> (±1.68)	1.35 <sup>a</sup> (±0.27)	1.89 <sup>a</sup> (±0.42)	0.49 <sup>a</sup> (±0.13)
T <sub>2</sub>	7.67 <sup>b</sup> (±1.2)	8.97 <sup>b</sup> (±1.4)	1.69 <sup>b</sup> (±0.3)	0.29 <sup>c</sup> (±0.08)	0.52 <sup>c</sup> (±0.1)	0.14 <sup>c</sup> (±0.02)	8.89 <sup>c</sup> (±1.27)	9.48 <sup>b</sup> (±1.44)	1.64 <sup>c</sup> (±0.08)	0.63 <sup>b</sup> (±0.12)	0.49 <sup>c</sup> (±0.17)	0.16 <sup>b</sup> (±0.04)
T <sub>3</sub>	7.33 <sup>b</sup> (±0.9)	5.81 <sup>c</sup> (±1.6)	1.28 <sup>b</sup> (±0.2)	0.23 <sup>c</sup> (±0.1)	0.34 <sup>c</sup> (±0.08)	0.10 <sup>c</sup> (±0.02)	8.01 <sup>c</sup> (±1.9)	8.09 <sup>b</sup> (±1.6)	0.88 <sup>c</sup> (±0.2)	0.35 <sup>b</sup> (±0.1)	0.22 <sup>c</sup> (±0.08)	0.12 <sup>b</sup> (±0.03)
T <sub>4</sub>	19.86 <sup>a</sup> (±3.3)	17.41 <sup>a</sup> (±1.9)	4.20 <sup>a</sup> (±0.9)	0.58 <sup>b</sup> (±0.1)	1.11 <sup>b</sup> (±0.25)	0.30 <sup>b</sup> (±0.2)	15.86 <sup>b</sup> (±2.4)	17.39 <sup>a</sup> (±2.9)	5.59 <sup>b</sup> (±1.2)	1.10 <sup>a</sup> (±0.4)	1.30 <sup>b</sup> (±0.23)	0.35 <sup>a</sup> (±0.08)
CD (5%)	2.31	2.69	1.10	0.10	0.27	0.08	2.61	4.43	0.88	0.32	0.23	0.15

**Table 1c:** Influence of biochar on growth parameters of arabica and robusta seedlings (6 months old seedlings)

Treatments	Arabica						Robusta					
	SL	RL	FSW	FRW	DSW	DRW	SL	RL	FSW	FRW	DSW	DRW
T <sub>1</sub>	21.46 <sup>b</sup> (±4.7)	17.21 <sup>a</sup> (±3.9)	7.84 <sup>b</sup> (±2.9)	1.68 <sup>a</sup> (±0.05)	1.92 <sup>b</sup> (±0.65)	0.37 <sup>b</sup> (±0.03)	26.44 <sup>a</sup> (±4.45)	17.74 <sup>a</sup> (±2.61)	11.54 <sup>a</sup> (±2.58)	2.30 <sup>b</sup> (±0.42)	3.22 <sup>a</sup> (±1.05)	0.95 <sup>a</sup> (±0.19)
T <sub>2</sub>	7.01 <sup>c</sup> (±1.8)	6.19 <sup>b</sup> (±2.1)	1.22 <sup>c</sup> (±0.75)	0.34 <sup>b</sup> (±0.15)	0.30 <sup>c</sup> (±0.13)	0.11 <sup>c</sup> (±0.1)	8.65 <sup>b</sup> (±1.2)	12.52 <sup>b</sup> (±1.5)	0.61 <sup>c</sup> (±0.1)	0.28 <sup>c</sup> (±0.08)	0.21 <sup>b</sup> (±0.03)	0.15 <sup>b</sup> (±0.04)

T <sub>3</sub>	10.47 <sup>c</sup> (±1.8)	13.81 <sup>c</sup> (±216)	1.89 <sup>c</sup> (±0.34)	0.81 <sup>b</sup> (±0.09)	0.52 <sup>c</sup> (±0.07)	0.21 <sup>c</sup> (±0.04)	11.48 <sup>b</sup> (±1.8)	13.19 <sup>b</sup> (±2.1)	3.25 <sup>b</sup> (±0.9)	0.64 <sup>c</sup> (±0.1)	0.98 <sup>b</sup> (±0.25)	0.30 <sup>b</sup> (±0.06)
T <sub>4</sub>	29.72 <sup>a</sup> (±3.6)	19.92 <sup>a</sup> (±3.2)	11.18 <sup>a</sup> (±2.3)	1.84 <sup>a</sup> (±0.6)	2.84 <sup>a</sup> (±0.6)	0.67 <sup>a</sup> (±0.15)	27.45 <sup>a</sup> (±3.8)	19.67 <sup>a</sup> (±2.7)	12.69 <sup>a</sup> (±2.2)	2.84 <sup>a</sup> (±0.9)	3.56 <sup>a</sup> (±0.8)	0.96 <sup>a</sup> (±0.12)
CD (5%)	3.88	3.05	3.70	0.63	0.61	0.17	4.54	2.82	2.77	0.40	0.88	0.18

SL - Shoot length (cm); RL- Root length (cm); FSW-Fresh shoot weight (gm); FRW- Fresh root weight (gm); DSW-Dry shoot weight (gm); DRW-Dry root weight (gm)

Analysis of soil samples collected from all the four growth mediums after six months of seedling growth under nursery condition revealed that the soil sample from T<sub>2</sub> growth medium showed the highest pH values (arabica-8.3; robusta-7.6) while the soil sample from T<sub>1</sub> growth medium recorded the least pH values (arabica-5.6; robusta-5.7). Similar trend was seen even in case of electrical conductivity (EC) in both arabica and robusta coffee varieties. Regarding organic carbon (OC) content, the soil samples from T<sub>2</sub> growth medium showed the highest OC values (arabica-7.3%; robusta-6.3%). In the remaining growth mediums (T<sub>1</sub>, T<sub>3</sub>, T<sub>4</sub>), the OC content ranged from 2.4% to 4.8% in arabica variety and from 2.9% to 4.2% in robusta variety (Table 2). The data on pH, EC and OC recorded in the current study matches to the results of Gopinandhan [et al., \(2022\)](#) and Lehmann et al., (2011) who reported that addition of biochar makes the soil alkaline, exhibits higher EC value and contains comparatively higher amount of organic carbon.

**Table 2.** Influence of biochar on soil properties of nursery potting mixture

Treatments	Arabica			Robusta		
	pH	EC (dS/m)	Organic carbon ((%)	pH	EC (dS/m)	Organic carbon (%)
T <sub>1</sub>	5.6 ± 0.57	0.43 ± 0.06	2.4 ± 0.63	5.7 ± 0.60	0.38 ± 0.03	2.9 ± 0.40
T <sub>2</sub>	8.3 ± 0.9	1.14 ± 0.1	7.3 ± 0.8	7.6 ± 0.4	0.97 ± 0.1	6.3 ± 0.5
T <sub>3</sub>	7.7 ± 0.8	0.86 ± 0.07	4.8 ± 0.55	7.2 ± 0.4	0.71 ± 0.1	4.2 ± 0.4
T <sub>4</sub>	6.2 ± 0.67	0.67 ± 0.09	2.9 ± 0.58	6.4 ± 0.7	0.57 ± 0.02	3.5 ± 0.20



Several authors have reported that optimal addition of biochar in the growth medium increased the overall biomass of the coffee seedlings (Samuel et al., 2019; Herviyanti et al., 2020; Jessica et al., 2022; Leta et al., 2022) and also in other crops such as soybean (Zamriyetti and Mufriah, 2020) and asparagus (Elmer and Pignatello, 2011). Coffee grows better at a pH value of between 5.5 and 6. In the present study, the poor performance of coffee seedlings in T<sub>2</sub> and T<sub>3</sub> growth mediums probably due to the presence of excess levels of biochar in the growth medium (T<sub>2</sub> -14%; T<sub>3</sub> -10%) which led to alkalisation of the growth mediums. Barotni et al., (2010) and Samuel et al., (2019) reported that that excess addition of biochar resulted in a general reduction of biomass due to changes in the physico-chemical properties of soil.

#### 4. CONCLUSION

The findings of current study clearly revealed that the overall growth performance of coffee seedlings was equally good in both T<sub>1</sub> growth medium (jungle soil: farmyard manure: sand in 6:2:1 proportion) and T<sub>4</sub> growth medium (jungle soil: farm yard manure: biochar in 6:2:1 proportion) when compared to T<sub>2</sub> growth medium (biochar: farmyard manure: sand in 6:2:1 proportion) and T<sub>3</sub> growth medium (jungle soil: biochar: sand in 6:2:1 proportion). The results of study suggests that jungle soil and farm yard manure (compost) are essential ingredients for the growth of coffee seedlings and sand can be replaced with the biochar. Biochar at the rate of 2% in the coffee nursery mixture is ideal dosage for the better performance of coffee seedlings. Previous reports along with the results generated in the current study clearly indicated that the excess addition biochar increases the pH of the growth medium and thus optimization of biochar application rate is paramount importance for the better performance of coffee seedlings.

#### REFERENCES

- Anonymous. (2023). International Coffee Organization. Historical data on the global coffee trade. London, England.
- Anonymous. (2024). Database on coffee. Coffee Board of India, Karnataka, India.
- Baronti, S., Alberti, G., Delle Vedove, G., Di Gennaro, F., Fellet, G., Genesio, L., Miglietta, F., Pere Sotti, A. & Vaccari, F. P. (2010). The biochar option to improve plant yields: first results from some field and pot experiments in Italy. *Italian Journal of Agronomy*, 5, 3-11.
- Dai, Y., Zheng, H., Jiang, Z. & Xing, B. (2020). Combined effects of biochar properties and soil conditions on plant growth: A meta-analysis. *Science Total Environment*, 713, 136635. <https://doi.org/10.1016/j.scitotenv.2020.136635>.

Dispenza, V., De Pasquale, C., Fascella, G., Mammano, M. M. & Alonzo, G. (2016). Use of biochar as peat substitute for growing substrates of *Euphorbia × lomi* potted plants. *Spanish Journal of Agriculture Research*, 14, 4 e0908. <http://dx.doi.org/10.5424/sjar/2016144-9082>.

Dwi, S. R. & Niken, P. S. (2017). Development of *Pratylenchus coffeae* in biochar applied soil, coffee roots and its effect on plant growth. *Pelita Perkebunan*, 33, 24-32.

Effa, E. B. & Otie, V. O. (2023). Biochar-A mechanism of soil amendment for agricultural productivity. *Global Journal of Agriculture Science*, 22, 147-152.

Elmer, W. H. & Pignatello, J. J. 2011. Effect of biochar amendments on mycorrhizal associations and *Fusarium* crown and root rot of asparagus in replant soils. *Plant Diseases*, 95, 960-966.

Gopinandhan, T N., Channabasamma. B. B., Sandeep. T. N., Chandrasekar. N., Shruthi, H. & Nagaraja, J. S. (2022). Production and physio-chemical characterization of biochar obtained from coffee processing wastes and crop residues of coffee plantation. *Asian Journal of Microbiology Biotechnology Environmental Science*, 24, 494-501.

Herviyanti, H., Maulana, A., Prima, S., Aprisal, A., Crisna, S. D. & Lita, A. L. (2020). Effect of biochar from young coconut waste to improve chemical properties of ultisols and growth coffee [*Coffea arabica* L.] plant seeds. *Earth and Environmental Science*, 497, 012038. doi:10.1088/1755-1315/497/1/012038.

Jackson, M L. (1973). Soil chemical analysis. Prentice Hall of India Pvt. Ltd. New Delhi

Jessica. E.C.C., Andrea, K.L.T., Holger, G.S., Galo, A.C.G. & Ángel, F.C.S. (2022). Growth and quality of arabica coffee plants with the application of biochar and biofertilizers in the nursery. *Chilean Journal of Agriculture Animal Science ex Agro-Ciencia*, 38, 3-14.

Lehmann, J., Rillig, M. C., Thies, J., Masiello, C. A., Hockaday. W. C. & Crowley, D. (2011). Biochar effects on soil biota-A review. *Soil Biology Biochemistry*, 43, 1812-1836.

Leta, A., Gezahegn, B. & Taye, K. (2022). Effects of pot sizes and biochar base media composition on nutrient uptake of coffee (*Coffea arabica* L.) seedlings in South Ethiopia. *World Journal of Agriculture and Soil Science*, 8, DOI: 10.33552/WJASS.2022.08.000678.

Li, Q., Chen, J., Caldwell, R.D. & Deng, M. (2009). Cowpeat as a substitute for peat in container substrates for foliage plant propagation. *Horticulture Technology*, 19, 340-345.

Navdeep, S. and Adesh, K. (2020). Plant disease management through biochar-A review. *International Journal of Current Microbiology Applied Science*, 11, 3499-3510.

Nieto, A., Gasco, G., Paz-Ferreiro, J., Fernandez, J.M., Plaza, C. & Mendez, A. (2016). The effect of pruning waste and biochar addition on brown peat based growing media properties. *Science Horticulture*, 199, 142- 148.

- Oliveira, F. R., Patel, A. K., Jaisi, D. P., Adhikari, S., Lu, H. & Khanal, S. K. (2017). Environmental application of biochar: Current status and perspectives. *Bioresource Technology*, 246, 110–122.
- Ostos, J. C., Lopez-Garrido, R., Murillo, J. M. & Lopez, R. (2008). Substitution of peat for municipal solid waste and sewage sludge based composts in nursery growing media: Effects on growth and nutrition of the native shrub *Pistacia lentiscus* L. *Bioresource Technology*, 99, 1793-1800.
- Picca, G., Goni-Urtiaga, A., Gomez-Ruano, C., Plaza, C. & Panettieri, M. (2023). Suitability of co-composted biochar with spent coffee grounds substrate for tomato (*Solanum lycopersicum*) fruiting stage. *Horticulture*, 9, 89, <https://doi.org/10.3390/horticulturae9010089>.
- Samuel, F. B., Ajebesone, F. N., Tata, P. N. & Tsi, E, A. (2019). Influence of biochar and poultry manure on weed infestation and growth of arabica coffee (*Coffea arabica*) seedlings. *Current Research in Agriculture Science*, 6: 9 -19.
- Sanchez-Reinoso, A. D., Lombardini, L. & Restrepo-Díaz, H. (2022). Physiological behaviour and nutritional status of coffee (*Coffea arabica* L. var. Castillo) trees in response to biochar application. *The Journal of Agriculture Science*, 160, 220–234.
- Sanchez-Reinoso, A.D., Avila-Pedraza, E.A., Leonardo, L. & Restrepo-Diaz, H. (2023<sup>a</sup>). The application of coffee pulp biochar improves the physical, chemical, and biological characteristics of soil for coffee cultivation. *Journal of Soil Science and Plant Nutrition* <https://doi.org/10.1007/s42729-023-01208-4>.
- Sánchez-Reinoso, A. D, Colmenares-Jaramillo, A., Lombardini, L. & Restrepo-Díaz, H. (2023<sup>b</sup>). Physiological response of Castillo el Tambo coffee plants to biochar and chemical fertilization applications. *Chilean Journal of Agriculture Research*, 83, 307-319.
- Sheng, Y. & Zhu, L. (2018). Biochar alters microbial community and carbon sequestration potential across different soil pH. *Science Total Environment*, 622, 1391–1399.
- Tian, Y., Sun, X., Li, S., Wang, H., Wang, L., Cao, J. & Zhang, L. (2012). Biochar made from green waste as peat substitute in growth media for *Calathea rotundifolia* cv. *Fasciata*. *Science Horticulture*, 143, 15-18.
- Vaughn, S. F., Kenar, J. A., Thompson, A. R. & Peterson, S. C. (2013). Comparison of biochars derived from wood pellets and pelletized wheat straw as replacements for peat in potting substrates. *Industrial Crops Production*, 51: 437-443.

Walkley, A. J. & Black, C. A. (1934). An examination of the method for determination of soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37, 29-38.

Zamriyetti, S. M. & Mufriah, D. (2020). Effects of type of biochar and dolomite-Coffee compost application on characteristic of soybean germination. International Conference on Agriculture, Environment and Food Security. *Earth and Environmental Science*, 454, 012177. doi:10.1088/1755-1315/454/1/012177.

Zhang, L., Sun, X., Tian, Y & Gong, X. (2014). Biochar and humic acid amendments improve the quality of composted green waste as a growth medium for the ornamental plant *Calathea insignis*. *Science Horticulture*, 176, 70-78.

\*\*\*\*\*