

Prevalence and antibiotic susceptibility pattern of bacteria isolated from Nigerian currency note in Enugu, Enugu State, Nigeria.

ABSTRACT

Background: Nigerian currency notes acts as fomites and thus can transmit pathogenic microorganisms from one person to another

Aim: This study was conducted to evaluate the prevalence and antibiotic susceptibility pattern of bacteria isolated from Nigerian currency note circulating in Enugu East local government area of Enugu.

Method: Isolation, characterization and antibiotic sensitivity tests were done using standard methods.

Result: A total of 632 bacteria strains comprising 13 different bacterial species were isolated and identified from the Nigerian currency notes. The respective isolation rates of the isolates were *Staphylococcus aureus*(74.3%), *E.coli* (31.6%), *Pseudomonas aeruginosa* (5.3%), *Bacillus*spp (28.6%), *Salmonella* spp (6.0%), *Klebsiella*spp(21.3%), *Streptococcus*spp(15.3%), *Acinetobacter*spp (3.3%), *Serratia marcescens* (3.6%), *Proteus mirabilis* (8.0%), *Enterococcus faecalis* (6.0%), *Citrobacter*sp(6.3%) and *Actinomycetes*sp(0.6%). The Gram positive and Gram negative bacteria showed resistance to cloxacilline (87%) and amoxicillin (84%). Further, 79.5% of all the bacterial isolates had an index >2 while 20.5% had a multi-antibiotic resistance index < 2.

Comment [WU1]: bacterial

Conclusion:Nigerian currency notes harbour bacteria. Many of these bacteria are multidrug resistant organisms.

Comment [WU2]: harbour

Key words: Prevalence, Bacteria, Antibiotic resistance, MARI.

INTRODUCTION

It has been well documented that bacteria is cosmopolitan in distribution. Money, which is one of the most commonly used commodities in life, provides a niche for the microbial community. They are used for the purpose of goods and services worldwide. The word "money" has its origin in Rome (Snehalatha *et al.*, 2016). These papers, most frequently passed from hand to hand can be contaminated with pathogenic bacteria by different ways during handling, thus increases the possibility of the transmission of potential pathogenic bacteria, it is fact that we buy day to day commodities transferring microorganisms from one location to another location, bacteria adherence the notes during coughing or sneezing, and by placement on dirty surface (Elsharief *et al.*, 2018).

In Nigeria, the currencies exist as naira and kobo and various denominations of the naira notes have been minted by the Central Bank of Nigeria (CBN). They are released to the public, through the commercial banks. Currently, there are eight denominations of the naira notes: ₦5, ₦10, ₦20, ₦50, ₦100, ₦200, ₦500 and ₦1000 notes. The ₦5, ₦10, ₦20, ₦50, ₦100 and ₦200 naira notes are the most common and are more involved in daily cash transactions. They are common especially among the populace while the ₦500 are used in corporate transactions (Moses *et al.*, 2018). In day to day transactions, money is handled by persons of varying health and hygienic standards and also stored under varying environmental and personal hygienic conditions. Poor currency handling practices are common globally. People often exhibit unhygienic habits like store money in unsanitary places (socks, bras, shoes) or handle it carelessly, introducing germs. Unhygienic habits like licking fingers while counting money or handling bills with dirty hands significantly increase contamination (Mousa & Idress, 2023). Contaminated surfaces, dust, soil, and even the handler's own body can contribute to the spread of microbes on currency notes. These practices not only contaminate money but also increase the risk of infection for those who handle it (Awe *et al.*, 2010). This has been implicated in serious health hazards such as impairment of lung function. The contamination of the notes can be traced to dust, soil, water and microflora of the body of handlers (hand, skin, etc.). *Citrobacter* spp, *Salmonella* spp, *Shigella* spp, *Escherichia coli*, *Staphylococcus aureus* and *Pseudomonas aeruginosa* have been isolated from naira notes (Muhumuza *et al.*, 2018). Thus, the purpose of this study is to ascertain the degree of bacterial contamination and antibiotic resistance pattern of the bacteria isolated from Nigerian currency notes circulating in Enugu East Local Government Area of Enugu State.

Materials and Method

Materials

Study Area: This research project work was carried out in Enugu east local government of Enugu state.

Test sample: Nigerian naira (currency notes) (Figure 1)

Test Organisms: The test organisms were the bacteria isolated and they include: *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Salmonella spp*, *Klebsiella aerogenes*, *Streptococcus spp*, *Acinetobacter spp*, *Serratia marcescens*, *Proteus mirabilis*, *Enterococcus faecalis*, *Citrobacter spp* and *Actinomycetesspp*

Culture media: Media for isolation of bacteria were purchased from Titan Biotech LTD (India) and they include Nutrient agar, MacConkey agar, Eosin methylene blue (EMB) agar and Mannitol salt agar.

Standard antibiotic disk

The selected antibiotics discs (Oxoid, UK) used for the study include meropenem (10 µg), nitrofurantoin (30 µg), cephalixin (30 µg), ciprofloxacin (10 µg), gentamicin (10µg), ofloxacin (5µg), clindamycin (10 µg), erythromycin (10µg), ceftriaxone (30µg), ampicillin (30µg), levofloxacin (5µg) amoxicillin (30µg), streptomycin (30µg), cloxacilline (10µg), perfloxacin (10µg), chloramphenicol (10µg).



Fig 1: Nigerian currency notes

Methods

Preparation and sterilization of media

The media used were prepared according to manufacturers' specification

Sample Collection: A total of 300 currency samples of different denominations of ₦5, ₦10, ₦20, ₦50, ₦100, ₦200, ₦500, ₦1000 notes were randomly collected using an adapted method from Igumbore *et al.* (2007). Samples were obtained from different artisan groups of meat sellers, food sellers, palm oil sellers, mechanics and non artisans in the study area. Samples were collected from the surface of the naira currency note using a dampened cotton wool swab stick and kept in a refrigerator for future use.

Cultivation, isolation and characterization: Each of the samples collected was inoculated into 5 ml of brain heart infusion (BHI) broth for enrichment and incubated for 12 h at 37°C. Following the enrichment, sub-culturing of the broth culture was done onto different selective media for presumptive isolation of different bacteria. Due incubation was done at optimum temperature of 37°C for 24 h. Then, organisms isolated were characterized based on morphology, biochemical, cultural and metabolic characteristics.

Antibiotic susceptibility test: Antibiotic susceptibility of all bacterial species were done using the Kirby Bauer disc agar diffusion method and interpreted according to the guidelines of Clinical Laboratory Standards Institute (CLSI, 2023). Sterile molten Mueller-Hinton agar plates were allowed to set and solidify on a level surface. An 18- 24 h old broth culture of the all the test isolates were standardized by adjusting its turbidity to 0.5 Mcfarland's standard. The sterile swab sticks were used for this cultivation by dipping it into a tube of standardized test isolates, drained to remove excess inoculum and inoculated by streaking on the surface of prepared Mueller-Hinton agar plate. After this, the inoculated Mueller-Hinton agar plate was allowed to dry for a few minutes at room temperature with the lid closed. After drying, antibiotic impregnated discs (Oxoid, UK) of known concentrations were placed on the inoculated plates and incubated overnight at 37°C. The IZD (inhibition zone diameter) were measured using a graduated meter rule and results recorded and interpreted following the guideline of CLSI (2023).

Comment [WU3]: delete

Comment [WU4]: tested

Comment [WU5]: were recorded ...

Comment [WU6]: How MARI was calculated ?

Result

Ethical approval: The samples were collected with the consent of all the artisan and non artisan groups.

Isolation of bacteria from the samples

Out of the 300 currency notes that were randomly sampled, a total of 632 bacteria were cultured. There was no bacterial growth detected on the control samples collected from First bank of Nigeria (N=5). The number of currency note according to source were shown in (Table 1) and the number in each denominations represented (65) hospital/laboratory, (27) students, (61) meat sellers, (20) food sellers, (33) palm oil sellers, (54) bus conductors/drivers and (40) mechanic. Table 2 shows the frequency of occurrence of bacteria isolate of naira note from whole sample collection. This shows that 13 different microorganism were isolated from the naira note and 632 microorganisms in total were isolated from 300 samples. (Table 3) shows the total percentage prevalence from each isolate in all the denomination and out of 632 bacteria were isolated there were 223(74.3%) *Staphylococcus* sp, 95(31.6%) *E.coli*, 86(28.6%) *Bacillus subtilis*, 64(21.3%) *Klebsiella aerogenes*, 46(15.3%) *Streptococcus* sp, (8.0%) *Proteus mirabilis*, other organisms that were isolated include 16(5.3%) *Pseudomonas aeruginosa*, 18(6.0%) *Salmonella* sp, 10(3.3%) *Acinetobacter* sp, 11(3.6%) *Serratia marcescens*, 18(6.0%) *Enterococcus faecalis*, 19(6.3%) *Citrobacter* sp, 2(0.6%) Actinomycete. Gram positive isolates made up 53% of the total isolate while the rest were gram negative (47%). The antibiotic resistance pattern(%) of the bacterial isolated from all denomination are shown in (Fig 2) all Gram positive bacteria isolated from naira currency notes showed high susceptibility to cefuroxime, ciprofloxacin, and ofloxacin. However, Clindamycin and Erythromycin are not effective against *Staphylococcus* and *Streptococcus*. (Fig 3) Ciprofloxacin and Ofloxacin consistently demonstrate high resistance rates across multiple Gram-negative species and Meropenem and Ceftriaxone show relatively higher susceptibility rates compared to other antibiotics for several genera. The multiple antibiotic resistant indexes are shown in Table 4.

Comment [WU7]: need to be checked

Comment [WU8]: were

Table 1: Number of Currency note according to source

Denomination (₹)	hospital/laboratories	Students	Meat vendors	Food vendors	Palm oil vendors	Bus conductor s/ Driver	Mechanics	Total
1000	24	20	40	4	0	0	8	96
500	20	6	2	2	14	16	15	75
200	3	4	2	2	2	2	3	18
100	4	5	1	2	3	2	1	18
50	9	0	6	2	4	1	7	29
20	5	11	7	4	7	2	4	39
10	3	8	2	1	2	1	2	19
5	0	0	1	1	1	3	0	6
Total	65	27	61	20	33	54	40	300

Table 2: Frequency of occurrence of bacterial isolates in the test Naira note.

Denomination (₦)	<i>EC</i>	<i>Sa</i>	<i>Pa</i>	<i>Bs</i>	<i>Sal</i>	<i>Ka</i>	<i>St s</i>	<i>Acto</i>	<i>Sm</i>	<i>Pm</i>	<i>Ef</i>	<i>Ct</i>	<i>Ac sp,</i>	Total
1000	35	76	5	16	1	20	17	1	1	12	6	5	1	196
500	10	48	1	30	7	13	13	0	2	0	1	4	0	129
200	8	13	0	9	2	3	1	0	0	0	1	1	1	39
100	7	16	1	4	0	9	2	1	3	2	0	0	0	45
50	12	25	6	8	6	5	4	1	2	6	1	2	0	78
20	12	30	1	8	1	11	5	5	2	3	5	5	0	88
10	9	10	2	7	1	2	3	0	1	1	3	1	0	40
5	2	5	0	4	0	1	1	2	0	0	1	1	0	17
Total	95	223	16	86	18	64	46	10	11	24	18	19	2	632

Key: *Ec*, *Escherichia coli*, *Sa*, *Staphylococcus sp*, *Pa*, *Pseudomonas aeruginosa*, *Bs*, *Bacillus subtilis*, *Sal*, *Salmonella sp*, *Ka*, *Klebsiella aerogenes*, *St s*, *Streptococcus sp*, *Acto*, *Acinetobacter sp*, *Sm*, *Serratia marcescens*, *Pm*, *Proteus mirabilis*, *EF*, *Enterococcus faecalis*, *Ct*, *Citrobacter sp*, *Ac sp*, *Acinomycetessp*

Table 3: Total percentage prevalence of each isolates from all the different denomination

Denomination (N)	EC	Sa	Pa	Bs	Sal	Ka	St s	Acto	Sm	Pm	Ef	Ct	Ac sp,	Total
1000	35	76	5	16	1	20	17	1	1	12	6	5	1	196
500	10	48	1	30	7	13	13	0	2	0	1	4	0	129
200	8	13	0	9	2	3	1	0	0	0	1	1	1	39
100	7	16	1	4	0	9	2	1	3	2	0	0	0	45
50	12	25	6	8	6	5	4	1	2	6	1	2	0	78
20	12	30	1	8	1	11	5	5	2	3	5	5	0	88
10	9	10	2	7	1	2	3	0	1	1	3	1	0	40
5	2	5	0	4	0	1	1	2	0	0	1	1	0	17
Total	95	223	16	86	18	64	46	10	11	24	18	19	2	632
% Prevalence	31.6	74.3	5.3	28.6	6.0	21.3	15.2	3.3	3.6	8.0	6.0	6.3	0.6	

Key: *Ec*, *Escherichia coli*, *Sa*, *Staphylococcus* sp, *Pa*, *Pseudomonas aeruginosa*, *Bs*, *Bacillus subtilis*, *Sal*, *Salmonella* sp, *Ka*, *Klebsiella aerogenes*, *St s*, *Streptococcus* sp, *Acto*, *Acinetobacter* sp, *Sm*, *Serratia marcescens*, *Pm*, *Proteus mirabilis*, *EF*, *Enterococcus faecalis*, *Ct*, *Citrobacter* sp, *Ac sp*, *Acinomycetessp*

Fig 2: Percentage antibiotic resistance pattern of Gram positive Bacteria.




Fig 3: Percentage antibiotic resistance of Gram negative Bacteria.




Table 4: Determination of Multiple Antibiotic Resistant Index (MARI)

Bacteria	Range of MARI
<i>Staphylococcus</i> sp	0.1 – 0.8
<i>Proteus mirabilis</i>	0.1 – 0.5
<i>Streptococcus</i> sp	0.1 – 0.7
<i>Bacillus subtilis</i>	0.3 – 0.5
<i>Serratia marcescens</i>	0.4 – 0.9
<i>Klesiella aerogenes</i>	0.3 – 0.6
<i>Pseudomonas aeruginosa</i>	0.2 – 0.9
<i>Acinetobacter</i> sp	0.1 – 0.5
<i>Citrobacter</i> sp	0.1 – 0.6
<i>Salmonella</i> sp	0.2 – 0.5

Discussion

Bacteria colonize both living and non living materials including food, stuff, meat, different fishes, wines, inert surfaces, currency notes and can be transferred from person to person through unethical practices (Adonuet *al.*, 2024). Whenever there is a presence of microorganisms on fomites, it suggests that the minimum conditions for their presence have been met. In the case of Nigerian naira notes, due to high frequency of use, one may question: 'How microbiologically safe are our naira notes in circulation'? In the present study, we isolated different bacteria that contaminate Naira currency notes. More so, we determined the antibiotic susceptibility pattern of the bacteria isolated from the test currency notes. This study made use of a total of 300 samples of naira notes (209 paper notes and 93 polymer notes) ranging from ₦5 to ₦1000 collected from hospital/laboratory, students, meat vendors (butchers), mechanics, bus conductors/drivers, palm oil vendors and food vendors. Some freshly printed naira (clean mint) notes were used as a control. All the naira notes that had bacterial growths were rough and dirty in appearance. The control group obtained from a commercial bank in the same study area did not show any bacterial growth. Of all the 300 samples used, the total of 632 isolates was isolated. *Staphylococcus aureus* was the prevalent bacterium (74.3%) followed by *Escherichia coli*

Comment [WU9]: italic

(31.6%) and *Bacillus subtilis* (28.6%), which is in agreement with results of other researchers (Umeh *et al.* 2007; Shekarforoushet *al.* 2009; Abdumoniemet *al.* 2010; Alwakeel and Nasser. 2011; Elememet *al.* 2016; Firoozeh *et al.* 2018; Nanda *et al.* 2019)). *Staphylococcus aureus* forms part of the normal flora of the skin of human hands and this accounts for high prevalence of this organism in the test currency notes. Similarly, *Escherichia coli*, which form part of the normal flora of the gut, frequently contaminate hands in individuals that do not maintain proper hygiene. The prevalence of *Klebsiella spp* was 21.3 %, which is in agreement with the findings of researchers (Chigozie *et al.*, 2021). Other bacteria which we isolated contaminated the currency notes in a manner that poses threats to public health. The results of this research shows that lower denomination was the most contaminated, this is consistent with a previous report (Tswana *et al.* 2000; Igumbore *et al.* 2007) and this is understandable because a good number of individuals currently have higher denominations in exchange for lower denomination when trading. It is important to note that bacterial growth was not detected in 5 samples of mint banknotes and this might be attributed to the fact that they had not been in circulation, which normally exposes them to usage and contamination. However, some researchers believed that some currency notes that are not yet in circulation could still be contaminated with fastidious organisms (Ahmed *et al.*, 2018).

The majority of the test bacteria showed resistance to most of the antibiotics used. The highest resistance was seen against Gram positive bacteria. Many of the bacterial isolates showed resistance to amoxicillin (78.4%), cloxacillin (68.6%), ampicillin (63.4%), cefuroxime (56.0%) and erythromycin (53.6%), which was in agreement with the result of work done elsewhere (Obajuluwa *et al.*, 2023). The high resistance pattern recorded against these antibiotics implies inappropriate use of antimicrobial agents in the study area. Most of the test bacteria had a MAR index > 0.2 indicating their source to where antibiotics are commonly used or previous exposure of the organism to antimicrobial agents as reported by other researchers (Gunasekar, 2017, Mthembu *et al.* 2019, Afunwa *et al.* 2020). In other words, the test isolates were obtained from high risk sources of antibiotic resistance. These isolates were from artisans and non-artisan groups, pointing to the fact that antibiotic resistance and development of superbugs are not limited to hospital acquired pathogens only, but can be from community acquired pathogens too. It has been shown that $MARI \geq 0.4$ or higher is associated with human faecal sources of contamination (Joseph *et al.*, 2017). This demonstrates a highly compromised hygienic environment for selling and processing of food.

Conclusion

This study showed that Nigerian currency notes harbour bacteria. Many of these bacteria are multidrug resistant and this poses a serious threat to public health. Inappropriate use and antibiotic misuse might have contributed to the presence of superbugs in the study area.

Comment [WU10]: correct it

References

- Abdulmoniem, M., Saadabi, M., Alhussaini, A., Al-Ghanayem, B., Mohammed, S. & Al Shuriam.(2017). Isolation and Identification of Pathogenic Bacteria and Fungi from some Saudi Bank Note Currency. *Bioscience Biotechnology Research, Asia*. 14(2):715-720.
- Adonu, C., Ujam, T., Ali, I., Okorie, H., Adonu, M., Restus, O. & Offiah, R.(2024). Phenotypic characterization of bacteria isolated from crayfish (*Procambrusclarkii*) and fresh catfish (*Clariasgariepinus*) in Nsukka, Nigeria. *Journal of Advances in Microbiology Research*. 5(2): 230-235.
- Afunwa, R. A., Ezeanyinka, J., Afunwa, E. C., Udeh, A. S., Oli, N. A. &Unachukwu, M.(2020).Multiple antibiotic resistant index of Gram-Negative Bacteria from Bird Droppings in Two Commercial Poultrys in Enugu, Nigeria.*Open Journal of Medical Microbiology*. 10(04):171-181.
- Ahmed, S., Parveen, S., Nasreen, T. & Feroza,B. (2010). Evaluation of the Microbial Contamination of Bangladesh Paper Currency Notes (Taka) in Circulation. *Advances in Biological Research*. 4 (5): 266-271.
- Alwakeel, S.S. & Nasser, L.A. (2011). Bacterial and fungal contamination of Saudi Arabian paper currency and cell phones. *Asian Journal of Biological Sciences*.4(7):556-562.
- Awe, S., Eniola, K. I. T., Ojo, F. T., & Sani, A. (2010). Bacteriological quality of some Nigerian currencies in circulation. *African Journal of Microbiology Research*,4(21): 2231–2234.
- Chigozie, E., Jude, O., Ijeoma, M., Perpetual, Z., Charles, O. &Malgorzata, K. (2021). Bacterial contamination of Nigerian currency notes: A comparative analysis of different denominations recovered from local food vendor. *National Library of Medicine*. e10795. 2021; doi:10.7717/peerj.10795
- CLSI. Performance standard for antimicrobial susceptibility testing. (2023). 33rd ed. CLSI Supplement M100. *Clinical and Laboratory Standard Institute*; c2023.
- Elemem, M. M., Dhawi, A.,Shaban, M.B. &Dahmani, K. A. (2016).Study of bacterial contamination on Libyan paper banknotes in circulation. *American Journal of Microbiology and Biotechnology*. 3(1):1-6.
- Elsharief, M., Haider, J. & Waly, S. (2018). A study of bacterial contamination of paper currency notes circulating in zliten area and their antibiotic resistance. *Journal of Basic Sciences*. 31: 99-115.
- Firoozeh, F., Dadgostar, E., Akbari, H., Zibaei, M., Sadjjadian, S., Moshtaghi, M. &Shakib, A. (2017). Bacterial Contamination of Iranian Paper Currency and Their Antibiotic Resistance Patterns. *International Journal of Enteric Pathogens*. 5(4): 106-110.
- Gunasekar, B. (2017). Multiple Antibiotic Resistance Indexing of non- fermenting gram negative bacilli. *Asian Journal of Pharmaceutical and Clinical Research*. (10): 78-80.
- Igumbor, E.O., Obi, C.L., Bessong, P.O., Potgieter, N.&Mkasi, T.C. (2007). Microbiological analysis of banknotes circulating in the Venda region of Limpopo province, South Africa. *South African Journal of Science*.103(2):365-366.

- Joseph, A., Odimayo, M., Olokoba, L., Olokoba, A. & Popoola, G. (2017). Multiple Antibiotic Resistance Index of *Escherichia Coli* Isolates in a Tertiary Hospital in South-West Nigeria. *Medical Journal of Zambia*. Vol. 44, No.4: 225-232.
- Moses, I., Ugbo, E., Iroha, I., Ukpai, E., Eluu, S., Ilang. & Otozi, S. (2018). Public Health Implications of Salmonella species Contamination of Naira notes Obtained from Butchers in Abakaliki meat Market, Ebonyi state. *African Journal of Clinical and Experimental Microbiology*. 9 (3): 165-170
- Mousa, M. & Idress, H. (2023). Evaluating the Level of Bacteria Contamination on Libyan Banknotes and Coins in Circulation City. *Mediterranean Journal of Basic and Applied Sciences*. 7(4):134-149,
- Mthembu, T.P., Zishiri, O.T. & El Zowalaty, M.E. (2019). Molecular Detection of Multidrug-Resistant *Salmonella* Isolated from Livestock Production Systems In South Africa. 14(12):3537-3548.
- Muhumuza, A., Catherine, A., Musisi, N., Francis, E. & Samuel, N. (2018). Bacterial contamination of Ugandan paper Currency notes possessed by food vendors around Mulago Hospital complex, Uganda. *Pan African Medical Journal*. 31:143.
- Nanda, B., Abdullah, A.S, Zobayda, F.H, Manisha, D., Muhammad, T. H. & Sukumar S. (2019). Survey of bacterial contamination and antibiotic resistance pattern of Bangladeshi paper currency notes in Mymensingh city. *African Journal of Microbiology Research*. 13(10):206-213
- Obajuluwa, A. F., Igwe, J. C., Durowaiye, M. T., Garba, S. & Bamai, D. Z. (2023). Microbial load and antibiotic resistance patterns of bacterial isolates from Nigeria naira notes. *FUDMA Journal of Sciences (FJS)*. 7(6):261 -265.
- Shekarforoush, S., Khajeh, A.E. & Zarei, M. (2009). Evaluation of the bacterial contamination of the Iranian currency notes. *Iranian Journal of Health and Environment*. 1(2):81-8.
- Snehalatha, V., Malashree, R. & Preeti, S. (2016). Isolation, Enumeration and Antimicrobial Susceptibility of Predominant Microbes Associated with Currency Notes. *International Journal of Current Microbiology and Applied Sciences*. 5(8):650-657.
- Tswana, S.A., Chihota, V.N., Moyo, S.R., Chigonde, S.C. & Nzara, M.J. (2000). Contamination of Zimbabwean banknotes by microorganisms – a pilot study. *South African Journal of Epidemiology and Infection*. 15(1):14–18.
- Umeh, E., Juluku, J. & Ichor, T. (2007). Microbial contamination of Naira (Nigerian currency) notes in circulation. *Research Journal of Environmental Sciences*. 1:336-339.

UNDER PEER REVIEW