

# Original Research Article

## Human Health Risk Assessment of Heavy Metals and Polycyclic Aromatic Hydrocarbons in Seafood from Southern Nigeria's Coastal Regions

### ABSTRACT

**Aim:** The levels of heavy metals and polycyclic aromatic hydrocarbons (PAHs) in tilapia fish, prawn, oyster and periwinkle samples from southern Nigeria's coastal regions were evaluated to determine possible human health risk associated with consumption.

**Study design:** Ten (10) samples each of the selected seafood were collected from mangrove swamps along Bakana River in the study area. The samples were analyzed for nutrient levels and concentrations of toxic metals and PAHs.

**Methodology:** The concentrations of heavy metals and PAHs in the selected seafood samples were determined using Gas Chromatography Flame Ionization Detector and Atomic Absorption Spectrophotometry. Probable human health risk associated with the seafood consumption for adult and children populations was evaluated using Estimated Daily Intake (EDI), Target Hazard Quotient (THQ) and Carcinogenic Risk (CR) assessment for heavy metals and PAHs.

**Results:** The seafood samples showed the presence of heavy metals (Cr, Cd, Cu, Pb and Zn) but values were all below regulatory permissible limits of 0.6, 0.05, 3.0, 0.2 and 5.0 mg/kg respectively. PAH compounds (fluorene and anthracene) were below detectable limits in the seafood samples; phenanthrene was detected in all the seafood samples but highest in Tilapia fish, recording  $28.32 \pm 0.05$  mg/kg. Benzo (a) pyrene was below detectable limit in the investigated seafood samples. Carcinogenic Risk (CR) assessment values for heavy metals in the seafood samples were significantly ( $p \leq 0.05$ ) high for Cd, Pb, Cr and Ni in both adult and children populations. CR values for PAHs were within regulatory threshold of  $10E-6 - 10E-4$ .

**Conclusion:** This study revealed that individual carcinogenic potencies for heavy metals (Cd, Pb, Ni, Cr) in the seafood samples exceed the guideline screening value for human consumption, indicating high potential for carcinogenic risk. Our findings suggest that Bakana river is exposed to substantial amount of carcinogenic environmental pollutants associated with deleterious environmental and epidemiological effects.

*Keywords: Heavy metals, Polycyclic aromatic hydrocarbons, Seafood, Risk assessment*

### 1. INTRODUCTION

Seafood has been a significant food source for humans, especially those in the coastal region, because of their flavor and nutritional value. Seafood describes various ocean creatures categorized into fish and shellfish (e.g. Oysters, squid, mussel, lobster, prawns, crab and shrimp). The low calorific value and level of protein in seafood makes it a healthier alternative to red meats or poultry. Seafood is reported to be rich in calcium, iron, and

vitamins A, E, C and D [1, 2]. Seafood consists of filter-feeding aquatic organisms, with high bioaccumulation rate for toxins in their tissues [3, 4]. They are vulnerable to ingesting chemical contaminants from sediment, water, micro- and macro-organisms [1]. These contaminants can further bioaccumulate in their tissues and subsequently be biomagnified in the food chain [5, 6].

Anthropogenic influences on aquatic habitats, especially arising from increased industrialization has heightened research on the safety of seafood. Anthropogenic sources include stubble burning and dispersion of contaminated sewage sludge on farmlands, exhausts from automobiles and oil pollution of surface waters [7 - 9]. Polycyclic aromatic hydrocarbons (PAHs) and heavy metals, among others have continued to pose environmental hazard [10]. Contamination of seafood by PAHs is as a result of deposition and transfer. PAHs are fat-soluble compounds that are carcinogenic, mutagenic, and capable of disrupting hormonal functions [11-13]. Sixteen PAHs compounds (Naphthalene, Benzo[k]fluoranthene, Acenaphthylene, Acenaphthene, Benzo[a]anthracene, Fluorene, Benzo[b]fluoranthene, Phenanthrene, Anthracene, Dibenzo[a,h]anthracene, Pyrene, Chrysene, Benzo[a]pyrene, Indeno[1,2,3-cd]pyrene, Fluoranthene, and Benzo[g,h,i]perylene) are considered by the U.S. Environmental Protection Agency (EPA) as priority pollutants as they are the most toxic, carcinogenic, and commonly found PAHs in the environment. These PAHs compounds are often used as markers for assessing PAH contamination and exposure. Meanwhile, Benzo(a)pyrene (BaP) a known carcinogen, is often used as a benchmark compound when evaluating the potential carcinogenicity of PAH mixtures [14]. High exposure to some heavy metals have been reported to cause persistent brain damage, kidney failure cognitive decline, memory issues, mood swings, interference with enzyme activity leading to organ damage, neurological abnormalities and several other health issues [15 -17]. Akpabio et al. [18] also reported that excessive amounts of heavy metals in food have potentials to be fatal and cause serious health issues. Consuming seafood tainted with heavy metals has been linked to a number of diseases that are known to have negative health consequences on humans

Pollution of the environment by crude oil spillage can be a source of PAH contamination mostly in fish, fishery products and seafood. In recent times, chemical contamination from oil spillage has shown to be a major source of pollution within the coastal areas of Nigeria's Niger Delta Region (NDR) [7, 19, 20]. The environment in the NDR has been seriously contaminated by oil production, according to numerous studies. Many illicit refineries and bunker operations exist in the region; these operations are not sufficiently regulated and thus, contaminants are released water bodies within the region. The aquatic system is most commonly contaminated component of the environment. This is due to the fact that pollutants in air, soil, or on land eventually find their way into the aquatic system due to a variety of factors, including local precipitation, water runoff, rock leaching, and solid waste. This poses a significant threat to marine ecosystems, with potentially devastating consequences for seafood safety. The bioaccumulation and toxic consequences associated with these marine creatures, as well as their subsequent transfer to people through the food chain, present considerable concerns due to their negative repercussions [21, 22]. Seafood Consumers are therefore gravely concerned about the health effects of consuming seafood from the Niger Delta Region.

This research examined the health risks of consuming seafood (Prawn, Oyster, Periwinkles and Tilapia fish) obtained from the study area. The study's objectives include; (i) to assess the concentrations of PAHs and heavy metals (HMs) in tissues of selected seafood obtained from Bakana River (ii) Determine the safety of the seafood samples under study for human consumption (iii) To evaluate the potential health hazards of consuming seafood from the study area.

## 2. MATERIAL AND METHODS

### 2.1 Study Location

The study area, shown in Fig. 1, is located within the pollution-prone mangrove swamp of Bakana River in Bakana Community in the coastal region of Southern Nigeria. Situated in the Degema Local Government Area of Rivers State, Bakana is home to the Kalabari ethnic Community. The Bakana River is situated within the Niger Delta basin. The basin spans 20,000 km<sup>2</sup> and includes all of the territories between latitudes 4° 14'N and 5° 35'N, and longitudes 5° 26'E and 7° 37'E. The location is characterized by silt and mud deposits, swampy areas, and mangrove plants. Bakana Community, as well as other coastal Communities within the Niger Delta region are known for fishing and serve as a source of seafood for marketers engaged in subsequent distribution.

Bakana River is constantly exposed to organic waste, litter, petroleum hydrocarbon pollutants, and toxic chemicals. The area lies in the Niger Delta wet equatorial climatic which experience extensive-rainy season from March to November, with mean annual rainfall range from 1500 mm around the northern fringe to 4500 mm around the coastal margin [23, 24].




Fig. 1 Map of the study area

### 2.2 Seafood Sample Collection

Ten (10) samples each of Tilapia fish, Prawn, Oyster and Periwinkle were collected between June and July 2024 in plastic containers from mangrove swamps along Bakana River. These species were selected considering their consumption rate and availability. Periwinkles and oysters were hand-picked at low tide from the mud in the creeks. Mangrove oysters were harvested from mangrove roots. Collected seafood samples were immediately transferred in

sterile isothermal container for onward delivery to the laboratory for analysis. The samples were analyzed for proximate composition, mineral content, PAHs and heavy metals levels.

### 2.3 Proximate Analysis

Proximate analysis was carried out according to the procedure of Association of Official Analytical Chemist (AOAC) [25] for moisture, ash, crude fibre and crude protein content. Carbohydrate component was calculated by the Difference Method [25] by subtracting the sum (g/100g dry matter) of crude protein, crude fat, ash and fibre from 100g. The caloric value was determined based on the Atwater Factor [26].

### 2.4 PAHs and Heavy metal determination

Sixteen (16) priority PAHs listed by the USEPA and cited by the International Agency for Research on Cancer (IARC) were analyzed [27, 28]. Sample analysis followed the method described by Odesa and Olanye [1]. The samples were homogenized and 25 ml of dichloromethane was used for the extraction after it has been dehydrated with anhydrous sodium sulfate ( $\text{Na}_2\text{SO}_4$ ). The samples were further evaporated and a portion of the solution was examined using Gas Chromatography Flame Ionisation Detector (GC model: Agilent 6890N) and Atomic Absorption Spectrophotometry (AAS model: SP-AA4530). The congeners were detected via flame ionization detection. Heavy metal levels in seafood were determined using an atomic absorption spectrophotometer (ASTM-E594-96).

### 2.5 Health Risk Assessment for PAHs and Heavy metals

Health risk evaluation was performed for children and adult population using Estimated Daily Intake (EDI), Target Hazard Quotient (THQ) and Carcinogenic Risk (CR) assessment.

#### 2.5.1 Estimated Daily Intake (EDI)

Approximate daily consumption of the seafood was determined using the formula below, taking into account the levels of PAHs and heavy metals in the chosen sea food samples.

$$EDI = \frac{C_m \times IR}{BW \times 10^{-3}} \quad (1)$$

Where BW is the average adult and child body weight who consume the seafood, IR is the ingestion rate of heavy metals and PAHs in kg/day, and  $C_m$  is the saturation of metals and PAHs in each chosen seafood in mg/kg.

#### 2.5.2 Target Hazard Quotient (THQ)

The Target Hazard Quotient (HQ), which is the ratio of the computed chronic intake (CDI) to the ingestion reference dose (RfD) of the chosen heavy metals, is typically used to highlight the degree of non-carcinogenic concerns [29]. USEPA [30] provides the formula, which is displayed in equation 2. If  $HQ > 1$ , it suggests that the exposed population is more likely to experience bad health impacts. Conversely, if  $HQ < 1$  then there is no possibility of negative health effects with the ingestion reference dose for heavy metals to be that set by the WHO 2017, with PAHs having the ingested reference dose to be Acy  $6.0 \times 10^{-2}$ , Acp  $6.0 \times 10^{-2}$ , Flr  $4.0 \times 10^{-2}$ , Ant  $3.0 \times 10^{-1}$ , Phe  $3.0 \times 10^{-2}$ , Flt  $4.0 \times 10^{-2}$  and Pyr  $3.0 \times 10^{-1}$  [30].

$$HQ = \frac{CDI}{RfD} \quad (2)$$

The ratio of estimated daily intake (EDI) to RfD was used to compute the health risks associated with consuming seafood. Equation 4 was used to determine the EDI (Ding et al. 2012).

$$\text{THQ} = \frac{\text{EDI} \times \text{EF} \times \text{ED}}{\text{AT} \times \text{RFD}} \times 10^{-3} \quad (3)$$

$$\text{RFD}_{\text{derm}} = \text{RFD}_{\text{oral}} \times \text{ABS}_{\text{gi}} \quad (4)$$

The ABS<sub>gi</sub> value is the gastrointestinal absorption factor. It has no unit of its own, Cr (0.25), Pb, (0.1), Cd (0.08), Cu (0.3) and Zinc (0.61) with Ni not assigned and is 0.89 for PAHs [31].

### 2.5.3 Carcinogenic Risk Assessment

The malignant growth slant factor (SF) (USEPA 2000) was duplicated by the CDI or EDI to gauge the HQs for cancer-causing risk from ingestion/dermal openness to seafood, as indicated in equation 5. The Incremental Lifetime Cancer Risk (ILCR) is calculated using potential cancer risk when the ratios are larger than 1.

$$\text{CR} = \text{EDI} \times \text{CSF}_{\text{for seafood sample}} \quad (5)$$

According to the toxicological assessments and risk system created by the USEPA, WHO, and International Agency for Research on Cancer (IARC), the following heavy metals have carcinogenic slopes coefficients of 0.38, 0.84, and 0.5 that indicate they are recognized human carcinogens: Cd, Ni, Cr, and Pb. and  $8.5 \times 10^{-3}$  (mg L<sup>-1</sup> day<sup>-1</sup>)[29] while verified cancer slope factor for PAHs is 11.5 mg/kg/day [32].

### 2.6 Statistical analysis

Statistical analysis of data All values were expressed as mean  $\pm$  SD and then subjected to analysis of variance (ANOVA) using the Statistical Package for Social Sciences (SPSS) version 17.0 (SPSS Inc., Chicago Illinois). Statistical significance was considered at P=0.05.

## 3. RESULTS AND DISCUSSION

### 3.1 Proximate Composition of Seafood samples

Results for proximate analysis of Tilapia fish, Prawn, Oyster and Periwinkle are presented in Figure 2. The results indicate that Prawn has the highest percentage of crude protein (19.71%) and carbohydrate (8.38%), when compared with other seafood in this study. Ash content was highest in Periwinkle, recording  $6.00 \pm 0.78\%$ .

### 3.2 Mineral composition of the selected seafood

Analytical results for mineral composition of seafood samples in this study are shown in Figure 3. Prawn and Oysters showed the highest calcium contents, recording 72.14 mg/100g and 86.32 mg/100g respectively. Tilapia fish recorded the least calcium content (54.05 mg/100g). The highest potassium level is recorded in Periwinkle (33.60 mg/100g), followed by Prawn (27.61 mg/100g). The highest level of sodium is found in Oyster (18.2 mg/100g), closely followed by Periwinkle (16.70 mg/100g).

### 3.3 Heavy metal composition of seafood samples

Table 3 shows the results for heavy metal levels in the seafood samples under study. Seafood samples studied showed the presence of heavy metals namely Cr, Cd, Cu, Pb, Zn, Ni, but values were all below WHO permissible limits. Tilapia fish recorded  $0.009 \pm 0.001$  mg/kg of chromium (Cr), while Prawn showed  $0.001 \pm 0.000$  mg/kg, Oyster recorded  $0.001 \pm 0.000$  mg/kg, and Periwinkle has  $0.001 \pm 0.000$  mg/kg.

### 3.4 PAH content in seafood samples

Table 4 shows the composition of PAHs in the selected seafood samples. Fluorene and anthracene was below the detectable limit for all the selected seafood samples

while phenanthrene was detected in all selected seafood samples and highest in Tilapia fish, recording  $28.32 \pm 0.05$  mg/kg. Benzo (a) pyrene was below detection limit (BDL) in the investigated seafood samples. Benzo (g,h, i) perylene was also not detected in all the samples, implying that the concentrations were too small to be precisely measured.




Figure 2. Proximate composition of selected seafood samples




Figure3. Mineral composition of selected seafood samples

Table 1. Heavy metal content of selected seafood samples

SEAFOOD SAMPLE	Tilapia fish	Prawn	Oyster	Periwinkle	Permissible Limits: FAO [26]; WHO [33]
PARAMETER					

Cd (mg/kg)	0.001±0.000 <sup>ac</sup>	0.003±0.001 <sup>b</sup>	0.001±0.00 <sup>ac</sup>	0.001±0.00 <sup>ac</sup>	0.05
Pb (mg/kg)	0.008±0.000 <sup>abc</sup>	0.008±0.001 <sup>abc</sup>	0.008±0.001 <sup>abc</sup>	0.075±0.001 <sup>c</sup>	0.2
Cr (mg/kg)	0.009±0.001 <sup>a</sup>	0.001±0.000 <sup>bcd</sup>	0.001±0.000 <sup>bcd</sup>	0.001±0.000 <sup>bcd</sup>	0.6
Zn (mg/kg)	0.101±0.000 <sup>ab</sup>	0.127±0.001 <sup>ab</sup>	0.205±0.01 <sup>cd</sup>	0.218±0.002 <sup>cd</sup>	5.0
Ni (mg/kg)	0.163±0.000 <sup>a</sup>	0.207±0.001 <sup>bc</sup>	0.211±0.01 <sup>bc</sup>	0.339±0.002 <sup>d</sup>	NS
Cu (mg/kg)	0.003±0.000 <sup>a</sup>	2.461±0.003 <sup>b</sup>	0.296±0.00 <sup>c</sup>	1.301±0.001 <sup>d</sup>	3.0

Values are Mean ± Standard Deviation. Data with the same alphabets (a,b,c,d) as superscript shows non-significant differences ( $p \geq 0.05$ ), while that with different alphabets as superscript shows significant differences ( $p \leq 0.05$ ).

Table 2. PAH content of selected seafood samples

PAH Compound (mg/kg)	Tilapia fish	Prawn	Oyster	Periwinkle
Naphthalene	13.112 ± 0.1108 <sup>a</sup>	6.116 ± 0.141 <sup>b</sup>	BDL	4.116 ± 0.017 <sup>c</sup>
Acenaphthylene	5.404 ± 0.220 <sup>a</sup>	8.441 ± 0.081 <sup>b</sup>	11.411 ± 0.053 <sup>c</sup>	BDL
Acenaphthene	11.411 ± 0.012 <sup>a</sup>	2.153 ± 0.022 <sup>b</sup>	17.667 ± 0.032 <sup>c</sup>	BDL
Fluorene	BDL	BDL	BDL	BDL
Phenanthrene	28.323 ± 0.054 <sup>a</sup>	12.111 ± 0.004 <sup>b</sup>	20.183 ± 0.014 <sup>c</sup>	7.004 ± 0.013 <sup>d</sup>
Anthracene	BDL	BDL	BDL	BDL
Fluoranthene	BDL	15.550 ± 0.017 <sup>a</sup>	BDL	4.627 ± 0.028 <sup>b</sup>
Pyrene	BDL	BDL	BDL	8.111 ± 0.061
Benz (a) anthracene	7.008 ± 0.012 <sup>a</sup>	11.105 ± 0.021 <sup>b</sup>	BDL	BDL
Chrysene	BDL	BDL	BDL	3.077 ± 0.006 <sup>a</sup>
Benzo (b) fluoranthene	11.441 ± 0.016 <sup>a</sup>	8.408 ± 0.051 <sup>b</sup>	7.653 ± 0.017 <sup>c</sup>	BDL
Benzo (k) fluoranthene	BDL	BDL	BDL	BDL
Benzo (a) pyrene	BDL	BDL	BDL	BDL
Indeno (1,2,3-cd) pyrene	BDL	3.115 ± 0.042	BDL	BDL
Dibenz (a,h) anthracene	BDL	BDL	BDL	BDL
Benzo (g,h,i) perylene	BDL	BDL	BDL	BDL

Values are Mean ± Standard Deviation. Data with the same alphabets (a,b,c,d) as superscript shows non-significant differences ( $p \geq 0.05$ ), while that with different alphabets as superscript shows significant differences ( $p \leq 0.05$ ); BDL – Below Detection Limit (measured concentration is below measurable detection limit).

Table 3. Estimated Dietary Intake for Heavy Metals in selected seafood samples

	Tilapia fish		Prawn		Oyster		Periwinkle		USEPA [34]
	Adults	Children	Adults	Children	Adults	Children	Adults	Children	
Cd (mg/kg)	2.60E-10	3.13E-10	3.12E-03	3.75E-03	2.60E-10	3.13E-10	1.04E-03	1.25E-03	1
Pb (mg/kg)	1.66E-08	2.00E-08	1.04E-03	1.25E-03	1.66E-08	2.00E-08	9.75E-03	1.17E-02	1
Cr (mg/kg)	7.02E-06	8.44E-06	3.47E-07	4.17E-07	7.80E-07	9.38E-07	3.47E-07	4.17E-07	1



Zn (mg/kg)	1.58E-05	1.89E-05	2.20E-04	2.65E-04	3.20E-05	3.84E-05	3.78E-04	4.54E-04	1
Ni (mg/kg)	1.70E-06	2.04E-06	5.38E-03	6.47E-03	2.19E-06	2.64E-06	8.81E-03	1.06E-02	1
Cu (mg/kg)	6.24E-08	7.50E-08	3.20E-02	3.85E-02	6.16E-06	7.40E-06	1.69E-02	2.03E-02	1
THI	2.46E-05	2.95E-05	4.18E-02	5.02E-02	4.11E-05	4.94E-05	3.69E-02	4.43E-02	1

Table 4. Estimated Dietary Intake for PAHs in selected seafood samples

	Tilapia fish		Prawn		Oyster		Periwinkle	
	Adults	Children	Adults	Children	Adults	Children	Adults	Children
Naphthalene	6.82E-06	8.20E-05	3.18E-06	3.82E-05	0.00E+00	0.00E+00	2.14E-06	2.57E-05
Acenaphthylene	2.81E-06	3.38E-05	4.39E-06	5.28E-05	5.93E-06	7.13E-05	0.00E+00	0.00E+00
Acenaphthene	5.93E-06	7.13E-05	1.12E-06	1.35E-05	9.19E-06	1.10E-04	0.00E+00	0.00E+00
Fluorene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Phenanthrene	1.47E-05	1.77E-04	6.30E-06	7.57E-05	1.05E-05	1.26E-04	3.64E-06	4.38E-05
Anthracene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Fluoranthene	0.00E+00	0.00E+00	8.09E-06	9.72E-05	0.00E+00	0.00E+00	2.41E-06	2.89E-05
Pyrene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.22E-06	5.07E-05
Benz (a) anthracene	3.64E-04	4.38E-03	5.77E-04	6.94E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Chrysene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.60E-05	1.92E-04
Benzo (b) fluoranthene	5.95E-04	7.15E-03	4.37E-04	5.26E-03	3.98E-04	4.78E-03	0.00E+00	0.00E+00
Benzo (k) fluoranthene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Benzo (a) pyrene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Indeno (1,2,3-cd) pyrene	0.00E+00	0.00E+00	1.62E-04	1.95E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Dibenz (a,h) anthracene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Benzo (g,h,i) perylene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00

Table 5. THQ for Heavy Metals in selected seafood samples

	Tilapia fish		Prawn		Oyster		Periwinkle		USEPA [34]
	Adults	Children	Adults	Children	Adults	Children	Adults	Children	
Cd (mg/kg)	2.60E-10	3.13E-10	3.12E-03	3.75E-03	2.60E-10	3.13E-10	1.04E-03	1.25E-03	1
Pb (mg/kg)	1.66E-08	2.00E-08	1.04E-03	1.25E-03	1.66E-08	2.00E-08	9.75E-03	1.17E-02	1
Cr (mg/kg)	7.02E-06	8.44E-06	3.47E-07	4.17E-07	7.80E-07	9.38E-07	3.47E-07	4.17E-07	1
Zn (mg/kg)	1.58E-05	1.89E-05	2.20E-04	2.65E-04	3.20E-05	3.84E-05	3.78E-04	4.54E-04	1
Ni (mg/kg)	1.70E-06	2.04E-06	5.38E-03	6.47E-03	2.19E-06	2.64E-06	8.81E-03	1.06E-02	1
Cu (mg/kg)	6.24E-08	7.50E-08	3.20E-02	3.85E-02	6.16E-06	7.40E-06	1.69E-02	2.03E-02	1
THI	2.46E-05	2.95E-05	4.18E-02	5.02E-02	4.11E-05	4.94E-05	3.69E-02	4.43E-02	1

Table 6. THQ for PAHs in selected seafood samples

	Tilapia fish		Prawn		Oyster		Periwinkle		USEPA [34]
	Adults	Children	Adults	Children	Adults	Children	Adults	Children	
Naphthalene	3.41E-04	4.10E-03	1.59E-04	1.91E-03	0.00E+00	0.00E+00	1.07E-04	1.29E-03	1
Acenaphthylene	-	-	-	-	-	-	-	-	1
Acenaphthene	9.89E-05	1.19E-03	1.87E-05	2.24E-04	1.53E-04	1.84E-03	0.00E+00	0.00E+00	1



phenanthrene	1.08E-04	1.85E-04	4.60E-05	7.89E-05	7.66E-05	1.32E-04	2.66E-05	4.57E-05	10E-6 – 10E-4
Anthracene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	-	-	10E-6 – 10E-4
Fluoranthene	0.00E+00	0.00E+00	5.90E-05	1.01E-04	0.00E+00	0.00E+00	1.76E-05	3.02E-05	10E-6 – 10E-4
Pyrene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.08E-05	5.29E-05	10E-6 – 10E-4
Benz (a) anthr	2.66E-03	4.57E-03	4.22E-03	7.24E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4
Chrysene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.17E-04	2.01E-04	10E-6 – 10E-4
Benzo (b) fluor	4.34E-03	7.46E-03	3.19E-03	5.48E-03	2.91E-03	4.99E-03	0.00E+00	0.00E+00	10E-6 – 10E-4
Benzo (k) fluor	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4
Benzo (a) pyrene	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4
Indeno (1,2,3-cd) p	0.00E+00	0.00E+00	1.18E-03	2.03E-03	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4
Dibenz (a,h) anthr	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4
Benzo (g,h,i) per	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	10E-6 – 10E-4

Results for proximate analysis of the seafood samples showed that prawn has the highest percentage of crude protein (19.71%) and carbohydrate (8.38%) when compared with other seafood samples in this study. This confirms that prawn is a good source of dietary protein. Since prawn has high carbohydrates content, its consumption could provide the body with fuel and energy that is required for daily activities and exercise [35]. Ofor *et al.* [36] noted that adequate carbohydrate is also required for optimum function of the brain, heart, nervous, digestive and immune system while carbohydrate deficiency causes depletion of body tissue.

Results for mineral composition revealed that prawn and oyster had the next highest calcium contents, recording 72.14 mg/100g and 86.32 mg/100g respectively. The least calcium-containing seafood in this study is Tilapia fish which recorded 54.05 mg/100g. The highest potassium level was found in periwinkle (33.60 mg/100g), followed by prawn (27.61 mg/100g). Calcium deficiency can lead to osteoporosis in which the bone deteriorates and there is an increased risk of fractures. If there is more phosphorus than calcium in the diet, the body will start to take calcium from its own reserves (the bones) to compensate. Over a long or short period of time, this may affect the bones in a negative way. It is therefore necessary that a good source of calcium is used to complement diets [37].

The levels of six (6) heavy metals (Zn, Cu, Ni, Pb, Cr and Cd) were investigated in the selected seafood samples. Results showed that heavy metal content of the selected seafood samples were all below permissible limits as set by the Food and Agriculture Organization (FAO) [26] and World Health Organisation (WHO) [33]. Results from previous similar studies indicated a low to moderate level of heavy metals in seafood collected from the Niger Delta Region. Also, few studies from this region reported very high levels of some heavy metals in seafood collected from sites that have been severely impacted by decades of crude oil pollution and had suffered from various environmental impacts [38], [39], [40], [41]. Similarly, Ezemonye *et al.* [42] reported that hazardous levels of Pb, Ni and Co were detected by in seafood sampled from the Benin River in Southern Nigeria.

Sixteen (16) PAH compounds recognized PAHs by USEPA [43] (acenaphthene, acenaphthylene, anthracene, fluoranthene, fluorene, naphthalene, phenanthrene, pyrene, benz[a]anthracene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[ghi]perylene, benzo[a]pyrene, chrysene, dibenz[a,h]anthracene, and indeno[1,2,3-cd]pyrene) were investigated in the seafood samples under study. GCMS analysis detected nine (9) out of the sixteen (16) PAH compounds. Fluorene, Anthracene, Benzo (k) fluoranthene, Benzo (a) pyrene, Dibenz (a,h) anthracene and Benzo (g,h,i) perylene were not detected in any of the seafood samples. Established guidelines suggest the use benzo[a]pyrene as a marker for occurrence of carcinogenic PAHs in food [44]. The below-detecton levels of benzo(a)pyrene as observed in tilapa fish, prawn, oyster and periwinkles is at variance with the report by Odesa and Olannye [1], and Tongo *et al.* [45], who both assessed the concentrations of

polycyclic aromatic hydrocarbons (PAHs) in various types of seafood sourced from crude oil-contaminated coastal regions of Southern Nigeria. Both Authors in separate submissions noted that high levels of benzo(a)pyrene were detected in periwinkles and crabs.

The results of Lifetime Carcinogenic risk assessment for Heavy metals and the PAHs in children and adults are presented in Tables 7 & 8. No traces of the potential carcinogenic PAHs were recorded for Fluorene, Anthracene, Benzo (k) fluoranthene, Benzo (a) pyrene, Dibenz (a,h) anthracene and Benzo (g,h,i) perylene in both adult and children. The Lifetime Carcinogenic risk assessment of other PAH compounds in the selected seafood samples were estimated and observed to show mean values within the regulatory threshold of  $10E-6$  –  $10E-4$ . However, values reported for Carcinogenic Risk (CR) assessment of heavy metals were significantly ( $p \leq 0.05$ ) high for Cd, Pb, Cr and Ni in both adult and children populations. It follows that the risk of intake of carcinogenic heavy metals is high for consumers of the investigated seafood obtained from Bakana River.

#### **4. CONCLUSION**

The results of this study clearly demonstrate that there are significant impact of human activities that could lead to serious contamination of seafood in the study area, and the resultant risks to human health. The study has revealed that the presence of PAHs and heavy metals, in this coastal environment in the Niger Delta Region of Nigeria, as reported in other studies, can contaminate the food chain, and endanger both human and aquatic life within the study area. The individual carcinogenic potencies for heavy metals (Cd, Pb, Ni, Cr) in the seafood samples exceed the guideline screening value for human consumption, indicating high potential for carcinogenic risk. Therefore, consumers of the studied seafood samples may be exposed to the risk of consuming carcinogenic environmental pollutants. Our findings also suggest that Bakana river is exposed to substantial amount of carcinogenic environmental pollutants associated with deleterious environmental and epidemiological effects and if crude oil pollution in southern Nigeria's coastal region continues, severe health implications will be inevitable. It is therefore pertinent to implement measures to mitigate hydrocarbon pollution and monitor seafood safety regularly to protect public health.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **AUTHORS' CONTRIBUTIONS**

Author SEO designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. 'Author ESI' and 'Author SCO' managed the analyses of the study. 'Author SCO' managed the literature searches. All authors read and approved the final manuscript.

#### **Disclaimer (Artificial intelligence)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### **REFERENCES**

1. Odesa, G. E., & Olanye, D. U. (2025). Health risk assessment of PAHs and heavy metal levels in periwinkles (*Pachymelania fuscumutans*) and crabs (*Scylla serrata*) consumed in crude oil-contaminated coastal regions of Southern Nigeria. *Toxicology Reports*, 14, 101852.

2. Boylston, T., Chen, F., Coggins, P., Hydlig, G., McKee, L. H., & Kerth, C. (2012). *Handbook of meat, poultry and seafood quality*. John Wiley & Sons.
3. Shumway S. E., Davis, C., Downey, R., Karney, R., Kraeuter, J., Parsons, J., & Wikfors, G. (2003). Shellfish aquaculture—in praise of sustainable economies and environments. *World Aquac.*, 34(4), 8-10.
4. Berry, J. P., & Lind, O. (2010). First evidence of “paralytic shellfish toxins” and cylindrospermopsin in a Mexican freshwater system, Lago Catemaco, and apparent bioaccumulation of the toxins in “tegogolo” snails (*Pomacea patulacatemacensis*). *Toxicon*, 55(5), 930-938.
5. Alam, M. G. M., Allinson, G., Stagnitti, F., Tanaka, A., & Westbrooke, M. (2002). Arsenic contamination in Bangladesh groundwater: a major environmental and social disaster. *International Journal of Environmental Health Research*, 12(3), 235-253
6. Alemdaroğlu, T., Onur, E., & Erkakan F. (2003). Trace metal levels in surficial sediments of lake Manyas, Turkey and Tributary Rivers. *International Journal of Environmental Studies*, 60(3), 287-298.
7. Onojake, M. C., Nwokonko, V. N., & Osakwe, J. O. (2020). Human Health Risks Assessment of Polycyclic Aromatic Hydrocarbons in Selected Seafood from Niger Delta Nigeria. *Nigerian Journal of Chemical Research*, 25(2), 32–50.
8. Beck, H., Dross, A., & Mathar, W. (1992). PCDDs, PCDFs, and related contaminants in the German food supply. *Chemosphere*, 25(7-10), 1539-1550.
9. IPCS, W. (1998). *Assessment of the health risks of dioxins: Re-evaluation of the Tolerable Daily Intake (TDI)*. Geneva: World Health Organization International Programme on Chemical Safety.
10. Bowles, K. C., Apte, S. C., Maher, W. A., Kawei, M., & Smith, R. (2001). Bioaccumulation and biomagnification of mercury in lake Murray, Papua New Guinea. *Canadian Journal of Fisheries and Aquatic Sciences*, 58(5), 888-897.
11. Ibezue, V.C., Odesa, E. G., Ndukwe, J.O., & Nwabneli. E.O. (2018). Hydrocarbon concentration levels in groundwater in Jesse and Environ, Ethiope West LGA Delta State, Nigeria. *International Journal of Computer Applications Technology and Research*, 7(7), 241-254.
12. Odesa, G. E., Eyankware, M.O., & Elozona, O. E. (2024). Assessing the total petroleum hydrocarbon content in groundwater in parts of Eleme, Southern Nigeria. *Reading Time*, 01-05.
13. Odesa, G. E., Ozulu, G. U., Eyankware, M. O., Mba-Otike, M. N., & Okudibie, E. J. (2024). A holistic review of three-decade oil spillage across the Niger Delta Region, with emphasis on its impact on soil and water. *Reading Time*, 2024b, 02-02.
14. US Environmental Protection Agency (U.S.E.P.A.) (2001). *Integrated Risk Information System: Benzo[a]pyrene (BaP) (CASRN 50-32-8)*, Available from: U.S. Environmental Protection Agency, <<http://www.epa.gov/iris/subst/0136.htm>>
15. Balali-Mood, M., Naseri, K., Tahergorabi, Z., Khazdair, M.R., & Sadeghi, M. (2021). Toxic mechanisms of five heavy metals: mercury, lead, chromium, cadmium, and arsenic. *Frontiers in Pharmacology*, 12, 643972.
16. Charkiewicz, A. E., & Backstrand J. R. (2020). Lead toxicity and pollution in Poland. *International Journal of Environmental Research and Public Health*, 17(12), 4385.
17. Salnikow, K., & Zhitkovich, A. (2008). Genetic and epigenetic mechanisms in metal carcinogenesis and cocarcinogenesis: nickel, arsenic, and chromium. *Chemical Research in Toxicology*, 21(1), 28-44.
18. Akpabio, E.M., & Subramanian, S.V. (2012). *Traditional Ecological Knowledge: an Emerging framework for Understanding Water and Sanitation practices in Nigeria*.

ZEF working paper series 94. Zentrum für Entwicklungsforschung, University of Bonn, Germany.

19. Moslen, M., Miebaka, C.A., & Boisa, N. (2019). Bioaccumulation of polycyclic aromatic hydrocarbon (PAH) in a bivalve (*Arca senilis*-blood cockles) and health risk assessment. *Toxicology Reports*, 6, 990-997.
20. Akakuru, O.C., Eyankware, M. O., Akakuru, O. U., Nkwoada, A. U., & Agunanne, V. C. (2023). Quantification of contamination, ecological risk index, and health risk assessment of groundwater using artificial neural network and multi-linear regression modeling approaches within Egbema, Nigeria. *Arabian Journal of Geosciences*, 16(9), 507.
21. Eyankware, M. O., Nnabo P. N., &Ogwah, C. (2020). Impact of past mining activity on water resources around active and abandoned mines in Ebonyi State South-Eastern Nigeria. *Hydro Science & Marine Engineering*, 2(2), 29-35.
22. Shamsollahi, H.R., Alimohammadi, M., Momeni, S., Naddafi, K., Nabizadeh, R., Khorasgani, F. C., & Yousefi, M. (2019). Assessment of the health risk induced by accumulated heavy metals from anaerobic digestion of biological sludge of the lettuce. *Biological Trace Element Research*, 188, 514-520.
23. Adejuwon, J. O. (2012). Rainfall seasonality in the Niger delta belt, Nigeria. *Journal of Geography and Regional Planning*, 5(2), 51-60.
24. Igweze, A. H., Amagoh, M. N. & Ashinze, A. N. (2014). Analysis of rainfall variations in the Niger Delta region of Nigeria. *Journal of Environmental Science, Toxicology and Food Technology*, 8(1), 25-30.
25. AOAC (1990): Official Methods of Analysis of the Association of Official's Analytical Chemists, 14th Ed. Association of Official Analytical Chemist, Washington D.C., pp: 223-225, 992-995.
26. FAO (2003): Food energy-methods of analysis and conversion factors. FAO Food and nutrition paper 77.
27. Andersson, J. T., &Achten. C. (2015). Time to say goodbye to the 16 EPA PAHs? Toward an up-to-date use of PACs for environmental purposes. *Polycyclic Aromatic Compounds*, 35(2-4), 330-354.
28. Muñoz, B., & Albores, A. (2011). DNA damage caused by polycyclic aromatic hydrocarbons: mechanisms and markers. *DNA Repair*, 20(1), 125-143.
29. Orosun, M. M., Adewuyi, A. D., Salawu, N. B., Isinkaye, M. O., Orosun, A. R., &Oniku, A.S. (2020). Monte Carlo approach to risks assessment of heavy metals at automobile spare part and recycling market in Ilorin, Nigeria. *Scientific Reports*, 10, 22084.
30. USEPA. (1999). Guidance for performing aggregate exposure and risk assessments. Office of Pesticide Programs, Environmental Protection Agency (EPA), Washington, DC.
31. Behera, B. K., Das, A., Sarkar, D. J., Weerathunge, P., Parida, P. K., Das, B. S., Thavamani, P., Ramanathan, R., & Bansal, V. (2018). Polycyclic Aromatic Hydrocarbons (PAHs) in inland aquatic ecosystems: Perils and remedies through biosensors and bioremediation. *Environmental Pollution*, 241, 212-233.
32. Karyab, H., Yunesian, M., Nasser, S., Rastkari, N., Mahvi, A., & Nabizadeh, R. (2016). Carcinogen Risk Assessment of Polycyclic Aromatic Hydrocarbons in Drinking Water, Using Probabilistic Approaches. *Iranian Journal of Public Health*, 45(11), 1455-1464.
33. World Health Organization (WHO) (2017). Guidelines for Drinking Water Quality, 4th ed., first Addendum Geneva. pp. 1 – 631.
34. United States Environmental Protection Agency (2011). Exposure Factors Handbook. (Final Edition). U.S. Environment Protection Agency, Washington, DC.
35. Udousoro, I., & Ekanem, P. (2013): Assessment of proximate composition of twelve edible vegetables in Nigeria: *International Journal of Modern Chemistry*, 4(2), 79-89.

36. Offor, I. F., Ehiri, R. C., & Njoku, C. N. (2014). Proximate nutritional analysis and heavy metal composition of dried *Moringa oleifera* leaves from Oshirionich L.G.A, Ebonyi State Nigeria. *Journal of Environmental Science, Toxicology and food technology (IOSR-JESTFT)*, 8(1), 57- 62.
37. Pravina, P., Sayaji, D., & Avinash, M. (2013): Calcium and its role in human body. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 4(2), 659-668.
38. Ihunwo, O.C., Dibofori-Orji, A.N., Olowu, C., & Ibezim-Ezeani, M.U. (2020). Distribution and risk assessment of some heavy metals in surface water, sediment and grey mullet (*Mugil cephalus*) from contaminated creek in Woji, southern Nigeria. *Marine Pollution Bulletin*, 154, Article 111042.
39. Ihunwo, O. C., Ibezim, M. U., Ezeani, V.O., Wekpe, A. R., Shahabinia, E., Bonnail, P. C., & Mmom, M. H. Jnr. (2022). Risk assessment of human exposure to lead and cadmium in tissues of Blackchin Tilapia (*Sarotherodon melanotheron*) from an intertidal creek in the Niger Delta region of Nigeria. *Environmental Research Communications*, 4(7) 75007.
40. Nkpaa, K. W., Onyeso, G. I., & Achugasim, O. (2017). Heavy metals levels in shellfish from Bodo City and B-Dere, Ogoniland, Rivers State, Nigeria, and evaluation of possible health risks to consumers. *Sustainable Water Resources Management*, 3, 83-91.
41. Nkpaa, K. W., Patrick-Iwuanyanwu, K. C., Wegwu, & M. O., Essien, E. B. (2016). Health risk assessment of hazardous metals for population via consumption of seafood from Ogoniland, Rivers State, Nigeria; a case study of Kaa, B-Dere, and Bodo City. *Environmental Monitoring and Assessment*, 188(1), 9, 10.1007/s10661-015-5006-4.
42. Ezemonye, L. I., Adebayo, P. O., Enuneku, A. A., & Tongo, I., Ogbomida, E. (2019). Potential health risk consequences of heavy metal concentrations in surface water, shrimp (*Macrobrachium macrobrachion*) and fish (*Brycinus longipinnis*) from Benin River, Nigeria. *Toxicology Reports*, 6, 1-9.
43. US Environmental Protection Agency (2005). US EPA Guidelines for Carcinogen Risk Assessment. EPA/630/P-03/001F., Risk Assessment Forum, Washington, DC.
44. Zelinkova, Z., & Wenzl, T. (2015). The Occurrence of 16 EPA PAHs in Food - A Review. *Polycyclic Aromatic Compounds*, 35(2-4), 248-284.
45. Tongo, I., Etor, E. E., & Ezemonye, L. I. N. (2018). Human health risk assessment of PAHs in fish and shellfish from Amariaria Community, Bonny River, Nigeria. *Journal of Applied Sciences and Environmental Management*, 22(5), 731-736.