

# Enriched Vermicompost made through bio waste of soybean stover (dry matter) and fresh cow dung using earthworms (*Eisenia fetida* spp.)

## ABSTRACT

This investigation was associated with different materials and techniques on bio-fertilizer application in soybean stover (dry matter) and fresh Cow dung for effect of bio fertilizer and earthworms (*Eisenia fetida* sp.) on bio waste decomposition. This study was analyzed by RBD (Randomized Block Design) with 6 treatments have to bio fertilizers combination *i.e.* (E<sub>1</sub>-control, E<sub>2</sub>- Rhizobium, E<sub>3</sub>- Rhizobium+ PSB, E<sub>4</sub>-Rhizobium+ KSB, E<sub>5</sub>-Rhizobium+ PSB+KSB, and E<sub>6</sub>-Rhizobium+ PSB+ KSB+ Trichoderma) and earthworms with 4 replications. The research was conducted at MRPC in Department of Soil Science and Agricultural Chemistry, JNKVV, Jabalpur during the Rabi season of 2021-22 and 2022-23. The result interpreted by partial decomposition, duration of completion, conversion rate and recovery percentage of vermicomposting. The result revealed that partial decomposition (kg) found in enriched vermicompost higher to lower sequence *i.e.* E<sub>5</sub> (7.11 kg) > E<sub>3</sub> (7.18 kg) > E<sub>2</sub> (7.19 kg) > E<sub>4</sub> (7.20 kg) = E<sub>6</sub> (7.20 kg) > E<sub>1</sub> (7.28 kg) from initial weight (10 kg). The production of enriched vermicompost maximum found in E<sub>6</sub> (5.58, 5.63 and 5.61 kg pot<sup>-1</sup>) within respective duration (42, 40 and 41 days) of vermicomposting in sequent years and statically pooled analysis. Conversion rate and recovery percentage of enriched vermicompost were increase with the combination of bio-agents. This investigation useful for making vermicompost using of agricultural waste through bio-fertilizer.

**Keyword:** *Rhizobium*, *Trichoderma*, *Phosphorus Solubilizing bacteria*, *Potassium solubilizing bacteria*, microorganisms and vermicompost *etc.*

## 1. INTRODUCTION

Waste management (solid, liquid and gaseous waste) is a major global challenge with increasing demand for protecting human health and the environment. In this research review, the classification and overview of research will help us identify the most important research areas for waste management. Furthermore, to promote the transition from a linear to a circular economy, waste management should be supported by policy-based initiatives and management policies (Maqsoodi *et al.*, 2023). The case of vermicomposting, or with mixed agricultural and eco-vermicomposting, the earthworms can assist in detoxifying commercial and ecological wastes and their very last product is useful for plant increase as natural fertilizers (Raza *et al.*, 2024). Millions of tons of agricultural and industrial waste are discarded every year at considerable financial and environmental cost. Instead of discarding the food scraps and waste, we can recycle these with the help of earth worms (Fayaz *et al.*, 2016). The Vermicompost is an eco-friendly, low cost, and effective way to recycle agricultural and kitchen waste. It is a mixture of earthworm castings, organic materials, humus, and other organisms. It has been advocated in integrated nutrient management systems in field crops due to its rich source of macro and micronutrients, vitamins, enzymes, antibiotics, and growth hormones. It also improves fertilizer and water use efficiency even better than FYM (Singh and Agarwal, 2005). Organics, inorganic and bio-fertilizers are essential to raise the crop yield. Vermicomposting of non-toxic biodegradable matter produces a stabilized humus like product known as vermicompost, which has a great potential as soil amendment. Vermicompost is a good soil conditioner that is rich in NPK, micronutrients, and growth hormones. Vermicompost application to soil also increases microbial populations and activities that further influence nutrient cycling, production of plant growth-regulating materials, and build up plant resistance to pathogen and nematode attacks (Gopal *et al.*, 2009). Composting waste and using it in agriculture is the most economical way to deal with it. Worldwide, there is a systematic improvement in the methods of treating waste and then recycling it for use in technological processes. This approach can be applied in a circular economy in which the value of raw materials and finished products can be preserved for as long as possible, minimizing the amount of waste (Szulc *et al.*, 2021). In agriculture, production is massive every year, but millions of tons of agricultural products are lost during the agricultural process (Serpil *et al.*, 2012). To achieve maximum and quality production, fertilizers are used. The necessary elements are absorbed by plants from fertilizers. This is called plant nutrition. In permanent agriculture, the nutrient content is reduced to compensate for this excessive use of artificial fertilizers [6]. With excessive use, soils are salinized, heavy metals come into contact with humans, and nitrates accumulate in water sources. They have harmful effects on the environment. Instead of artificial

57 fertilizers, bio-fertilizers are used for mass and quality food production. In this case, agricultural waste  
 58 produces bio-fertilizer and energy. Composting is a biological process in which microorganisms  
 59 decompose organic matter and lower the carbon-nitrogen ratio of the substrate. It is generally prepared  
 60 from organic waste material such as crop residue, household waste etc. this research focus the using  
 61 of different bio agent on agricultural waste.

## 62 2. METHODOLOGY:

63 The experiment was conducted in the Department of soil science and agriculture chemistry,  
 64 JNKVV, Jabalpur. Enriched vermicompost made through earthen pots their dimension was height  
 65 (120cm), upper diameter (60cm) and lower diameter (30cm) and the enriched vermicompost material  
 66 composted made through biowaste of soybean stover (dry matter) and fresh cow dung with earthworms  
 67 (*Eisenia fetida sp.*) under 6 treatments of biofertilizer combination with 4 replications and the data was  
 68 statically analysis by RBD design (Gomez and Gomez, 1984). Vermicompost made by continue two  
 69 years (2022 or 2023) completely in *Rabi* season. This experimental trial has using different materials  
 70 and technology that is following heading listed below.

### 71 2.1 Collection of bio wastes, bio agents and earthworms

72 Soybean stover and cow dung were gathered from study campus Breeder Seed Research Unit  
 73 and Dairy Research Farm respectively, for use in the current study as vermicomposting substrates.  
 74 However, bio fertilizers and earthworms were obtained from the Microbes Research Production Centre  
 75 and Dairy Research Farm of the vermicompost unit at JNKVV, Jabalpur, Madhya Pradesh respectively.  
 76 The dry matter contents of soybean stover and fresh cow dung were determined using the oven dry  
 77 method at 105 °C and were 67.8, 22.3%, respectively.

### 78 2.2 Partial decomposition after adding of bio fertilizers and earthworms in pot<sup>-1</sup>

79 The earthworms have a completely unique capacity to transform degradable bio wastes into  
 80 precious composts. However, those wastes need to be partly decomposed bio inoculate and launch of  
 81 earthworms throughout the start of decomposition technique can be survival and improvement of  
 82 earthworms. Before pre decomposition, dung was turned into delivered as additive in reputable  
 83 substrata in an identical proportion (1:1) which allowed to decompose for 28 days. After pre-  
 84 decomposition technique Rhizobium, PSB, KSB and Trichoderma have been injected discretely @ 10  
 85 g kg<sup>-1</sup> every pot. Desired moisture stage of 70-80% turned into maintained with inside the decomposing  
 86 bio-waste with normal watering on the price of 4 liter water for 8 kg of substrata at 7 days periods  
 87 throughout the partial decomposition (28 days) duration the partial decomposition was calculated  
 88 through following formula.

89 
$$\text{Partial Decomposition (kg)} = \text{Fresh weight of substrata (at initial)} - \text{Oven dry weight of substrata (after 28 days)}$$

### 90 2.4 Duration of vermicomposting

91 The total number of days required to complete the composting process, including 28-days for  
 92 partial decomposition, were recorded as vermicompost duration for every treatment.

### 93 2.5 Conversion rate of vermicompost

94 The vermicomposting rate is the daily conversion of bio waste into decomposed organic matter  
 95 by bio inoculants and earthworms were recorded in the final stage of vermicomposting. In fact, the  
 96 amount of vermicompost produced during a treatment period is known as the conversion rate and is  
 97 determined for each treatment according to a given method.

98 
$$\text{Conversion rate (g/day)} = \frac{\text{Quantity of vermicompost (weight)}}{\text{Number of days it took to convert the substrate into vermicompost}}$$

### 99 2.6 Recovery of vermicompost

100 Vermicompost recovery was calculated based on the final dry matter obtained from each  
 101 treatment from the total dry matter used for decomposition according to the following formula:

102 
$$\text{Recovery of vermicompost (\%)} = \frac{\text{Dry weight of vermicompost}}{\text{Dry weight of substrata}} \times 100$$

## 103 3. RESULT AND DISCUSSION

### 104 3.1 Partial decomposition of bio-waste

105 The data table 1 revealed that the partial decomposition of bio-waste shows non-significantly  
 106 in the both year and statistical pooled data. However, The partial decomposition maximum observed  
 107 in treatment E<sub>4</sub> (Rhizo.+PSB+KSB) *i.e.* 7.11 kg found in both year and pooled data followed by E<sub>3</sub> -  
 108 Rhizobium + PSB (7.12, 7.24 and 7.18 kg) letter on E<sub>4</sub>-Rhizobium + PSB (7.13, 7.27 and 7.20 kg) and  
 109 the minimum partial decomposition found in E<sub>1</sub> - control *i.e.* 7.25, 7.31 and 7.28 in the year 2021,2022  
 110 and pooled data respectively. Wiharyanto *et.al.*, 2018 reported that the partial decomposition in addition  
 111 of local microorganisms mixture of food waste can accelerate the process of compost maturity.

112 **Table 1. Partial Decomposition (kg) of bio-waste under different bio agent with earthworm after**  
 113 **28 days**

Treatments	2021	2022	Pooled
------------	------	------	--------

<b>E<sub>1</sub> (Control)</b>	7.25	7.31	7.28
<b>E<sub>2</sub> (Rhizobium)</b>	7.16	7.22	7.19
<b>E<sub>3</sub> (Rhizobium + PSB)</b>	7.12	7.24	7.18
<b>E<sub>4</sub> (Rhizobium + KSB)</b>	7.13	7.27	7.20
<b>E<sub>5</sub> (Rhizo. + PSB + KSB)</b>	7.11	7.11	7.11
<b>E<sub>6</sub> (Rhizo.+PSB+KSB+Tricho)</b>	7.18	7.21	7.20
<b>SEm±</b>	0.23	0.13	0.19
<b>CD (P=0.05)</b>	0.69	0.37	0.55


### 3.2 Production of vermicompost under different treatments with duration of completion of vermicomposting

Production of vermicompost and vermicompost completion under different treatments were presented in Table 2 and Figure 1. The results were found significantly on production and duration of completion of vermicompost. The production of vermicompost found in E<sub>6</sub> - Rhizo. + PSB+ KSB+ Tricho (5.58, 5.63 and 5.61 kg plot<sup>-1</sup>) within respective duration of vermicomposting (42, 40 and 41 days) followed by E<sub>5</sub> - Rhizo. + PSB + KSB (5.20, 5.25 and 5.23 kg plot<sup>-1</sup>) within 52, 54 and 53 days after completion of decomposition of bio- waste significantly superior to E<sub>2</sub> (Rhizobium) *i.e.* 4.39, 4.45 and 4.42 kg plot<sup>-1</sup> within 66, 63 and 64.50 days and E<sub>1</sub> (Control) *i.e.* 4.23, 4.25 and 4.24 kg plot<sup>-1</sup> within 79, 74 and 76.50 days of bio-waste decomposition. While, the treatments E<sub>3</sub>-Rhizobium + PSB (4.90, 4.95 and 4.93 kg plot<sup>-1</sup>) and E<sub>4</sub>-Rhizobium + KSB (4.92, 4.87 and 4.90 kg plot<sup>-1</sup>) within E<sub>3</sub> (55, 57 and 56 days) and E<sub>3</sub> (57, 55 and 56 days) respective duration of vermicomposting. Addition of the bio-fertilizers altered the soil physicochemical properties due to the microbial activity from the bio-fertilizers. Zea Mays showed an enhanced growth and reproduction rate upon application of the bio-fertilizers. Vermicomposting can be used as a waste corn pulp management strategy and at the same time obtain bio-fertilizers reported by Manyuchi, 2013. Under paddy straw based vermicomposting reported by Vijaya *et al.*, 2008 and similarly observation obtained in different agricultural crops reported by researcher that is Kalantari *et al.*, 2010, Kumari *et al.*, 2011 and Ansari and Sukhraj, 2010.

**Table 2. Production of Vermicompost (Kg pot<sup>-1</sup>) under various treatments.**

Treatments	Vermicompost Production (kg pot <sup>-1</sup> )			Duration (days) of completion vermicomposting		
	2021	2022	Pooled	2021	2022	Pooled
<b>E<sub>1</sub> (Control)</b>	4.23	4.25	4.24	79.00	74.00	76.50
<b>E<sub>2</sub> (Rhizobium)</b>	4.39	4.45	4.42	66.00	63.00	64.50
<b>E<sub>3</sub> (Rhizobium + PSB)</b>	4.90	4.95	4.93	55.00	57.00	56.00
<b>E<sub>4</sub> (Rhizobium + KSB)</b>	4.92	4.87	4.90	57.00	55.00	56.00
<b>E<sub>5</sub> (Rhizo. + PSB + KSB)</b>	5.20	5.25	5.23	52.00	54.00	53.00
<b>E<sub>6</sub> (Rhizo.+PSB+KSB+Tricho)</b>	5.58	5.63	5.61	42.00	40.00	41.00
<b>SEm±</b>	0.28	0.26	0.27	2.62	2.55	2.58
<b>CD (P=0.05)</b>	0.82	0.77	0.79	7.72	7.51	7.62

Figure 1. Production of Vermicompost (Kg pot<sup>-1</sup>) and Duration (days) of completion vermicomposting under various treatments.



### 3.3 Vermicompost recovery percentage under various treatments

The data Table 3 and Figure 2 shows that vermicompost recovery percentage on E<sub>6</sub>- Rhizo.+ PSB+ KSB+ Tricho (69.75, 70.38 and 70.06 %) and E<sub>5</sub> - Rhizo. + PSB + KSB (65.00, 65.63 and 65.31%) were significantly superior to E<sub>2</sub> –Rhizobium (54.88, 55.63 and 55.25 %) and E<sub>1</sub>-Control (52.88, 53.13 and 53.00 %). While, the treatments E<sub>3</sub> -Rhizobium + PSB (61.25, 61.88 and 61.56 %) and E<sub>4</sub> - Rhizobium + KSB (61.50, 60.88 and 61.19 %) were found partly in the year of 2021 and 2022 as well as statically pooled analysis. The recovery percentage depend on dry vermicompost production *i.e.* increase with the increasing the different bio-fertilizer numbers.

**Table 3. Vermicompost recovery percentage and Conversion rate of bio-waste (g day<sup>-1</sup>) under various treatments.**


Treatments	Vermicompost Recovery (%)			Conversion rate of waste (g day <sup>-1</sup> )		
	2021	2022	Pooled	2021	2022	Pooled
E <sub>1</sub> (Control)	52.88	53.13	53.00	55.00	59.15	57.07
E <sub>2</sub> (Rhizobium)	54.88	55.63	55.25	69.80	73.76	71.78
E <sub>3</sub> (Rhizobium + PSB)	61.25	61.88	61.56	92.34	89.70	91.02
E <sub>4</sub> (Rhizobium + KSB)	61.50	60.88	61.19	90.97	93.67	92.32
E <sub>5</sub> (Rhizo. + PSB + KSB)	65.00	65.63	65.31	103.16	99.85	101.50
E <sub>6</sub> (Rhizo.+PSB+KSB+Tricho)	69.75	70.38	70.06	134.90	142.88	138.89
SEm±	3.47	3.26	3.37	6.77	6.51	6.64
CD (P=0.05)	10.24	9.62	9.93	19.96	19.20	19.58

### 3.4 Conversion rate of bio-waste (g day<sup>-1</sup>) under different treatments

The data have present in Table 3 and Figure 2 Bio-waste Conversion rate estimated that significantly in the both year and pooled analysis. The heights conversion rate computed in E<sub>6</sub>- Rhizo.+PSB+KSB+Tricho (134.90, 142.88 and 138.89 g day<sup>-1</sup>) *i.e.* was significantly superior to all bio agent based treatments. While, E<sub>5</sub> - Rhizo. + PSB + KSB (103.16, 99.85 and 101.50 g day<sup>-1</sup>), E<sub>4</sub> - Rhizo.+ KSB (90.97, 93.67 and 92.32 g day<sup>-1</sup>) E<sub>3</sub>- Rhizo.+ PSB (92.34, 89.70 and 91.02 g day<sup>-1</sup>) were found significant to E<sub>2</sub>- Rhizo. (69.80, 73.76 and 71.78 g day<sup>-1</sup>) and E<sub>1</sub>- control (55.00, 59.15 and 57.07 g day<sup>-1</sup>) in the year of 2021, 2022 and statically pooled analysis data. The conversation rate might be dependent on the bio-fertilizer effected to deteriorate the cellulose tissue in soybean straw so that earthworm easy to digest the soybean straw. Similar finding reported by Tsai *et al.*,2007, Yong *et al.*,2021, Dikko *et al.*,2019, Srivastava *et al.*,2024.

161  
162

**Figure 2. Graph of Vermicompost recovery percentage and Conversion rate of bio-waste (g day<sup>-1</sup>) under various treatments.**



163  
164

#### 4. CONCLUSION

The present studies concluded on the basis of following observation taken during experimental period *i.e.* partial decomposition, recovery percentage and production of vermicompost with recovery percentage. The partial decomposition under statically pooled data found in higher on treatment E<sub>6</sub> (Rhizo.+PSB+KSB+Tricho) and E<sub>4</sub> (Rhizobium + KSB) after 28 days after application enriched biofertilizer. Moreover, vermicompost production maximum in the E<sub>6</sub> (Rhizo.+PSB+KSB+Tricho) *i.e.* 5.58, 5.63, and 5.61 with lowest days (42, 40 and 41) and minimum production (4.23, 4.25, 4.24) found with higher days (79, 74, 76.50) on the E<sub>1</sub>(Control) after complete decomposition of bio waste respective in 2021, 2022 and statically pooled data. Simultaneously, vermicompost recovery percentage and biowaste conversion rate maximum in the enriched biofertilizers of E<sub>6</sub> (Rhizo.+PSB+KSB+Tricho) followed by E<sub>5</sub> (Rhizo. + PSB + KSB) in compare to E<sub>1</sub>(Control). This experiment helpful to maintain the agriculture bio-waste material for decomposition and it also increased the fertility levels on the soil.

#### ACKNOWLEDGEMENTS

I express my gratitude and honors to my chairmen and my advisory committee for his able guidance, constant inspiration, and valuable suggestion and critical during entire course of present studies and preparation manuscript. I extend my heartiest thanks to HOD sir Dr. P.S. Kulhare, Dr. B. K. Dixit, Dr. H. K. Rai, Dr. Y. M. Sharma, Dr. S. S. Baghel, Dr. B. S. Dubedi, Dr. R. K. Sahu, Dr. A. K. Upadhyay, Dr. G. S. Tegore and all department members soil science and agricultural chemistry, CoA, JNKVV, Jabalpur. For provide valuable resources for vermicompost production, field works, research works and laboratory works. There is no funding agency for support this research.

#### AUTHORS' CONTRIBUTIONS

The K.K.P. (1<sup>st</sup> author) carried out the experimental during the research period, D.S.S.B. (2<sup>nd</sup> author) given guidance for above investigation, the D.S.B.A. (3<sup>rd</sup> author), D.H.K.R. (4<sup>th</sup> author), D.R.K.S. (5<sup>th</sup> author), and D.U.S. (6<sup>th</sup> author) supporting for laboratory guidance and valuable suggestion for the research after that A.K.S. (7<sup>th</sup> author) and P. (8<sup>th</sup> author) were supported in data collection, vermicompost testing analysis and lot of moral supported for research.

191  
192  
193

194 **Disclaimer (Artificial intelligence)**

195 Author(s) hereby declares that NO generative AI technologies such as Large Language  
196 Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or  
197 editing of this manuscript.

## 198 REFERENCES

- 199 Ansari, A. A., and Jaikishun, S., (2010) "An investigation into the vermicomposting of sugarcane  
200 bagasse and rice straw and its subsequent utilization in cultivation of *Phaseolus vulgaris* L. In  
201 Guyana", *American-Eurasian J. Agriculture and Environmental Science*, 8 (6), 666-671.
- 202 Dikko, A., Mahmud, Mustapha, Y., Abubakar, D., and Yusuf, A., (2019) STUDIES ON CONVERSION  
203 OF SOLID WASTE TO BIOFERTILIZER BY VERMICOMPOSTING. *FUW Trends in Science &*  
204 *Technology Journal*, www.ftstjournal.com e-ISSN: 24085162; p-ISSN: 20485170; 4 (1): 137 –  
205 139.
- 206 Fayaz, A., Ahmad, L., and Singh, P. (2016). Vermi-composting: an effective technique to recycle wastes  
207 into valuable organic fertilizer: a review. *Journal of Pure and Applied Microbiology*, 10(2), 1109.  
208 [https://link.gale.com/apps/doc/A481650197/AONE?u=anon~5c8ff683&sid=googleScholar&xid](https://link.gale.com/apps/doc/A481650197/AONE?u=anon~5c8ff683&sid=googleScholar&xid=6b493c7a)  
209 [=6b493c7a](https://link.gale.com/apps/doc/A481650197/AONE?u=anon~5c8ff683&sid=googleScholar&xid=6b493c7a)
- 210 Gomez, K.A., and Gomez, A.A., (1984) *Statistical Procedures for Agricultural Research*. 2nd Edition,  
211 John Wiley and Sons, New York, 680 p.
- 212 Gopal, M., Gupta, A., Sunil, E., Thomas, V.G., (2009) Amplification of plant beneficial microbial  
213 communities during conversion of coconut leaf substrate to vermicompost by *Eudrilus* sp. *Curr*  
214 *Microbiol.* 59:15–20. doi: 10.1007/s00284-009-9388-9.
- 215 Kalantari, S., Hatami, S., Ardalan, M. M., Alikhani, H. A., and Shorafa, M., (2010) "The effect of compost  
216 and vermicompost of yard leaf manure on growth of corn", *African Journal of Agricultural*  
217 *Research*, 5 (11), 1317-1323.
- 218 Kumari, M., Kumar, S., Chauhan, R.S., and Ravikanth, K., (2011) "Bioconversion of herbal industry  
219 waste into vermicompost using an epigeic earthworm *Eudrilus Eugeniae*", *Waste Management*  
220 *and Research*, 29 (11), 1205-1212.
- 221 Maghsoudi, M.S., Shokouhyar, S., Khanizadeh, S., and Shokoohyar, (2023). Towards a taxonomy of  
222 waste management research: An application of community detection in keyword network J.  
223 *Clean. Prod.*, 401 Article 136587.
- 224 Manyuchi, M., (2013). Production of Bio-Fertilizers from Vermicomposting of Waste Corn Pulp Blended  
225 with Cow Dung as a Solid Waste Management Approach.
- 226 Raza, S.T., Zhu, B., Yao, Z.Y., Wu, J.P., Chen, Z., Ali, Z., and Tang, J.L., (2023) Impacts of  
227 vermicompost application on crop yield, ammonia volatilization and greenhouse gases  
228 emission on upland in Southwest China *Sci. Total. Environ.*, 860 Article 160479.
- 229 Serpil S., and Scribe B., (2012) *International Journal of Environmental Science and Development*, 3 (1)  
230 An Agricultural Pollutant: Chemical.
- 231 Singh, R., and Agarwal, S.K., (2005). Effect of levels of farmyard manure and nitrogen fertilization on  
232 grain yield and use efficiency of nutrients in wheat (*Triticum aestivum*). *Indian Journal of*  
233 *Agricultural Sciences*. 75. 408-413.
- 234 Srivastava, S., Singh, P., Barbora, L., Baruah, D., Saikia, R., Mohanty, K., and Kalita, P., (2024)  
235 Performance assessment of innovative waste management system developed for the  
236 production of bio-fertilizer, *Sustainable Chemistry for the Environment*, 7, 100148, ISSN 2949-  
237 8392, <https://doi.org/10.1016/j.scenv.2024.100148>. ([https://www.sciencedirect.com/science/arti](https://www.sciencedirect.com/science/article/pii/S2949839224000919)  
238 [cle/pii/S2949839224000919](https://www.sciencedirect.com/science/article/pii/S2949839224000919))
- 239 Szulc, W., Rutkowska, B., Gawroński, S., Wszelaczyńska, E., (2021) Possibilities of Using Organic  
240 Waste after Biological and Physical Processing-An Overview. *Processes*, 9, 1501.  
241 <https://doi.org/10.3390/pr9091501>
- 242 Tsai, S., Liu, C., and Yang, S., (2007) Microbial conversion of food wastes for biofertilizer production  
243 with thermophilic lipolytic microbes, *Renewable Energy*, 32 (6): 904-915, ISSN 0960-1481,  
244 <https://doi.org/10.1016/j.renene.2006.04.019>. ([https://www.sciencedirect.com/science/article/pi](https://www.sciencedirect.com/science/article/pii/S0960148106001030)  
245 [i/S0960148106001030](https://www.sciencedirect.com/science/article/pii/S0960148106001030))
- 246 Vijaya, D., Padmadevi, S.N., Vasandha, S., Meerabhai, R.S., and Chellapandi, P., (2008) "Effect of  
247 vermicomposting coirpith on the growth of *Andrographis Paniculata*", *Journal of Organic*  
248 *Systems*, 3 (2), 51-56.
- 249 Wiharyanto, O., Mochtar, H., Purwono, P., (2018) Decomposition of food waste using bulking agent and  
250 bio-drying technology. *E3S Web of Conferences*. 73. 05013. 10.1051/e3sconf/20187305013.
- 251 Yong, Z. J., Bashir, Mohammed J.K., and Hassan, M. S., (2021) Biogas and biofertilizer production from  
252 organic fraction municipal solid waste for sustainable circular economy and environmental  
253 protection in Malaysia, *Science of The Total Environment*, 776,145961, ISSN 0048-9697,

254 <https://doi.org/10.1016/j.scitotenv.2021.145961>.([https://www.sciencedirect.com/science/article](https://www.sciencedirect.com/science/article/pii/S0048969721010287)  
255 [/pii/S0048969721010287](https://www.sciencedirect.com/science/article/pii/S0048969721010287))

256 **ABBREVIATIONS**

257 Rhizo= Rhizobium, PSB= Phosphorus solubilizing bacteria, KSB= Potassium solubilizing  
258 bacteria, Tricho.= Trichoderma, *i.e.* = that is

UNDER PEER REVIEW