

Infestation of mulberry leaves by Leaf roller/webber-*Diaphania pulverulentalis* (Hampson) and *Glyphodes pyloalis*(Walker) and its management strategies

Abstract

The mulberry is a perennial, fast growing plant mainly cultivated worldwide for rearing of mulberry silkworm, *Bombyx mori*. The *B. mori* which is a monophagous insect feeds only on mulberry leaves for its proper growth and development. The mulberry plant during its growth is susceptible to various diseases viz., leaf spot, powdery mildew, leaf rust etc. and pests viz., leaf roller, Bihar hairy caterpillar, mealy bugs etc. in the field which deteriorates its leaf quality and quantity. Among these pests, defoliators attain significant importance as they cause damage up to 12-25% thereby making mulberry leaves unfit for silkworm rearing. The leaf rollers are major pests currently causing devastation to mulberry plants both in India and other parts of the world. Thus, in this review an attempt has been made regarding new insights about the leaf roller pests and devise more sustainable management of this pest through integrated pest management (IPM) strategy to control it, so that the overall cocoon crop production will be enhanced worldwide.

Keywords: leaf roller, *D. pulverulentalis*, infestation, pest, IPM, mulberry

Introduction

The mulberry (*Morus spp.*) is an important valuable plant and has commercial significance as it is the only sole food source of *B. mori*. Its contribution is around 38.20% for the successful harvesting of cocoon crop (Miyashita, 1986). It is fast growing plant with deep root system present in order Urticales/Rosales and family Moraceae. The feeding of *B. mori* on good quality mulberry leaves results in its healthy growth and development and eventually enhanced silk production (Islam *et al.*, 2022a, 2022b; Islam *et al.*, 2023; Islam *et al.*, 2024). During its growth it also faces many challenges as it is subjected to various kinds of diseases and pests which hampers its leaf production both qualitatively and quantitatively. The leaf roller, *D. pulverulentalis* is a serious pest in Southern India and also is a potential carrier of *Nosema bombycis*, which causes deadly pebrine disease to the silkworms as pebrine spores can get easy entry in the rearing rooms through contaminated mulberry leaves (Ifat *et al.*, 2011). Similarly, Watanabe *et al.* (1988) found leaf roller *G. pyloalis* as the alternate hosts

of densoviruses and picornaviruses. The *D. pulverulentalis* was reported from Karnataka in 1995 and then spread to Tamil Nadu, Andhra Pradesh and Kerala and its infestation causes leaf yield loss up to 12.8% having average incidence of 21.77% causing severe damage to sericulture industry (Siddegowda *et al.*, 1995; Geetha Bai *et al.*, 1997; Rajadurai *et al.*, 1999). Manjunath Gowda *et al.* (2005) stated that 37 parasitoids and 6 predators are known naturally controlling the leaf roller infestation at different stages, also 10 entomopathogenic fungi were reported regulating the incidence of leaf roller naturally (Srinivasa Gowda *et al.*, 2000). Around 300 insect and non-insect pest species are known causing attack to mulberry. The insect pests may be sapsuckers, defoliators, borers and residing in soil like termites. The infestation by major insect pests like leaf roller, pink mealy bug and thrips severely reduces nutrient content of leaves by hampering photosynthesis which eventually reflects in the poor growth of plant (Kotikal *et al.*, 1982; Biradar, 1989; Srinivasa Gowda, 2004; Sakthivel *et al.*, 2019). The yield loss caused by major insect pests of mulberry namely Bihar hairy caterpillar (*Spilosoma obliqua*), leaf roller (*D. pulverulentalis*) and mealy bugs (*Maconellicoccus hirsutus*) is up to 25-30%, 12-15% and 10-15% respectively (Manjunath *et al.*, 2000). Mulberry leaf roller is a major dangerous pest of mulberry plant in India causing heavy infestation in the states of Karnataka, Andhra Pradesh, Tamil Nadu, Kashmir, Assam etc. (Mathur, 1980; Chakraborty, 2005; Dandin and Giridhar, 2010; Illahi *et al.*, 2013). Keeping in view the status of this pest the current review elucidated this pest in more detail like its biology, nature of damage, period of occurrence and subsequent management strategies for controlling this pest so that the overall silk production will be increased.

Life cycle and management strategy of *D. pulverulentalis* through IPM

D. pulverulentalis is a key seasonal defoliating pest and its maximum population buildup is recorded during September-November and there is a linear relationship between the decrease in temperature and increase in relative humidity resulting in increased pest population (Kumari *et al.*, 2021). The biochemical components in six mulberry varieties viz., M5, Mysore local, MR2, S54, S36 and V1 were analysed infested by *D. pulverulentalis*. It was revealed that the biochemical composition of mulberry leaves was reduced almost in all the varieties (Mahadeva and Nagaveni, 2011). Bhagyamma and Vijaya Kumari (2022) after carrying out study on three mulberry varieties viz., V1, Mysore local and S36 infested by *D. pulverulentalis* found that the photosynthetic pigments in the leaves like total chlorophyll and

carotenoids decreased by 40.17% and 40.85% respectively. Further, the biomolecule components like proteins, carbohydrates and starch decreased by 24.62%, 47.09% and 16.09% respectively. However, the phenol content increased by 57.39% than control. **Life Cycle:** The eggs of *D. pulverulentalis* (Order-Lepidoptera, Family-Pyralidae) are pale yellow in colour laid singly on the lower surface along the leaf vein of the mulberry leaves having fecundity around 80-150. The egg period lasts for 3-4 days. The total larval period is 12-15 days having 5 larval instars. The first larval instar is minute and is fluorescent yellow in colour. The second larval instar is light yellowish orange in colour having small sub-median black spots enclosed by white patches. The third larval instar is deep orange in colour and its dorsal, mid dorsal lines and spots are clearly visible. The fourth larval instar is dark greenish brown in colour and the fifth larval instar is dark pinkish brown in colour. The pupal period lasts for 7-9 days and are dark brown in colour and pupates in dried leaves/soil. The adults are yellowish grey in colour having longevity of 8 and 11 days for male and female respectively. The total lifecycle is completed in 17-24 days (Mahadeva, 2018). **Symptoms:** The early instar larvae occupy the apical parts of the mulberry shoot and feed on the young, tender leaves. The larvae reside on the leaves by forming web and fold the leaves and thereby making shelter for itself. The leaf margins of apical leaves are rolled and tied by the larval web wherein they live. Occasionally 1 or 2 leaves are rolled into cup shape with the web secreted from the larvae which remains inside and hence this pest is called as leaf roller/leaf webber. The lower leaves of the infested plants become soiled due to faecal matter of larva.

IPM of *D. pulverulentalis*: (a) **Mechanical control:** The infested portions are trimmed and collected in a polythene bag and then destroyed by burning. Flood irrigation and deep ploughing helps in killing of pupae of the pest residing in the soil. Light traps can be used to attract and kill adults of pest. (b) **Chemical control:** Foliar application of 0.076% DDVP 76% EC (one ml in one litre of water) 10 days after pruning/leaf harvest is helpful. If the infestation continues then further 2-3 sprays are needed. The leaf for silkworm rearing can be used after 7 days of DDVP application. The commercial neem pesticide (0.03% AZ) 0.05% can be sprayed having safe period of 10 days. (c) **Biological control:** The release of egg parasitoid, *Trichogramma chilonis* @ 1 lakh adults/acre in 4 split doses, larval parasitoid, *Bracon brevicornis* @ 200 adult wasps and ectopupal parasitoid, *Tetrastichus howardii* @ 1 pouch /acre is effective for controlling leaf roller *D. pulverulentalis* (Sakthivelet al., 2019; Kumar et al., 2024). The *Apanteles* sp. and *Chelonus* sp. are also reported to be parasitic on leaf roller (Geetha Bai et al., 1997; Srinivasa Gowda et al., 2001), also the

Calosoma sp. which is a larval predator predaes on leaf webber larvae (Annon., 1997). Furthermore, the aqueous leaf extracts of *Catharanthus roseus*, *Piper betle*, *Ocimum sanctum*, *Tagetes patula* and *Mentha piperita* are found to be effective against leaf roller larva. The insecticidal properties of various medicinal plants viz., *Lantana camara*, *Allium sativum*, *Zingiber officinale*, *Azadirachta indica* and *Vitex negundo* was studied against *D. pulverulentalis* and among these, lantana extract at lower concentration of 11% was found to be most effective against leaf roller (Maheswari and Govindaiah, 2017, 2018).

Life cycle and management strategy of *G. pyloalis* through IPM

The *G. pyloalis* (lesser mulberry pyralid) is a serious pest of mulberry in India, China, Japan, Malaysia, Pakistan, Uzbekistan, Burma and Korea (Madyarov *et al.*, 2006). Borgohain *et al.* (2015) carried out study on mulberry leaves infested by *G. pyloalis* and revealed that early first and second instar larvae fed by scraping green tissues and mesophyll layer leaving behind only transparent epidermis layer. The larvae (first and second instar) damaged up to 0.11-0.33 cm² area and full-grown larvae damaged up to 0.69-1.75 cm² area of mulberry leaf. **Life Cycle:** *G. pyloalis* (Order-Lepidoptera, Family-Pyralidae) is a holometabolic insect consisting of egg, larva, pupa and adult stages having much similarity to silkworm. The moth is nocturnal (Khosravia and Sendi., 2010) and lays around 200 eggs under side of leaf, the egg is pale yellow in colour, round and its size measures about 0.2mm. The eggs hatch within 5-6 days into larva. The larvae measure around 0.2-2cm in length and are slender, fusiform and segmented. Each larva has 3 pair of forelegs, 4 pairs of hind legs and a pair of caudal legs and then mature, transforming into brown pupae and then into moth. The whole life cycle takes about a month to complete. **Symptoms:** Under Kashmir climatic conditions, infestation is observed around July-October. After hatching the larva spin fine silky net of threads around themselves and eat leaves leaving behind transparent cuticular layer. The larval excreta are held in between fine silky threads, thus making leaves unsuitable for silkworms. After the pest damage, the nutrients in the leaves are lost and this in turn inhibits the growth of plants. **Mainly**, much damage is caused by 4th and 5th instar larva. The larvae feed inside the leaves after webbing the leaves together. The larvae skeletonise the mulberry leaf after eating all the green portions. The infested part of the leaf becomes dark brown in colour and is devoid of chlorophyll, moisture content, proteins, sugar etc. (Hassan and Mir., 2018). **IPM of *G. pyloalis*:** (a) **Mechanical control:** Hand pick the larval stages of pest and destroy it. Ensure deep ploughing and weeding in order to destroy the hibernating larva. Burn the diseased and fallen leaves in September-October. Ensure straw banding of trees and light trapping for

moths to control this pest. (b) **Chemical control:** Spraying of 0.04% DDVP on leaves can significantly reduce the infestation level of the pest. (c) **Biological control:** The *G. pyloalis* was reported as the most predominant defoliator pest and its natural enemies like hymenopteran parasitoids viz., *Apanteles obliquae*, Ichneumonid wasp, *Chelonus carbonator* and the coccinellid predator, *Cheilomeness exmaculata* proved to be active natural enemies against it (Sultana et al., 2023) (Table 1 & 2). Nighat et al. (2002) recorded two braconid larval parasitoids viz., *Apanteles glomeratus* and *Chelonus sp.* parasitizing the larvae of *G. pyloalis*, further parasitoids namely *Habrobacon hebetor*, *C. carbonator*, *Glyptapanteles sp.*, *Pristomerus sulci*, *Xanthopimpla sp.*, *Perilampus sp.*, *Campoletis sp.* and *Brachymeria lasus* could be used for the control of *G. pyloalis* under temperate conditions of Kashmir (Mitta et al., 2011, 2016) (Table 3). In a recent finding *G. pyloalis* was found as a host of *P. sulci* (Bhat et al., 2020).

Table 1: Natural enemies present on mulberry

| Natural enemies | Order | Family |
|-------------------------------------|-------------|---------------|
| <i>Apanteles obliquae</i> Wilkinson | Hymenoptera | Braconidae |
| <i>Chelonus carbonator</i> Marshall | Hymenoptera | Braconidae |
| <i>Megaselia scalaris</i> | Diptera | Phoridae |
| <i>Cheilomeness exmaculata</i> | Coleoptera | Coccinellidae |
| <i>Disophryssp.</i> | Hymenoptera | Braconidae |
| Tachinid fly | Diptera | Tachinidae |
| Ichneumonid wasp | Hymenoptera | Ichneumonidae |
| Spider | Araneae | Salticidae |

Table 2: Defoliator pests present on mulberry

| Common name | Scientific name | Order | Family |
|----------------------|--|-------------|--------------------------|
| Mulberry leaf roller | <i>Glyphodes pyloalis</i> Walker | Lepidoptera | Pyalidae |
| Common cutworm | <i>Spodoptera litura</i> Fabricius | Lepidoptera | Noctuidae |
| Tussock caterpillar | <i>Euproctis fraterna</i> Moore | Lepidoptera | Erebidae |
| Spanworm | <i>Hemerophillaa trilineata</i> Butler | Lepidoptera | Geometridae |
| Green weevil | <i>Myloccerus viridanus</i> Fabricius | Coleoptera | Curculionidae |

Sultana et al., 2023

Table 3: Parasitoids of *G. pyloalis*

| Name | Family | Order | Stage attacked | Period of activity |
|--|---------------|-------------|-----------------------------|--------------------|
| <i>Apanteles obliquae</i> Wilkinson | Braconidae | Hymenoptera | Larval instars | July - Oct. |
| <i>Bracon hebetor</i> Say | Braconidae | Hymenoptera | Larval instars | Sept. |
| <i>Chelonus carbonator</i> Marshall | Braconidae | Hymenoptera | Late instars | July - Oct. |
| <i>Pristomerus sulci</i> M. & K. | Ichneumonidae | Hymenoptera | Late instars / Pupae | Aug. |
| <i>Xanthopimpla sp.</i> | Ichneumonidae | Hymenoptera | Late instars / Pupae | Aug. |
| <i>Campoletis sp.</i> | Ichneumonidae | Hymenoptera | Late larval instars / Pupae | Aug. - Sept. |
| <i>Brachymeria lasus</i> (Walker) | Chalcididae | Hymenoptera | Pupal | July - Aug. |

Mittal *et al.*, 2016

Conclusion

The incidence of pests poses a great threat to mulberry plants and deteriorates its leaf quality and eventually results in poor cocoon production. The outbreak of pest attack like defoliators on mulberry plants incur heavy losses to silkworm farmers and in turn it becomes imperative to design sustainable management strategies to control it through IPM. As the wide use of chemicals is not a long-term solution to control pests as they harm the environment by leaving more residues, so utilising biological means like its natural enemies is best for naturally maintaining the balance in the mulberry ecosystem. So, in order to check the leaf roller population like *D. pulverulentalis* and *G. pyralis*, multipronged approach using IPM should be followed for better efficiency to boost the silk industry.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Competing interests

Authors declare that no competing interests exist

References

- Anonymous(1997).Bionomics and integrated control of of *Spilosoma obliqua* and other important lepidopteran mulberry pests.Annual report,Central Sericultural Research and Training Institute,Pampore,India pp:86-91.
- Bhagyamma, C.T.,& Kumari, N.V. (2022). Studies on the impact of leaf webber *Diaphania pulverulentalis* on different mulberry varieties and silkworm *Bombyx mori* L. *Indian Journal of Agricultural Research*, 56(3), 351-356. <http://dx.doi.org/10.18805/IJARE.A-5887>
- Bhat, D.M., Sheikh, M.A.,& Khan, S.A. (2020). An updated checklist of Ichneumonid parasites (Hymenoptera: Ichneumonidae) associated with insects/insects pests of Agricultural, Horticulture and other Economically important plants in the union territory of Jammu & Kashmir. *Int.J.Curr.Microbial.App.Sci.*, 9(4), 2266-2277. <https://doi.org/10.20546/ijcmas.2020.904.271>
- Biradar, NA. (1989). Faunistic study of arthropods infesting mulberry (*Morus* spp.) and biology of *Euproctis fraterna* (Lepidoptera: Lymantridae) M. Sc. (Seri.) Thesis, UAS, Bengaluru, 120.
- Borghain,A., Battacharjee,J., Dutta,L.C., Bhattacharya, B., Singha,A. (2015).Mulberry Leaf Damage Caused by Leaf Roller, *Glyphodes Pyloalis* Walker.*J. Exp. Zool. India*, 18(1), 313-314.
- Chakraborty, I.R. (2005). Package and Practices of Muga, Eri and Mulberry Sericulture for North Eastern region of India (CM & RTI), CSB –Lahdoigarh, Assam.
- Dandin, S.B., & Giridhar, K. (2010). Mulberry diseases and pest control measures. Handbook of Sericulture Technologies, CSB, pp 175-177.
- Geethabai,M.,Marimadaiah,B.,Narayana Swamy,K.C., &Rajgopal.(1997).An outbreak of leaf roller pest *Diaphania* (*Margorania*)*pulverulentalis*Hampson)on mulberry in Karnataka *Geobios news reports*,16(2), 73-79.
- Hassan, F., & Mir, M.A.(2018). *Glyphodes* Infestation in Mulberry, *Int. J. Pure App. Biosci*, 6(1), 1195-1197. <http://dx.doi.org/10.18782/2320-7051.6039>
- Ifat, B., Sharma, S.D.,& Shabir, A.B. (2011). Screening of different insect pests of mulberry and other agricultural crops for microsporidian infection. *Int. J. Biotechnol. Mol. Biol. Res.*, 2(8), 138-142.

- Illahi, I., Mittal, V., Ramegowda, G.K., & Dhar, A. (2013). Leaf webber a serious insect pest of mulberry in Kashmir. *Indian Silk*, 3(51), 11, 8-10.
- Islam, T., Bhat, S.A., Malik, F.A., Khan, F.A., Mir, S.A., Nazir, N., & Wani, S.A. (2022b). Evaluation of some mulberry genotypes for nutritional consumption parameters of silkworm, *Bombyx mori* L. under temperate conditions of Kashmir, India. *Plant Archives*, 22(2), 136-139. <http://dx.doi.org/10.51470/PLANTARCHIVES.2022.v22.no2.024>
- Islam, T., Bhat, S.A., Malik, F.A., Khan, F.A., Mir, S.A., Nazir, N., & Wani, S.A. (2022a). Rearing of silkworm, *Bombyx mori* L. on different mulberry genotypes and its impact on post cocoon parameters. *Plant Archives*, 22(2), 380-382. <http://dx.doi.org/10.51470/PLANTARCHIVES.2022.v22.no2.065>
- Islam, T., Bhat, S.A., Malik, F.A., Wani, S.A., Khan, F.A., Mir, S.A., & Nazir, N. (2023). Feeding of different Mulberry Varieties and its Impact on Silk Gland of Silkworm, *Bombyx mori* L. *Biological Forum – An International Journal*, 15(1), 488-492.
- Islam, T., Bhat, S.A., Qadir, J., Malik, F.A., Wani, S.A., Khan, F.A., Mir, S.A., & Nazir, N. (2024). Impact of leaves of different mulberry genotypes on nutritional efficiency of conversion parameters of silkworm, *Bombyx mori* L. *Journal of Scientific Agriculture*, 8, 66-68. doi: 10.25081/jsa.2024.v8.9051
- Kotikal, Y.K. (1982). Studies on pests of mulberry *Morus alba* L. with special reference to the black headed caterpillar, *Spilosoma obliqua* Walker (Lepidoptera: Arctiidae). M. Sc.(Agri.) Thesis, University of Agricultural Sciences, Bangalore, India.
- Kumar, K., Sanappa, B., Tanvi Rahman., Mahesh, R., Lalhmangaizuali., Malswamdwangi. (2024). *Bombyx mori* L.: Acts as host for *Tetrastichus howardi* for mass multiplication and its utilization for controlling leaf roller in mulberry. *JETIR*, 11(8), 797-803.
- Kumar, S., Ramakrishna Naika., & Nithya, B.N. (2021). Population dynamics of mulberry leaf roller, *Diaphania pulverulentalis* (Hampson) (Lepidoptera: Pyralidae). *Journal of Entomology and Zoology Studies* 2021, 9(4), 281-285.
- Madyarov, S.H.R., Khamraev, A.S.H., Otalbaev, D.O., Kamita, S.G. & Hammock, B.D. (2006). Comparative effects of wild and recombinant baculoviral insecticides on mp *Glyphodes pyloalis* wlk. and mulberry silkworm *Bombyx mori* L. *International*

Workshop on Silk Handcrafts Cottage Industries and Silk Enterprises Development in Africa, Europe, Central Asia and the Near East, & Second Executive Meeting of Black, Caspian seas and Central Asia Silk Association (BACSA) Bursa, Turkey.

- Mahadeva, A. (2018). Insect Pest Infestation, an Obstacle in Quality Mulberry Leaves Production. *Asian J. Biol. Sci.*, 1-12. <https://scialert.net/abstract/?doi=ajbs.2018.41.52>
- Mahadeva., & Nagaveni. (2011). Alterations in the biochemical components and photosynthetic pigments of mulberry (*Morus Spp.*) attacked by leaf – roller (*Diaphania pulverulentalis*) pest. *African Journal of Biochemistry Research*, 5(14), 365-372. <http://dx.doi.org/10.5897/AJBR11.091>
- Maheswari, M., & Govindaiah, G. (2017). Evaluation of different plant extracts for management of mealy bugs and leaf rollers in the mulberry field. *Journal of Entomology and Zoology Studies*, 5(6), 2081-2085.
- Maheswari, M., & Govindaiah, G. (2018). Larvicidal activity of a few botanical extracts against leaf roller in the mulberry. *Journal of Entomology and Zoology Studies*, 6(3), 1667-1671.
- Manjunath Gowda., Narayanaswamy, K.C., & Ninge Gowda, K.N. (2005). Status of leaf roller pests of mulberry. *J Ecobiol*, 17(3), 201-216.
- Manjunath, D., Satyaprasad, K., Katiyar, R. L., Rajaduari, S., Shekhar, M. A., Sen, A. K., & Data, R.K. (2000). Integrated pest management in sericulture. National Conference on Strategies for Sericulture Research and Development, 5 to 7 Nov. C. S. R. T. I. Mysore.
- Mathur, R.N. (1980). Biology of the mulberry defoliator-*Glyphodes pyloalis* (Wlk.) (Lepidoptera: Pyralidae). *Indian Forest Bulletin*, 273 1-6.
- Mittal, V., Irfan Illahi., Anil Dhar., & Khan, M.A. (2011). Natural enemies of mulberry pyralid, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae) in temperate climate of Kashmir. *Journal of Biological Control*, 25(1), 55-57. <https://doi.org/10.18641/JBC%2F25%2F1%2F40100>
- Mittal, V., Ramegowda, G.K., Irfan, I., Babulal., & Sharma, S.P. (2016). Biodiversity of Natural Enemies of Mulberry Leaf Webber, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae) in Kashmir Valley. *Indian Horticulture Journal*, 6(special), 50-53.

- Miyashita, Y. (1986). A report on mulberry cultivation and training methods suitable to bivoltine rearing in Karnataka. pp. 7-30.
- Nighat, M., Malik, M.A., Dar, H.U., Sofi, M.A., & Malik Farooq. (2002). Preliminary studies on the natural enemies of *Glyphodes pyloalis* Walker (Pyrilidae: Lepidoptera), a serious pest of mulberry trees in Kashmir. *Oriental Science*, 7, 111-113.
- Rajadurai, S., Manjunath, D., Katiyar, R.L., & Kumar, V. (1999). Leaf roller: A serious pest of mulberry. *Indian Silk*, 37, 9-11.
- Roya Khosravia., & Jalal Jalali Sendi. (2010). Biology and demography of *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae) on mulberry. *Journal of Asia-Pacific Entomology*, 13, 273-276. <https://doi.org/10.1016/j.aspen.2010.04.005>
- Sakthivel, N., Narendra Kumar, J.B., Dhahira Beevi, N., Devamani, M., & Teotia, R.S. (2019). Mulberry pests current status and management practices, Central Sericultural Research & Training Institute Central Silk Board, Ministry of Textiles, Government of India, pp, 17-18.
- Siddegowda, K., Gupta, V.K., Sen, A.K., Benchamin, K.V., & Manunath, D. (1995). *Diaphania sp.* infests mulberry in South India. *Indian Silk*, 34, 6-8.
- Srinivas Gowda, T.K., Narayana Swamy, & Sanaveerappanavar, V.T. (2001). Natural enemies of *Diaphania pulverulentalis* (Hampson) (Lepidoptera: Pyralidae) infesting mulberry. *Sericologia*, 41(4), 621-625.
- Srinivasa Gowda, R. (2004). Bio-Ecological and integrated management of mulberry leaf roller, *Diaphania Pulverulentalis* (Hampson) Lepidoptera: Pyralidae. Ph.D Thesis, University of Agricultural Sciences, Bangalore, India. P.87.
- Srinivasa Gowda, R., Narayanaswamy, T.K., & Rajegowda. (2000). Management of mulberry leaf roller, *Diaphania pulverulentalis* (Hampson) (Lepidoptera: Pyralidae). *Environment & Ecology*, 23(3), 485-487.
- Sultana, M., Brahma, D., Chakraborty, R., Borah, R., & Sarma, J. (2023). Diversity of Defoliators and Natural Enemies Associated with the Mulberry Ecosystem. *Int.J.Curr.Microbiol.App.Sci*, 12(12), 160-180. <https://doi.org/10.20546/ijcmas.2023.1212.020>
- Watanabe, H., Kurihara, Y., Wang, Y.X., & Shimizu, T. (1988). Mulberry pyralid, *Glyphodes pyloalis*: habitual host of nonoccluded viruses pathogenic to the silkworm, *Bombyx*

UNDER PEER REVIEW