

## Original Research Article

### Effect of mexican sunflower (*Tithonia diversifolia*) extract on the growth and herbage yield of lemon grass (*Cymbopogon flexuosus*) var. krishna

#### ABSTRACT

A field experiment was conducted in the Medicinal plant unit, Department of Horticulture, Faculty of Agriculture, Annamalai university to study the effect of mexcian sunflower (*Tithonia diversifolia*) leaf extract on the growth and herbage yield of lemon grass (*Cymbopogon flexuosus* var. krishna). The experiment was laid out in a Randomized Block Design with nine treatments and replication thrice. The treatments consisted of FYM (25 t ha<sup>-1</sup>) and Tithonia extract at different concentration viz., 5, 10, 15, 20, 25, 30, 35 and 40 per cent as foliar application on 15, 45, 75, 105, and 135 days after transplanting. The results of the experiment revealed that the combined application of FYM 25 t ha<sup>-1</sup> + Tithonia extract per cent 40 per cent (T<sub>9</sub>) produced the highest values for growth parameters such as plant height, number of tillers, plant spread (East-West and North-South), number of leaves plant<sup>-1</sup> and yield parameters such as fresh herbage, dry herbage yield and dry matter production (per plant, per plot and per hectare).

#### INTRODUCTION

*Cymbopogon flexuosus* (Nees ex Steud) Wats. belonging to the family Poaceae is known as Cochin or East Indian or Malabar lemongrass. It is a perennial aromatic cum medicinal herb native to the Indian subcontinent (Adinarayana *et al.*, 2012). The crop is extensively cultivated in the poor, marginal and waste lands and also along the bunds as live mulch. The well ramified root system of the plant helps in soil and water conservation. Lemongrass was one of the herbs to travel along the spice route from Asia to Europe.

Lemongrass oil of commerce is popularly known as Cochin oil in the world trade, since 90 per cent of it is shipped from Cochin port. It is grown in several countries particularly West Indies, China, Indonesia, Brazil, Congo, Republic of Malagasy, Sri Lanka, Zambia and other countries. In India it is cultivated along Western Ghats, Karnataka and Tamil Nadu states besides foothills of Arunanchal Pradesh and Sikkim (Srivastava *et al.*, 2013).

Lemon grass is commonly known for its higher citral content. The major constituents of root, stem and leaves are geraniol 30.5 per cent, citronellol 24.1 per cent, neral 10.3 per cent and geranial 13.6 per cent. Lemon grass oil is mainly used in the manufacture of perfumes for soaps, hair oils scents and medicines. It also has antibacterial properties. Ionone prepared from citral present in lemon grass oil is one of the most important raw material for the preparation of vitamin A. The use of lemon grass is part of a competitive market which includes pharmaceuticals, food, cosmetics, and perfumery markets. There are numerous examples of the application of lemongrass for health remedies by different ethnic groups (Ravinder *et al.*, 2010).

The yield and quality of lemongrass is mostly influenced by the climatic and agronomic management factors. The management of plant nutrition is a significant aspect that influences plant output. Organic farming system emphasise on the use of organic matters for maintaining soil health, growth and multiplication of beneficial microbes and minimizing health hazards associated with food (Badalingappanavar *et al.*, 2018). As conventional method of farming uses the chemical based fertilizers which are not useful in case of medicinal plants as use of chemical fertilizers had adverse effect on nature of secondary

metabolites. Organic method of cultivation is found to effectively enhancing the production quantitatively and qualitatively over chemical fertilizers. Beside their effectiveness these are easily available material in low cost and also helps in reduction in farm waste.

Farmyard manure (FYM) is a decomposed mixture of dung, urine, litter, and leftover materials from roughages and fodder fed to animals. A well-decomposed FYM contains 0.5–1.5 per cent Nitrogen, 0.2–0.4 per cent Phosphorus, and 0.5–1.0 per cent potassium. (Kumar *et al.*, 2021).

*Tithonia diversifolia* commonly known as Mexican sunflower was probably introduced into West Africa as an ornamental plant. *Tithonia diversifolia* leaves are rich in essential nutrients such as nitrogen, phosphorus, potassium, and other micronutrients that are beneficial for plant growth. Chemical composition of *Tithonia diversifolia* is Nitrogen 1.76 per cent, Phosphorous 0.82 per cent, Potassium 3.92 per cent, Calcium 3 per cent and Magnesium 0.005 per cent (Olabode *et al.*, 2007). Using *Tithonia diversifolia* as a foliar spray is an organic method to provide nutrients to plants.

In view of the above the present study was carried out on the Standardization and effect of Mexican sunflower (*Tithonia diversifolia*) extract on the growth and herbage yield of lemon grass (*Cymbopogon flexuosus*) var. krishna.

## Materials and methods

The field experiment was carried at Department of Horticulture, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu during January – July, 2024. Geographical location of land and climatic condition of experimental site is geographically situated at 11° 24' North latitude and 79° 44' East longitude at an altitude of + 5.72 m above MSL. The experiment was laid out in a Randomized Block Design with nine treatments and replicated thrice. The treatments consisted of T<sub>1</sub>- FYM 25 t ha<sup>-1</sup> (Control), T<sub>2</sub>- FYM 25 t ha<sup>-1</sup> + Tithonia Extract 5%, T<sub>3</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 10%, T<sub>4</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 15%, T<sub>5</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 20%, T<sub>6</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 25%, T<sub>7</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 30%, T<sub>8</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 35%, T<sub>9</sub>-FYM 25 t ha<sup>-1</sup> + Tithonia Extract 40%. Hundred gram of fresh shoot of *Tithonia diversifolia* were harvested and chopped into 2-3 cm length and finely ground with mortar and pestle. The ground plant material was soaked in one litre of distilled water in a large beaker for 12 hours. The solution was filtered through cheese cloth to remove debris and finally filtered using Whatman No. 1 filter paper. The final filtrate served as the leaf extract of *Tithonia diversifolia*. The chlorophyll content index was recorded with the help of chlorophyll content index meter (Model CCM – 200 001 300. CE).

## Result and discussion

### Growth attributes

In the present study, it was observed that applying FYM 25 t ha<sup>-1</sup> + Tithonia Extract 40 per cent (T<sub>9</sub>) recorded the maximum plant height, number of tillers plant<sup>-1</sup>, number of leaves plant<sup>-1</sup>, plant spread (East-West and North-South) at 30, 60, 90, 120 and 150 DAP. The perusal of data furnished in Table 1 shows that the maximum plant height with the values of 59.73, 101.03, 136.24, 78.36 and 115.29 cm at 30, 60, 90, 120 and 150 days after planting and the least plant height of 41.38, 71.40, 95.30, 55.45 and 89.36 cm respectively, were registered in the Control (T<sub>1</sub>) which received only FYM 25 t ha<sup>-1</sup>. The influence of spraying tithonia extract on number of tillers plant<sup>-1</sup> was found to be insignificant at 30 DAP, while the difference were found significant among various treatments at later growth of the stage. The maximum number of tillers plant<sup>-1</sup> registered with the values of 4.60, 16.93, 34.71, 42.54 and 58.02 at 30, 60, 90,

120 and 150 DAT. The least number of tillers (3.72, 7.23, 21.23, 26.31 and 35.17) recorded in the control ( $T_1$ ).

FYM increases soil macro elements, such as nitrogen (N), phosphorous (P) and potassium (K), resulting in maximum NPK uptake by the crop and improved soil organic carbon as well as available N, P, and K for increased growth of forage grasses (Walie *et al.*, 2022). The increase in plant height might be due to greater availability of macronutrients from organic sources which helped in acceleration of various metabolic processes (Momin and Gajbhiye 2016). The present results are in conformation with the findings of Anuja and Jayasri (2011) in sweet basil, Punam *et al.* (2012) in lemongrass. The highest plant height and number of tillers is due to amount of N, P, and K content present in tithonia. The N concentration in tithonia leaves is higher than the critical level of 2.0 to 2.5 per cent, below which net immobilization of N, would be expected. The P concentration is also higher than the critical level of 0.25 per cent for net P mineralization (Ademiluyi *et al.*, 2007). Phosphorus plays a crucial role in enhancing vegetative growth in plants by affecting soil nitrogen dynamics and promoting plant and microbial immobilization, ultimately leading to increased soil total nitrogen levels (Wang *et al.*, 2022). 'K' helps in photosynthesis, favors high energy status, maintains cell turgour, regulates opening of leaf stomata, promotes water uptake, regulates nutrients translocation in plant, favours carbohydrate transport, enhances N uptake, helps in protein synthesis and promotes starch synthesis (Mitra and Dhaliwal 2009). The present results are in conformation with the findings of Babajide *et al.* (2008) in tomato and Shokalu *et al.* (2009) in *Celosia argentea*.

It can be inferred from the data presented in Table. 2 shows the maximum number of leaves recorded with the values of 14.04, 55.95, 152.52, 192.17 and 261.11 at 30, 60, 90, 120 and 150 DAP and the least value for leaf production was noticed in the treatment  $T_1$  (control) with the values of 9.32, 28.92, 89.21, 102.32 and 161.94 respectively. It can be observed from the data presented in Table 3 that the maximum plant spread of East-West direction was registered the values of (39.82, 73.25, 98.08, 46.57 and 70.81 cm) at 30, 60, 90, 120 and 150 days after planting, while the least plant spread was registered in the control  $T_1$  (FYM 25 t ha<sup>-1</sup>) with the values of 25.36, 45.36, 68.36, 27.35 and 46.35 cm respectively. the maximum plant spread in North-South direction with the values of 19.93, 57.19, 87.99, 39.43 and 60.05 cm at 30, 60, 90, 120 and 150 days after planting respectively, Treatment  $T_1$  (control) recorded the minimum plant spread 12.25, 38.45, 61.25, 17.36 and 37.36 cm respectively.

The highest number of leaves plant<sup>-1</sup> and maximum plant spread (East-West and North South) was due to the soil application of FYM and foliar application of tithonia extract. Farmyard manure (FYM) plays a crucial role in enhancing soil fertility due to its optimal C: N ratio, which facilitates the release of nitrogen in readily available forms during decomposition (Vinayak *et al.*, 2022). The maximum number of leaves plant<sup>-1</sup> and plant spread (East-West and North-South) may be due to increase in soil nitrogen levels. This is in accordance with the finding of Shinde *et al.* (2013) in ashwagandha and Rashid *et al.* (2013) in stevia. The growth promotion could be explained by the high level of macro and micronutrients in *T. diversifolia* leaves and stems, essential for crop growth and positively involved as organic amendments in many physiological processes such as the molecular structure of nucleic acid, the storage functions, the energy transfer processes within the plant and the roots developmet (Ewane *et al.*, 2020). This is in accordance with the findings of Aguyoh *et al.*, (2010) in watermelon.

### Physiological attributes

The highest Chlorophyll Content Index (Table 2) at I Harvest (90 DAP) and II Harvest (150 DAP) with values of 25.85 and 28.77 and the least in control ( $T_1$ ) with the values of 12.83

and 13.29. Chlorophyll is a vital pigments in plant photosynthesis, absorbing sunlight and transferring energy efficiently. They have diverse absorption bands in the visible and near-infrared regions, crucial for plant growth and development (Paterson 2023). The increasing chlorophyll content was due to the presence of microorganisms in the FYM that colonizes in the rhizosphere and stimulate the plant growth and biochemical contents (Ravimycin 2016). This is in line with the findings of (Chaurasia and Singh (2021) in Ashwagandha. Organic manures are good sources of nitrogen, which favour chlorophyll production. Nitrogen supply affects leaf growth because it increases the leaf area of plants thus influencing photosynthesis. Photosynthetic proteins represent a large proportion to total leaf N (Devkota and Jha 2013). The green leaf biomass of tithonia is high in nutrients, averaging about 3.5% Nitrogen, 0.37% Phosphorous and 4.1% potassium on dry matter basis (Jama *et al.*, 2000), Which helps in photosynthesis.

### Yield attributes

It can be inferred from the data presented in Table. 4&5 shows that the maximum fresh herbage production at I Harvest and II Harvest (280.30 g plant<sup>-1</sup>, 2.14 kg plot<sup>-1</sup> and 13.24 t ha<sup>-1</sup> and 320.81 g plant<sup>-1</sup>, 2.45 kg plot<sup>-1</sup> and 15.15 t ha<sup>-1</sup>) and dry herbage yield I Harvest and II Harvest (56.06 g plant<sup>-1</sup>, 0.43 Kg plot<sup>-1</sup> and 2.65 t ha<sup>-1</sup> and 64.16 g plant<sup>-1</sup>, 0.49 Kg plot<sup>-1</sup> and 3.03 t ha<sup>-1</sup>) was recorded in plants supplied with FYM 25 t ha<sup>-1</sup> + Tithonia extract 40 per cent (T<sub>9</sub>), which could have attributed to the better growth of the plants as reflected in yield contributing characters such as plant height, number of tillers plant<sup>-1</sup>, plant spread (East-West and North-South) and number of leaves plant<sup>-1</sup>. The increased in yield parameters was due to increase in plant height, number of tillers plant<sup>-1</sup>, number of leaves plant<sup>-1</sup> and largest leaves leading to higher dry matter accumulation in plants and translocation of photosynthates from source to sink which might due to sufficient availability of major and micronutrients from FYM, helped in the uptake of more nutrients (Muruganandam *et al.*, 2021).

Organic manures, as noted by Krishnamoorthy and Ravikumar (1973), can enhance plant growth through various mechanisms. Their high humus content promotes microbial nitrogen fixation, ensuring a steady supply of nitrogen to plants. Additionally, organic manures may stimulate the production of plant growth hormones. This combination of factors can lead to extended leaf area duration, allowing for increased photosynthetic rates and, ultimately, higher dry matter accumulation (Padmanaban, 2003). Similar findings have been reported for sweet basil by Yadav *et al.* (2023). FYM also help to induce soil physical conditions which make the plant roots to proliferate better for utilisation of nutrients to boost up the yield (Rathore 2021). Increased yield parameters on application of increased level of FYM might also be due to enhanced canopy photosynthesis through increased vegetative growth of crop (Rahman *et al.*, 2004). This is in accordance with the findings of Priya *et al.* (2018) in sweet flag.

Increase in yield of lemongrass was due to steady decomposition of FYM and release of nutrients throughout the crop growth period coupled with better assimilation of nutrients (singh *et al.*, 2017). This results are in agreement with the findings of Umesh *et al.* (2007) in medicinal solanum and Sharaf *et al.* (2013). Additionally, Previous crops and tillage practices in an experimental field can impact soil nitrogen availability and microbial activity, which subsequently influence the mineralization rate of FYM.

Foliar fertilization has the ability to improve the efficiency and rapidity of utilization of a nutrient urgently required by the plant for various metabolic activities. (Alshaal *et al.*, 2017). The spraying of tithonia extract @ 40 per cent was found to be the best treatment due to the effect of increased nitrogen, phosphorous and potassium in the tithonia extract (Abeyrathne *et al.*, 2021). This finding is consistent with the reports of Parman (2007) in

potato and Zhai *et al.* (2009) in tomato. Increasing or decreasing trend of growth parameters and yield parameters corresponds to increase in concentration of Mexican Sunflower extract as well as with period of transplanting.

### Conclusion

From the findings of the current study, it can be inferred that farm yard manure 25 t ha<sup>-1</sup> + Tithonia extract 40 % applied every month could be the best management practice to get the highest possible herbage yield in lemongrass. The present finding clearly shows that Mexican sunflower extract has potential to serve as a natural biostimulant for the sustainable lemongrass cultivation providing an eco-friendly substitute in place of synthetic inputs by promoting environmental health.

**Table 1. Effect of Tithonia extract on Plant height (cm) and Number of tillers plant<sup>-1</sup> of Lemon grass (*Cymbopogon flexuosus*) var. krishna**

TREATMENTS	Plant height (cm)					Number of tillers plant <sup>-1</sup>				
	30 DAP	60 DAP	90 DAP	120 DAP	150 DAP	30 DAP	60 DAP	90 DAP	120 DAP	150 DAP
T <sub>1</sub> - FYM 25 t ha <sup>-1</sup> (Control)	41.38	71.40	95.30	55.45	89.36	3.72	7.23	21.23	26.31	35.17
T <sub>2</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 5%	41.44	71.94	96.19	55.92	90.38	3.89	7.41	21.65	26.80	35.80
T <sub>3</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 10%	42.06	72.57	97.43	56.44	91.64	4.10	7.61	22.14	27.32	36.52
T <sub>4</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 15%	45.32	77.83	104.95	60.60	92.00	4.23	9.07	24.26	29.84	40.38
T <sub>5</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 20%	46.18	78.62	106.36	61.23	93.36	4.36	9.29	24.77	30.40	41.20
T <sub>6</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 25%	49.50	83.93	113.70	65.50	98.75	4.43	10.96	27.11	32.97	45.31
T <sub>7</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 30%	52.84	89.41	121.11	69.72	104.17	4.54	12.79	29.54	35.86	49.50
T <sub>8</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 35%	56.25	95.14	128.64	74.01	109.70	4.58	14.80	32.10	38.98	53.73
T <sub>9</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 40%	59.73	101.03	136.24	78.36	115.29	4.60	16.93	34.71	42.54	58.02
<b>S. Ed</b>	1.18	2.02	2.73	1.57	2.40	NS	0.27	0.65	0.80	1.09
<b>CD (P=0.05)</b>	3.07	4.29	5.78	3.34	5.10	NS	0.58	1.38	1.70	2.31

**Table 2. Effect of Tithonia extract on number of leaves plant<sup>-1</sup> and chlorophyll content index (CCI) of Lemon grass (*Cymbopogon flexuosus*) var. Krishna**

TREATMENT	Number of leaves plant <sup>-1</sup>					chlorophyll content index (CCI)		
	30 DAP	60 DAP	90 DAP (I Harvest)	120 DAP	150 DAP (II Harvest)	90 DAP (I Harvest)	150 DAP (II Harvest)	MEAN
T <sub>1</sub> - FYM 25 t ha <sup>-1</sup> (Control)	9.32	28.92	89.21	102.32	161.94	12.83	13.29	13.06
T <sub>2</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 5%	9.48	30.12	91.53	103.44	165.07	13.07	13.50	13.29
T <sub>3</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 10%	9.68	31.41	93.92	104.63	168.39	13.36	13.76	13.56
T <sub>4</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 15%	10.38	35.23	104.89	120.88	185.60	15.39	16.22	15.81
T <sub>5</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 20%	10.61	36.59	107.58	122.10	189.10	15.72	16.55	16.14
T <sub>6</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 25%	11.38	40.82	118.60	138.79	206.62	17.98	19.27	18.63
T <sub>7</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 30%	12.19	45.34	129.81	156.02	224.48	20.27	22.16	21.21
T <sub>8</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 35%	13.08	49.96	141.13	173.94	242.67	22.59	25.18	23.88
T <sub>9</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 40%	14.04	55.95	152.52	192.17	261.11	25.85	28.77	27.31
<b>S. Ed</b>	NS	0.98	2.84	3.38	4.95	0.43	0.47	-
<b>CD (P=0.05)</b>	NS	2.09	6.02	7.18	10.51	0.93	1.01	-

**Table 3. Effect of Tithonia extract on Plant spread (East-West) and (North- South) (cm) of Lemon grass**  
*(Cymbopogon flexuosus) var. krishna*

TREATMENT	Plant spread (East-West) (cm)					Plant spread (North- South) (cm)				
	30 DAP (cm)	60 DAP (cm)	90 DAP (I Harvest) (cm)	120 DAP (cm)	150 DAP (II Harvest) (cm)	30 DAP (cm)	60 DAP (cm)	90 DAP (I Harvest) (cm)	120 DAP (cm)	150 DAP (II Harvest) (cm)
T <sub>1</sub> - FYM 25 t ha <sup>-1</sup> (Control)	25.36	45.36	68.36	27.35	46.35	12.25	38.45	61.25	17.36	37.36
T <sub>2</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 5%	25.56	46.56	69.65	27.65	47.24	12.45	39.05	62.27	17.71	38.10
T <sub>3</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 10%	26.36	48.02	71.29	28.33	48.26	12.78	39.93	63.73	18.39	39.08
T <sub>4</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 15%	27.96	51.61	75.82	30.83	51.55	13.80	42.43	67.69	21.64	42.31
T <sub>5</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 20%	28.31	54.13	77.74	31.85	52.79	14.25	43.73	69.38	22.87	43.52
T <sub>6</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 25%	30.50	59.01	82.42	34.74	56.68	15.57	46.49	73.58	26.52	47.21
T <sub>7</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 30%	33.36	64.19	87.40	38.39	60.91	16.98	49.69	78.12	30.49	51.19
T <sub>8</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 35%	36.48	69.85	92.63	42.28	65.60	18.40	53.29	82.98	34.78	55.40
T <sub>9</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 40%	39.82	73.25	98.08	46.57	70.81	19.93	57.19	87.99	39.43	60.05
<b>S. Ed</b>	0.78	1.47	2.05	0.88	1.42	0.36	1.17	1.84	0.67	1.18
<b>CD (P=0.05)</b>	1.59	2.99	4.18	1.80	2.90	0.79	2.38	3.74	1.38	2.41

**Table 4. Effect of Tithonia extract on fresh herbage yield of lemon grass (*Cymbopogon flexuosus*) var. krishna**

<b>TREATMENT</b>	<b>90 DAP I HARVEST (g Plant<sup>-1</sup>)</b>	<b>90 DAP I HARVEST (kg plot<sup>-1</sup>)</b>	<b>90 DAP I HARVEST (t ha<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (g Plant<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (kg plot<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (t ha<sup>-1</sup>)</b>
T <sub>1</sub> - FYM 25 t ha <sup>-1</sup> (Control)	123.62	0.9457	5.84	170.32	1.30	8.04
T <sub>2</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 5%	126.83	0.9702	5.99	173.34	1.33	8.19
T <sub>3</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 10%	130.36	0.9973	6.16	176.55	1.35	8.34
T <sub>4</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 15%	158.01	1.2088	7.46	201.69	1.54	9.52
T <sub>5</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 20%	161.90	1.2385	7.65	205.16	1.57	9.69
T <sub>6</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 25%	189.93	1.4530	8.97	231.52	1.77	10.93
T <sub>7</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 30%	218.49	1.6714	10.32	258.91	1.98	12.23
T <sub>8</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 35%	249.28	1.9070	11.77	288.55	2.21	13.63
T <sub>9</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 40%	280.30	2.1443	13.24	320.81	2.45	15.15
<b>S. Ed</b>	4.65	0.04	-	5.65	0.03	-
<b>CD (P=0.05)</b>	9.86	0.10	-	11.97	0.07	-



**Table 5. Effect of Tithonia extract on dry herbage yield of lemon grass (*Cymbopogon flexuosus*) var. krishna**

<b>TREATMENT</b>	<b>90 DAP I HARVEST (g Plant<sup>-1</sup>)</b>	<b>90 DAP I HARVEST (kg plot<sup>-1</sup>)</b>	<b>90 DAP I HARVEST (t ha<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (g Plant<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (kg plot<sup>-1</sup>)</b>	<b>150 DAP II HARVEST (t ha<sup>-1</sup>)</b>
T <sub>1</sub> - FYM 25 t ha <sup>-1</sup> (Control)	24.72	0.19	1.17	34.06	0.26	1.61
T <sub>2</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 5%	25.37	0.19	1.20	34.67	0.27	1.64
T <sub>3</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 10%	26.07	0.20	1.23	35.31	0.27	1.67
T <sub>4</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 15%	31.60	0.24	1.49	40.34	0.31	1.90
T <sub>5</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 20%	32.38	0.25	1.53	41.03	0.31	3.77
T <sub>6</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 25%	37.99	0.29	1.79	46.30	0.35	2.19
T <sub>7</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 30%	43.70	0.33	2.06	51.78	0.40	2.45
T <sub>8</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 35%	49.86	0.38	2.35	57.71	0.44	2.73
T <sub>9</sub> -FYM 25 t ha <sup>-1</sup> + Tithonia Extract 40%	56.06	0.43	2.65	64.16	0.49	3.03
<b>S. Ed</b>	1.38	0.004	-	0.37	0.02	-
<b>CD (P=0.05)</b>	2.93	0.01	-	0.79	0.06	-

### Disclaimer (Artificial intelligence)

#### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

### Reference:

1. Abeyrathne, T., K. D. Harris and A. M. K. D. M. Attanayaka. 2021. Effect of Foliar Application of Wild Sunflower (*Tithonia Diversifolia*) Leaf Extract on Yield Attributes of Vegetable Cowpea Cv. Bs-1 in Sandy Regosol. World J. Cur. Sci. Res., 1(2): 1-4.
2. Ademiluyi, B. O and S. O. Omotoso. 2007. Comparative evaluation of *Tithonia diversifolia* and NPK fertilizer for soil improvement in maize (*Zea mays*) production in Ado Ekiti, southwestern Nigeria., American-Eurasian J. Sustain. Agric., 1(1): 32-36.
3. Adinarayana, G., G. Rahul, R. S. Kiran, K. V. Syamsundar and B. R. Rajeswara. 2012. Evaluation of antimicrobial potential of field distilled and water-soluble essential oils of *Cymbopogon flexuosus*. J Pharmacogn, 3(2), 142-6.
4. Aguyoh, J. N., W. Audi, M. Saidi and L. Gao-Qiong. 2010. Growth, yield and quality response of watermelon (*Citrullus lanatus* [thunb] mansf. & nakai) cv. crimson sweet) subjected to different levels of tithonia manure. Int. J. Nat. Sci., 1(1): 7-11.
5. Alshaal, T. and H. El-Ramady. 2017. Foliar application: from plant nutrition to biofortification. Environ., Biodiv. & Soil Security, 1: 71-83.
6. Amadalo. 2000. *Tithonia diversifolia* as a green manure for soil fertility improvement in western Kenya: A review. Agrofor. Syst., 49: 201-221.
7. Anuja S. and P. Jayasri. 2011. Effect of soil and foliar application of organic nutrients on NPK uptake, herbage yield and inflorescence yield of sweet basil (*Ocimum basilicum* L.). Asian J. Hort., 6(1):195-198.
8. Babajide, P. A., O. S. Olabode, W. B. Akanbi, O. O. Olatunji, and E. A. Ewetola. 2008. Influence of composted Tithonia biomass and N-mineral fertilizer on soil physico-chemical properties and performance of Tomato (*Lycopersicon lycopersicum*). Res. J. Agronomy, 2(4): 101-106.

9. Badalingappanavar, R., M. Hanumanthappa, H. K. Veeranna, S. Kolakar and G. Khidrapure. 2018. Organic fertilizer management in cultivation of medicinal and aromatic crops: a review. *J. Pharm. and Phytochem.*, 7(3S): 126-129.
10. Chaurasia, S., and R. Singh. 2021. Effect of integrated nutrient management system on chlorophyll a, chlorophyll b and carotenoids content of Ashwagandha (*Withania somnifera* L. Dunal). *Int. J. Adv. Res. Biol. Sci.*, 8(10): 154-165.
11. Devkota, A. and P. K. Jha. 2013. Effect of integrated manuring on growth and yield of *Centella asiatica* (L.) Urb. *Trop. Ecol.*, 54(1): 89-95.
12. Ewané, C. A., N. T. Mbanya and T. Boudjeko. 2020. *Tithonia diversifolia* leaves and stems use as substrate amendment promote the growth of plantain vivoplants in the nursery. *Agric. sci.*, 11: 849-859.
13. Gaur, B. L. and M. S. Rathore. 1991. Varietal response of opium poppy (*Papaver somniferum* L.) to nitrogen fertilization on vertisols. *Indian. J. Agron*, 36: 100-101.
14. Jama, B., C. A. Palm, R. J. Buresh, A. L. Niang, C. Gachengo, G. Nziguheba and B.
15. Krishnamoorthy, K.K. and V. Ravikumar. 1973. Comparative effectiveness of organic and mineral fertilizers on demopodzolic soils. *Soil. Fertl.*, 30: 2920.
16. Kumar, S., S. Dhar, S. Barthakur, M. V. S. Rajawat, S. A. Kochewad, S. Kumar and L. R. 2021. Farmyard manure as k-fertilizer modulates soil biological activities and yield of wheat using the integrated fertilization approach. *Front. Env Sci.*, 9: 764489.
17. Mbatha, A.N., 2008. Influence of organic fertilizers on the yield and quality of cabbage and Carrots. M. Sc. Thesis, University of the Free State Bloemfontein.
18. Mitra, S. K., and S. S. Dhaliwal. 2009. Effect of potassium on fruit quality and their storage life. In *Proceedings IPI-OUAT-IPNI International Symposium.*, 335: 327- 342.
19. Momin, Y. D. and B. R. Gajbhiye. 2016. Influence of FYM and NPK Fertilization on Growth Characters of Lemongrass (*Cymbopogon flexuosus*). *Adv. Life Sci*, 5(12): 5360-536.
20. Muruganandam, C., R. Ezhilnilavu and S. Sivasankar. 2021. Effect of integrated nutrient management on growth parameters of medicinal coleus (*Coleus forskohlii* Briq.). *Plant Arch.*, 21(1): 2525-2528.
21. Olabode, O. S., O. Sola, W. B. Akanbi, G. O. Adesina and P. A. Babajide. 2007. Evaluation of *Tithonia diversifolia* (Hemsl.) A Gray for soil improvement. *World J. of Agric. Sci.*, 3(4): 503-507.
22. Padmanaban, K. 2003. Effect of organic manures on growth, root yield and alkaloid content of ashwagandha (*Withania somnifera* (L.) dunal) 'jawahar'. M.Sc. (Hort.) Thesis, Tamilnadu Agricultural University, Coimbatore.
23. Parman, S. 2007. The effect of applying liquid organic fertilizer on the growth and production of potatoes (*Solanum tuberosum* L.). *Anatomy Physiol.*, 15(2): 21-31.
24. Paterson. 2023. Handbook of natural colorants. John Wiley & Sons. <https://doi.org/10.1002/9781119811749.ch10>
25. Priya, B. T., P. R. Devi and P. Sunitha. 2018. Effect of spacing and FYM on growth and yield of sweet flag (*Acorus calamus* L.). *J. Pharmacogn. Phytochem.*, 7(3S): 354-357.
26. Punam, P., R. Kumar, S. Sharma and D. Atu. 2012. The effect of organic management treatments on the productivity and quality of lemon grass (*Cymbopogon citratus*). *J. Org. Syst.*, 7(2): 36-48.
27. Rahman MS, Islam MA, Haque MS, Karim MA. Effects of planting dates and gibberellic acid on growth and yield of garlic. 2004. *Asian J Pl. Sci.* 344-352.

28. Rashid, Z., M. Rashid, S. Inamullah, S. Rasool and F. A. Bahar, F. A. 2013. Effect of different levels of farmyard manure and nitrogen on the yield and nitrogen uptake by stevia (*Stevia rebaudiana* Bertoni). African. J. Agric. Res., 8(29): 3941-3945.
29. Rathore, V. 2021. Effect of FYM and nitrogen levels on growth, dry matter accumulation, yield and nutrient uptake of brahmi (*Bacopa monnieri* L.). Pantnagar J. Res., 19(31): 151-158.
30. Ravimycin, T. 2016. Effects of Vermicompost (VC) and Farmyard Manure (FYM) on the germination percentage growth biochemical and nutrient content of Coriander (*Coriandrum sativum* L.). Int. J. Adv. Res., 3(6): 91-98.
31. Ravinder, K., K. Pawan, S. Gaurav, K. Paramjot, S. Gagan and K. Appramdeep. 2010. Pharmacogn. Invest of Cymbopogon citratus (DC) Stapf. Der Pharm. Lett., 2(2): 181-189.
32. Setyowati, N., S. Sudjatkiko, Z. Mukhtar, F. Fahrurrozi, M. Chozin and P. Simatupang. 2018. Growth and yield responses of cauliflower on Tithonia (*Tithonia diversifolia*) compost under organic farming practices. Int. J. Agric. Technol., 14(7): 1905-1914.
33. Sharaf, E. D., M. N. Shalan, R. A. Fouda and A. S. Dapour. 2013. Effect of some organic and bio-fertilizers on quality and quantity of *Rosmarinus officinalis* L. plants. J. Plant Prod., 4(7): 1061-1076.
34. Sharaf, E. D., M. N. Shalan, R. A. Fouda and A. S. Dapour. 2013. Effect of some organic and bio-fertilizers on quality and quantity of *Rosmarinus officinalis* L. plants. J. Plant Prod., 4(7): 1061-1076.
35. Shinde, A., P. Gahunge, P. Singh, S. K. Rath and N. Khemani. 2013. Effect of inorganic fertilizers and organic manures on growth, quality and yield of ashwagandha (*Withania somnifera* Dunal) cv. Jawahar Ashwagandha-20. Ann. Pharm. & Pharm. Sci., 4(1): 13-16.
36. Shokalu, A. O., A. O. Ojo, D. T. E.-Adewoyin, H. A. Akintoye and A. A. Idowu. 2009. August. Use of *Tithonia diversifolia* and Compost as Soil Amendments for Growth and Yield of Celosia argentea. In I All Afr. Horti. Congr., 911: 149-156.
37. Srivastava, V., S. Dubey, and A. Mishra. 2013. A review on lemon grass: agricultural and medicinal aspect. Int. Res. J. Pharm., 4(8): 42-44.
38. Umesh, K., S. P. Soumya, G. R. Smitha and B. S. Sreeramu. 2007. Effect of different organic manures on growth, yield and quality of Makoi (*Solanum nigrum*). In *National Seminar on Production, Processing and Marketing of Medicinal, Aromatic and Dye Yielding Crops* (pp. 10-16).
39. Vinayak, M., C. Ramana, B. Haribabu and K. M. Reddy. 2022. Determination of Physical Properties of FYM for the Development of FYM Shredder cum Spreader. Int. J. Stress Manag., 13: 1398-1402.
40. Walie, M., F. Tegegne, Y. Mekuriaw, A. Tsunekawa, N. Kobayashi, T. Ichinohe and E. Adegbo. 2022. Effects of farmyard manure and Desmodium intercropping on forage grass growth, yield, and soil properties in different agro-ecologies of Upper Blue Nile basin, Ethiopia. Cogent food agric., 8(1): 2082041.
41. Wang, R., B. Bicharanloo, E. Hou, Y. Jiang and F. A. Dijkstra. 2022. Phosphorus supply increases nitrogen transformation rates and retention in soil: a global meta-analysis. Earth's Future, 10(3): 1- 15.
42. Yadav, R., D. K. Meena, R. Dawar, S. T. Pandey and T. Singh. 2023. Enhancing sweet basil (*Ocimum basilicum*) production and quality with fermented organic manure. J. Pharm Innov., 12(3): 2845-2850.

43. Zhai, Z., D. L. Ehret, T. Forge, T. Helmer, W. Lin, M. Dorais and A. P. Papadopoulos. 2009. Organic fertilizers for greenhouse tomatoes: productivity and substrate microbiology. *Hort. Sci.*, 44(3): 800-809.
44. Singh, D. P and D. Singh. 2017. Effect of nitrogen and FYM on yield, quality and uptake of nutrients in wheat (*Triticum aestivum*). *Annals of Plant and soil Research*, 19(2), 232-236.

UNDER PEER REVIEW