

Advancing Groundwater Remediation: Efficacy of Slow-Release Permanganate Gels (SRPG) in Treating Contaminant Plumes

ABSTRACT

Aim: The aim of this study is to evaluate the efficiency of the slow-release permanganate gels (SRPG) as a technique for the remediation of contaminant plumes in aquifers to afford an improved degree of groundwater quality and reduced environmental risks.

Study Design: Experimental and modeling studies were reviewed to assess the oxidation capacity, release kinetics, and field applicability of SRPG for the remediation of groundwater. The review included literature from 2019 to 2024, focusing on the latest techniques and challenges in the field.

Methodology: A systematic search was conducted in several databases, including Google Scholar, PubMed, Scopus, and Web of Science. Inclusion criteria ensured that the selected articles were limited to those that were peer-reviewed discussing mechanisms, hydrogeologic applications, and contaminant mitigation involving SRPG. Experimental setup, numerical models, and field data were considered in order to assess performance and scalability for various usage of SRPG.

Results: The review identified 5 effective studies on SRPG in reducing contaminant concentrations in groundwater. Key factors for remediation success were increased oxidation capacity, longer release kinetics, and improved spatial distribution. One notable finding was a 75% reduction in trichloroethylene concentrations within six months of SRPG application in a pilot-scale study. Field applications reported significant contaminant reduction in heterogeneous aquifer systems, supporting the scalability of SRPG technology.

Conclusions: SRPG is a promising low-cost and sustainable method for groundwater remediation, particularly in complex aquifer systems. This study points to the need for future studies with regard to optimization of the gel formulations concerning long-term environmental impacts and the ability to biodegrade in environmental conditions toward globally sustainable water resources management.

Keywords: Slow-Release Permanganate Gels (SRPG), Contaminant Plumes, Groundwater Remediation, In-situ Chemical Oxidation (ISCO)

1. INTRODUCTION

Contamination of groundwater is considered one of the most widespread environmental hazards threatening the availability and quality of the most vital resources on earth[1]. Such contaminants include chlorinated solvents, petroleum hydrocarbons, pesticides, and heavy metals. The contamination is generally introduced into the aquifers from industrial discharge, agricultural runoff, and improper waste disposal as plumes of contaminants expand in space with time [2, 3]. Contaminated groundwater plumes are associated with high risks for

ecosystems, human health, and the long-term sustainability of this valuable resources[4]. These challenges, in turn, call for innovative remediation technologies that can ensure long-term effectiveness while being economically and environmentally viable. Traditional remediation techniques include pump-and-treat, in-situ chemical oxidation (ISCO), and bioremediation, among others, which have seen extensive application over the last decades [5, 6]. However, these methods often fall short in complex hydrogeologic environments where contaminants are heterogeneously distributed. For example, pump-and-treat systems are energy-intensive and ineffective in low-permeability areas, whereas conventional ISCO techniques, relying on liquid oxidants, have generally been plagued by a limited subsurface distribution and rapid oxidant depletion [7]. These limitations point to the need for more effective and sustainable solutions.

Slow-release permanganate gels (SRP-G) represent an exciting development in the evolution of groundwater remediation technologies. Unlike liquid oxidants, SRPG formulations provide a controlled and sustained release of potassium permanganate into the subsurface, enabling prolonged oxidation of contaminants and reducing the need for frequent applications [8]. This mechanism is particularly advantageous in heterogeneous aquifers where diffusion-limited processes dominate contaminant transport [9]. Furthermore, it is cost-effective for long-term remediation projects due to its capability to maintain reactive capacity over long periods. While pump-and-treat systems usually tend to be quite costly, power-consuming, and inefficient in low-permeability zones, the SRPG offers more specific and sustained oxidant delivery to reduce operational cost and energy consumption [7]. Whereas conventional ISCO approaches using liquid permanganate may face issues with distribution or the rapid consumption of oxidants, such challenges are overcome through the controlled, extended release with SRPG for far better contaminant treatment for extended periods. In contrast to bioremediation, which may be slow and less effective in heavily contaminated sites, the SRPG allows for faster, more predictable treatment performance, particularly in areas that may not easily reach optimum performance with other methods.[10].

Various research has proved that SRPG is effective in various contaminants, including trichloroethylene, perchloroethylene, and benzene-frequent groundwater pollutants [11,12]. Laboratory studies have shown that SRPG exhibits enhanced oxidative efficiency compared to liquid permanganate, particularly under conditions of high contaminant concentrations [13]. Field applications have further validated its efficacy, with case studies reporting significant reductions in contaminant levels and improvements in aquifer health within months of deployment [14, 15]. From an environmental sustainability perspective, SRPG aligns with the principles of green chemistry by minimizing secondary pollution and energy consumption [2]. The gels can be tailored to specific site conditions by modifying their composition and release kinetics, offering flexibility in addressing diverse remediation scenarios [8]. Their compatibility with other remediation strategies like bioremediation and natural attenuation further raises their potential for integrated approaches in the management of groundwater[4].

Despite these benefits, the effectiveness of SRPG still confronts a lot of challenges in their applications. The design of gels should consider site-specific factors: the rates of groundwater flow, types of contaminants, and geochemical conditions for optimum performance [7]. Additionally, there is a need to continue studies of the long-term impacts of SRPG in aquifer systems on possible byproduct formation and interactions with native microbial communities [10]. All these knowledge gaps should be bridged in scaling up the technology.

In this regard, slow-release permanganate gels represent one of the key developments in recent decades in the science of groundwater remediation. Combining the principles of

sustained release mechanisms with high oxidative capacity, SRPG offers an economically viable and environmentally acceptable method for remediating complex aquifer systems contaminated by diverse contaminant plumes. This review discusses state-of-the-art developments in the SRPG technology, inclusive of its applications, limitations, and future avenues for research.

2. METHODOLOGY

This review considers the systematic search, selection, and evaluation of literature with regard to effectiveness in using slow-release permanganate gels for the remediation of groundwater. The methodology includes a wide literature search, followed by a detailed selection of the studies and analysis of inherent limitations. Searching of relevant literature was done through four major academic databases: Google Scholar, PubMed, Scopus, and Web of Science. These databases were chosen for their wide coverage of peer-reviewed journals, conference proceedings, and technical reports. Searches were limited to publications from 2019 to 2024 to ensure the review incorporated the most recent advancements in SRPG technology. These include "slow-release permanganate gel", "groundwater remediation", "contaminant plume treatment", and "oxidation capacity". Boolean operators "AND" and "OR" were also used to refine the search results for comprehensive inclusion of relevant studies. Inclusion was limited to those that presented experimental, field, or modeling data related to SRPG applications in the English language. The exclusion criteria included publication before 2019, non-peer-reviewed articles, and studies that were not experimentally focused or were inaccessible in full-text format.




Figure 1: Flow diagram of the literature search and study selection for the review.

Of a total of 95 identified studies (see figure 2), 35 were from Google Scholar, 25 from PubMed, 20 from Scopus, and 15 from Web of Science. After removing duplicates, 70 unique studies remained. These were screened based on their titles and abstracts to exclude 55 either for being irrelevant or completely lacking in any focus on applications of SRPG. These left 15 studies that then underwent a detailed full-text review, and 10 others were excluded during this time. Common reasons for exclusion included insufficient methodological rigor, absence of experimental or field data on slow-release oxidants, lack of quantitative performance metrics, or inaccessibility of publications' format. The final selection included five studies that provided data on laboratory experiments, field trials, and numerical modeling and offered a wide range of perspectives on SRPG applications.

Notwithstanding the stringency of the above process, a number of limitations have been identified. Language restrictions posed a significant barrier, as non-English studies, potentially offering valuable regional insights, were excluded. Additionally, restricted access to full-text articles limited the inclusion of otherwise relevant studies, introducing potential

selection bias. The focus on SRPG specifically may have excluded broader insights from studies on other in situ chemical oxidation methods, potentially narrowing the scope of the review [1]. Furthermore, many of the included studies lacked long-term data, which limits understanding of the sustainability and environmental impacts of SRPG over extended periods. Variability in the methodological quality of the reviewed studies also presented challenges, as some offered limited quantitative data or inadequate statistical analyses, reducing the ability to draw consistent conclusions [11]. The systematic methodology adopted herein underlines the importance of transparency and reproducibility in literature reviews. Though there are certain limitations identified, this review underlines some key developments in SRPG technology and identifies areas of future research, including more extensive field trials, evaluation of long-term impacts, and broader inclusion of global studies to provide a comprehensive understanding of SRPG's role in groundwater remediation.

3. RESULTS AND DISCUSSION

The review identified key advances in the application of slow-release permanganate gels for the remediation of groundwater. These findings could indicate key aspects of their performance and field applicability:

Enhanced Oxidation Capacity

SRPG was very effective in oxidizing contaminants like chlorinated solvents, hydrocarbons, and organic compounds present in groundwater. Wang et al. [16], and Samborska-Goik Pogrzeba, [17] revealed that the supply of permanganate with SRPG is fairly stable for long periods, which improves the oxidation process in heterogeneous aquifers. Moreover, these gels provided effective treatment of low-permeability zones where conventional oxidants cannot easily reach.

Prolonged Release Kinetics

The formulation of permanganate in SRPG has been proven to ensure controlled release, thus maintaining effective concentrations for extended periods. For instance, Yen et al. [18] reported that SRPG can maintain its oxidation activity, thereby reducing the frequency of applications. The prolonged release diminishes operational costs while sustaining contaminant plume treatment.

Improved Spatial Distribution

Another important aspect in the reviewed studies is the potential of SRPG to permeate permanganate through heterogeneous and fractured aquifer systems. Wang et al. [19] investigated how SRPG delivers a more homogenous contaminant reduction compared to traditional liquid oxidants under varied hydrogeologic conditions. In this study, it was established that the gels conform to heterogeneities within the aquifer to enable appropriate treatment even in complex subsurface environments.

Field Applications and Scalability

Field trials gave strong evidence of the scalability and practical benefits of SRPG. Chang et al. [20] revealed that in a pilot-scale project, TCE concentrations were reduced by 75% within six months of the application of SRPG. Similarly, O'Connor et al. [21] demonstrated the feasibility of deploying SRPG in large-scale remediation projects and showed its effectiveness in mitigating contaminant plumes over a 2-hectare area. Both studies

emphasized the cost efficiency and environmental sustainability of SRPG technology when compared to other conventional methods.




Figure 2: Key Benefits of Slow-Release Permanganate Gels in Groundwater Remediation

Environmental Implications and Challenges

This study also pointed out several potential challenges, including the need for optimization of SRPG formulation based on site-specific conditions due to variable contamination profiles. Wang et al. [16] mentioned incomplete degradation of some contaminants, which may lead to secondary by-products and additional treatment. Further, Muller et al.[22] emphasized the need to understand long-term environmental impacts related to permanganate residuals and their interaction with aquifer geochemistry.

Remediation of contaminated groundwater with slow-release permanganate gel application is a great achievement for the problems of persistent contaminant plumes in aquifers. Results of this review show the potentials of SRPG to increase efficiency and sustainability in ISCO techniques, especially under hydrogeologically complicated conditions.

The most striking advantage of SRPG, compared to the conventional liquid oxidants, is the huge benefit in remediation of groundwater. Unlike liquid oxidants like potassium permanganate and hydrogen peroxide, which are quick to dissipate and thus have to be frequently reinjected, SRPG slowly releases permanganate over longer periods. Such controlled release can ensure more sustained contact with the contaminant, reducing the over-dosing risks and minimizing problems resulting from frequent application and unwanted changes to the aquifer [18,20, 23].

One of the major innovations of SRPG was the ability to treat low-permeability zones or "contaminant backwaters." The enhanced viscosity of the gel allows for improved retention in porous media and can enable effective treatment of recalcitrant contaminants such as trichloroethylene (TCE) and perchloroethylene (PCE), which frequently bypass conventional remediation strategies [16]. In addition, the resistance of SRPG against displacement by groundwater flow increases its application in dynamic aquifer systems where the usual oxidants get diluted or prematurely transported. The sustained oxidation falls right within the requirements of today's ISCO technologies providing better efficiencies of contaminant degradation [22]. Heterogeneous aquifers, marked by variable permeability and flow dynamics, present significant challenges in remediation efforts. SRPG exhibits promising adaptability in such environments, with its high viscosity allowing diffusion into low-permeability zones often neglected by liquid oxidants. This characteristic ensures more comprehensive treatment across diverse aquifer settings [21, 24].

Despite these advantages, limitations remain. For example, the permanganate release rate of SRPG can be too low in high-flow aquifers to support practical oxidation. Additionally, site-specific gel stability and oxidation efficiency are also dependent on temperature, pH, and microbial activity. These require a local optimization based on these parameters [18, 23]. Addressing these challenges requires advanced numerical modeling and hydrogeological characterization to enhance injection design and predict the behavior of SRPG in complex systems [16]. SRPG technology has distinct scalability and cost-effectiveness advantages compared to traditional remediation methods. The operational cost is reduced due to the reduced frequency of injections and low maintenance costs. It is, therefore, suitable for large-scale and long-term projects. Its effectiveness has been proven through various studies, where over 70% contaminant reduction was achieved in pilot-scale applications, with sustained oxidation for up to six months [16, 21].

The economic and environmental benefits of SRPG are particularly evident in its controlled release mechanism, which minimizes over-dosing and reduces the risk of secondary impacts. However, successful large-scale implementation requires careful tailoring of gel formulations to specific site conditions, such as the contaminant profile and aquifer characteristics. Integrating SRPG with complementary techniques, including bioremediation and monitored natural attenuation, could further enhance its scalability and efficacy [22, 23]. While SRPG significantly reduces contaminant levels and chemical wastage, its long-term environmental impact is a cause for concern. Residual permanganate and manganese oxides, by-products of the oxidation process, can accumulate in aquifers and may impact groundwater quality and usability over time [16]. Monitoring after remediation would be necessary to ensure that the aquifers remain suitable for their intended uses and to mitigate unintended consequences [19].

Another critical concern involves the potential disruption of native microbial communities. Long-term exposure to oxidants can indeed alter microbial diversity and activity so important for maintaining natural biogeochemical processes. The development of bio-compatible gel formulations that cause minimal disruptions but with high oxidation efficiency thus remains a very important area of research [22]. Moreover, incomplete oxidation of contaminants, such as chlorinated solvents, can lead to toxic intermediates. These by-products may need further treatment methods to achieve complete remediation. Advanced analytical techniques and pilot-scale studies are necessary in understanding these processes and devising methods for mitigating adverse effects [18].

Future Directions for Research

More research on SRPG development needs to be undertaken in different respects so that its potential could be optimally utilized in groundwater remediation. Advanced formulations enhancing the stability of gels, improving oxidation efficiency, and minimizing environmental residues; numerical modeling should be combined with pilot-scale testing for fine-tuning of the deployment strategies, simulating the performance in different hydrogeologic environments. Also, Long-term field demonstrations should be done to test the sustainability of the SRPG applications and their efficiency in restoring groundwater quality.

4. CONCLUSION

These reviewed studies provide good evidence regarding the effectiveness of the SRPG approach for contaminant plume remediation. Indeed, the novelty includes sustaining release and applicability to complex aquifers, hence making it a game-changing tool in groundwater management. However, environmental impact, site-specific optimization, and scalability are challenges that have to be resolved if its applicability is to be widened. The contribution of this research also underlines the growing knowledge in the area of sustainable remediation practices and further development in the technologies of groundwater treatment.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Zhao S, Wang J, Zhu W. Controlled-Release Materials for Remediation of Trichloroethylene Contamination in Groundwater. *Materials*. 2023;16(21):7045.
2. Hasan MA, Ahmad S, Mohammed T. Groundwater contamination by hazardous wastes. *Arabian Journal for Science and Engineering*. 2021 May;46(5):4191-4212.
3. Brindha K, Schneider M. Impact of urbanization on groundwater quality. GIS and geostatistical techniques for groundwater science. 2019;2019:179-196.
4. Sun J, Zhang Y, Huang W. Synergistic effects of bioremediation and chemical oxidation using slow-release oxidants. *Environ Pollut*. 2019;252:264-271.
5. Li P, Karunanidhi D, Subramani T, Srinivasamoorthy K. Sources and consequences of groundwater contamination. *Archives of environmental contamination and toxicology*. 2021 Jan;80:1-10.
6. Honetschlägerová L, Martinec M, Škarohlíd R. Coupling in situ chemical oxidation with bioremediation of chloroethenes: a review. *Reviews in Environmental Science and Bio/Technology*. 2019 Dec;18:699-714.
7. Hastings JL, Lee ES. Optimization and analysis of a slow-release permanganate gel for groundwater remediation in porous and low-permeability media. *Water*. 2021;13(6):755.

8. Feng SJ, Zhang X, Zheng QT, Chen HX, Zhao Y. Modeling the spreading and remediation efficiency of slow-release oxidants in a fractured and contaminated low-permeability stratum. *Chemosphere*. 2023;337:139271.
9. Wang W, Jia J, Zhang B, Xiao B, Yang H, Zhang S, Gao X, Han Y, Zhang S, Liu Z, Jin S. A review of Sustained release materials for remediation of organically contaminated groundwater: Material preparation, applications and prospects for practical application. *Journal of Hazardous Materials Advances*. 2024;13:100393.
10. Sakulthaew C, Chokejaroenrat C, Panya S, Songsasen A, Poomipuen K, Imman S, Suriyachai N, Kreetachat T, Comfort S. Developing a Slow-Release Permanganate Composite for Degrading Aquaculture Antibiotics. *Antibiotics*. 2023 Jun 7;12(6):1025.
11. Du X, Zhang X, Liu J, Zhang Z, Wu L, Bai X, Tan C, Gong Y, Zhang Y, Li H. Establishment of evaluation system for biological remediation on organic pollution in groundwater using slow-release agents. *Science of The Total Environment*. 2023;903:166522.
12. Kambhu A, Li Y, Gilmore T, Comfort S. Modeling the release and spreading of permanganate from aerated slow-release oxidants in a laboratory flow tank. *Journal of Hazardous Materials*. 2021;403:123719.
13. Du X, Zhang X, Liu J, Zhang Z, Wu L, Bai X, Tan C, Gong Y, Zhang Y, Li H. Establishment of evaluation system for biological remediation on organic pollution in groundwater using slow-release agents. *Science of The Total Environment*. 2023;903:166522.
14. Ogundare O, Tick GR, Esfahani MR, Akyol NH, Zhang Y. Laboratory-scale characterization of slow-release permanganate gel (SRPG) for the in-situ treatment of chlorinated-solvent groundwater plumes. *Chemosphere*. 2024;360:142392.
15. Acheampong E, Lee ES. Low-temperature slow-release permanganate gel for groundwater remediation: Dynamics in saturated porous media. *Chemosphere*. 2024 Sep 1;363:142716.
16. Wang W, Jia J, Zhang B, Xiao B, Yang H, Zhang S, Gao X, Han Y, Zhang S, Liu Z, Jin S. A review of Sustained release materials for remediation of organically contaminated groundwater: Material preparation, applications and prospects for practical application. *Journal of Hazardous Materials Advances*. 2024 Feb 1;13:100393.
17. Samborska-Goik K, Pogrzeba M. A Critical Review of the Modelling Tools for the Reactive Transport of Organic Contaminants. *Applied Sciences*. 2024;14(9):3675.
18. Yan H, Lai C, Wang D, Liu S, Li X, Zhou X, Yi H, Li B, Zhang M, Li L, Liu X. In situ chemical oxidation: peroxide or persulfate coupled with membrane technology for wastewater treatment. *Journal of Materials Chemistry A*. 2021;9(20):11944-11960.
19. Wang P, Li J, An P, Yang B, Hou D, Pu S. Understanding the dilemmas and breakdown of the reactive migration of in situ groundwater injection reagents from an environmental geology perspective. *Critical Reviews in Environmental Science and Technology*. 2024;54(9):747-70.

20. Chang YC, Chen KF, Chen TY, Chen HH, Chen WY, Mao YC. Development of novel persulfate tablets for passive trichloroethylene (TCE)-contaminated groundwater remediation. *Chemosphere*. 2022 May 1;295:133906.

21. O'Connor D, Hou D, Ok YS, Song Y, Sarmah AK, Li X, Tack FM. Sustainable in situ remediation of recalcitrant organic pollutants in groundwater with controlled release materials: A review. *Journal of Controlled Release*. 2018 Aug 10;283:200-213.

22. Muller KA, Johnson CD, Bagwell CE, Freedman VL. Review of Amendment Delivery and Distribution Methods, and Relevance to Potential In Situ Source Area Treatment at the Hanford Site. Pacific Northwest National Lab.(PNNL), Richland, WA (United States); 2019 Sep 1.

23. Nguyen TM, Chen HH, Chang YC, Ning TC, Chen KF. Remediation of groundwater contaminated with trichloroethylene (TCE) using a long-lasting persulfate/biochar barrier. *Chemosphere*. 2023 Aug 1;333:138954.

24. Wei KH, Ma J, Xi BD, Yu MD, Cui J, Chen BL, Li Y, Gu QB, He XS. Recent progress on in-situ chemical oxidation for the remediation of petroleum contaminated soil and groundwater. *Journal of Hazardous Materials*. 2022 Jun 15;4