

(Original Research Article)

Osmopriming with Polyethylene Glycol (PEG-6000) improves the action of seed germination, growth, and physiology in carrot

ABSTRACT

In order to improve germination and vigour, the current study was conducted using three carrot varieties viz. Carrot Florence (G_1), Deb Kuroda-1 (G_2), and Deb Kuroda-3 (G_3), and various concentrations and durations of PEG-6000, including 0.1 MPa for 24 hours (T_2), 0.1 MPa for 48 hours (T_3), 0.25 MPa for 24 hours (T_4), 0.25 MPa for 48 hours (T_5), 0.40 MPa for 24 hours (T_6), and 0.40 MPa for 48 hours (T_7), non-primed seeds (T_1). A pre-sowing technique called seed priming produces a physiological environment that promotes more efficient seed germination. The experiment was carried out in the Department of Seed Science and Technology's seed testing lab at the BCKV, Mohanpur, Nadia, West Bengal, India. According to the experiment's results, seeds treated with 0.25 MPa PEG-6000 soaking for 48 hours produced the best results among treatments over genotype; these seeds showed notably greater potential than seeds treated with other priming concentrations and durations. Deb Kuroda-3 is the best from a germination perspective, and Deb Kuroda-1 is the best from a vigour perspective. The best results were clearly obtained with a 0.25 MPa PEG-6000 soaking duration of 48 hours for seed quality parameters like germination energy (47.273), seedling Vigour Index-I (639.032), and germination index (5.503). Therefore, to improve seedling establishment, PEG-6000 0.25 Mpa pre-sowing treatment for 48 hours is recommended for carrot.

Keywords: *Germination, PEG-6000, priming, vigour.*

INTRODUCTION

One of the most important vegetable crop in India is carrot (*Daucus carota* L.) ($2n=18$). This is a biennial plant in the Apiaceae family. Since seed is a key component of crop production, optimal seed germination is a prerequisite for a successful stand establishment. These days, the proportion of seed germination, emergence, and vigour of seedlings has been negatively impacted by many environmental and abiotic stressors, which eventually leads to low crop output. Numerous physiological and non-physiological methods are available to improve seed performance and overcome environmental limitations in order to speed up the germination process. Seed priming is a low-cost effective hydration technique to stimulate seed germination. During priming, seeds go through a physiological process, i.e. controlled

hydration and drying which results in enhanced and improved pre-germinative metabolic process for rapid germination. Seed priming can synchronize seed germination, and increase emergence (Heydecker, 1973).

Instead of using pure water, osmopriming entails soaking of seeds in an osmotic solution with a low water potential. The low water potential of osmotic solutions causes water to enter seeds slowly, allowing for progressive imbibition and the activation of early germination phases while preventing radicle protrusion. Various osmotic solutions, including sugar, polyethylene glycol (PEG), glycerol, sorbitol, and mannitol, are used depending on the type of plant, and they are then allowed to air dry before being sown (Slama *et al.*, 2007). Seed priming can improve crop performance under stress conditions, speed up germination, and reduce germination time (Basu, 1976; Chakraborty and Bordolui, 2021).

Benefits of seed priming include improved crop production, maturity, photo and thermo-dormancy release, nutrient uptake, and water use efficiency (Slama *et al.*, 2007). Therefore, our goal was to ascertain the proper PEG 6000 concentration and duration, which are crucial for carrot seed priming. Given the aforementioned factors, the current study investigated the effects of PEG-6000 seed priming at different doses and periods, along with dry seeds as a control, on vigour status, seedling growth, and germination in a laboratory setting.

MATERIALS AND METHODS

In the current study, three carrot genotypes and different osmo priming concentrations and durations were used which was carried out during 2022 at the Seed Testing Laboratory, Department of Seed Science and Technology, BCKV, Mohanpur, Nadia, West Bengal, using a completely randomised design with three replications. Three carrot genotypes were Carrot Florence (G₁), Deb Kuroda-1 (G₂), Deb Kuroda-3 (G₃). PEG-6000 was applied at 0.1 MPa for 24 hrs (T₂), 0.1 MPa for 48 hrs (T₃), 0.25 MPa for 24 hrs (T₄), 0.25 MPa for 48 hrs (T₅), 0.40 MPa for 24 hrs (T₆), 0.40 MPa for 48 hrs (T₇). The control (T₁) was non-primed seeds. AICRP Vegetable provided the seeds, which were analyzed in the Seed Testing Laboratory.

Time to 50% germination

The number of seeds that germinated each day was noted using the AOSA method. The following formulas from Coolbear *et al.* (1984), as modified by Farooq *et al.* (2005), were used to calculate the time of 50% germination (T₅₀):

$$T_{50} = t_i + \frac{\left(\frac{N}{2} - n_i\right)(t_j - t_i)}{(n_j - n_i)}$$

Where, N stands for final number of germination and n_i, n_j are cumulative number of seeds germinated by adjacent counts at times t_i and t_j when $n_i < N/2 < n_j$.

Mean germination time (MGT)

The Ellis and Roberts (1981) equation was used to calculate the mean germination time (MGT):

$$\text{MGT} = \frac{\sum Dn}{\sum n}$$

Where D is the number of days measured from the start of germination and n is the number of seeds that germinated on day D.

Germination percentage

Germination percentage (G) is computed as:

$$G = \frac{X}{Y} \times 100$$

Where X is the number of normal seedlings produced and Y is the total number of seeds taken for germination (ISTA, 1996). Percentage is used to illustrate it.

Germination index (GI)

According to Ruan *et al.*, 2002, the germination index (GI) was calculated using this formula:

$$GI = \frac{\text{Number of germinated seeds}}{\text{Day of first count}} + \dots + \frac{\text{Number of germinated seeds}}{\text{Day of last count}}$$

Germination Energy

On the fourth day after planting, the germination energy (GE) was noted. In relation to the total number of seeds tested, it is the percentage of seeds that germinated 4 days after planting (Ruan *et al.*, 2002).

Germination percentage

Cotton was placed in the petridish, and after that blotting paper was placed on it. Then it was wetted by distilled water. After the seeds were prepared, they were put on the blotting paper and covered with a lid. Such eight pairs of petridish as were kept in the germinator for each genotype and lot. The petridishes were removed from the seed germinator after fourteen days, and the numbers of normal seedlings were counted.

$$\text{Germination (\%)} = \frac{\text{Number of normal seedlings}}{\text{Total number of seeds}} \times 100$$

Seedling parameters: Root lengths and shoot lengths of ten seedlings were measured at 14 days after germination using the glass plate method in the laboratory with the help of a scale and graph paper and after that average was made out, expressed in centimetre (cm). A digital balance was used to measure the fresh weight of ten seedlings. After two hours of drying in a hot air oven at 80⁰C, the seedlings were weighed using a digital balance. The fresh weight and dry weight of the seedlings were both stated in grams (g).

Vigour index: Vigour index (VI) was computed by using the formula advised by Abdul-Baki and Anderson (1973): VI= G X L Where, 'G' stands for germination percentage and 'L' denotes average seedling length (cm).

RESULTS AND DISCUSSION

Time of 50% Germination (Days)

The highest time to 50% germination over genotypes (6.529) was observed to produce by T₁ on an average followed by T₂, T₃ and T₄; while it was of shortest length for T₅ preceded by T₆ and T₇. Elkoca *et al.* (2007) observed that Osmo- priming by PEG solution improved time of 50% germination after seed treatment in pea. Highest time to 50% germination (5.725) was observed for G₁ and lowest time to 50% germination was recognized for G₃, (4.020) over treatments (Table 1). When the interaction effect of genotypes and seed treatments were taken into consideration, G₁T₁ showed highest value (7.753) for this parameter. Kundu and Bordolui (2023) found a similar result in carrots primed with Ag-nano particles.

Table 1. Effect of osmo-priming on Time of 50% Germination (days) of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	7.753	6.367	6.000	5.733	4.233	4.767	5.220	5.725
G ₂	6.433	5.900	5.200	4.600	3.800	4.033	4.333	4.900
G ₃	5.400	4.797	4.067	3.667	3.207	3.450	3.550	4.020
Mean G	6.529	5.688	5.089	4.667	3.747	4.083	4.368	
		G	T	GXT				
SEm (±)		0.039	0.060	0.104				
LSD (0.05)		0.113	0.173	0.299				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40MPa PEG-6000 for 48 hrs.								




Fig.1. Graphical representation of Time of 50% Germination (days)
Mean Germination Time (Days)

Treatments over genotypes, highest mean germination time was observed in T₁ (7.922) followed by T₂, T₃ and T₄; while it was minimum for T₅ preceded by T₆ and T₇. Hasan *et al.* (2016) found that Osmo-priming improved mean germination time after seed treatment in rice. Over treatments the highest mean germination time was observed in G₁ (7.130) and lowest for G₃, (5.407) (Table 2). G₁T₁ showed highest value (9.147) for this parameter when interaction was made between genotypes and seed treatments. But interaction value was non-significant. G₂T₁, G₁T₂; G₃T₂, G₁T₆; G₃T₃, G₂T₆ were statistically at par. Ray and Bordolui found a similar result in tomatoes (2022a).

Table 2. Effect of osmo-priming on Mean Germination Time (days) of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G₁	9.147	7.793	7.393	7.147	5.660	6.160	6.613	7.130
G₂	7.827	7.290	6.593	5.993	5.187	5.427	5.707	6.289
G₃	6.793	6.140	5.460	5.060	4.623	4.860	4.910	5.407
Mean G	7.922	7.074	6.482	6.067	5.157	5.482	5.743	
		G	T	GXT				
SEm (±)		0.037	0.056	0.098				
LSD (0.05)		0.106	0.162	0.280				

Note: G = Genotypes, G₁ = Carrot Florence, G₂ = Deb Kuroda-1, G₃ = Deb Kuroda-3; T = Treatment, T₁ = Control, T₂ = 0.1 MPa PEG-6000 for 24 hrs, T₃ = 0.1 MPa PEG-6000 for 48 hrs, T₄ = 0.25 MPa PEG-6000 for 24 hrs, T₅ = 0.25 MPa PEG-6000 for 48 hrs, T₆ = 0.40 MPa PEG-6000 for 24 hrs, T₇ = 0.40 MPa PEG-6000 for 48 hrs.

Germination index

Highest germination index over genotypes was observed in T₅ (5.503) followed by T₆, T₇, and T₄, whereas T₁ (control) had the lowest germination index (2.089) preceded by T₂ and T₃. Sadeghi *et al.* (2011) found that Osmo-priming by PEG solution improved time of germination index after seed treatment in soybean. Over the treatments, G₃ (4.922) had the highest germination index and G₁ (4.130) had the lowest (Table 3). When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ (5.670) showed highest value for this parameter. But G₁T₁ G₂T₁; G₁T₅, G₂T₇ and G₃T₃; G₃T₄, G₃T₇ were statistically at par.

Table 3. Effect of osmo-priming on Germination index of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	2.087	3.700	4.067	4.250	5.247	5.033	4.530	4.130
G ₂	2.057	4.633	5.153	5.213	5.593	5.377	5.250	4.754
G ₃	2.123	5.157	5.280	5.353	5.670	5.500	5.370	4.922
Mean G	2.089	4.497	4.833	4.939	5.503	5.303	5.050	
		G	T	GXT				
SEm (±)		0.016	0.025	0.044				
LSD (0.05)		0.047	0.072	0.125				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40 MPa PEG-6000 for 48 hrs.								

Germination Energy (%)

The highest germination energy over genotypes was observed in T₅ (**47.273**) on an average followed by T₆, T₇ and T₄; while it was of lowest for T₁ (control) preceded by T₂ and T₃. Sadeghi *et al.* (2011) found that Osmo-priming by PEG solution improved time of germination energy after seed treatment in soybean. Highest germination energy (42.450) was observed for G₃ and lowest germination energy was recognized for G₁ (40.256) over treatments (Table 4). When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ showed highest value (47.687) for this parameter, though G₁T₃ and G₂T₂; G₃T₇ and G₂T₇; G₂T₆, G₃T₆, were statistically at par with each other.

Table 4. Effect of osmo-priming on Germination Energy (%) of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	18.833 (25.707)	40.467 (39.488)	42.377 (40.599)	43.680 (41.353)	46.700 (43.091)	45.367 (42.324)	44.367 (41.749)	40.256 (39.187)
G ₂	21.667 (27.721)	42.167 (40.477)	42.967 (40.940)	45.167 (42.209)	47.433 (43.511)	46.333 (42.880)	45.333 (42.305)	41.581 (40.006)

G₃	23.833 (29.208)	43.500 (41.248)	44.100 (41.595)	45.333 (42.305)	47.687 (43.656)	46.767 (43.139)	45.933 (42.650)	42.450 (40.542)
Mean G	21.444 (27.545)	42.044 (40.405)	43.148 (41.045)	44.727 (41.956)	47.273 (43.419)	46.156 (42.778)	45.211 (42.235)	
		G	T	GXT				
SEm (±)		0.087	0.133	0.230				
LSD (0.05)		0.249	0.380	0.659				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40MPa PEG-6000 for 48 hrs.								

Shoot Length (cm)

The longest shoot length over genotypes (3.759cm) was observed to produce by T₅ followed by T₆, T₇ and T₄; while it was of shortest length for T₁ (control) preceded by T₂ and T₃. Farooq *et al.* (2005) showed increased shoot length after seed treatment with Osmo-priming by PEG solution in rice. Highest shoot length (3.686 cm) was observed for G₃ and shortest shoot length (2.907 cm) was recognized for G₁, over treatments (Table 5). Though G₂ and G₃ over treatment were non-significantly differ. When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ showed highest value (4.392 cm) for this parameter, though G₁T₁ and G₂T₁; G₂T₂ and G₃T₂; G₁T₆, G₁T₇ were statistically at par with each other. Similar outcomes were noted by Choudhury and Bordolui (2022a) in Bengal gram when they used sodium molybdate (Na₂MoO₄) nutri-priming to increase shoot length.

Table 5. Effect of osmo-priming on Shoot Length (cm) of carrot genotypes

	T₁	T₂	T₃	T₄	T₅	T₆	T₇	Mean T
G₁	2.400	2.743	2.880	2.970	2.550	3.474	3.330	2.907
G₂	2.503	2.925	3.090	3.585	4.335	3.726	3.774	3.420
G₃	2.823	2.947	3.437	4.266	4.392	4.074	3.864	3.686
Mean G	2.576	2.872	3.136	3.607	3.759	3.758	3.656	
		G	T	GXT				
SEm (±)		0.037	0.057	0.099				
LSD (0.05)		0.107	0.164	0.284				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40MPa PEG-6000 for 48 hrs.								

Root length (cm)

T₅ (3.165 cm) produced highest root length (3.141 cm) over genotypes, followed by T₄, T₆, and T₃, whereas T₁ (control) had the least root length, preceded by T₂ and T₇. In pea,

Yanglem *et al.* (2021) found that Osmo-priming by PEG solution improved root length after seed treatment. Over the treatments, G₂ had the highest root length (3.124 cm), and G₁ had the smallest root length (2.210 cm) (Table 6). Despite the fact that G₁ and G₃, over treatments showed non-significant difference. The interaction between genotypes and seed treatments G₃T₅ showed highest value (3.165 cm) for this parameter, though G₁T₂ and G₂T₂; G₁T₃ and G₁T₄ were statistically at par with each other. Choudhury and Bordolui (2022b) used potassium nitrate to observe a similar kind of result in Bengal gram.

Table 6. Effect of osmo-priming on Root length (cm) of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	2.267	2.136	2.289	2.175	2.634	2.020	1.950	2.210
G ₂	2.523	3.193	3.247	3.381	3.623	3.057	2.844	3.124
G ₃	2.313	2.397	2.445	2.610	3.165	2.913	2.994	2.691
Mean G	2.368	2.575	2.660	2.722	3.141	2.663	2.596	
		G	T	GXT				
SEm (±)		0.041	0.062	0.108				
LSD (0.05)		0.117	0.178	0.309				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40 MPa PEG-6000 for 48 hrs.								

Seedling length (cm)

Among the treatments over genotypes, T₁ (control) observed shortest seedling lengths, which was preceded by T₂ and T₃, while T₅ produced seedlings with highest length of 6.900 cm, followed by T₆, T₄, and T₇. According to Singh *et al.* (2015), cowpea seeds treated with Osmo-priming by PEG solution produced longer shoots. Here, G₁ and G₂ are non-significantly differ. According to Singh *et al.*, cowpea seeds treated with Osmo-priming by PEG solution produced longer shoots (2014). G₂ had the longest shoots (6.544 cm) and G₁ had the shortest shoots (5.117 cm) over treatments (Table 7). When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ showed highest value (7.557 cm) for this parameter, though G₃T₁ and G₁T₄; G₁T₃ and G₁T₄ were statistically at par with each other.

Table 7. Effect of osmo-priming on Seedling length (cm) of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	4.667	4.879	5.169	5.145	5.184	5.494	5.280	5.117
G ₂	5.027	6.118	6.337	6.966	7.958	6.783	6.618	6.544
G ₃	5.137	5.343	5.882	6.876	7.557	6.987	6.858	6.377

Mean G	4.943	5.447	5.796	6.329	6.900	6.421	6.252	
		G	T	GXT				
SEm (±)		0.060	0.091	0.158				
LSD (0.05)		0.171	0.261	0.453				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40MPa PEG-6000 for 48 hrs.								

Germination percentage

In case of treatments over genotypes significantly differ with each other. But T₃ recorded highest germination percentage (93.644), followed by T₄, T₂, and T₆ whereas T₁ (control) produced the lowest germination percentage preceded by T₇ and T₅. T₃ and T₄ were non-significantly differing with each other. Lemmens *et al.* (2019) found that Osmo-priming by PEG solution improved germination percentage in wheat. Over the treatments, G₁ showed the lowest germination percentage (91.312) while G₃ had the highest germination percentage (92.126) (Table 8). Interaction between genotypes and seed treatments G₃T₃ observed highest value (93.967). G₁T₁, G₃T₁; G₂T₂, G₃T₂, G₁T₃, G₃T₂; G₂T₄ and G₂T₅ were statistically at per. Ray and Bordolui (2022b) discovered a similar kind of outcome in tomato.

Table 8. Effect of osmo-priming on Germination percentage of carrot genotypes

	T₁	T₂	T₃	T₄	T₅	T₆	T₇	Mean T
G₁	85.583 (67.661)	91.300 (72.833)	93.367 (75.128)	93.400 (75.091)	93.533 (75.245)	91.867 (73.404)	90.133 (71.666)	91.312 (73.004)
G₂	87.083 (68.910)	93.600 (75.323)	93.600 (75.380)	92.333 (73.904)	92.833 (74.444)	93.267 (74.933)	90.833 (72.356)	91.936 (73.607)
G₃	85.917 (67.932)	93.933 (75.748)	93.967 (75.768)	93.867 (75.634)	91.767 (73.307)	93.267 (74.954)	92.167 (73.726)	92.126 (73.867)
Mean G	86.194 (68.168)	92.944 (74.635)	93.644 (75.425)	93.200 (74.876)	92.711 (74.332)	92.800 (74.430)	91.044 (72.583)	
		G	T	GXT				
SEm (±)		0.192	0.294	0.509				
LSD (0.05)		0.551	0.842	1.458				
Note: G = Genotypes, G ₁ = Carrot Florence, G ₂ = Deb Kuroda-1, G ₃ = Deb Kuroda-3; T = Treatment, T ₁ = Control, T ₂ = 0.1 MPa PEG-6000 for 24 hrs, T ₃ = 0.1 MPa PEG-6000 for 48 hrs, T ₄ = 0.25 MPa PEG-6000 for 24 hrs, T ₅ = 0.25 MPa PEG-6000 for 48 hrs, T ₆ = 0.40 MPa PEG-6000 for 24 hrs, T ₇ = 0.40MPa PEG-6000 for 48 hrs.								




Fig. 2. Graphical representation of Germination (%)

Vigour Index

The highest vigour index over genotypes was observed in T₅ (639.032) followed by T₆, T₄ and T₇; while it was lowest for T₁ (control) preceded by T₂ and T₃. Rouhi *et al.* (2010) found that Osmo-priming by PEG solution improved vigour index in clover. Highest vigour index (602.773) was observed for G₂ and lowest vigour index (467.527) was recognized for G₁, over treatments (Table 9). Though G₁ and G₃ over treatment were non-significantly differ. When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ showed highest value (693.447) for this parameter, though G₂T₁ and G₁T₁; G₁T₂, G₃T₁; G₁T₄ and G₁T₅ were non-significant with each other.

Table 9. Effect of osmo-priming on Vigour Index of carrot genotypes

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G₁	399.333	445.413	482.173	480.330	484.873	504.630	475.937	467.527
G₂	437.747	572.743	593.107	643.340	738.777	632.587	601.113	602.773
G₃	441.340	501.860	552.717	645.407	693.447	651.760	631.950	588.354
Mean G	426.140	506.672	542.666	589.692	639.032	596.326	569.667	
		G	T	GXT				
SEm (±)		5.390	8.234	14.261				
LSD (0.05)		15.438	23.582	40.845				

Note: G = Genotypes, G₁ = Carrot Florence, G₂ = Deb Kuroda-1, G₃ = Deb Kuroda-3; T = Treatment, T₁ = Control, T₂ = 0.1 MPa PEG-6000 for 24 hrs, T₃ = 0.1 MPa PEG-6000 for 48 hrs, T₄ = 0.25 MPa PEG-6000 for 24 hrs, T₅ = 0.25 MPa PEG-6000 for 48 hrs, T₆ = 0.40 MPa PEG-6000 for 24 hrs, T₇ = 0.40MPa PEG-6000 for 48 hrs.




Fig. 3. Graphical representation of Vigour Index (%)

Seedling Fresh weight (mg) of 10 seedlings

Over genotypes, T₅ produced the highest fresh weight (107.556 mg) followed by T₆, T₇, and T₄; whereas T₂ showed the lowest fresh weight, preceded by T₁ and T₃. Ghiyasi *et al.* (2008) found that Osmo-priming by PEG solution improved fresh weight after seed treatment in case of wheat. Similarly, Chakraborty and Bordolui (2021) discovered that Ag nano priming increased the fresh weight of green grams seedlings compared to other treatments. Genotypes over treatments, G₃ had the highest fresh weight (93.714 mg) and G₁ observed the lowest fresh weight (79.667 mg) (Table 10). When the interaction effect of genotypes and seed treatments were taken into consideration, G₃T₅ showed highest value (108.333 mg) for this parameter but they were non-significantly differ with each other.

Table 10. Effect of osmo-priming on Seedling Fresh weight (mg) of carrot genotypes (10 seedlings)

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	62.333	49.000	76.000	87.333	102.000	95.000	86.000	79.667
G ₂	72.333	80.000	86.000	94.667	112.333	104.333	106.333	93.714
G ₃	74.667	79.000	82.667	92.000	108.333	103.000	103.333	91.857
Mean G	69.778	69.333	81.556	91.333	107.556	100.778	98.556	
		G	T	GXT				
SEm (±)		1.806	2.759	4.779				
LSD (0.05)		5.173	7.902	NS				

Note: G = Genotypes, G₁ = Carrot Florence, G₂ = Deb Kuroda-1, G₃ = Deb Kuroda-3; T = Treatment, T₁ = Control, T₂ = 0.1 MPa PEG-6000 for 24 hrs, T₃ = 0.1 MPa PEG-6000 for 48 hrs, T₄ = 0.25 MPa PEG-6000 for 24 hrs, T₅ = 0.25 MPa

PEG-6000 for 48 hrs, T₆ = 0.40 MPa PEG-6000 for 24 hrs, T₇ = 0.40MPa PEG-6000 for 48 hrs.

Seedling Dry Weight (mg) of 10 seedlings

The highest dry weight over genotypes was observed in T₅ (11.508) followed by T₆, T₇, and T₄. But, T₂ showed the lowest dry weight, preceded by T₁ and T₃. T₁ and T₂ were statistically at par. Ghiyasi *et al.* (2008) found that Osmo- priming by PEG solution improved dry weight after seed treatment in case of wheat. Over the treatments, G₃ had the highest dry weight (10.028), and G₁ had the lowest dry weight (8.522) (Table 11). G₁ and G₂ over treatments were non-significantly differing. The interaction effect of genotypes and seed treatments were non-significantly variation with each other but G₃T₅ showed highest dry weight (11.590).

Table 11. Effect of osmo-priming on Seedling Dry Weight (mg) of carrot genotypes (10 seedlings)

	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	Mean T
G ₁	6.670	5.223	8.133	9.343	10.913	10.167	9.203	8.522
G ₂	7.740	8.560	9.203	10.130	12.020	11.163	11.380	10.028
G ₃	7.990	8.453	8.843	9.847	11.590	11.020	11.057	9.829
Mean G	7.467	7.412	8.727	9.773	11.508	10.783	10.547	
		G	T	GXT				
SEm (±)		0.193	0.295	0.511				
LSD (0.05)		0.553	0.845	NS				

Note: G = Genotypes, G₁ = Carrot Florence, G₂ = Deb Kuroda-1, G₃ = Deb Kuroda-3; T = Treatment, T₁ = Control, T₂ = 0.1 MPa PEG-6000 for 24 hrs, T₃ = 0.1 MPa PEG-6000 for 48 hrs, T₄ = 0.25 MPa PEG-6000 for 24 hrs, T₅ = 0.25 MPa PEG-6000 for 48 hrs, T₆ = 0.40 MPa PEG-6000 for 24 hrs, T₇ = 0.40MPa PEG-6000 for 48 hrs.

Conclusion

Carrot seeds treated with PEG-6000 had better seed quality than the control. In comparison to other treatments, PEG-6000 @ 0.25Mpa soaking for soaking duration 48 hours was the most effective treatment over genotypes. Significantly highest germination index, germination energy, germination percentage and lowest mean germination time were noted for Deb Kuroda-3 (G₃) while highest seedling length, fresh weight, dry weight and vigour index were observed for Deb Kuroda-1(G₂) although these genotypes were statistically at par. So, in germination point of view, Deb Kuroda-3 is best and in vigour point of view, Deb Kuroda-1 is best. For seed quality parameters such as germination energy (47.273), seedling vigour Index-I (639.032), and germination index (5.503), PEG-6000 @ 0.25Mpa soaking for 48 hours shown noticeably the best results. Consequently, PEG-6000 @ 0.25Mpa for soaking

duration 48 hours is advised as a pre-sowing treatment for carrot seeds in order to improve seedling establishment.

Disclaimer (Artificial intelligence)

The authors of this manuscript hereby declare that no generative AI technologies, including text-to-image generators and Large Language Models (ChatGPT, COPILOT, etc.), were used in its writing or editing.

REFERENCES

- Abdul-Baki, A. and Anderson, J.D. (1973). Vigor Determination in Soybean Seed by Multiple Criteria. *Crop Science*. 13:630-633
- Basu, R.N. (1976). Physico-chemical control of seed deterioration. *Seed Research*. 4(1):15-23.
- Chakraborty, A. and Bordolui, S.K. (2021). Impact of Seed Priming with Ag-Nanoparticle and GA₃ on Germination and Vigor in Green gram. *Int.J.Curr.Microbiol.App.Sci*. 10(03): 941-950. doi: <https://doi.org/10.20546/ijcmas.2021.1003.119>
- Chakraborty, A. and Bordolui, S.K. (2021). Standardization of the Appropriate Doses of GA₃ and Ag-Nanoparticle in Green Gram for Quality Seed Production. *International Journal of Environmental & Agriculture Research*. 7(04), 1-11.
- Choudhury, A. and Bordolui S.K. (2022b). Inducement of Seed Priming with Potassium Nitrate on quality Performance of Chickpea (*Cicer arietinum* L.). *Biological Forum – An International Journal*. 14(4), 779-783.
- Choudhury, A. and Bordolui, S.K. (2022a). Seed invigoration treatment with sodium molybdate (Na₂MoO₄) nutri-priming for improvement of quality performance of Bengal gram (*Cicer arietinum* L.). *The Pharma Innovation Journal*. 11(12), 3381-3386.
- Coolbear, P., Francis, A., and Grierson, D. (1984). The effect of low temperature pre-sowing treatment on the germination performance and membrane integrity of artificially aged tomato seeds. *Journal of Experimental Botany*. 35(11): 1609-1617.
- Elkoca, E., Haliloglu, K. and Esitken, A. (2007). Hydro and osmopriming improve chickpea germination. *Acta Agriculturae Scandinavica Section B-soil and Plant Science year*. 57(3): 193-200.

- Ellis, R.H. and Roberts E.H.(1981).The quantification of ageing and survival in orthodox seeds. *Seed Science and Technology*. 9: 373-409.
- Farooq, M., Basra, S.M.A., Ahmad, N. and Hafeez, K. (2005). Thermal hardening: A new seed vigor enhancement tool in rice. *Journal of Integrative Plant Biology*. 47(2):187-193.
- Ghiyasi, M., Abbasi S.E., Mehdi, T., Amirnia, R. and Hojat, S. (2008). Effect of Osmopriming with Polyethylene Glycol (8000) on Germination and Seedling Growth of Wheat (*Triticum aestivum* L.) Seeds under Salt Stress. *Research Journal of Biological Sciences*.3
- Hasan, M.N. , Salam, M.A. ,Chowdhury, M.M.I. , Sultana, M. and Islam, N.(2016). Effect of osmopriming on germination of rice seed. *Bangladesh J. Agril.* 41(3): 451-460
- Heydecker, W. (1973). “Germination of an Idea: The Priming of Seeds,” School of Agriculture Research, University of Nottingham, Nottingham. 50-67.
- ISTA (1996). International Rules for Seed Testing 1996. The International Seed Testing Association, Zurich.
- Kundu, E. and Bordolui, S.K. (2023). Silver Nanoparticles –mediate Seed priming Improves Germination and Physiological Performance in Carrot. *Biological Forum – An International Journal*. 15(10): 1079-1085.
- Lemmens, E., Deleu, L.J., De Brier, N., De Man, W.L., De Proft, M., Prinsen, E. and Delcour, J.A. (2019). The Impact of Hydro-Priming and Osmo-Priming on Seedling Characteristics, Plant Hormone Concentrations, Activity of Selected Hydrolytic Enzymes, and Cell Wall and Phytate Hydrolysis in Sprouted Wheat (*Triticum aestivum* L.). *ACS omega*. 4(26): 22089–22100.
- Ray, J. and Bordolui, S.K. (2022a). Effect of seed priming as pre-treatment factors on germination and seedling vigour of tomato. *International Journal of Plant & Soil Science*. 34(20), 302-311.
- Ray, J. and Bordolui, S.K. (2022b). Seed quality deterioration of tomato during storage: Effect of storing containers and condition. *Biological Forum – An International Journal*. 14(2), 137-142.

- Rouhi, H.R., Afshari, R.T., Moosavi, S. and Gharineh, M.H. (2010). Effects of Osmopriming on Germination and Vigour Traits of Bersim Clover (*Trifolium alexandrinum* L.). *Notulae Scientia Biologicae*. 2(4): 59-63.
- Ruan, S., Xue, Q., and Tylkowska, K. (2002). The influence of priming on germination of rice (*Oryza sativa* L.) seeds and seedling emergence and performance in flooded soil. *Seed Sci Tech*. 30: 61-67.
- Sadeghi, H., Khazaei, F., Yari, L. and Sheidaei, S. (2011). Effect of seed osmopriming on seed germination behavior and vigor of soybean (*Glycine max* L.). *ARP Journal of Agricultural and Biological Science*. 6
- Singh, H., Jassal, R.K., Kang, J.S., Sandhu, S.S., Kang, H. and Grewal, K. (2015). Seed priming techniques in field crops - A review. *Agri. Review*. 36 (4): 251-264.
- Slama, I., Ghnaya, T., Hessini, K., Messedi, D., Savoure, A. and Abdely, C. (2007). Comparative study of the effects of mannitol and PEG osmotic stress on growth and solute accumulation in *Sesuvium portulacastrum*. *Environmental and Experimental Botany*. 61(1):10-17. DOI: 10.1016/j.envexpbot.2007.02.004
- Yanglem, S.D. and Ram, V. (2021). Effects of seed priming on root-shoot behaviour and stress tolerance of pea (*Pisum sativum* L.). *Bangladesh Journal of Botany*. 50(2): 199-208.