

Gross morphological and radiographic studies on the bones of the shoulder and arm in Indian grey mongoose or Asian grey mongoose (*Herpestes edwardsii*)

ABSTRACT

The scapula of the Indian grey mongoose was a flat and trapezoid shaped bone with two surfaces, four borders and four angles. The scapular spine divided its lateral surface into two unequal fossae and terminated as a pointed hamate process, the latter presented a triangular suprahamate process whose apex was pointed and directed caudally. The glenoid cavity was shallow with a cranially projected supraglenoid tubercle that possessed medially a small indistinct coracoid process. The humerus was a long bone with a distinct spirally twisted shaft and two extremities. Its proximal extremity had an ovoid head, lesser tuberosity placed medial to the head and a greater tuberosity lateral to the head. The bicipital groove was wide and shallow. The musculospiral groove was deep. The proximal half of the shaft had a medially located pectoral ridge and a laterally situated prominent sharp deltoid crest. The distal end had two condyles, two epicondyles and two fossae, a cranially placed shallow coronoid fossa and caudally placed deep olecranon fossa. The two fossae communicated cranially through the supratrochlear foramen. A sharp prominent lateral epicondylar crest was present and above the medial epicondyle there was the oval supracondyloid foramen.

Key words: Mongoose, scapula, hamate process, humerus, supratrochlear, supracondyloid foramen.

INTRODUCTION

Mongoose is a small terrestrial carnivorous mammal belonging to the family Herpestidae. The subfamily Herpestinae comprises 23 living species that are native to southern Europe, Africa and Asia [1]. The Indian grey mongoose or Asian grey mongoose (*Herpestes edwardsii*) is a mongoose species native to the Indian subcontinent and West Asia. This is one of 5-6 species in the Indian subcontinent and is the state animal of Chandigarh. This species is considered as the least concern in IUCN red list [2]. It is a good climber and can also swim. They have short limbs and are known for their speed and agility with their short limbs and are capable of combating venomous snakes. Scientific studies on the anatomy and morphology of this animal has become the area of recent interest for wildlife biologist. Research studies on various organs and organ systems such as its skull, brain, spinal cord, tongue, liver, pancreas, vertebral column and hindlimb bones etc., of the Indian Grey mongoose and related species in this Herpestinae family has begun to add up to the available pool of literature. Yet, studies are not yet done on the bones of the pectoral limb of this species. This current study on the gross morphological and radiographic anatomy of the bones of the shoulder and arm region in Indian grey mongoose (*Herpestes edwardsii*) is done to fill and enrich this knowledge gap.

MATERIALS AND METHODS

The cadaver of the Indian grey mongoose from the Puducherry region of India was utilized for the present study. A total of ten adult animals of both sexes that suffered natural death was utilized for this study. They were referred by the Department of Forest and Wildlife, Puducherry for their post mortem examination in the Department of Veterinary Pathology, Rajiv Gandhi Institute of Veterinary Education and Research, (RIVER) Puducherry. These cadavers were then subjected to radiographic inspection of the bones and joints in the radiology Unit, Teaching Veterinary Clinical complex, RIVER. After radiographic examination, they were carefully removed off their skin and muscles by careful dissection to the best possible extent and the bony remains were subjected to maceration without burying as recommended by [3] for smaller specimens and [4] for human cadavers. Necessary precautions were taken during processing with chemicals such as Potassium hydroxide and hydrogen peroxide for this study. The bones were then brushed with a soft bristled tooth brush to make them clean and to get rid of any adhering remnants. They were finally washed thoroughly with running tap water to get rid of the residual chemicals and then air dried as recommended by [5]. These bones were used to study for their gross morphological features and were photographed with Nikon digital camera for their features.

RESULTS AND DISCUSSION

Scapula

The scapula of the mongoose was flat and almost trapezoid in shape (Fig 1). Such quadrangular shaped scapula was reported by [6] in civet cat; [7] in Indian wild cat; [8] in tiger and [9] in Himalayan mongoose. Whereas, [10] described a nearly triangular scapula in New Zealand rabbit and domestic cat. In leopard cat and fishing cat it was rounded and an almost triangular scapula was present in the small Indian civet cat [11]. In Indian grey mongoose the scapula had 2 surfaces, 4 borders and 4 angles. The surfaces were lateral and medial (Fig 1). The lateral surface presented a sharp edged spine which gradually increased in height and presented in the middle a tuberosity. The scapular spine in its distal one third, presented a notch. This spine terminated as the pointed acromion or the hamate process, which was at the level of the glenoid cavity (Fig 1a).

In lion, the acromion process hanged over the glenoid cavity [12]. Contrarily, [13] observed that the scapular spine diminished from the tuber spine and did not form the acromion process. The spine in this species divided the lateral surface into two unequal fossae; supraspinous and infraspinous fossae as reported by [9] in the Himalayan mongoose. In contrast, the scapula was divided into two equal fossae in lion [7], domestic cat [10], great Indian rhino [13], dog [14], Hedge hog [15], Indian wild cat [16].

According to [17] in sloth bear, the scapula had an additional spine that divided the supraspinous fossa into cranial and caudal parts. In the present study, the scapular spine was inclined towards the infraspinous fossa as reported by [11] in fishing cat and small Indian civet cat. The acromion or hamate process in this species presented a triangular suprahamate process with its base attached to the spine. It deviated from the spine at an acute angle (Fig 1a). Similar to the above finding, [12] described that suprahamate process was a thick triangular backwardly

directed plate in lion and [6] termed it as metacromion process that was roughly triangular and caudally directed in civet cat. The apex of the suprahumeral process was pointed caudally towards the infraspinous fossa as observed by [9] in the Himalayan mongoose and [11] in fishing cat. The suprahumeral process was broad and semicircular in cat and was an irregular L shaped process in the rabbit [10]. In contrast, the metacromion process was found absent in dog [18].

In the present study, the cranial border of scapula was curved with a small indentation anteriorly, just below the cranial angle and a prominent notch distally as reported by [9] in the Himalayan mongoose. The dorsal border curved from the cranial angle and terminated caudally as a thick blunt projection. The caudal border was thick. From this thick blunt projection, it curved slightly till the middle, forming a convexity and then dropped straight down. In lion, a pointed projection was seen from the caudal angle of the scapula [12]. The ventral border included the curve of the cranial border and the rim of the glenoid cavity (Fig 1a). The cranial angle was obtuse. Caudal angle was acute and the thickest. The cranio-ventral angle was right angled and the caudo-ventral angle was slightly obtuse. A nutrient foramen was noticed at the base of the spine, towards the glenoid angle which coincides with the observation of [14]. The glenoid cavity in this species was in the shape of an oblong circle. But, it was rectangular in Indian elephants [6], oval to quadrangular in tiger [8] circular and deep in black Bengal goat [19], elongated in elephant [20]. In the present study, the rim of the glenoid cavity, near the posterior border of scapula was wide and round, while near the anterior border it was narrow. The glenoid cavity was shallow and a muscular process called the supraglenoid tubercle projected in front of the cavity that gave origin to the biceps brachii muscle (Fig 4a). These findings are similar to the dog [21]. This tubercle projected cranially and had a medial inclination as reported by [14] in dog. The glenoid notch was present along the ventromedial aspect of the rim of the glenoid cavity. Whereas, it was on the lateral aspect of the rim in the Himalayan mongoose [9] and on the anterolateral aspect of the rim in fishing cat [11].

The scapula of the Indian grey mongoose on its medial surface presented a relatively shallow subscapular fossa. Cranial and caudal to the fossa was a cranial and a caudal ridge as reported in tiger by [8], whereas four ridges were seen in civet cat [6]. On the other hand, the subscapular fossa was distinct with two prominent ridges in hippopotamus [22]. The coracoid process in this species was small and indistinct situated on the medial aspect of the supraglenoid tubercle (Fig 4a). It was hook like in civet cat [6], tiger [8], rounded in Himalayan mongoose [9], beak like in rabbit [10] and in domestic cat [10, 23]. The coracoid process is a remnant of the coracoid bone that is still distinct in monotremes [14].

Humerus:

The humerus of the mongoose was a long bone of the arm. It was placed obliquely downward and backward and articulated proximally with the glenoid fossa of the scapula and distally with the radius and ulna (Fig 5). It had a spirally twisted shaft and two extremities proximal and distal. The shaft presented 4 surfaces namely anterior, posterior, medial and lateral as studied in mongoose [24]. The proximal extremity consisted of a posteromedially placed smooth, ovoid head and two tuberosities, medial and lateral (Fig 4b) as in dog [18] and cat [10, 25]. Whereas, it

was spherical in civet cat [26] long and craniocaudally curved in lion [12]. This ovoid head in mongoose would permit a high degree of extension and flexion in this species. Lateral to the humeral head was the greater or lateral tuberosity. The lesser or medial tuberosity was located just medial to the head. (Fig 4b). Anteriorly, between the two tuberosities was placed the wide and shallow bicipital groove, through which the tendon of biceps brachii passed (Fig 2b, 4b) as reported by [24] in mongoose. A distinct well defined neck was noticed in the present study. The lateral tuberosity presented on its posterolateral aspect, a deep depression. But a grooved lateral tuberosity was reported by [27] in the Indian tiger and was a deep notch in mongoose [24].

The shaft of the humerus in this species was distinctly twisted with a deep musculospiral groove (Fig 2a) as reported by [24]. The proximal half of the shaft presented a medially placed pectoral ridge (Fig 2b) and a lateral prominent deltoid crest. (Fig 2a). The pectoral ridge originated from the greater tuberosity and terminated in the well-developed teres tubercle. A prominent teres tubercle was also seen in domestic cat [25]. The deltoid crest in the present study was prominent and sharp and was noticed on the lateral border as reported by [27] in tiger (Fig 2a). In contrast, [24] reported a sharp deltoid tuberosity in mongoose and a complete absence of it in domestic cat [10].

The medial surface of humerus was flat, wide above and narrow behind. The posterior surface was rough and narrow that widened distally. The distal end of the humerus had 2 condyles, 2 epicondyles and 2 fossae (Fig 4c, 4d). Of the 2 condyles the medial one was larger. The humeral condyles involved its articular area, together with the adjacent fossae. The radial or coronoid fossa was shallow and situated cranially above the trochlea. In dog coronoid fossa is absent [14]. In this study in Indian grey mongoose the epicondyles above the condyles enclosed the olecranon fossa caudally. The coronoid and olecranon fossa communicated through the supratrochlear foramen which was present in the bony plate that separated the 2 fossae (Fig 4c, 4d). The olecranon fossa was deep and in addition to the supratrochlear foramen it also had several small foramina. This finding is in agreement with [10] who reported that supratrochlear foramen of different size perforated the bony lamina in rabbit. The margin of the olecranon fossa in the present study was sharp on the lateral aspect, while on the medial aspect, the margin continued onto the medial epicondyle as in tiger [27]. The supratrochlear foramen may be absent when the humerus is small in dog [14]. The presence of supratrochlear foramen in mammals was variable and was linked to the range of mobility in the elbow joint [28].

A sharp prominent crest called the lateral epicondylar crest extended from the lateral epicondyle, on the posterior aspect, to end a little below the middle of the shaft (Fig 3, 5) as reported by [10] in cat, [14] in dog, [24] in mongoose and [26] in civet cat. In the present study, the distal $1/3^{\text{rd}}$ or $1/4^{\text{th}}$ of the humeral shaft was flattened cranio-caudally. Above the medial epicondyle was noticed the oval supracondyloid foramen roofed by a thick bony ledge (Fig 4c, d) as reported by [10] in cat, [12] in Indian lion, [24] in mongoose, [26] in civet cat, domestic cat and [29] in cheetah. It was absent in dog [18]. The supracondylar foramen was an ancestral structure in mammals which was lost during mammalian evolution [30]. In the African giant rats and cats, supracondyloid foramen is present and transmitted the median nerve and brachial artery [21]. Just above the lateral epicondylar crest, in mongoose on the posterior surface was present a nutrient foramen. It was on the caudal surface of the distal half in lion [12]. But, in civet cat it was on the medial surface [26]. In contrary, [24] reported the absence of nutrient foramen in

mongoose. However, numerous nutrient foramina in both the diaphysis and epiphysis was observed in the humerus of cat [23].

CONCLUSION

The acromion or hamate process of scapula in Indian grey mongoose presented a distinct suprahamate process similar to that of cat. The glenoid cavity was shallow with a supraglenoid tubercle that projected cranially in front of the cavity as in dog and cat. The ovoid humeral head was similar to canine and feline species. The teres tubercle was prominent as in cat. The presence of lateral epicondylar crest resembled both dog and cat. The presence of supracondyloid foramen was similar to that in cat. The supratrochlear foramen was seen as in dog. Thus the bones of the shoulder and arm in mongoose presented features that resembled either of these two carnivores and some that was in common for both carnivores. These findings signify that these morphological similarities in Indian grey mongoose are the adaptations of locomotion for predation by this carnivore.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES


1. Gilchrist JS, Jennings AP, Veron G, Cavallini P. "Family Herpestidae (Mongooses)". In Wilson, D. E. & Mittermeier, R. A. (eds.). *Handbook of the Mammals of the World*. Vol. 1. Carnivores. Barcelona: Lynx Edicions. 2009; 262–328.
2. Mudappa D, Choudhury A. "*Herpestes edwardsii*". *IUCN Red List of Threatened Species*. 2016:e.T41611A45206787. doi:10.2305/IUCN.UK.2016-1.RLTS.T41611A45206787.en
3. Soni A, Kumar A, Sharma A, Vohra H. Comparison of maceration techniques for retrieval of bones. *J Anat Soc India*. 2021;70:93-6
4. Savitri RC, Yakkundi, Matapathi N, Talawar V. A review of Bone Preparation Techniques for Anatomical Studies. *J Ayurveda Integr Med Sci* 2023;10:112-117. <http://dx.doi.org/10.21760/jaims.8.10.16>
5. Aggarwal N, Gupta M, Goyal PM, Jaswinder Kaur J. An alternative approach to bone cleaning methods for anatomical purposes. *Int J Anat Res*. 2016;4(2):2216-2221. DOI: 10.16965/ijar.2016.181
6. Sarma K, Suri S, Sasan JS. Gross Anatomical and morphometrical studies on scapula of civet cat (*Viverricula indica*). *Indian J Vet Anat*, 2023. 35(2): 162-164.
7. Palanisamy D, Tomar MPS, Ankem PB, Ullakula RS, Jonnalagadda N, Korampalli V. Gross morphology of scapula in Indian wild cat (*Felis silvestris ornata*). *Int J Curr Microbiol Appl Sci*, 2018;7:2473-2477.

8. Tomar MPS, Taluja JS, Vaish R, Shrivastav, AB, Shahi A, Sumbria D. Gross anatomy of scapula in tiger (*Panthera tigris*). Indian J Anim Res, 2018;52:547-550.
9. Rajput R, Pathak V, Shukla P, Gupta D. Morphological and morphometrical studies on the scapula of Himalayan mongoose. Indian J Vet Anat, 2022;34(1):43-45.
10. El-Ghazali HM, El-Beheri EI. Comparative macroanatomical observations of the appendicular skeleton of New Zealand rabbit (*Oryctolagus cuniculus*) and domestic cat (*Felis domestica*) thoracic limb. Int J Vet Sci, 2018;7(3):127-133.
11. Pathak SK, Archana M, Amarpal, Pawde AM. Morphological and morphometrical characterization of scapula bone of fishing cat, leopard cat and small Indian civet. Indian J Vet Anat, 2020;32 (1): 69-71.
12. Nzalak JO, Eki MM, Sulaiman MH, Umosen AD, Salami SO, Maidawa, S.M, Ibe CS, Gross Anatomical studies on the bones of the thoracic limbs of the lion (*Panthera leo*). J Vet Anat, 2010;3 (2): 65-71.
13. Bordoloi CC, Kalita HC, Kalita SN, Baishya G. Scapula of the Great Indian rhino (*Rhinoceros unicornis*). Indian Vet. J, 1993;70: 540-542.
14. Evans HW. Anatomy of the dog. W.B. Saunders Co., Philadelphia, USA. 1993
15. Ozkan ZE, Macro-Anatomical Investigations on the hedgehog skeleton (*Erinaceus europaeus*) I-Ossa Membri Thoracici. Turk. J. Vet. Anim. Sci, 2004;28: 271-274.
16. Pandey SP, Bhayani DM, Vyas YL. Gross anatomical study on the scapula of Asiatic lion (*Panthera persica*). Indian J Vet Anat, 2004;16:53-56.
17. Kalita PC, Bhattacharya R. Macroanatomy of the scapula of sloth bear (*Melursus ursinus*). Indian J. Vet. Anat, 2002;14; 77-79.
18. Lahunta ED, Miller's Anatomy of the dog. 4th edn. Elsevier Saunders, 2013;127-132.
19. Siddiqui MSI, Khan MZI, Sarma M, Islam MN, Jahan MR.. Macroanatomy of the bones of the limb of black Bengal goat (*Capra hircus*). Bangladesh J. Vet. Med, 2008;6(1):59-66.
20. Ahasan ASM, Quasem M, Rahman ML, Hasan RB, Kibria ASM, Shil SK. Macroanatomy of the bones of thoracic limb of an Asian elephant (*Elephas maximus*). Int J Morphol, 2016;34(3):909-917.
21. Dyce KM, Sack WO, Wensing CJG. Textbook of Veterinary Anatomy. Saunders-Elsevier Science, Philadelphia. 3rd ed. 2002; 840 pp.
22. Shuvo MJH, Shuvo RI, Emran AA, Rahman MT, Robin IH, Hasan MA, Jahid MT, Hussan M, Rahman S. Morphometric analysis of adult hippopotamus forelimb bones. Bangl. J. Vet. Med, 2023;21(1):7-15.
23. Sebastiani, AM, Fishbeck DW. Mammalian Anatomy: The Cat. 2nd edn. Morton Publishing Company, Colorado, 2005; pp:30.
24. Shunmugam R, Sundaram, M. Comparative morphology of the humerus of rabbit, guinea pig and mongoose. Ind. J. Vet. Sci and Biotech. 2022;18(4), 59-63.
25. Choudhury KB, Dev, Deka A, Rajkhowa J, Sinha S, Kachari J. Comparative morphological studies on appendicular skeleton of arm of Asian palm civet (*Paradoxurus hermaphroditus*) with domestic cat. J Entomol Zool Stud. 2020;8(4):1765-1767.
26. Sarma K, Sasan JS, Suri S. Gross and morphometrical studies on humerus of civet cat (*Viverricula indica*). Indian J Pure and Appl Biosci, 2017;5: 80-85.
27. Pathak SK, Archana M, Amarpal, Pawde AM. Morphological and certain morphometrical study of humerus bone of Indian tiger. Int J Curr Microbiol Appl Sci, 2017; 6(2), 546-551.

28. Witkowska A, Alibhai A, Hughes C, Price J, Klisch K, Sturrock CJ, Rutland CS. Computed tomography analysis of guinea pig bone: Architecture, bone thickness and dimensions throughout development. *Peer J*, 2014;2, 1-20.
29. Rosu PM, Predoi G, Belu C, Georgescu B, Dumitrescu I, Raita SM, Morphometric biodiversity in cheetah thoracic limb bones: a case study. *Sci works. Ser C. Vet Med*, vol. Lxii, issn 2065-1295, 2016; pp:41-45.
30. Polly PD. Limbs in mammalian evolution. In: Hall, B.K., ed. *Fins to Limbs : Evolution, Development and Transformation*. The University of Chicago Press. USA, 2007; pp.245-268.

FIGURES:

Fig 1: Photograph of the left Scapula of mongoose showing its features in its lateral and costal surface.



1- Scapular spine, 2- Infra spinous fossa, 3- Supra spinous fossa, 4- Acromion/hamate process, 5- Suprahamate metacromion process, 6- Candal angle, 7- Cranial angle, 8- Scapular notch, 9- Candal ridge, 10- Sub scapular fossa, 11- Cranial ridge, 12- Supra glenoid tubercle, 13- Glenoid cavity

a

b

Fig 2: Photograph of the left Humerus of mongoose showing its features in its lateral and medial sides.




Fig 3: Photograph of the left Humerus of mongoose showing its features in its anterior and posterior sides.






Fig 4: Photograph of the extremities of the scapula and Humerus of mongoose showing its features.



(a)- Distal extremity of scapula – 1- Coracoid process, 2- Supraglenoid tubercle, 3- Acromion process, 4- Suprahumeral process, 5- Glenoid cavity. (b)- Proximal extremity of humerus – 6- Head, 7- Lateral tuberosity, 8- Medial tuberosity, 9- Bicipital groove. (c)- Distal extremity of humerus (anterior) & (d) (posterior)- 10- supratrochlear foramen, 11- Supra condylar foramen, 12- Medial condyle, 13- Lateral condyle, 14- Lateral epicondyle, 15- Medial epicondyle

Fig 5: Radiographic anatomy of the bones of the shoulder and arm in mongoose.



Cr- Cranial ridge, Sfg-Supraspinous fossa, If- Infraspinous fossa, Ca- Caudal angle, Ss- Scapular spine, Sgt_ Supraglenoid tubercle, Shp- Suprahumeral process, Sj- Shoulder joint, Lt- Lateral tuberosity, Dc- Deltoid crest, Mg- Musculo-spiral groove, Lc- Lateral epicondylar crest, Op- Olecranon process, Ej- Elbow joint.