

Original Research Article

ANTI-HYPERGLYCEMIC AND ANTI-INFLAMMATORY EFFECTS OF A COCOYAM-BAMBARA GROUNDNUT-SOYA BEAN FLOUR BLEND IN STREPTOZOTOCIN-INDUCED DIABETIC RATS

ABSTRACT

Background: Diabetes mellitus has assumed a Public Health prominence as a result of the global prevalence, mortality and disability. With effective and affordable management not universally in focus, as a result of the huge financial burden associated with its management, attention has been shifted to medicinal food plants, most of which have been administered singly. The aim of this study is to determine the effect of flour blend of the commonly consumed cocoyam, Bambara groundnut and soya beans, on the serum lipid, bilirubin and inflammatory biomarkers' levels, in streptozotocin-induced diabetic rats (SIDRs), in order to build scientific evidence on the use of such food mix by diabetics. **Methodology:** Locally sourced cocoyam (CY) roots, Bambara groundnut (BGN) and Soya bean (SB) seeds were peeled in the case of cocoyam, washed, boiled and dried separately with oven-dryer, until constant weights were obtained, before grounding them and the formulations blended and pelletized. The three controls and seven intervention groups were appropriately placed on commercial rat feed, metformin and the pelletized flour blends for 28 days, after which, the blood specimen were collected for serum lipids, bilirubin and inflammatory biomarkers determination. **Results & Discussion:** The generated data were statistically analysed using SPSS, and one-way ANOVA, while the mean difference were determined by Turkey's post-hoc test. Formulation 1 (16.6%CY: 16.6%SB: 16.6%BGN: 50%RF)-treated rats had the least mean values of NFkB and IL-10, total and direct bilirubin levels, total cholesterol and LDLc, which were comparable to those of the normal control. **Formulation 1 reduced the blood glucose level by 13.02%, just like the anti-diabetic drug, which reduced same by 19.99%, when compared to the negative control. Formulation 1 and the antidiabetic drug, equally significantly reduced Total cholesterol (TC) and Low density lipoprotein cholesterol (LDLc) levels by 12.53% and 24.71% respectively when compared to the diabetic control.** **Conclusion:** Various formulations of the Cocoyam-bambara groundnut-soya bean flour blend, exerted varying effects on the inflammatory biomarkers, serum bilirubin and lipid profile in SIDRs, with formulation one showing more potent hypocholesterolamic and anti-inflammatory activities than the antidiabetic drug. **These findings underscores the possibility of using such plant food mix in the management and prevention of T2DM, and hence opens the gate for clinical trials in human subjects.**

Key words: Medicinal plant foods, Serum lipids, diabetic rats

1. INTRODUCTION

Diabetes mellitus affects various age-groups globally and at a rate considered of Public Health importance due to the prevalence, disability and mortality [1]. As a pancreatic organ disorder, it makes insulin production inadequate and ineffective [2]. Its prevalence in the Asian and African continents far surpasses that in the Western nations and Europe [3]. Arising from dilapidated health infrastructure, shortage of human resource for health, and limited funding, the disease outcome is poor, thereby affecting productivity and worsening disability [4, 5, 6]. In finding solutions to these menace a critical alternative to the expensive and ineffective orthodox management strategies has led to the choice of dietary and other lifestyle modifications as treatment and prophylactic adjuncts [7], with the diabetic diets used in the Western Nations proving effective in T2DM management [8, 9] and its use and the utilization of nutraceuticals and functional foods recommended by World Health Organization [10]. The presence of bioactive compounds in these functional foods and their biochemical effects *in vivo*, have been the basis for their usage in managing non-communicable diseases [11, 12]. Several bioactive compounds have been identified in these plant foods, including the isoflavones, stilbene, and flavonoids, with reported actions on the pancreatic cells [13]. For instance, the hypoglycemic bioactivity of the chemical constituents in cocoyam [14-16], antioxidant and antidiabetic actions of Bambara groundnut (*vigna subterranean*), attributed to the chemical constituents [17-31], and the use of Soya Bean (*Glycine max. (L) Merrill*), in non-communicable diseases management, linked to the same bioactive compounds and their biochemical actions [32-46]. Rarely are these medicinal food plants consumed singly without mixing two or more of them as food mixtures. Consequently, understudying the effects of such flour blends on biochemical parameters such as the serum lipids and inflammatory biomarkers is the focus of this study.

2.0 Materials and Methods

2.1 Collection and Preparation of the Plant Food Material

The Bambara groundnut, soya bean and cocoyam were purchased from the local market and prepared by washing, peeling and soaking (cocoyam), washing and rinsing (Bambara groundnut and soya bean). All the three plant foods were then boiled, oven-dried at a temperature of 60⁰C until a constant weight was achieved, before being ground and formulated as pellets. The pelletized blends were oven-dried and stored for future use.

2.2 Experimental Animals

A total of eighty-five (85) male albino rats (134 – 249 grams), were purchased locally and segregated in groups of eight per cage, with each rat marked on the tail, in circles of 1-8 as identification marks. Ten groups, A- J were mapped out, and fed with commercial rat feed and water administered *ad libitum* for seven days to allow for acclimatization. The body weights of the rats were recorded and the protocols for handling experimental animals followed [47].

2.3 Induction of Insulin Resistance Using Low Fructose Diet

Once the acclimatization period was completed, 30 g of fructose was dissolved in 300 mL of water and administered *ad libitum* as low dose fructose diet for six days. The commercial rat feeds were maintained throughout this period.

2.4 Induction of Type 2 Diabetes Mellitus using STZ

The rats in the ten groups were fasted overnight at the 6th day of the fructose diet and the fasting blood glucose recorded. All the normo-glycemic rats were administered with streptozotocin (STZ) prepared as 1 g in 50 mL of freshly prepared sodium citrate buffer and given by intra-peritoneal route using the formula for extracts to experimental animal subjects to estimate the dose [48, 49]. Post STZ treatment, blood samples were collected from the rats and estimated for blood glucose levels at days 6 and 12. The rats that were hyperglycemic were randomly assigned to standard control, negative control and intervention groups. The rats that were not administered with STZ and normoglycemic, acted as the normal control. The various groups were assigned to interventions as shown in Table 1, according to Nnadi and colleagues [50].

Table 1: The Formulations and Their Compositions given to Each Rat Group

Rat Groups	Formulations	Formulation Compositions
A	1	16.6%CY: 16.6%SB: 16.6%BGN: 50%RF
B	Standard Control	Metformin + 100% Rat Feed (RF)
C	Normal Control	100% RF
D	Negative Control	100% RF
E	2	12.5%CY: 12.5%SB: 25%BGN: 50%RF
F	4	12.5%CY: 25%SB: 12.5%BGN: 50%RF
G	7	0%CY: 50%SB: 0%BGN: 50%RF
H	5	0%CY: 0%SB: 50%BGN: 50%RF
I	6	50%CY: 0%SB: 0%BGN: 50%RF
J	3	25%CY: 12.5%SB: 12.5%BGN: 50%RF

RF =Rat Feed

CY = Cocoyam

SB = Soya Bean

BGN = Bambara Groundnut

2.5 Estimation of Serum Lipids and Inflammatory Biomarkers

The interventions were conducted for 28 days, following which, blood samples were collected for lipid using the Randox assay diagnostic kits [51], with the LDLc and VLDLc concentration obtained using the formula according to Friededwald and colleagues [52]: $LDL-C = Total\ cholesterol - (HDL-C + [TG/5])$.

For the serum inflammatory biomarkers determination, the blood specimen, was collected and allowed to clot, then centrifuged at 2000 RPM for 10 min at 4⁰C and the serum obtained, used to determine the interleukin-6 and IL-8 with the aid of human Enzyme-linked Immunosorbent Assay (ELISA) kits (Elabscience, Quangzhou, China) [53]. The C-reactive proteins, NFk β , and IL-10 were measured using a highly sensitive commercial method in Hitachi 917 autoanalyzer [54].

2.6 Statistical Analysis

The Statistical Package for Social Sciences (SPSS Inc., Chicago, IL) version 20.0, one-way analysis of variance (ANOVA) and the Turkey's post-hoc test were

employed for data analysis, standard error of mean and the mean differences respectively. The statistically significant was pegged at $p < 0.05$.

3.1 RESULTS

3.1.1 Inflammatory Markers Results




Figure 1: Bar Charts showing NFκB, Interleukin (IL)-10 and C - reactive protein (CRP) for the various diabetic and non-diabetic rat groups showing the lowest values for Group A in the pro-inflammatory and anti-inflammatory biomarkers.

Figure 1 showed that Group A were least while Group B and D were the highest for both markers NFκB and IL-10. For CRP, Groups A, C and F had least mean CRP while Group B had highest mean CRP.

3.1.2 Serum bilirubin and Protein Results




Figure 2: Bar Charts showing Serum bilirubin and Serum Protein for the various diabetic and non-diabetic rat groups showing that the diabetic control, had the least values of the bilirubin and serum total protein.

Figure 2 revealed that T-BILL and D-BILL has no significant ($p>0.05$) mean difference among the groups, however mean values for TP, ALB and GLB were significantly ($p<0.05$) different. Mean TP and ALB were minimum in group D while mean GLB were least in group J.

3.1.3 Results of Serum Lipid Profile




Figure 3: Bar Charts showing Serum Lipid Profile for the various diabetic and non-diabetic rat groups showing the comparable values of Groups A and C.

Figure 3 is the result of mean analysis of serum lipid profile among various groups. Significant ($p < 0.05$) difference among the groups were observed for the TCHOL, HDL-c and LDL-c only.

3.2 DISCUSSION

3.2.1 Effects on Inflammatory Biomarkers

In acute and chronic inflammation, cytokines which are soluble factors are engaged in recruiting leucocytes [55] and exhibit synergistic and antagonist effects [56]. NF- κ B is activated in prolonged hyperglycemia, with resultant expression of genes for TNF- α , IL-6, IL-8, MCP-1, MIP-1 α , MIP-2, resistin, ICAM-1 and VCAM-1 [57]. These expressed cytokines, regulate growth factor, apoptotic genes, cell-cycle programme and inflammation [58]. Circulating cytokines can provide the general state of inflammation in the individual. Inflammatory markers –CRP, IL-6 and TNF- α are raised in patients with metabolic syndrome and sub-clinically overt T2DM [59]. Transforming growth factor β 1 [60], MCP-1 [61] or Lipoprotein Associated Phospholipase A2 (LP-PLA2) [62] also increase in T2DM.

When the Acute phase proteins are chronically raised, it promotes IR in skeletal muscles and endothelial dysfunction and release of CRP from the liver [63]. There is significant correlation between CRP level and features of metabolic syndrome such as hyper-insulinaemia and IR [64].

In this study, the mean values of NF κ B and IL-10 among the groups, showed significant differences, with Group A having the least and the standard and negative controls having the highest values, and these higher values were significantly different that of the normal control and the groups on intervention formulation. The normal control and intervention groups had no significant difference. Similar results have been gotten when whole grains, and foods containing high amounts of antioxidants and phytochemicals, are consumed with resultant improved glycemic index [65, 66].

In terms of the c-reactive proteins, Groups A,C and F had the least values, meaning that the formulations 1 and 4 respectively given to these groups, had as good anti-

inflammatory protection as the normoglycemic group which was better than the effect of the standard control. Other intervention formulations used in groups E, G, H, I and J were not as protective as those used in Groups A and F as they had higher CRP values.

Group A rats were given formulation 1 (Table 1) which contained mainly phenolic compounds and stilbene [67]. Among the several mechanisms of action of phenolic are inhibition of transcription factors [68], pro-inflammatory cytokines release [68, 69] and enzymes associated with inflammation such as cyclo-oxygenase [70, 71], lipoxygenases, inducible nitric oxide synthase (iNOS) [72]. To exert anti-inflammatory activities, polyphenols, block MAPK-mediated pathway, an action that has been linked to their structure [70, 73].

Stilbene in the same vein, inhibits nucleotide-binding and oligomerization domain (NOD)-Like receptors (NLRP)-3 [74], which is activated by myeloid cells in obesity [75] and by Danger Associated Molecular Patterns (DAMPs) in chronic inflammation and insulin resistance [76], with subsequent release of IL-1 β and IL-18 [77]. Studies have shown that absence of NLRP3 gene, protects against obesity-induced inflammation is prevented [78]. The actions of Stilbenes have been linked to silent information regulator 1 (SIRT1) activation [79] or by thioredoxin-interacting protein (TXNIP) inhibition [80, 81]. Activation of SIRT1, promotes the deacetylation of peroxisome proliferator activator gamma co-activator 1 alpha (PGC-1 α), critical in regulating mitochondrial biogenesis [82]. Thirdly, stilbene modulates pro-inflammatory cytokines, which occurs in the presence of infectious agents, with resultant IL-1 β release [83]. Hence by arresting its activation stilbene inhibits the development of T2DM [84]. Fourthly, stilbene inhibits the overexpression of IL-6 through downregulation of NF- κ B and activator protein-1 (AP-1) in different cells [85] and by so acting, there is non-induction of the intercellular adhesion molecule (ICAM-1) expression, non-phosphorylation of signal transducer and activator of transcription 3 (STAT3) [86] and downregulation of suppressor of cytokine signaling (SOCS3) expression in Hep G2 cells in rat liver cells resulting in improved insulin signaling [87]. Additionally, Stilbene, downregulates the p38 MAPK pathway

thereby modulating NF κ B that causes TNF- α and IL-1 β release and upregulates heme-oxygenase-1 (HO-1),, with the later resulting in the upregulation of enzymatic antioxidants such as catalase (CAT), superoxide dismutase (SOD), and glutathione peroxidase (GPX) [88].

Also stilbene reduces the endoplasmic reticulum (ER) stress, through the activation of transcription factor 4 (ATF-4) and Tribbles Pseudokinase 3 (TRIB3) [89, 90], downregulation of protein kinase-like ER kinase (PERK) and eukaryotic initiation factor 2 alpha (eIF2 alpha) [91] and hence improve insulin sensitivity [91]. Stilbene also acts on innate and adaptive immunity. Stilbene inhibits the proliferation of CD4⁺ and CD8⁺ T cells. When this happens, cytokines are not released [92] thereby affecting the type 1 (IL2, IFN-g, and lymphotoxin) cell mediated inflammatory response and type 2 (IL4, IL5, IL10, and IL13) antibody-mediated immune response T cells.

3.2.2 Effect on Bilirubin

The bilirubin can be indirect bilirubin or direct bilirubin, and each of these subtypes having clinical relevance. When moderate hyperbilirubinemia involve all subtypes, physiological benefit and disease prevention and management occurs [93] because their antioxidant and anti-inflammation properties manifest [94].

In this study, the bilirubin mean values were not significantly different among the groups though the total bilirubin and direct bilirubin were least in the normal control, and in Groups A, E and F rats. The diabetic and standard control groups were among the groups with highest values of total and direct bilirubin, in line with previous study that reported raised level of direct bilirubin in T2DM [95]. Presence of elevated levels of direct bilirubin is more related to metabolic syndrome and stroke [96] than total bilirubin and indirect bilirubin. However in chronic hyperglycemia,,more free radicals are and more bilirubin consumed, resulting in decreased bilirubin [95, 97]. In view of the short duration of this study and the non-chronicity of the raised glucose levels, low concentration of the bilirubin as reported in the intervention groups could be attributed to the presence of bioactive compounds

in the formulations which might have not only regulated the blood glucose but also reduced the free radicals.

3.2.3 Effect on Serum Lipid Profile

Groups A, E and I had the least mean values of the Total Cholesterol (TC) in this study and this was comparable with values in the normal control. For the triglyceride, there was no significant difference among the groups, while in the case of HDLc, Groups A, B, C and D showed significant difference when compared to those in Groups F, G, H, I and J. The LDLc, values in Groups A, C and E were the least and were significantly different from the diabetic control and other groups. For the VLDLc, there was no significant difference among the group means. Overall, in this study, there were elevated plasma values of the TC and LDLc in the diabetic control when compared to the normal control and this was in agreement with earlier reports by other researchers [98]. The remarkably low values of TC and LDL in Group A relative to the diabetic control, was equally in tandem with the work of Wang and colleagues [99].

T2DM is often associated with dyslipidemia, in which there is elevated levels of triglycerides and low levels of high-density lipoproteins (HDL) and elevated low-density lipoproteins (LDL) [100 - 102]. The more plasma triglycerides increases, the more, HDL is broken down, resulting in reduced level of HDL which in turn leads to raised LDL [103]. A hypertriglyceridemia state with low levels of HDL aids elevated fatty acids, which stimulate insulin resistance and cause β -cell dysfunction [104, 105]. Such alterations in lipid profile, impairs the ability of insulin to inhibit hormone-sensitive lipase and hence cause an increase in serum concentrations of free fatty acids [105, 106], which further induce insulin resistance in adipose tissue and skeletal muscle [107]. The resultant DNA damage activates DNA repair enzyme which inhibits metabolic enzymes such as glyceraldehyde-3-phosphate dehydrogenase which ultimately leads to *de novo* synthesis of diacylglycerides, up-regulation of PKC pathway, stimulation of free radical generating-enzymes like lipooxygens and the development of deteriorating toxic environment [105, 108].

Studies have reported significant correlation between serum triglyceride and blood sugar levels [109] and the association in diabetics between high levels of LDL cholesterol and higher risk of diabetic complications such as cardiovascular diseases [98]. When compounds with potent antioxidant properties are given, there is improved lipid profile. Wang and colleagues reported that taurine-treated diabetic rats showed a significant improvement in lipid profile through reduction in triglyceride, cholesterol and LDL and elevated HDL [99]. Other studies linking isoflavones and other bioactive compounds to regulation of blood glucose level and improving lipid profile have been documented as found in this study [110 -112]

4. CONCLUSION

Various formulations of the cocoyam-bambara groundnut-soya bean flour blend, exerted varying effects on the inflammatory biomarkers, serum bilirubin and lipid profile in **STZ**-induced diabetic rats. Formulation 1 composed of 16.6%cy: 16.6%sb: 16.6%bgn: 50%rf, and administered to rats in group a, had the least mean values of **NFKB** and **IL-10**, total and direct bilirubin levels, total cholesterol and **LDLC**. These values were comparable to values obtained in the normal control. **Formulation 1 reduced the blood glucose level by 13.02%, just like the anti-diabetic drug, which reduced same by 19.99%, when compared to the negative control. Formulation 1 and the antidiabetic drug, equally significantly reduced Total cholesterol (TC) and Low density lipoprotein cholesterol (LDLc) levels by 12.53% and 24.71% respectively when compared to the diabetic control.**The implication is that formulation 1 had better hypocholesterolamic and anti-inflammatory activities than metformin antidiabetic drug. However, this blend formulation had no effect on the triglyceride and VLDLc levels. **These findings underscores the possibility of using such plant food mix in the management and prevention of T2DM, and hence opens the gate for clinical trials in human subjects.**

ETHICAL APPROVAL

Ethical approval was sought and obtained from the Research Ethical Committee of the Faculty of Basic Medical Sciences, Rivers State University, Port Harcourt, Nigeria.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

- [1] Sun, H., Saeedi, P., Karuranga, S., Pinkepank, M., Ogurtsova, K., Duncan, B.B., Stein, C., Basit, A., Chan, J.C.N., Mbanya, J.C., Pavkov, M.E., Ramachandaran, A., Wild, S.H., James, S., Herman, W.H., Zhang, P., Bommer, C., Kuo, S., Boyko, E.J., Maqliano, D. J., (2022). IDF Diabetes Atlas: Global, Regional and Country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Research and Clinical Practice*, 183, 109119. <https://doi.org/10.1016/j.diabres.2021.109119>
- [2] Van Belle, T.L., Coppieters, K.T. and Von Herrath, M.G. (2011).Type 1 diabetes: etiology, immunology and therapeutic strategies. *Physiological Reviews*, 91:1.
- [3] Lin, X., Xu, Y., Pan, X., *et al.*, (2020) Global, regional, and national burden and trend of diabetes in 195 countries and territories: an analysis from 1990 to 2025. *Sci Rep*, 10, 14790
- [4] Kirigia JM, Sambo HB, Sambo LG and Barry SP. (2009).Economic burden of diabetes mellitus in the WHO African region. *BMC International Health and Human Rights*, 9:6.
- [5] Ashcroft, F.M. and S.J.H. Ashcroft, (1992). *Insulin, Molecular Biology to Pathology*. Oxford University Press. pp 266-284.
- [6] Sobngwi E, Mauvais-Jarvis F, Vexiau P, Mbanya JC and Gautier JF. (2001).Diabetes in Africans. Epidemiology and clinical specificities. *Diabetes Metab.*, 27(6):628-634.
- [7] Piero MN. (2006) Hypoglycemic effects of some Kenyan plants traditionally used in management of diabetes mellitus in eastern province, Msc thesis, Kenyatta University
- [8] Wylie-Rosett, J and Delahanty, L.M (2017) *Nutrition in the Prevention and Treatment of Disease (Fourth Edition)* Pp 691-707. <https://doi.org/10.1016/B978-0-12-802928-2.00031-X>
- [9] Forouhi, G.N., Misra, A., Mohan, V., Taylor, R., Yancy, W (2018) Dietary and nutritional approaches for prevention and management of type 2 Diabetes, *Science and Politics of Nutrition*, BMJ 361 Doi:<https://doi.org/10.1136/bmj.k2234>
- [10] Piero MN, Njagi JM, Kibiti CM, Ngeranwa JJN, Njagi ENM and Miriti PM. (2012).The Role of Vitamins and Mineral Elements in Management of Type 2 Diabetes Mellitus: A Review *South As. J. Biol.Sci.*, 2(Supp.1):107 –115.
- [11] Merida L.A., Mattos E.B., Correa A.C., Pereira P.R., Paschoalin V.M., Pinho M.F., Vericimo M.A (2018). Tarin stimulates granulocyte growth in bone marrow cell cultures and minimizes immunosuppression by cyclophosphamide in mice. *PLoS ONE*. 13:e0206240. doi: 10.1371/journal.pone.0206240
- [12] Chukwuma C.I., Islam M.S., Amonsou E.O. (2018) A comparative study on the physicochemical, anti-oxidative, anti-hyperglycemic and anti-lipidemic properties of amadumbe (*Colocasia esculenta*) and okra (*Abelmoschus esculentus*) mucilage. *J. Food Biochem*. 42:e12601. doi: 10.1111/jfbc.12601

- [13] Prabhakar, P.K., and Doble, M (2008) A Target Based Therapeutic Approach Towards Diabetes Mellitus Using Medicinal Plants. *Current Diabetes Reviews*, 291-308. Bentham Science Publishers Ltd.
- [14] Kumawat, N.S., Chaudhari, S.P., Wani, N.S., Deshmukh, T.A., & Patil, V.R. (2010). Antidiabetic activity of ethanol extract of *Colocasia esculenta* leaves in alloxan induced diabetic rats. *International Journal of Pharmaceutical Technique Research*, 2, 1246–1249.
- [15] Eleazu CO, Iroaganachi M & Eleazu KC (2013). Ameliorative potentials of cocoyam (*Colocasia esculenta* L.) and unripe plantain (*Musa paradisiaca* L.) on the relative tissue weights of streptozotocin-induced diabetic rats. *J Diabetes Res.* 1–8 <https://doi.org/10.1155/2013/160964>
- [16] Uro-Chukwu, H.C., Ezekwe, A.S., Roberts, F., Okari, K.A & Uro-Chukwu, F.N.C (2024) Anti-hyperglycemic Effects and Histological Changes in Streptozotocin-induced diabetic rats fed with high concentration of *Colocasia esculenta* flour. *South Asian Research Journal of Natural Products*, 7(2), 123–134.
- [17] Arise, A. K., Amonsou, E. O. & Ijabadeniyi, O. A. (2015) Influence of extraction methods on functional properties of protein concentrates prepared from South African Bambara groundnut landraces. *International Journal of Food Science and Technology*. 50, 1095–1101. <https://doi.org/10.1111/ijfs.12746>.
- [18] Oyeyinka, S.A. & Oyeyinka, A.T. (2018), A review on isolation, composition, physicochemical properties and modification of Bambara groundnut starch. *Food Hydrocolloid.*, 75, 62–71.
- [19] Oyeyinka, S.A. Tijani, S.T., Oyeyinka, A.T., Arise, A.K., Balogun, M.A., Kolawole, F.L., Obalowu, M.A. & Joseph, J.K (2018). Value added snacks produced from Bambara groundnut (*Vigna subterranea*) paste or flour. *LWT-Food Science & Technology*, 88, 126-131.
- [20] Halimi, R.A., Barkla, B.J., Mayes, S. & King, G.J. (2019). The potential of the underutilized pulse bambara groundnut (*Vigna subterranea* (L.) Verdc.) for nutritional food security. *The Journal of Food Composition and Analysis*, 77(2), 47–59.
- [21] Olaleye, A.A., Adeyeye E.I., and Adesina, A.J (2013) Chemical Composition of Bambara groundnut (*Vigna subterranea* L. Verdc) *Bangladesh Journal of Scientific & Industrial Research*, 48(3): 167-178.
- [22] Salawu, (2016). Comparative study of the antioxidant activities of methanolic extract and simulated gastrointestinal enzyme digest of Bambara nut (*Vigna subterranea*) FUTA J. Res. Sci., 1: 107-120.
- [23] Harris, T., Jideani, V., and Le Rose-Hill, M (2018) Flavonoids and tannin composition of Bambara groundnut (*Vigna subterranea*) of Mpumalanga, South Africa. *Heliyon*, 4: e00833. Doi: 10.1016/j.heliyon.2018.e00833
- [24] Baptista, A., Pinho, O., Pinto, E., Casal, S., Mota, C. & Ferreira, I.M. (2016). Characterization of protein and fat composition of seeds from common beans (*Phaseolus vulgaris* L.), cowpea (*Vigna unguiculata* L. Walp) and Bambara

- groundnuts (*Vigna subterranean* L. Verde) from Mozambique. *Journal of Food Measurement and Characterization*, 12, 1-9
- [25] Uro-Chukwu, H.C., Ezekwe, A.S., Okorie, U., Eleazu, C.O., Uro-Chukwu, F.C. U., Ajah, D.O.J., Nwokpoku, B.I, Okari, K.A &Ekwe N.P (2024) Hypoglycemic and Antioxidant Effects of High Concentration of Bambara Groundnut (*Vigna Subterranean*) Flour Fed on Streptozotocin-induced Diabetic Rats. *International Journal of Basic and Clinical Toxicology*, 3(5):10-20.
- [26] Yao, D.N., Kouassi, K.N., Erba, D., Scazzina, F., Pellegrini, N. & Casiraghi, M.C. (2015). Nutritive Evaluation of the Bambara Groundnut Ci12 Landrace [*Vigna subterranea* (L.) Verdc. (*Fabaceae*)] Produced in Côte d'Ivoire. *International Journal of Molecular Sciences*, 16(9), 21428-21441.
- [27] Barbieri, R., Coppo, E., Marchese, A., Daglia, M., Sobarz-Sánchez, E., Nabavi, S.F.& Nabavi, S.M. (2017). Phytochemicals for human disease: An update on plant-derived compounds antibacterial activity. *Microbiology Research*, 196, 44-68.
- [28] Saxena, M., Saxena, J., Nema, R., Singh, D. & Gupta, A. (2013). Phytochemistry of medicinal plants. *Journal of Pharmacognosy & Phytochemistry*, 1(6), 168–182.
- [29] Uro-Chukwu, H.C., Okoli, E.C., Okpala, L.C, Uro-Chukwu, F.N.C.(2024) Antioxidant and Cytotoxic Activities of Protein Hydrolysates from Shrimp Shell Wastes, germinated soybean and pigeon pea blends: A Mixture Response Surface Methodology Approach. *Journal of Drug Delivery & Therapeutics*, 14 (6):7-14
- [30] Iwai, K., Kim, M., Onodera, A. & Matsue, H. (2006). α -Glucosidase Inhibitory and Antihyperglycemic Effects of Polyphenols in the Fruit of *Viburnum dilatatum* Thunb. *Journal of Agricultural and Food Chemistry*, 54(13), 4588-4592
- [31] Ruzaidi, A., Abbe, M., Amin, L., Nawalyah, A.G., Muhajir, H., Pauliena, M.B.S.M.J., Muskinah, M.S (2008) Hypoglycemic properties of Malaysia cocoa (*Theobromacacao*) polyphenols-rich extract. *Int Food Res. J.*, 15:305-312
- [32] Esteves, E. A., Martino, H. S. D., Oliveira, F. C. E., Bressan, J., and Costa, N. M. B. (2010). Chemical composition of a soybean cultivar lacking lipoxygenases (LOX2 and LOX3). *Food Chemistry*, 122 (1): 238-242
- [33] Lebiezinska, A., &Szefer, P. (2006). Vitamins B in grain and cereal-grain food, soyproducts and seeds. *Food Chemistry*, 95 (1): 116-122
- [34] Martino, H. S. D., Martin, B. R., Weaver, C. M., Bressan, J., Esteves, E. A., and Costa, N. M. B. (2007). Zinc and iron bioavailability of genetically modified soybeans in rats. *Journal of Food Science*, 72 (9): 689-695
- [35] Esteves, E. A., Bressan, J., Costa, N. M. B., Martino, H. S. D., Donkin, S. S., & Story, J. A. (2011). Modified soybean affects cholesterol metabolism in rats similarly to a commercial cultivar. *Journal of Medicinal Food*, Vol.
- [36] Uro-Chukwu, H.C.,Ezekwe, A.S., Roberts, F., Okari, K.A., &Uro-Chukwu, F.N.C. (2024) Comparative Analysis of the Anti-diabetic Effects of Cocoyam,

- Soya Bean & Bambara groundnut flour Fed differently to Streptozotocin (STZ)-induced Diabetic Rats. *Asian Journal of Research & Reports in Endocrinology*, 7(1), 60–68
- [37] Duranti, M., Lovati, M. R., Dani, V., Barbiroli, A., Scarafoni, A., Castiglioni, S., Ponzzone, C., and Morazzoni, P. (2004). The alpha' Subunit from Soybean 7S Globulin Lowers Plasma Lipids and Upregulates Liver Beta-VLDL Receptors in Rats Fed a Hypercholesterolemic Diet. *The Journal of Nutrition*, 134(6):1334-1339
- [38] Lee, S. H., Parka, H. J., Chuna, H. K., Choa, S. Y., Junga, H. J., Choa, S. M., Kimb, D. Y., Kangb, M. S., and Lillehojc, H. S. (2007). Dietary phytic acid improves serum and hepatic lipid levels in aged ICR mice fed a high-cholesterol diet. *Nutrition Research*, 27 (8): 505-510
- [39] Kim, S. M., Rico, C. W., Lee, S. C., and Kang, M. Y. (2010). Modulatory Effect of Rice Bran and Phytic Acid on Glucose Metabolism in High Fat-Fed C57BL/6N Mice. *Journal of Clinical Biochemistry and Nutrition*, 47 (1): 12-17
- [40] Teixeira Damasceno, N., Apolinário, E., Dias Flauzino, F., Fernandes, I., and Abdalla, D. (2007). Soy isoflavones reduce electronegative low-density lipoprotein (LDL-) and anti-LDL- autoantibodies in experimental atherosclerosis. *European Journal of Nutrition*, 46 (3):125-132.
- [41] Rios, D. R., Rodrigues, E. T., Cardoso, A. P., Montes, M. B., Franceschini, S. A., and Tolo, M. R. (2008). Lack of effects of isoflavones on the lipid profile of Brazilian postmenopausal women. *Nutrition*, 24 (11-12):1153-1158
- [42] Uro-Chukwu, H.C., Ezekwe, A.S., Roberts, F., Okari, K.A., & Uro-Chukwu, F.C.U (2024) Biochemical Effects of High Concentration of Colocasia Esculenta Flour Fed on Streptozotocin-Induced Diabetic Rats. *International Research Journal of Gastroenterology & Hepatology* 7(1): 44-60
- [43] Saito, K., Jin, D. H., Ogawa, T., Muramoto, K., Hatakeyama, E., Yasuhara, T., and Nokihara, K. (2003). Antioxidative Properties of Tripeptide Libraries Prepared by the Combinatorial Chemistry. *Journal of Agricultural and Food Chemistry*, 51 (12): 3668-3674
- [44] Takahashi, T., Nakamura, A., Kato, M., Maeda, H., Mandella, R. C., Broadmeadow, A., and Ruckman, S. A. (2003). Soluble soybean fiber: a 3-month dietary toxicity study in rats. *Food Chem Toxicol.*, 41 (8):1111-1121
- [45] Takenaka, Y., Doyama, N., Maruyama, N., Utsumi, S., and Yoshikawa, M. (2004). Introduction of DPR, an enterostatin fragment peptide, into soybean beta-conglycinin alpha' subunit by site-directed mutagenesis. *Bioscience, Biotechnology, and Biochemistry*, 68 (1):253-256
- [46] Ascencio, C., Torres, N., Isoard-Acosta, F., Gímez-Pérez, F. J., Hernández-Pando, R., & Tovar, A. R. (2004). Soy Protein Affects Serum Insulin and Hepatic SREBP-1 mRNA and Reduces Fatty Liver in Rats. *The Journal of Nutrition*, 134 (3):522-529
- [47] NRC (National Research Council) (1985) Guide for the care and use of laboratory Animals. Bethesda (MD): *National Institute of Health*, 8523

- [48] Nair AB, Jacob S. (2016) A simple practice guide for dose conversion between animals and human. *J Basic Clinical Pharmacology*, 7(2):27-31. doi: 10.4103/0976-0105.177703.
- [49] Pneu-Dart (2023) Dosage Calculation. <https://www.pneudart.com>. Accessed 14th April 2024
- [50] Nnadi, N.N., Ezekwesili, C.N., Ezeigwe, O.C (2022) Effects of Formulated Unripe Plantain and Millet Dietary Feeds in Alloxan-Induced Diabetic Albino Rats. *International Journal of Innovative Research and Advanced Studies (IJIRAS)*, 9 (6)
- [51] Tietz, N.W.(1995). *Clinical Guide to Laboratory Tests*. 3rd Ed. Philadelphia, PA: WB Saunders Company.
- [52] Friedewald WT, Levy RI & Fredrickson DS (1972). Estimation of the concentration of low density lipoprotein cholesterol in plasma without use of preparative ultracentrifuge. *Clin Chem.* **18**, 499-505
- [53] Mussa BM, Srivastava A, Al-Habshi A, Mohammed AK, Halwani R, Abusnana S (2021). Inflammatory Biomarkers Levels in T2DM Emirati Patients with Diabetic Neuropathy. *Diabetes, Metabolic Syndrome and Obesity*, 14:3389-3397. doi: 10.2147/DMSO.S319863.
- [54] Vinagre, I., Sánchez-Quesada, J.L., Sánchez-Hernández, J., Santos, D., Ordoñez-Llanos, J., De Leiva, A., Pérez, A.(2014). Inflammatory biomarkers in type 2 diabetic patients: effect of glycemic control and impact of LDL subfraction phenotype. *Cardiovascular Diabetology*, 13:34. doi: 10.1186/1475-2840-13-34.
- [55] Mak TW , Saunders ME (2006) The immune response. Basic and clinical principles . In: Press EA, editor, San Diego, CA Elsevier Academic Press, 464–516
- [56] Feghali CA & Wright TM (1997). Cytokines in acute and chronic inflammation *Front Biosci.*, 12 – 26
- [57] Shoelson, S. E., Lee, J., and Goldfine, A. B. (2006). Inflammation and insulin resistance. *J. Clin. Invest.* 116 (7), 1793–1801. doi:10.1172/jci2906.
- [58] Sultuybek, G.K., Soydas, T., Yenmis, G.(2019). NF-κB as the mediator of metformin's effect on ageing and ageing-related diseases. *Clinical and experimental pharmacology & physiology*, 46: 413–422.
- [59] Mirza S, Hossain M, Mathews C, Martinez P, Pino P, Gay JL, Rentfro A, McCormick JB, Fisher-Hoch SP. (2012) Type 2-diabetes is associated with elevated levels of TNF-alpha, IL-6 and adiponectin and low levels of leptin in a population of Mexican Americans: a cross-sectional study. *Cytokine*. 57(1):136-

42. doi: 10.1016/j.cyto.2011.09.029. Epub 2011 Oct 28. PMID: 22035595; PMCID: PMC3270578

- [60] Azar, S. T, Salti, I., Zantout, M.S., & Major S., (2000) Alterations in plasma transforming growth factor beta in normoalbuminuric type 1 and type 2 diabetic patients. *Journal of Clinical Endocrinology Metabolism*. 85(12), 4680–4682. doi,10.1210/jcem.85.12.7073
- [61] Sell H, Eckel J (2007) Monocyte chemotactic protein-1 and its role in insulin resistance. *Curr OpinLipidol*, 18(3):258–262.
- [62] Serban G, Stanasel O, Serban E, Bota S.(2018) 2-Amino-1,3,4-thiadiazole as a potential scaffold for promising antimicrobial agents. *Drug Des Devel Ther*. May 31;12:1545-1566. doi: 10.2147/DDDT.S155958.
- [63] Calle, M. C.,& Fernandez, M. L (2012) Inflammation and type 2 diabetes. *Diabetes Metabolism.*, 38(3),183–191.
- [64] Nakano S, Kuboki K, Matsumoto T, Nishimura C, Yoshino G: (2010) Small, dense LDL and high-sensitivity C-reactive protein (hs-CRP) in metabolic syndrome with type 2 diabetes mellitus. *Journal of Atherosclerosis and Thrombosis*, 17(4):410–415
- [65] Golan R, Tirosh A, Schwarzfuchs D, Harman-Boehm I, Thierry J, Fiedler GM, Blüher M, Stumvoll M & Shai I (2012). Dietary intervention induces flow of changes within biomarkers of lipids, inflammation, liver enzymes, and glycemic control. *Nutrition* **28**, 131–137.
- [66] Sanchez-Rosales, A.I., Guadarrama-Lopez, A.L., Gaona-Valle, L.S., Martinez-Carrillo, B.E., Valdes-Ramos, R(2022) The Effect of Dietary Patterns on Inflammatory Biomarkers in Adults with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutrients*, 14(21)4577 <https://doi.org/10.3390/nu14214577>.
- [67] Uro-Chukwu, H.C (2024) Anti-hyperglycemic Effects of Cocoyam (*Colocasia esculenta*)-Bambara Groundnut (*Vigna subterranean*)-Soya bean (*Glycine max. (l.) Merril*) (CY-SB-BGN) Flour Blend Fed on Steptozotocin-induced Diabetic Rats, a Ph.D Dissertation submitted to the Department of Medical Biochemistry, Rivers State University, Nkporlu, Port Harcourt, Nigeria.
- [68] Karlsen A, Retterstol L, Laake P, Paur I, Kjolsrud-Bohn S, Sandvik L, Blomhoff R, Anthocyanins inhibit nuclear factor-kappaB activation in monocytes and reduce plasma concentrations of pro-inflammatory mediators in healthy adults. *Journal of Nutrition*, 2007; 137: 1951–1954

- [69] Gauliard B, Grieve D, Wilson R, Crozier A, Jenkins C, Mullen WD & Lean M (2008). The effects of dietary phenolic compounds on cytokine and antioxidant production by A549 cells. *J. Med. Food*. **11**, 382–384. doi: 10.1089/jmf.2007.593.
- [70] Hou, D.X., Masuzaki, S., Hashimoto, F., Uto, T., Tanigawa, S., Fujii, M., & Sakata, Y. (2007). Green tea proanthocyanidins inhibit cyclooxygenase-2 expression in LPS-activated mouse macrophages: molecular mechanisms and structure-activity relationship. *Archives of Biochemistry and Biophysics*. 460,67–74. doi: 10.1016/j.abb.2007.01.009.
- [71] Hou, D.X., Yanagita, T., Uto, T., Masuzaki, S., Fujii, M. (2005). Anthocyanidins inhibit cyclooxygenase-2 expression in LPS-evoked macrophages: structure-activity relationship and molecular mechanisms involved. *Biochemical journal. Pharmacology*, 70, 417–425. doi: 10.1016/j.bcp.2005.05.003.
- [72] Pergola, C., Rossi, A., Dugo, P., Cuzzocrea, S. & Sautebin, L. (2006). Inhibition of nitric oxide biosynthesis by anthocyanin fraction of blackberry extract. *Nitric Oxide*, 15(1) :30–39.
- [73] Herath, H.M., Takano-Ishikawa, Y., Yamaki, K. (2003). Inhibitory effect of some flavonoids on tumor necrosis factor- α production in lipopolysaccharide-stimulated mouse macrophage cell line J774.1. *Journal of Medical Food*, 6,365–370. doi: 10.1089/109662003772519930.
- [74] Wang, H., Sun, X., Zhang, N., Ji, Z., Ma, Z., Fu, Q., Qu, R., & Ma, S. (2017) Ferulic acid attenuates diabetes-induced cognitive impairment in rats via regulation of PTPIB and insulin signalling pathway. *Physiology & Behaviour*, 182, 93 – 100.
- [75] Vandanmagsar B, Youm YH, Ravussin A, Galgani JE, Stadler K, Mynatt RL, Ravussin E, Stephens JM, Dixit VD. (2011) The NLRP3 inflammasome instigates obesity-induced inflammation and insulin resistance. *Nat Med*.17(2):179-88. doi: 10.1038/nm.2279. Epub 2011 Jan 9. PMID: 21217695; PMCID: PMC3076025
- [76] Shao, Y., Nanayakkara, G., Cheng, J., Cueto, R., Yang, W.Y., Park, J., Wang, H and Yang, X (2018) Lysophospholipids and Their Receptors Serve as Conditional DAMPs and DAMP Receptors in Tissue Oxidative and Inflammatory Injury. *Antioxidant & Redox Signaling*, 28(10), <https://doi.org/10.1089/ars.2017.7069>
- [77] Ballak DB, Stienstra R, Tack CJ, Dinarello CA, van Diepen JA.(2015) IL-1 family members in the pathogenesis and treatment of metabolic disease: Focus on adipose tissue inflammation and insulin resistance. *Cytokine*. 75(2):280-90. doi: 10.1016/j.cyto.2015.05.005. Epub 2015 Jul 17. PMID: 26194067; PMCID: PMC4553099.
- [78] Meier, D.T., Souza, J.D., Donath, M.Y (2024) Targeting the NLRP3 inflammasome-IL-1 β pathway in type 2 Diabetes and Obesity. *Diabetologia*, 68 (1): 3-16.

- [79] Li X., Watanabe K., Kimura I. (2017) Gut Microbiota Dysbiosis Drives and Implies Novel Therapeutic Strategies for Diabetes Mellitus and Related Metabolic Diseases. *Frontiers in Immunology*, 8:1882. doi: 10.3389/fimmu.2017.01882.
- [80] Borra MT, Smith BC, Denu JM, Mechanism of human SIRT1 activation by resveratrol. *Journal of Biological Chemistry*, 2005; 280 (17): 17187-17195, doi: 10.1074/jbc.M501250200
- [81] Olcum M, Toston B, Ercan I, Eitutan IB, Genc S, Inhibitory effects of phytochemicals on NLRP3 inflammasome activation: a review. *Phytomedicine*, 2020; 75: 153238. <https://doi.org/10.1016/j.phymed.2020.153238>
- [82] Ungvari, Z., Sonntag, W.E., de Cabo, R., Baur, J.A., and Csiszar, A (2011) Mitochondrial Protection by Resveratrol. *Exercise Sport Science Review*, 39(3): 128 – 132. Doi: 10.1097/JES.ob013e3182141f80.
- [83] He, Y., Hara, H., & Nunez, G. (2016) Mechanism and regulation of NLRP3 inflammasome activation. *Trends in Biochemical Sciences*, 41(12), 1012 -1021. <https://doi.org/10.1016/j.tibs.2016.09.002>
- [84] Zou P., Liu, X., Li, G., & Wang, Y. (2018). Resveratrol pretreatment attenuates traumatic brain injury in rats by suppressing NLRP3 inflammasome activation via SIRT1. *Molecular Medicine Reports*, 17 (2): 3212 – 3217
- [85] Yang, Z., Wang, M., Zhang, Y., Cai, F., Jiang, B., Zha, W., Yu, W. (2020). Metformin ameliorates diabetic cardiomyopathy by activating the PK2/PKR pathway. *Frontiers in Physiology*, 11:425.
- [86] Dziemidowicz, M., Bonda, T.A., Litvinovich, S., Taranta, A., Winnicka, M. M., & Kaminski, K.A (2019) The role of interleukin-6 in intracellular signal transduction after chronic beta-adrenergic stimulation in mouse myocardium, *Archives of medical science: AMS*, 15(6), 1565.
- [87] Rehman, K., Akash, M.S. H., Liaqat, A., Kamal, S., Qadir, M.I., & Rasul, A., (2017) Role of interleukin-6 in development of insulin resistance and type 2 diabetes mellitus. *Critical Reviews in Eukaryotic Gene Expression*, 27(3): 229 – 236.
- [88] Bellaver B., Bobermin, L.D., Souza, D.G., Rodrigues, M.D.N., de Assis, A.M., Wajner, M., Goncalves, C., Souza, D.O., Quincozes-Santos, A (2016) Signaling mechanisms underlying glioprotective effects of resveratrol against mitochondrial dysfunction. *Biochimica et Biophysica Acta (BBA) – Molecular Basis of Disease*, 1862 (9): 1827-1838**
- [89] Ko, Y-J., Kim, H-H., Kim, E-J., Katakura, Y., Lee, WW-S., Kim, G-S., & Ryu, C-H. (2013) Piceatannol inhibits mast cell-mediated allergic inflammation. *International Journal of Molecular Medicine*, 31(4): 951-958. <https://doi.org/10.3892/ijmm.2013.1283>.
- [90] Yang, J., Leng, J., Li, J.J., Tang, J-F., Li, Y., Liu, B-L., & Wen, X-D. (2016) Corsolic acid inhibits adipose tissue inflammation and ameliorates insulin

resistance via AMPK activation in high-fat fed mice. *Phytomedicine*, 23(2), 181-190. <https://doi.org/10.1016/j.phymed.2015.12.018>

- [91] Zhao, H., Zhang, Y., Shu, L., Song, G., and Ma, H (2019) Resveratrol reduces liver endoplasmic reticulum stress and improves insulin sensitivity in vivo and in vitro. *Drug Design, Development and Therapy*, 1473-1485, Doi: 10.2147/DDDT.S203833 Wang, G., Li W; Lu, X., Zhao, X., Xu, L. (2013). Taurine attenuates oxidative stress and alleviates cardiac failure in type I diabetic rats. *Croatian medical journal*, 54:171-9 doi: 10.3325/cmj.2013.54.171.
- [92] Lai, X., Pei, Q., Song, X.U., Zhuo, X., Yin, Z., Jia, R., Zou, Y., Li, L., Yue, G., Liang, X., Yin, L., Lv, C., & Jing, B.O. (2016) The enhancement of immune function and activation of NF-Kb by resveratrol-treatment in immunosuppressive mice. *International immunopharmacology*, 33, 42-47. <https://doi.org/10.1016/j.intimp.2016.01.028>
- [93] Chan, K. H., O'Connell, R. L., Sullivan, D. R., Hoffmann, L. S., Rajamani, K., Whiting, M., Donoghoe, M. W., Vanhala, M., Hamer, A., Yu, B., Stocker, R., Ng, M. K & Keech, A. C. (2013) FIELD Study Investigators. Plasma total bilirubin levels predict amputation events in type 2 diabetes mellitus: the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. *Diabetologia*. 56(4):724-36. doi: 10.1007/s00125-012-2818-4. Epub 2013 Jan 17. PMID: 23322233.
- [94] Dong H, Huang H, Yun X, Kim D, Yue Y, Wu H, Sutter A, Chavin KD, Otterbein LE, Adams DB, Kim Y & Wang H (2014). "Bilirubin increases insulin sensitivity in leptin-receptor deficient and diet-induced obese mice through suppression of ER stress and chronic inflammation," *Endocrinology*, 155(3), 818–828,.
- [95] Wang, J, Li Y., Han X., Hu H., Wang, F., et al. (2017). Serum bilirubin levels and risk of type 2 diabetes: results from two independent cohorts in middle-aged and elderly Chinese, *Scientific Reports*, 7: 41338.
- [96] Pineda, S., Bang OY, Saver JL, Starkman S, Yun SW, Liebeskind DS, Kim D, Ali LK, Shar SH, Ovbiagele B. (2008), Association of serum bilirubin with ischemic stroke outcomes. *J. Stroke Cerebrovasc. Dis.* 17, 147–152, doi: 10.1016/j.jstrokecerebrovasdis.2008.01.009.
- [97] Rochette, L; M. Zeller, Y. Cottin, and C. Vergely,(2018) Redox functions of heme oxygenase-1 and biliverdin reductase in diabetes, *Trends in Endocrinology & Metabolism*, 29(2): 74–85.
- [98] Racheck LI. Free fatty acids and skeletal muscle insulin resistance.(2014), *Prog Mol Biol Transl Sci.*; 121: 267–92.

- [99] Wang, G., Li W; Lu, X., Zhao, X., Xu, L. (2013). Taurine attenuates oxidative stress and alleviates cardiac failure in type I diabetic rats. *Croatian medical journal*, 54:171-9 doi: 10.3325/cmj.2013.54.171.
- [100] Wu, L. and Parhofer, K.G.(2014). Diabetic dyslipidemia. *Metabolism*, 63(12): 1469-79.
- [101] Taskinen M. R. (2002) Diabetic dyslipidemia. *Atheroscler Suppl.* 2002;3(1):47–51. doi: 10.1016/S1567-5688(01)00006-X.
- [102] Ozder A. (2014) Lipid profile abnormalities seen in T2DM patients in primary healthcare in Turkey: a cross-sectional study. *Lipids in Health & Disease*, 6 (13):183. doi: 10.1186/1476-511X-13-183. PMID: 25481115; PMCID: PMC4271485.
- [103] Guérin M, Le Goff W, Lassel TS, Van Tol A, Steiner G & Chapman MJ (2001). Atherogenic role of elevated CE transfer from HDL to VLDL(1) and dense LDL in type 2 diabetes : impact of the degree of triglyceridemia. *ArteriosclerThrombVasc Biol.* 21(2), 282–288.
- [104] Briaud, .I, Harmon, J. S., Kelpe, C. L., Segu, V. B.,&Poitout, V. (2001) Lipotoxicity of the pancreatic betacell is associated with glucose-dependent esterification of fatty acids into neutral lipids. *Diabetes*. 50(2), 315–321.
- [105] Wolfrum C, Asilmaz E, Luca E, Friedman JM, Stoffel M. (2004) Foxa2 regulates lipid metabolism and ketogenesis in the liver during fasting and in diabetes. *Nature* 432: 1027–1032.
- [106] Schenk, S.; Saberi, M.; Olefsky, J.M.(2008), Insulin sensitivity: Modulation by nutrients and inflammation. *J. Clin Investig*, 118, 2992–3002.
- [107] Petersen, M.C.; Shulman, G.I. Mechanisms of insulin action and insulin resistance. (2018) ,*Physiol. Res.* 98(4), 2133–2223.
- [108] Kang, Q., & Yang, C. (2020). Oxidative stress and diabetic retinopathy: Molecular mecha-nisms, pathogenetic role and therapeutic implications. *Redox Biology*, 37, 101-799.
- [109] Hamzah, S. A. (2019). Association between Lipid Profiles and Renal Functions among Adults with Type 2 Diabetes. *International Journal of Diabetes & Metabolic Disorders*, 25,234-138.
- [110] Barańska A, Błaszczuk A, Polz-Dacewicz M, Kanadys W, Malm M, Janiszewska M, Jędrych M. (2021) Effects of Soy Isoflavones on Glycemic Control and Lipid Profile in Patients with Type 2 Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials.

Nutrients, 13(6):1886. doi: 10.3390/nu13061886. PMID: 34072748; PMCID: PMC8229139.

- [111] Zuo, X., Zhao, R., Wu, M., Wan, Q., Li, T (2023) Soy Consumption and the Risk of Type 2 Diabetes and Cardiovascular Diseases: A Systematic Review and Meta-Analysis. *Nutrients*. 2023; 15(6):1358. <https://doi.org/10.3390/nu15061358>, <https://www.mdpi.com/2072-6643/15/6/1358>
- [112] Al-Khayri JM, Mascarenhas R, Harish HM, Gowda Y, Lakshmaiah VV, Nagella P, Al-Mssallem MQ, Alessa FM, Almaghasla MI, Rezk AA. (2023) Stilbenes, a Versatile Class of Natural Metabolites for Inflammation-An Overview. *Molecules*, 28(9):3786. doi: 10.3390/molecules28093786. PMID: 37175197; PMCID: PMC10180133