

Studies of Spacing on Growth and Yield of Different Varieties of Field Pea (*Pisum sativum* L.)

ABSTRACT

A field experiment entitled " **Studies of spacing on growth and yield of different varieties of field pea (*Pisum sativum* L.)**" was undertaken during the *rabi* season of 2023 - 2024 at Pandit Deen Dayal Upadhyay Institute of Agricultural Sciences, Utlou, Bishnupur District, Manipur, India. The treatment comprised of three different spacing (S₁- 20×10 cm, S₂- 30×10 cm, and S₃- 40×10 cm,) and three varieties V₁- Rachna, V₂ –Aman and V₃- Prakash with a total of nine treatment combinations. The experiment was laid out in a Factorial Randomized Block Design (FRBD) with three replications. The results revealed that the individual effect of spacing S₃ - 40×10 cm (S₃) and variety S₃V₂ (V₂) significantly enhanced the growth attributes i.e. plant height, number of branches, fresh and dry weight, number of nodules, dry weight of nodules for all the growth stages recorded. The treatment combination S₃V₂ (40 × 10 cm + Aman) recorded maximum plant height, number of branches, fresh and dry weight, number of nodules and dry weight of nodules for all the growth stages recorded. The different planting spaces and varieties significantly enhanced the yield attributes of pea. The spacing(S₃) and varieties (V₂) significantly increased the number of pods per plant, seeds per pod, pod length, test weight, seed yield, stover yield of pea. The treatment combination S₃V₂ (40 × 10 cm + Aman) gave the maximum seed yield (20.69 kg/ha) and stover yield (25.19 kg/ha). The highest gross return, net return and highest benefit-cost ratio were obtained from the treatment S₃V₂ (40 × 10 cm + Aman). Thus, from the experiment it can be concluded that the application of S₃V₂ (40 × 10 cm + Aman) is more favorable for attaining sustainable higher profits and productivity in the cultivation of pea during Rabi season of Manipur.

Keywords: Pea, spacing, varieties, growth, yield.

1. INTRODUCTION

Pea (*Pisum sativum* L.) is a very common leguminous crop grown in the rabi season throughout the world. Pea crop own a strategic position in Indian agriculture as it is an excellent source of dietary protein and a mini-nitrogen plant having ameliorative effect on soil. It helps in improving physical, chemical and biological properties of soil and also utilize natural resources in a better way (Kolb *et al.*, 2017). Less inputs particularly the irrigation and fertilizer are needed in cultivation of pea. It improves soil fertility due to fixation of nitrogen by rhizobium bacteria. Nutritionally, pea contains, 7.2 g, fats 0.1 g, minerals 0.8 g, carbohydrates 15.8 g, calcium 20 mg, magnesium 34 mg, copper 0.23 mg, iron 1.5 mg and vitamin C 9.0 mg/100 g of edible portion (Sepehya *et al.*, 2015).

Spacing is also one of the important parameters, which ultimately affected nutrients uptake, growth and yield of plant. Increase in spacing, the total population decrease, but with more nutrition the individual plant grows better and get more yield and vice-versa. The increase or decrease of row spacing's and plant population has definite pattern in relation to the yield. Among various agronomic yield limiting factors, planting pattern is considered of great importance. Lone *et al.* (2009) stated that the optimum plant density with proper geometry of planting is dependent on variety, its growth habit and agroclimatic conditions. Optimum spacing is necessary to obtain maximum yield in any crop by reducing the competition among the plants for light, nutrient, moisture, etc. Optimum spacing for any crop varies considerably due to environment under which it is grown and different variety.

In agriculture, varieties are essential for enhancing productivity and are chosen based on their ability to grow in specific environments and maximize yield potential. The growth and yield of a crop are influenced by several factors, and selecting the right variety plays a crucial role in improving both. Each variety has a genetic potential for growth, which defines its maximum size, growth rate, and the ability to produce leaves, flowers, and roots. This potential is determined by genes that regulate cell division, elongation, and differentiation. Varieties with superior genetic growth potential can achieve better vegetative growth, which supports higher yield outcomes. Keeping these points in views, the present investigation entitled "Studies of Spacing on Growth and Yield of Different varieties of Field Pea (*Pisum sativum* L.) was conducted during *rabi* 2023-24 at the farm of Pandit Deen Dayal Upadhyay Institute of Agriculture Science, Utlou, Bishnupur, Manipur.

2. MATERIAL AND METHODS

The field experiment was conducted at the experimental site of the Pandit Deen Dayal Upadhyay Institute of Agricultural Sciences situated in Utlou, Bishnupur District, Manipur, during the Rabi season of 2023–2024 which is at 24°43'23"N latitude & 93°51'33"E longitude and at an altitude of 790 above mean sea level (MLS). The soil of the experimental site was clayey, the soil pH was acidic in reaction (5.2) with high organic carbon content (1.9%). The available nitrogen (188 kg/ha) is low and phosphorus (20.0 kg/ha) is medium and potassium (324.0 kg/ha) is high in range according to TNAU soil rating chart. During the period of experimentation, the monthly maximum and minimum temperature were between 22.3° C - 28.9°C and 4.6°C - 8.8°C, and the maximum and minimum relative humidity were recorded between 93% - 94% and 32% - 57%, respectively. There are nine treatments and three replications laid out in a Factorial Randomized Complete Block Design (FRBD). The treatments were: T₁ S₁V₁ 20×10 cm + Rachna T₂ S₁V₂ 20×10 cm + Aman T₃ S₁V₃ 20×10 cm + Prakash T₄ S₂V₁ 30×10 cm + Rachna T₅ S₂V₂ 30×10 cm + Aman T₆ S₂V₃ 30×10 cm + Prakash T₇ S₃V₁ 40×10 cm + Rachna T₈ S₃V₂ 40×10 cm + Aman T₉ S₃V₃ 40×10 cm + Prakash. A uniform dose of 20 kg nitrogen (as urea), 60 kg phosphorus (SSP) and 40 kg potash (MOP) were applied to all the treatments during the time of sowing. The biometric observation on different growth and yield attributes were recorded at various crop growth period

3. RESULTS AND DISCUSSION

3.1 Effect of spacing and varieties on plant height (cm)

The data on plant height as influenced by spacing and varieties recorded during 30, 60, 90 DAS and at harvest are presented in Table 1. The individual effect of spacing on plant height of field pea has been found to be significant for all the growth stages. In general, the spacing, S₃ (40×10 cm) recorded the maximum plant height i.e. 11.69 cm, 30.13 cm, 41.62 cm and 42.84 cm respectively during 30, 60, 90 DAS and at harvest, and S₁ (20×10 cm) recorded minimum plant height i.e. 10.89 cm, 28.88 cm, 39.31 cm and 46.29 cm. Again, the varieties, V₂ (Aman) recorded maximum plant height i.e. 12.98 cm, 30.68 cm, 46.29 cm and 47.22 cm, respectively during 30, 60, 90 DAS and at harvest and V₁ (Rachna) recorded minimum plant height i.e. 9.77 cm, 27.92 cm, 35.61 cm, and 36.62 cm respectively during 30, 60, 90 DAS and at harvest. The combined effect of spacing and varieties on plant height of field pea has been found to be non-significant for all the growth stages. Significant enhancement in plant height under different spacing and varieties seems to be due to increase in cell division which results in rapid growth of plants obtained by Yadav (2003) in cowpea and Sen *et.al.* (2005) in dwarf field pea. These findings are in good lines with those obtained by Khan *et al.* (2021), reported that increased row spacing increase the plant height.

Table 1. Effect of spacing and varieties on plant height (cm)

Treatments	Plant height (cm)			
	30 DAS	60 DAS	90 DAS	At harvest
Spacing				
S ₁ : 20×10 cm	10.89	28.88	39.31	40.98
S ₂ : 30×10 cm	11.24	29.51	40.53	42.04
S ₃ : 40×10 cm	11.69	30.13	41.62	42.84
S.Ed (±)	0.10	0.36	0.42	0.32
C.D. (P = 0.05)	0.21	0.77	0.88	0.67
Varieties				
V ₁ : Rachna	9.77	27.92	35.61	36.62
V ₂ : Aman	12.98	30.68	46.29	47.22
V ₃ : Prakash	11.07	29.92	39.55	42.03
S.Ed (±)	0.10	0.36	0.42	0.32
C.D. (P = 0.05)	0.21	0.77	0.88	0.67

3.2 Effect of spacing and varieties on number of branches per plant of field pea.

The data on number of branches per plant as influenced by spacing and varieties recorded during 30, 60, 90 DAS and at harvest are presented in Table 2. The individual effect of spacing on number of branches per plant of field pea has been found to be significant for all the growth stages. In general, the spacing, S₃ (40×10 cm) recorded the maximum number of branches per plant i.e. 1.30, 2.53, 2.95 and 3.93 which was at par with S₂ again S₂ was also at par with S₁ during 30 DAS and 60 DAS but superior at 90 DAS and at harvest. Minimum number of branches per plant was recorded in S₁ (20×10 cm) i.e.1.19, 2.42, 2.74 and 3.7. Again, the varieties, V₂ (Aman) recorded maximum number of branches per plant i.e. 1.37, 3.12, 3.52 and 4.62, respectively during 30, 60, 90 DAS and at harvest and V₁ (Rachna) recorded minimum number of branches per plant i.e. 1.12, 1.94, 2.31, and 3.20 respectively during 30, 60, 90 DAS and at harvest. The combined effect of spacing and varieties on number of branches per plant of field pea has been found to be non-significant for all the growth stages. Significant enhancement in number of branches per plant might be due to

different row spacing which had sufficient space, nutrients, moisture and sunlight for better overall development of individual plant. The variation in production of branches per plant in varieties may be due to genetically makeup of individual varieties. This result is also obtained by Sajib *et al.* (2012) and Kumari *et al.* (2021) in pea.

Table 2. Effect of spacing and varieties on number of branches per plant of field pea.

Treatments	Number of branches per plants			
	30 DAS	60 DAS	90 DAS	At harvest
Spacing				
S₁: 20×10 cm	1.19	2.42	2.74	3.71
S₂: 30×10 cm	1.27	2.48	2.89	3.84
S₃: 40×10 cm	1.30	2.53	2.95	3.93
S.Ed (±)	0.04	0.03	0.02	0.02
C.D. (P = 0.05)	0.08	0.07	0.03	0.03
Varieties				
V₁: Rachna	1.12	1.94	2.31	3.20
V₂: Aman	1.37	3.12	3.52	4.62
V₃: Prakash	1.27	2.37	2.74	3.65
S.Ed (±)	0.04	0.03	0.02	0.02
C.D. (P = 0.05)	0.08	0.07	0.03	0.03

3.3 Effect of spacing and varieties on number of pods per plant

The data on number of pods per plants as influenced by spacing and varieties are presented in Table 3. The individual effect of spacing and varieties on number of pods per plants of field pea has been found to be significant. Among the different spacing S₃ (40×10 cm) recorded the maximum number of pods per plants i.e. 13.01 and S₁ (20×10 cm) recorded minimum pods length per plant i.e. 10.87. Among the different variety, V₂ (Aman) recorded maximum number of pods per plants i.e. 13.53 and V₁ (Rachna) recorded minimum number of pods per plants i.e. (10.46). The combined effect of spacing and varieties on number of pods per plant of field pea has been found to be non-significant for all the growth stages. This might be due to wider row spacing which give the sufficient space of individual plant for better reproductive growth and increase the pod bearing ability because easily provide essential plant nutrients in this row spacing. Significant variation in pods per plant may be correlated with the number of branches. Significant interaction between row spacing on number of pods plants was also reported by Sajid *et al.* (2012) and Shaukat *et al.* (2012) in field pea.

3.4 Pods length (cm)

The data on pods length of plants as influenced by spacing and varieties are presented in Table 3. The individual effect of spacing and varieties on number of pods length of plants of field pea has been found to be significant. Among the different spacing S₃ (40×10 cm) recorded the maximum number of pods length i.e. 6.01 cm and S₁ (20×10 cm) recorded minimum pods length per plant i.e. 4.71 cm. Among the different variety, V₂ (Aman) recorded maximum number of pods length i.e. 5.65 cm and V₁ (Rachna) recorded minimum number of pods length i.e. (4.98 cm). The interaction of spacing and varieties on number of pods length of field pea was found to be significant for all the growth stages of field pea. The maximum number of pods length per plants was found to be for the treatment S₃V₂ (40×10 cm + Aman), i.e. (6.11). The treatments combination of (S₂V₂, S₃V₁, S₃V₂ and S₃V₃) and (S₁V₂, S₁V₃, S₂V₁ and S₂V₃) was recorded to be at par with each other. The lowest number of pods

length was for the treatment S₁V₁(20×10 cm + Rachna), i.e. (4.23 cm). Enhancement in number of pod length under different spacing and varieties seems to be due to the variation in pod length among varieties which accounted for varietals inheritance. Significant interaction between row spacing on pod length was also observed by Alizai *et al.* (2005) in pea. Significant effect on variety were also reported by Bhutia *et al.* (2017) in pea.

3.5 Seed yield (q/ha)

The data on seed yield as influenced by spacing and varieties are presented in Table 3. The individual effect of spacing and varieties on seed yield of field pea has been found to be significant. Among the different spacing S₃ (40×10 cm) recorded the maximum seed yield i.e. 17.60 q/ha and S₁ (20×10 cm) recorded minimum seed yield i.e.13.81 q/ha. Among the different variety, V₂ (Aman) recorded maximum seed yield i.e.18.46 q/ha and V₁ (Rachna) recorded minimum seed yield i.e. (13.06 q/ha). The interaction of spacing and varieties on seed yield of field pea was found to be significant for all the growth stages of field pea. The seed yield ranged from 11.21 q/ha to 21.11 q/ha. The maximum seed yield was found to be for the treatment S₃V₂ (40×10 cm + Aman), i.e. (20.69 q/ha) followed by treatment S₂V₂ (30×10 cm + Aman) i.e. 18.34 q/ha. The lowest seed yield was for the treatment S₁V₁ (20×10 cm + Rachna) i.e. (11.21 q/ha). The variation in seed yield in varieties may be due to maximum number of nodules per plant, pods per plant, seed yield per plant and better seed index. This favorable phenomenon resulted in higher yield. Significant interaction between row spacing on seed yield was observed by Hussain *et al.* (2017) in pea. Significant effect on variety were also reported by Kumar *et al.* (2018) in field pea. Significant interaction between row spacing and variety on seed yield/plant was also reported by Malek *et al.* (2012) and Mondal *et al.* (2014).

208 **Table 3.** Effect of spacing and varieties on number of pods, pod length (cm) and seed yield
 209 (q/ha) of field pea.
 210

Treatments	Number of pods	Pod length (cm)	Seed yield (q/ha)
Spacing			
S_1	10.19	4.71	11.26
S_2	10.50	5.25	12.33
S_3	10.72	6.01	12.86
S.Ed(±)	0.01	0.11	0.04
C.D	0.03	0.23	0.08
Varieties			
V_1	9.74	4.98	10.60
V_2	11.08	5.65	13.56
V_3	10.58	5.34	12.29
S.Ed(±)	0.01	0.11	0.04
C.D	0.03	0.23	0.08
Spacing x varieties			
$S_1 V_1$	9.60	4.23	9.41
$S_1 V_2$	12.50	5.03	10.86
$S_1 V_3$	11.01	4.87	10.30
$S_2 V_1$	10.45	4.81	9.81
$S_2 V_2$	13.66	5.82	11.05
$S_2 V_3$	12.24	5.12	10.64
$S_3 V_1$	11.32	5.90	10.01
$S_3 V_2$	14.43	6.11	11.34
$S_3 V_3$	13.27	6.04	10.81
S.Ed(±)	0.11	0.18	0.03
C.D	0.22	0.39	0.05

211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222

4. CONCLUSION

Based on the results from the experiment it can be concluded that the effect of spacing and varieties on growth and yield of field pea (*Pisum sativum* L.) significantly increases the growth attributes, yield and yield attributes of field pea. The spacing (40×10 cm) and variety (Aman) was found best for field pea cultivation. The treatment combination S_3V_2 (40×10 cm + Aman) was found most effective from all the other treatment. From this research outputs we can conclude that the spacing i.e. 40×10 cm and variety i.e. S_3V_2 (70,810) leads to better net returns.

REFERENCES

- Bhutia, T. L., Shivani. and Saurabh, K. (2017). Evaluation of different varieties of pea (*Pisum sativum* L.) for yield and quality under late sown conditions in Eastern region. *C.R.*, 52(4-5):176-179.
- Hussain, M., Qasim, M. and Ali, S. (2017). Optimal Plant Spacing for Growth and Yield in Peas. *J. Agric. Sci.*, 9(1): 55-62.
- Khan, A., Ahmad, M. and Rahman, S. (2021). Soil Structure and Plant Height Growth in Peas: The Role of Spacing. *Soil Biol. Biochem.*, 156, 108171.
- Kolb AY, Kolb DA. Experiential learning theory as a guide for experiential educators in higher education. *Experiential Learning & Teaching in Higher Education*. 2017;1(1):7-44
- Kumar, R., Singh, J. and Sharma, P. (2018). Comparative Growth and Yield performance of different pea varieties in Punjab. *Indian J. Agric. Sci.*, 88(5): 765-770.
- Kumari, P., Singh, R. and Tripathi, A. (2021). Soil Nutrient Availability and Its Impact on Branch Growth of Peas at Varying Spacings. *S.S.P.N.*, 21(2): 1500-1508.
- Lone, Bilal Ahmad, Hasan, Badrul, Singh Amarjeet, Haq S. A. and Sofi, R. (2009). Effects of seed rate, row spacing and fertility levels on yield attributes and yield of soya bean under temperate condition. *J. Agric. Bio. Sci.* 4 (2): 19-25.
- Malek, M.A., Shafiquzzaman, M., Rahman, M.S., Ismail M.R. and Mondal, M.M.A. (2012). Standardization of soya bean row spacing based on morpho-physiological characters. *Legume Res.* 35(2): 138-143.
- Mondal, M.M.A., Puteh, A.B., Kashem, M.A. and Hasan, M>M (2014). Effect of plant density on canopy structure and dry matter partitioning into plant parts of soy bean [*Glycine max* (L). *Life Sci. J.*, 11(3): 67-74
- Sajib, M., Rab, A., Amin, N.U., Fazaliwahid, Jan, 1., Ahmad, 1., Khan, I.A. and Khan, MA (2012). Effect of herbicides and row spacing on growth and yield of pea. *Pak. J. Weed sci. Res.*, 18 (1): 1-13.
- Sen, K.C., Prasad, S.M. and Sinha, S.P. (2005). Effect of plant population and nitrogen level on growth, yield and yield attributes of dwarf field pea in North Bihar. *J. Appl. Biol.*, 15 (1): 25-27.
- Sepehya S, Bhardwaj SK, Dhiman S. Quality Attributes of Garden Pea (*Pisum sativum* L.) as influenced by Integrated Nutrient Management under Mid Hill Conditions. *J. Krishi Vigyan.*, 2015, 3(2): 78-83.
- Shaukat SA, Ahmad Z, Choudhary YA, Shaukat SK. Effect of different sowing dates and row spacing on the growth, seed yield and quality of off-season pea (*Pisum sativum* L. Cv. Climax) under temperate conditions of Rawalkot Azad Jammu and Kashmir. *J. Sci. Agric.* 2012, (15): 117-125.
- Yadav, G.L. (2003). Effect of sowing time, row spacing and seed rate on yield of cowpea under rainfed condition. *Indian J. Pulses Res.*, 16 (2): 157-158.