

A Study On Gaussian Generalized Edouard Numbers

Abstract-In this study, we define Gaussian generalized Edouard numbers in detail, and focus on two specific cases: Gaussian Edouard numbers, Gaussian Edouard-Lucas numbers.

In addition, we present some identities and matrices related to these sequences, as well as recurrence relations, Binet's formulas, generating functions, Simpson's formulas, and summation formulas.

Keywords: Gaussian Edouard numbers, Gaussian Edouard-Lucas numbers.

1. Introduction

In this section, firstly, we give some preliminary result on Edouard numbers.

The generalized Edouard sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2)\}_{n \geq 0}$ is defined by the third-order recurrence relation as

$$W_n = 7W_{n-1} - 7W_{n-2} + W_{n-3} \quad (1.1)$$

with the initial values W_0, W_1, W_2 not all being zero.

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = 7W_{-(n-1)} - 7W_{-(n-2)} + W_{-(n-3)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n .

Characteristic equation of $\{W_n\}$ is

$$z^3 - 7z^2 + 7z - 1 = (z^2 - 6z + 1)(z - 1) = 0$$

whose roots are

$$\alpha = 3 + 2\sqrt{2},$$

$$\beta = 3 - 2\sqrt{2},$$

$$\gamma = 1.$$

Note that

$$\begin{aligned}\alpha + \beta + \gamma + \delta &= 7, \\ \alpha\beta + \alpha\gamma + \beta\delta &= 7, \\ \alpha\beta\gamma &= 1.\end{aligned}$$

For $n = 1, 2, 3, \dots$. Hence, recurrence (1.1) is true for all integer n . Soykan has conducted a study on this particular generalized Edouard sequence and its three spacial cases, for more details, see [20]

Third order recurrence relations has been studied by many authors, for more detail see [2,3,5,6, 7,12,13,15,16,18,19,24,28,29].

Next, we present Binet's formula of generalized Edouard numbers.

THEOREM 1.1. [20] *Binet formula of generalized Edouard numbers can be presented as follows:*

$$W_n = \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha^n}{(\alpha - \beta)(\alpha - 1)} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta^n}{(\beta - \alpha)(\beta - 1)} - \frac{(W_2 - 6W_1 + W_0)}{4} \right).$$

Now we define two particular cases of the sequence $\{W_n\}$ as follows: the edouard sequence $\{E_n\}_{n \geq 0}$, the Edouard-Lucas sequence $\{K_n\}_{n \geq 0}$, respectively, by the third-order recurrence relations,

$$E_n = 7E_{n-1} - 7E_{n-2} + E_{n-3}, \quad E_0 = 0, E_1 = 1, E_2 = 7, \tag{1.2}$$

$$K_n = 7K_{n-1} - 7K_{n-2} + K_{n-3}, \quad K_0 = 3, K_1 = 7, K_2 = 35. \tag{1.3}$$

The sequences $\{E_n\}_{n \geq 0}$, $\{K_n\}_{n \geq 0}$ can be extended to negative subscriptys by defining,

$$\begin{aligned}E_{-n} &= 7E_{-(n-1)} - 7E_{-(n-2)} + E_{-(n-3)}, \\ K_{-n} &= 7K_{-(n-1)} - 7K_{-(n-2)} + K_{-(n-3)},\end{aligned}$$

for $n = 1, 2, 3, \dots$ respectively. As a result, recurrences (1.2)-(1.3) hold for all integer n .

For all integers n , Edouard and Edouard-Lucas numbers can be expressed using Binet's formulas as

$$\begin{aligned}E_n &= \frac{\alpha^{n+1}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^{n+1}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}, \\ K_n &= \alpha^n + \beta^n + 1.\end{aligned}$$

Note that, Gaussian numbers, generally known as Gaussian integers, are a subset of the complex numbers. A complex number is expressed in the form $a + bi$ where a and b are arbitrary real numbers, and i is the imaginary unit such that $i^2 = -1$. Gaussian integers are a specific type of complex number. In other word, z is a Gaussian integers such that $z = a + bi$ where a and b are arbitrary integers.

Next, we give some information about Gaussian sequences from literature.

First, we give some Gaussian numbers with second order recurrence relations.

- Horadam [11] introduced Gaussian Fibonacci numbers and defined by

$$GF_n = F_n + iF_{n-1}$$

where $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$ (in fact, he defined these numbers as $GF_n = F_n + iF_{n+1}$ and he called them as complex Fibonacci numbers.).

- Pethe and Horadam [14] introduced Gaussian generalized Fibonacci numbers by

$$GF_n = F_n + iF_{n-1},$$

where $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$.

- Halıcı and Öz [10] studied Gaussian Pell and Pell Lucas numbers by written , respectively,

$$GP_n = P_n + iP_{n-1},$$

$$GQ_n = Q_n + iQ_{n-1}$$

where $P_n = 2P_{n-1} + P_{n-2}$, $P_0 = 0$, $P_1 = 1$ and $Q_n = 2Q_{n-1} + Q_{n-2}$, $Q_0 = 2$, $Q_1 = 2$.

- Aşçı and Gürel [1] presented Gaussian Jacobsthal and Gaussian Jacobsthal Lucas numbers given by, respectively,

$$GJ_n = J_n + iJ_{n-1},$$

$$Gj_n = j_n + ij_{n-1}$$

where $J_n = J_{n-1} + 2J_{n-2}$, $J_0 = 0$, $J_1 = 1$ and $j_n = j_{n-1} + 2j_{n-2}$, $j_0 = 2$, $j_1 = 1$.

- Taşçı [25] introduced and studied Gaussian Mersenne numbers defined by

$$GM_n = M_n + iM_{n-1}$$

where $M_n = 3M_{n-1} - 2M_{n-2}$, $M_0 = 0$, $M_1 = 1$.

- Taşçı [26] introduced and studied Gaussian balancing and Gaussian Lucas Balancing numbers given by, respectively,

$$GB_n = B_n + iB_{n-1},$$

$$GC_n = C_n + iC_{n-1}$$

where $B_n = 6B_{n-1} - BJ_{n-2}$, $B_0 = 0$, $B_1 = 1$ and $C_n = 6Cj_{n-1} - C_{n-2}$, $C_0 = 1$, $C_1 = 3$.

- Ertaş and Yılmaz [8] studied Gaussian Oresme numbers and defined them as

$$GS_n = S_n + iS_{n-1}$$

where oresme numbers are given by $S_n = S_{n-1} - \frac{1}{4}S_{n-2}$, $S_0 = 0$, $S_1 = \frac{1}{2}$.

Now, we present some gaussian numbers with third order recurrence relations.

- Soykan, et al. [22] presented Gaussian generalized Tribonacci numbers given by

$$GW_n = W_n + iW_{n-1}$$

where $W_n = W_{n-1} + W_{n-2} + W_{n-3}$, with the initial condition W_0, W_1, W_2 .

- Taşcı [27] studied Gaussian Padovan and Gaussian Pell- Padovan numbers by written, respectively,

$$GP_n = P_n + iP_{n-1}$$

$$GR_n = R_n + iR_{n-1}$$

where $P_n = P_{n-2} + P_{n-3}$, $P_0 = 1$, $P_1 = 1$, $P_2 = 1$, and $R_n = 2R_{n-2} + R_{n-3}$, $R_0 = 1$, $R_1 = 1$, $R_2 = 1$.

- Cerda-Morales [4] defined Gaussian third-order Jacobsthal numbers as

$$GJ_n = J_n + iJ_{n-1}$$

where $J_n = J_{n-1} + J_{n-2} + 2J_{n-3}$, $J_1 = 0$, $J_2 = 1$, $J_3 = 1$.

2. Gaussian Generalized Edouard Numbers

In this section, we define Gaussian generalized Edouard numbers and present some properties such as Binet's formula and generating function.

Gaussian generalized Edouard numbers $\{GW_n\}_{n \geq 0} = \{GW_n(GW_0, GW_1, GW_2)\}_{n \geq 0}$ are defined by

$$GW_n = 7GW_{n-1} - 7GW_{n-2} + GW_{n-3}, \tag{2.1}$$

with the initial conditions

$$GW_0 = W_0 + i(7W_0 - 7W_1 + W_2),$$

$$GW_1 = W_1 + iW_0,$$

$$GW_2 = W_2 + iW_1,$$

not all being zero. The sequences $\{GW_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$GW_{-n} = 7GW_{-(n-1)} - 7GW_{-(n-2)} + GW_{-(n-3)} \tag{2.2}$$

for $n = 1, 2, 3, \dots$. Thus, recurrence (2.1) holds for all integer n . Note that for all integers n , we get,

$$GW_n = W_n + iW_{n-1}, \tag{2.3}$$

$$GW_{-n} = W_{-n} + iW_{-n-1}.$$

The first few generalized Gaussian Edouard numbers with positive subscript and negative subscript are presented in the following table.

Table 1. The first few generalized Gaussian Edouard numbers.

n	GW_n
0	$W_0 + i(7W_0 - 7W_1 + W_2)$
1	$W_1 + iW_0$
2	$W_2 + iW_1$
3	$W_0 - 7W_1 + 7W_2 + iW_2$
4	$7W_0 - 48W_1 + 42W_2 + i(W_0 - 7W_1 + 7W_2)$
5	$42W_0 - 287W_1 + 246W_2 + i(7W_0 - 48W_1 + 42W_2)$
6	$246W_0 - 1680W_1 + 1435W_2 + i(42W_0 - 287W_1 + 246W_2)$
7	$1435W_0 - 9799W_1 + 8365W_2 + i(246W_0 - 1680W_1 + 1435W_2)$
8	$8365W_0 - 57120W_1 + 48756W_2 + i(1435W_0 - 9799W_1 + 8365W_2)$

and with negative subscript

n	GW_{-n}
0	$W_0 + i(7W_0 - 7W_1 + W_2)$
1	$7W_0 - 7W_1 + W_2 + i(42W_0 - 48W_1 + 7W_2)$
2	$42W_0 - 48W_1 + 7W_2 + i(246W_0 - 287W_1 + 42W_2)$
3	$(246W_0 - 287W_1 + 42W_2) + i(1435W_0 - 1680W_1 + 246W_2)$
4	$(1435W_0 - 1680W_1 + 246W_2) + i(8365W_0 - 9799W_1 + 1435W_2)$
5	$(8365W_0 - 9799W_1 + 1435W_2) + i(48756W_0 - 57120W_1 + 8365W_2)$
6	$(48756W_0 - 57120W_1 + 8365W_2) + i(284172W_0 - 332927W_1 + 48756W_2)$
7	$(284172W_0 - 332927W_1 + 48756W_2) + i(1656277W_0 - 1940448W_1 + 284172W_2)$
8	$(1656277W_0 - 1940448W_1 + 284172W_2) + i(9653491W_0 - 11309767W_1 + 1656277W_2)$

Gaussian Edouard numbers, $GW_n : GW_n(0, 1, 7 + i) = GE_n$, are defined by

$$GE_n = 7GE_{n-1} - 7GE_{n-2} + GE_{n-3} \tag{2.4}$$

with the initial conditions

$$GE_0 = 0, GE_1 = 1, GE_2 = 7 + i.$$

Gaussian Edouard-Lucas numbers, $GW_n(3 + 7i, 7 + 3i, 35 + 7i) = GK_n$, are defined by

$$GK_n = 7GK_{n-1} - 7GK_{n-2} + GK_{n-3} \tag{2.5}$$

with the initial conditions

$$GK_0 = 3 + 7i, GK_1 = 7 + 3i, GK_2 = 35 + 7i.$$

Note that for all integers n , we have

$$GE_n = E_n + iE_{n-1},$$

$$GK_n = K_n + iK_{n-1}.$$

The first few values of Gaussian Edouard numbers, Gaussian Edouard-Lucas numbers with positive and negative subscript are given in the Table 2.

Table 2. Special cases of Gaussian generalized Edouard numbers with positive and negative subscripts.

n	0	1	2	3	4	5	6
GE_n	0	1	$7 + i$	$42 + 7i$	$246 + 42i$	$1435 + 246i$	$8365 + 1435i$
GE_{-n}		i	$1 + 7i$	$7 + 42i$	$42 + 246i$	$246 + 1435i$	$1435 + 8365i$
GK_n	$3 + 7i$	$7 + 3i$	$35 + 7i$	$199 + 35i$	$1155 + 199i$	$6727 + 1155i$	$39203 + 6727i$
GK_{-n}		$7 + 35i$	$35 + 199i$	$199 + 1155i$	$1155 + 6727i$	$6727 + 39203i$	$39203 + 228487i$

Next, we present The Binet's formula for the Gaussian generalized Edouard numbers.

THEOREM 2.1. *The Binet's formula for the Gaussian generalized Edouard numbers is*

$$\begin{aligned}
 GW_n = & \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha^n}{(\alpha - \beta)(\alpha - 1)} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta^n}{(\beta - \alpha)(\beta - 1)} - \frac{(W_2 - 6W_1 + W_0)}{4} \right) \\
 & + i \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha^{(n-1)}}{(\alpha - \beta)(\alpha - 1)} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta^{(n-1)}}{(\beta - \alpha)(\beta - 1)} - \frac{(W_2 - 6W_1 + W_0)}{4} \right).
 \end{aligned}$$

Proof. The proof follows from 1.1 and 2.3. \square

The previous Theorem gives the following results, as special cases.

COROLLARY 2.2. *For all integers n , we have following identities,*

$$\begin{aligned}
 \text{(a): } GE_n &= \left(\frac{\alpha^{n+1}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^{n+1}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4} \right) + i \left(\frac{\alpha^n}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^n}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4} \right). \\
 \text{(b): } GK_n &= (\alpha^n + \beta^n + 1) + i(\alpha^{n-1} + \beta^{n-1} + 1).
 \end{aligned}$$

The next Theorem presents the generating function of Gaussian generalized Edouard numbers.

THEOREM 2.3. *Let $f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n x^n$ denote the generating function of Gaussian generalized Edouard numbers. Then,*

$$f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n x^n = \frac{GW_0 + (GW_1 - 7GW_0)x + (GW_2 - 7GW_1 + 7GW_0)x^2}{1 - 7x + 7x^2 - x^3}. \tag{2.6}$$

Proof. Using the definition of Gaussian Edouard numbers, and subtracting $7xf(x)$, $-7x^2f(x)$ and $x^3f(x)$ from $f(x)$ we obtain

$$\begin{aligned}
 (1 - 7x + 7x^2 - x^3)f_{GW_n}(x) &= \sum_{n=0}^{\infty} GW_n x^n - 7x \sum_{n=0}^{\infty} GW_n x^n + 7x^2 \sum_{n=0}^{\infty} GW_n x^n - x^3 \sum_{n=0}^{\infty} GW_n x^n, \\
 &= \sum_{n=0}^{\infty} GW_n x^n - 7 \sum_{n=0}^{\infty} GW_n x^{n+1} + 7 \sum_{n=0}^{\infty} GW_n x^{n+2} - \sum_{n=0}^{\infty} GW_n x^{n+3}, \\
 &= \sum_{n=0}^{\infty} GW_n x^n - 7 \sum_{n=1}^{\infty} GW_{n-1} x^n + 7 \sum_{n=2}^{\infty} GW_{n-2} x^n - \sum_{n=3}^{\infty} GW_{n-3} x^n, \\
 &= (GW_0 + GW_1 x + GW_2 x^2) - 7(GW_0 x + GW_1 x^2) + 7GW_0 x^2 \\
 &\quad + \sum_{n=3}^{\infty} (GW_n - 7GW_{n-1} + 7GW_{n-2} - GW_{n-3}) x^n, \\
 &= GW_0 + GW_1 x + GW_2 x^2 - 7GW_0 x - 7GW_1 x^2 + GW_0 x^2, \\
 &= GW_0 + (GW_1 - 7GW_0)x + (GW_2 - 7GW_1 + 7GW_0)x^2,
 \end{aligned}$$

and rearranging above equation, we get 2.6. \square

Theorem 2.3 gives the following results as special cases,

$$f_{GE_n}(x) = \frac{x + ix^2}{1 - 7x + 7x^2 - x^3}, \quad f_{GK_n}(x) = \frac{(7 + 3i)x^2 - (14 + 42i)x + 3 + 7i}{1 - 7x + 7x^2 - x^3}.$$

3. Some Identities About Recurrence Relations of Gaussian Generalized Edouard Numbers

In this section, we present some identities on Gaussian Edouard, Gaussian Edouard-Lucas numbers.

THEOREM 3.1. *The following equations hold for all integer n*

$$GE_n = \frac{7}{64}GK_{n+3} - \frac{11}{16}GK_{n+2} + \frac{21}{64}GK_{n+1}, \tag{3.1}$$

$$GK_n = 35GE_{n+3} - 238GE_{n+2} + 199GE_{n+1}. \tag{3.2}$$

Proof. To proof identity (3.1), we can write $GE_n = aGK_{n+3} + bGK_{n+2} + cGK_{n+1}$ and solve the system of equations we get,

$$GE_0 = aGK_3 + bGK_2 + cGK_1,$$

$$GE_1 = aGK_4 + bGK_3 + cGK_2,$$

$$GE_2 = aGK_5 + bGK_4 + cGK_3.$$

Then, we obtain $a = \frac{7}{64}, b = -\frac{11}{16}, c = \frac{21}{64}$. The other identities can be found similarly. \square

We can write $GE_n = aGK_{n+3} + bGK_{n+2} + cGK_{n+1}$ and solve the system of equations we get,

$$\begin{aligned} GK_0 &= aGE_3 + bGE_2 + cGE_1, \\ GK_1 &= aGE_4 + bGE_3 + cGE_2, \\ GK_2 &= aGE_5 + bGE_4 + cGE_3. \end{aligned}$$

Then, we obtain $a = 35, b = -238, c = 199$. \square

LEMMA 3.2. ([9]) We assume that $f(x) = \sum_{n=0}^{\infty} a_n x^n$ is the generating function of the sequence $\{a_n\}_{n \geq 0}$. Then the generating functions of the sequences $\{a_{2n}\}_{n \geq 0}$ and $\{a_{2n+1}\}_{n \geq 0}$ are stated as

$$f_{a_{2n}}(x) = \sum_{n=0}^{\infty} a_{2n} x^n = \frac{f(\sqrt{x}) + f(-\sqrt{x})}{2}$$

and

$$f_{a_{2n+1}}(x) = \sum_{n=0}^{\infty} a_{2n+1} x^n = \frac{f(\sqrt{x}) - f(-\sqrt{x})}{2\sqrt{x}}$$

respectively.

The generating functions of the even and odd-indexed generalized Edouard sequences are provided by the following theorem.

THEOREM 3.3. The generating functions of the sequence GW_{2n} and GW_{2n+1} are provided by

$$f_{GW_{2n}}(x) = \frac{GW_0 + (GW_2 - 35GW_0)x + (42GW_0 - 48GW_1 + 7GW_2)x^2}{1 - 35x + 35x^2 - x^3}. \tag{3.3}$$

$$f_{GW_{2n+1}}(x) = \frac{GW_1 + (GW_0 - 42GW_1 + 7GW_2)x + (7GW_0 - 7GW_1 + GW_2)x^2}{1 - 35x + 35x^2 - x^3} \tag{3.4}$$

Proof. We only proof 3.3. From Theorem 2.3 we can obtain following identities:

$$\begin{aligned} f_{GW_n}(\sqrt{x}) &= \frac{GW_0 - \sqrt{x}(GW_1 - 7GW_0) + x(GW_2 - 7GW_1 + 7GW_0)}{7x - 7\sqrt{x} - \sqrt{x^3} + 1}, \\ f_{GW_n}(-\sqrt{x}) &= -\frac{GW_0 + \sqrt{x}(GW_1 - 7GW_0) + x(GW_2 - 7GW_1 + 7GW_0)}{7\sqrt{x} + 7x + \sqrt{x^3} - 1}. \end{aligned}$$

Thereby, using Lemma 3.2 identity 3.3 can be proved . The other identity can be found similarly. \square

From Theorem 3.3, we get the following corollary.

COROLLARY 3.4. **a):**

$$\begin{aligned} f_{GE_{2n}}(x) &= \frac{(1 + 7i)x^2 + (7 + i)x}{-x^3 + 35x^2 - 35x + 1}, \\ f_{GE_{2n+1}}(x) &= \frac{ix^2 + (7 + 7i)x + 1}{-x^3 + 35x^2 - 35x + 1}. \end{aligned}$$

b):

$$\begin{aligned} f_{GK_{2n}}(x) &= \frac{(35 + 199i)x^2 - (70 + 238i)x + (3 + 7i)}{-x^3 + 35x^2 - 35x + 1}, \\ f_{GK_{2n+1}}(x) &= \frac{(7 + 35i)x^2 - (46 + 70i)x + (7 + 3i)}{-x^3 + 35x^2 - 35x + 1}. \end{aligned}$$

From Corollary 3.4 we can obtain the following corollary which presents the identities on Gaussian Edouard sequences.

COROLLARY 3.5. **(a):** $(7 + i)GK_{2n-2} + (1 + 7i)GK_{2n-4} = (3 + 7i)GE_{2n} - (70 + 238i)GE_{2n-2} + (35 + 199i)GE_{2n-4}$.

(b): $iGK_{2n-4} + (7 + 7i)GK_{2n-2} + GK_{2n} = -(70 + 238i)GE_{2n-1} + (35 + 199i)GE_{2n-3} + (3 + 7i)GE_{2n+1}$.

(c): $(5383 + 931i)GE_{2n-4} - (46 + 70i)GE_{2n-2} + (7 + 3i)GE_{2n} = (7 + i)GK_{2n-1} + (1 + 7i)GK_{2n-3}$.

(d): $(7 + 3i)GE_{2n+1} - (46 + 70i)GE_{2n-1} + (7 + 35i)GE_{2n-3} = iGK_{2n-3} + (7 + 7i)GK_{2n-1} + GK_{2n+1}$.

Proof. From Corollary 3.4 we obtain

$$((1 + 7i)x^2 + (7 + i)x)f_{GK_{2n}} = ((3 + 7i) - (70 + 238i)x + (35 + 199i)x^2)f_{GE_{2n}}.$$

The LHS (left hand side) is equal to

$$\begin{aligned} LHS &= ((1 + 7i)x^2 + (7 + i)x) \sum_{n=0}^{\infty} GK_{2n}x^n \\ &= (7 + i)x \sum_{n=0}^{\infty} GK_{2n}x^n + (1 + 7i)x^2 \sum_{n=0}^{\infty} GK_{2n}x^n \\ &= (7 + i) \sum_{n=0}^{\infty} GK_{2n}x^{n+1} + (1 + 7i) \sum_{n=0}^{\infty} GK_{2n}x^{n+2} \\ &= (7 + i) \sum_{n=1}^{\infty} GK_{2n-2}x^n + (1 + 7i) \sum_{n=2}^{\infty} GK_{2n-4}x^n \\ &= (7 + i)(3 + 7i)x + \sum_{n=2}^{\infty} ((7 + i)GK_{2n-2} + (1 + 7i)GK_{2n-4})x^n \end{aligned}$$

whereas the RHS (right hand side) is equal to

$$\begin{aligned} RHS &= ((3 + 7i) - (70 + 238i)x + (35 + 199i)x^2) \sum_{n=0}^{\infty} GE_{2n}x^n \\ &= (3 + 7i) \sum_{n=0}^{\infty} GE_{2n}x^n - (70 + 238i)x \sum_{n=0}^{\infty} GE_{2n}x^n + (35 + 199i)x^2 \sum_{n=0}^{\infty} GE_{2n}x^n \\ &= (3 + 7i) \sum_{n=0}^{\infty} GE_{2n}x^n - (70 + 238i) \sum_{n=0}^{\infty} GE_{2n}x^{n+1} + (35 + 199i) \sum_{n=0}^{\infty} GE_{2n}x^{n+2} \\ &= (3 + 7i) \sum_{n=0}^{\infty} GE_{2n}x^n - (70 + 238i) \sum_{n=1}^{\infty} GE_{2n-2}x^n + (35 + 199i) \sum_{n=2}^{\infty} GE_{2n-4}x^n \\ &= (3 + 7i)(7 + i)x + \sum_{n=2}^{\infty} ((3 + 7i)GE_{2n} - (70 + 238i)GE_{2n-2} + (35 + 199i)GE_{2n-4})x^n. \end{aligned}$$

Comparing the coefficients and the proof of the first identity (a) is done. We can present other identity similarly. \square

We can get an identity related to Gaussian Edouard numbers and Edouard-Lucas numbers given below.

THEOREM 3.6. *For all integers m, n the following identity holds:*

$$GW_{m+n} = E_{m-1}GW_{n+2} + (E_{m-3} - 7E_{m-2})GW_{n+1} + E_{m-2}GW_n.$$

Proof. First, we assume that $m, n \geq 0$. The proof can be given by mathematical induction on m . If $m = 0$ we get,

$$GW_n = E_{-1}GW_{n+2} + (E_{-3} - 7E_{-2})GW_{n+1} + E_{-2}GW_n$$

which is true since $E_{-1} = 0, E_{-2} = 1, E_{-3} = 7$. We assume that the identity given holds for $m \leq k$. For $m = k + 1$, we get,

$$\begin{aligned} GW_{(k+1)+n} &= 7GW_{n+k} - 7GW_{n+k-1} + GW_{n+k-2} \\ &= 7(E_{k-1}GW_{n+2} + (E_{k-3} - 7E_{k-2})GW_{n+1} + E_{k-2}GW_n) \\ &\quad - 7(E_{k-2}GW_{n+2} + (E_{k-4} - 7E_{k-3})GW_{n+1} + E_{k-3}GW_n) \\ &\quad + (E_{k-3}GW_{n+2} + (E_{k-5} - 7E_{k-4})GW_{n+1} + E_{k-4}GW_n) \\ &= (7E_{k-1} - 7E_{k-2} + E_{k-3})GW_{n+2} + ((7E_{k-3} - 7E_{k-4} + E_{k-5}) \\ &\quad - 7(7E_{k-2} - 7E_{k-3} + E_{k-4}))GW_{n+1} + (7E_{k-2} - 7E_{k-3} + E_{k-4})GW_n \\ &= E_kGW_{n+2} + (E_{k-2} - 7E_{k-1})GW_{n+1} + E_{k-1}GW_n \\ &= E_{(k+1)-1}GW_{n+2} + (E_{(k+1)-3} - 7E_{(k+1)-2})GW_{n+1} + E_{(k+1)-2}GW_n. \end{aligned}$$

Consequently, by mathematical induction on m , this proves Theorem 3.6. The other case of m, n can be proved similarly. \square

Taking $GW_n = GE_n$ or $GW_n = GK_n$ in above theorem, respectively, we get.

COROLLARY 3.7. *For all integers m, n , we get,*

$$\begin{aligned} GE_{m+n} &= E_{m-1}GE_{n+2} + (E_{m-3} - 7E_{m-2})GE_{n+1} + E_{m-2}GE_n, \\ GK_{m+n} &= E_{m-1}GK_{n+2} + (E_{m-3} - 7E_{m-2})GK_{n+1} + E_{m-2}GK_n. \end{aligned}$$

4. Simson's Formula

In this section, we present Simpson's formula of generalized Gaussian Edouard numbers. This is a special cases of [17, Theorem 4.1].

THEOREM 4.1 (Simpson’s formula of generalized Gaussian Edouard numbers). *For all integers n , we can write following equality*

$$\begin{aligned} \begin{vmatrix} GW_{n+2} & GW_{n+1} & GW_n \\ GW_{n+1} & GW_n & GW_{n-1} \\ GW_n & GW_{n-1} & GW_{n-2} \end{vmatrix} &= \begin{vmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{vmatrix} \\ &= (6GW_1 - GW_0 - GW_2)(GW_2^2 + 8GW_1^2 + GW_0^2 - 8GW_1GW_2 \\ &\quad + 6GW_0GW_2 - 8GW_0GW_1). \end{aligned}$$

Proof. Using [17, Theorem 4.1] we can obtain the required result. \square

From the Theorem 4.1 we get the following Corollary.

COROLLARY 4.2. *For all integers n , we get the following identities:*

$$\begin{aligned} \text{(a): } \begin{vmatrix} GE_{n+2} & GE_{n+1} & GE_n \\ GE_{n+1} & GT_n & GE_{n-1} \\ GE_n & GE_{n-1} & GE_{n-2} \end{vmatrix} &= 6 - 6i. \\ \text{(b): } \begin{vmatrix} GK_{n+2} & GK_{n+1} & GK_n \\ GK_{n+1} & GK_n & GK_{n-1} \\ GK_n & GK_{n-1} & GK_{n-2} \end{vmatrix} &= -3072 + 3072i. \end{aligned}$$

5. SUM FORMULAS

In this section we identify some sum formulas of generalized Gaussian Edouard numbers.

THEOREM 5.1.

$$\begin{aligned} \text{a): } \sum_{k=0}^n W_k &= \frac{1}{4}(-n + 3)W_n + (n + 2)(7W_{n+1} - W_{n+2}) - (n + 1)W_{n+1} + 2W_2 - 13W_1 + 7W_0. \\ \text{b): } \sum_{k=0}^n W_{2k} &= \frac{1}{32}(-n + 3)W_{2n} + (n + 2)(-7W_{2n+2} + 48W_{2n+1} - 7W_{2n}) - (n + 1)W_{2n+2} + 15W_2 - \\ &\quad 96W_1 + 49W_0. \\ \text{c): } \sum_{k=0}^n W_{2k+1} &= \frac{1}{32}(-n + 3)W_{2n+1} + (n + 2)(-W_{2n+2} + 42W_{2n+1} - 7W_{2n}) - (n + 1)(7W_{2n+2} - \\ &\quad 7W_{2n+1} + W_{2n}) + 9W_2 - 56W_1 + 15W_0. \end{aligned}$$

Proof. It is given in Soykan [23, Theorem 3.3]. \square

In this section, we identify some sum formulas of generalized Gaussian Edouard numbers. \square

THEOREM 5.2. *For all integers $n \geq 0$, we have sum formulas given below:*

$$\begin{aligned} \text{(a): } \sum_{k=0}^n GW_k &= \frac{1}{4}(-n + 3)GW_n + (n + 2)(7GW_{n+1} - GW_{n+2}) - (n + 1)GW_{n+1} + 2GW_2 - 13GW_1 + \\ &\quad 7GW_0. \\ \text{(b): } \sum_{k=0}^n GW_{2k} &= \frac{1}{32}(-n + 3)GW_{2n} + (n + 2)(-7GW_{2n+2} + 48GW_{2n+1} - 7GW_{2n}) - (n + 1)GW_{2n+2} + \\ &\quad 15GW_2 - 96GW_1 + 49GW_0. \end{aligned}$$

$$(c): \sum_{k=0}^n GW_{2k+1} = \frac{1}{32}(- (n+3)GW_{2n+1} + (n+2)(-GW_{2n+2} + 42GW_{2n+1} - 7GW_{2n}) - (n+1)(7GW_{2n+2} - 7GW_{2n+1} + GW_{2n}) + 9GW_2 - 56GW_1 + 15GW_0).$$

Proof. Use Theorem 5.1 and the definition of GW_n . \square

As a special case of the Theorem 5.2, we present the following corollary.

COROLLARY 5.3. *For all integers $n \geq 0$, we have sum formulas given below:*

$$\begin{aligned} a): \sum_{k=0}^n GE_k &= \frac{1}{4}(- (n+3)GE_n + (n+2)(7GE_{n+1} - GE_{n+2}) - (n+1)GE_{n+1} + 1 + 2i). \\ b): \sum_{k=0}^n GE_{2k} &= \frac{1}{32}(- (n+3)GE_{2n} + (n+2)(-7GE_{2n+2} + 48GE_{2n+1} - 7GE_{2n}) - (n+1)GE_{2n+2} + 9 + 15i). \\ c): \sum_{k=0}^n GE_{2k+1} &= \frac{1}{32}(- (n+3)GE_{2n+1} + (n+2)(-GE_{2n+2} + 42GE_{2n+1} - 7GE_{2n}) - (n+1)(7GE_{2n+2} - 7GE_{2n+1} + GE_{2n}) + 7 + 9i). \end{aligned}$$

As a special case of the Theorem 5.2 we present the following corollary.

COROLLARY 5.4. *For all integers $n \geq 0$, we have sum formulas given below:*

$$\begin{aligned} a): \sum_{k=0}^n GK_k &= \frac{1}{4}(- (n+3)GK_n + (n+2)(7GK_{n+1} - GK_{n+2}) - (n+1)GK_{n+1} + 24i). \\ b): \sum_{k=0}^n GK_{2k} &= \frac{1}{32}(- (n+3)GK_{2n} + (n+2)(-7GK_{2n+2} + 48GK_{2n+1} - 7GK_{2n}) - (n+1)GK_{2n+2} + 160i). \\ c): \sum_{k=0}^n GK_{2k+1} &= \frac{1}{32}(- (n+3)GK_{2n+1} + (n+2)(-GK_{2n+2} + 42GK_{2n+1} - 7GK_{2n}) - (n+1)(7GK_{2n+2} - 7GK_{2n+1} + GK_{2n}) - 32). \end{aligned}$$

6. Matrix Formulation of GW_n

In this section, we examine some matrices related to Gaussian Edouard and Gaussian Edouard-Lucas numbers. We define the square matrix A of order 3 as

$$A = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

such that $\det A = 1$. Note that

$$A^n = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^n = \begin{pmatrix} E_{n+1} & -7E_n + E_{n-1} & E_n \\ E_n & -7E_{n-1} + E_{n-2} & E_{n-1} \\ E_{n-1} & -7E_{n-2} + E_{n-3} & E_{n-2} \end{pmatrix}.$$

Then we give the following Lemma.

LEMMA 6.1. For $n \geq 0$ the following identity is true:

$$\begin{pmatrix} GW_{n+2} \\ GW_{n+1} \\ GW_n \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}$$

Proof. The proof can be given by mathematical induction on n . If $n = 0$ we obtain

$$\begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}$$

which is true. We assume that the identity given holds for $n = k$. Thus, the following identity is true.

$$\begin{pmatrix} GW_{k+2} \\ GW_{k+1} \\ GW_k \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}.$$

For $n = k + 1$, we get,

$$\begin{aligned} \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} &= \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} \\ &= \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} GW_{k+2} \\ GW_{k+1} \\ GW_k \end{pmatrix} \\ &= \begin{pmatrix} 7GW_{k+2} - 7GW_{k+1} + GW_k \\ GW_{k+2} \\ GW_{k+1} \end{pmatrix} \\ &= \begin{pmatrix} GW_{k+3} \\ GW_{k+2} \\ GW_{k+1} \end{pmatrix}. \end{aligned}$$

Consequently, by mathematical induction on n , the proof is completed. \square

For the proof see [19].

We define

$$N_{GW} = \begin{pmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{pmatrix}, \tag{6.1}$$

$$S_{GW} = \begin{pmatrix} GW_{n+2} & GW_{n+1} & GW_n \\ GW_{n+1} & GW_n & GW_{n-1} \\ GW_n & GW_{n-1} & GW_{n-2} \end{pmatrix}. \quad (6.2)$$

Now, we have the following theorem with N_{GW} and S_{GW} .

THEOREM 6.2. *Using N_{GW} and S_{GW} , we get,*

$$A^n N_{GW} = S_{GW}.$$

Proof. Note that we get,

$$\begin{aligned} A^n N_{GW} &= \begin{pmatrix} E_{n+1} & -7E_n + E_{n-1} & E_n \\ E_n & -7E_{n-1} + E_{n-2} & E_{n-1} \\ E_{n-1} & -7E_{n-2} + E_{n-3} & E_{n-2} \end{pmatrix} \begin{pmatrix} GW_2 & GW_1 & GW_0 \\ GW_1 & GW_0 & GW_{-1} \\ GW_0 & GW_{-1} & GW_{-2} \end{pmatrix}, \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \end{aligned}$$

where

$$\begin{aligned} a_{11} &= GW_2 E_{n+1} + GW_1 (E_{n-1} - 7E_n) + GW_0 E_n, \\ a_{12} &= GW_1 E_{n+1} + GW_0 (E_{n-1} - 7E_n) + GW_{-1} E_n, \\ a_{13} &= GW_0 E_{n+1} + GW_{-1} (E_{n-1} - 7E_n) + GW_{-2} E_n, \\ a_{21} &= GW_2 E_n + GW_1 (E_{n-2} - 7E_{n-1}) + GW_0 E_{n-1}, \\ a_{22} &= GW_1 E_n + GW_0 (E_{n-2} - 7E_{n-1}) + GW_{-1} E_{n-1}, \\ a_{23} &= GW_0 E_n + GW_{-1} (E_{n-2} - 7E_{n-1}) + GW_{-2} E_{n-1}, \\ a_{31} &= GW_2 E_{n-1} + GW_1 (E_{n-3} - 7E_{n-2}) + GW_0 E_{n-2}, \\ a_{32} &= GW_1 E_{n-1} + GW_0 (E_{n-3} - 7E_{n-2}) + GW_{-1} E_{n-2}, \\ a_{33} &= GW_0 E_{n-1} + GW_{-1} (E_{n-3} - 7E_{n-2}) + GW_{-2} E_{n-2}. \end{aligned}$$

Using Theorem (3.6) the proof is done. \square

By taking, $GW_n = GE_n$ with $GE_0 = 0, GE_1 = 1, GE_2 = 7 + i$ in (6.1) and (6.2), $GW_n = GK_n$ with $GK_0 = 3 + 7i, GK_1 = 7 + 3i, GK_2 = 35 + 7i$ in (6.1) and (6.2), respectively, we get,

$$N_{GE} = \begin{pmatrix} 7+i & 1 & 0 \\ 1 & 0 & i \\ 0 & i & 1+7i \end{pmatrix}, \quad S_{GE} = \begin{pmatrix} GE_{n+2} & GE_{n+1} & GE_n \\ GE_{n+1} & GE_n & GE_{n-1} \\ GE_n & GE_{n-1} & GE_{n-2} \end{pmatrix}$$

$$N_{GK} = \begin{pmatrix} 35+7i & 7+3i & 3+7i \\ 7+3i & 3+7i & 7+35i \\ 3+7i & 7+35i & 35+199i \end{pmatrix}, \quad S_{GK} = \begin{pmatrix} GK_{n+2} & GK_{n+1} & GK_n \\ GK_{n+1} & GK_n & GK_{n-1} \\ GK_n & GK_{n-1} & GK_{n-2} \end{pmatrix}$$

From Theorem 6.2, we get the following corollary.

COROLLARY 6.3. *The following identities are holds*

(a): $A^n N_{GE} = S_{GE}.$

(b): $A^n N_{GK} = S_{GK}.$

References

- [1] Aşçı M., Gürel E., Gaussian Jacobsthal and Gaussian Jacobsthal Lucas numbers, *Ars Combinatoria*, 111, 2013.
- [2] Bruce, I., A modified Tribonacci sequence, *Fibonacci Quarterly*, 22(3), 244–246, 1984.
- [3] Catalani, M., Identities for Tribonacci-related sequences, arXiv:math/0209179, 2012.
- [4] Cerda-Morales, G., On Gauss third-order Jacobsthal numbers and their applications, *Annals of the Alexandru Ioan Cuza University-Mathematics*, 67(2),231-241, 2022.
- [5] Choi, E., Modular Tribonacci Numbers by Matrix Method, *Journal of the Korean Society of Mathematical Education Series B: Pure and Applied. Mathematics.* 20(3), 207–221, 2013.
- [6] Elia, M., Derived Sequences, The Tribonacci Recurrence and Cubic Forms, *Fibonacci Quarterly*, 39 (2), 107-115, 2001.
- [7] Er, M. C., Sums of Fibonacci Numbers by Matrix Methods, *Fibonacci Quarterly*, 22(3), 204-207, 1984.
- [8] Ertaş A., Yılmaz F., On Quaternions with Gaussian Oresme Coefficients, *Turk. J. Math. Comput. Sci.*, 15(1), 192-202, 2023.
- [9] Frontczak, R., Convolutions for Generalized Tribonacci Numbers and Related Results, *International Journal of Mathematical Analysis*, Vol. 12, 2018, no. 7, 307 - 324.
- [10] Halcı S., Öz S., On Some Gaussian Pell And Pell-Lucas Numbers, *Ordu Univ. J. Sci. Tech.*, 6(1), 8-18, 2016.
- [11] Horadam, A.F., Complex Fibonacci Numbers and Fibonacci Quaternions, *Amer. Math. Monthly*, 70 , 289-291,1963.
- [12] Lin, P. Y., De Moivre-Type Identities For The Tribonacci Numbers, *Fibonacci Quarterly*, 26, 131-134, 1988.
- [13] Pethe, S., Some Identities for Tribonacci sequences, *Fibonacci Quarterly*, 26(2), 144–151, 1988.
- [14] Pethe, S., Horadam, A.F., Gaussian Generalized Fibonacci Numbers, *Bull. Austral. Math. Soc.*, 33 , 37-48., 1986.
- [15] Scott, A., Delaney, T., Hoggatt Jr., V., The Tribonacci sequence, *Fibonacci Quarterly*, 15(3), 193–200, 1977.
- [16] Shannon, A., Tribonacci numbers and Pascal’s pyramid, *Fibonacci Quarterly*, 15(3), pp. 268 and 275, 1977.
- [17] Soykan, Y., Simson Identity of Generalized m-step Fibonacci Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 45-56, 2019.
- [18] Soykan, Y. Tribonacci and Tribonacci-Lucas Sedenions. *Mathematics* 7(1), 74, 2019.
- [19] Soykan Y., A Study On Generalized (r,s,t)-Numbers, *MathLAB Journal*, 7, 101-129, 2020.
- [20] Soykan, Y., Generalized Edouard Numbers, *International Journal of Advances in Applied Mathematics and Mechanics*, 9(3), 41-52, 2022.

- [21] Soykan, Y., Generalized Tribonacci Polynomials, *Earthline Journal of Mathematical Sciences*, 13(1), 1-120, 2023.
- [22] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, *Journal of Progressive Research in Mathematics*, 14(2), 2018
- [23] Soykan, Y., Sums and Generating Functions of Special Cases of Generalized Tribonacci Polynomials, *International Journal of Advances in Applied Mathematics and Mechanics*, 11(2), 80-173, 2023.
- [24] Spickerman, W., Binet's formula for the Tribonacci sequence, *Fibonacci Quarterly*, 20, 118-120, 1982.
- [25] Taşçı, D., On Mersenne Numbers, *Journal of Science and Arts*, 4(57), 1021-1028, 2021.
- [26] Taşçı, D., Gaussian Balancing and Gaussian Lucas Balancing Numbers, *Journal of Science and Arts*, 3(44), 661-666, 2018.
- [27] Taşçı, D., Gaussian Padovan and Gaussian Pell- Padovan numbers, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 67(2), 82-88, 2018.
- [28] Yalavigi, C. C., Properties of Tribonacci numbers, *Fibonacci Quarterly*, 10(3), 231-246, 1972.
- [29] Yılmaz, N., Taskara, N., Tribonacci and Tribonacci-Lucas Numbers via the Determinants of Special Matrices, *Applied Mathematical Sciences*, 8(39), 1947-1955, 2014.