

Gaussian Numbers with Generalized Pandita Numbers Components

Abstract: In this study, we define Gaussian generalized Pandita numbers in detail, and focus on two specific cases: Gaussian Pandita numbers, Gaussian Pandita-Lucas numbers.

We present some identities and matrices related to these sequences, as well as recurrence relations, Binet's formulas, generating functions, exponential generating functions, Simson's formulas, and summation formulas.

Keywords: Pandita numbers, Pandita-Lucas numbers, Gaussian Pandita numbers, Gaussian Pandita-Lucas numbers, Binet's formulas, generating functions, exponential generating functions.

1. Introduction

In this section, we give some preliminary result on Pandita numbers.

The generalized Pandita sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, W_3)\}_{n \geq 0}$ is defined by the fourth-order recurrence relations as

$$W_n = 2W_{n-1} - W_{n-2} + W_{n-3} - W_{n-4}. \quad (1.1)$$

with the initial values W_0, W_1, W_2, W_3 are not all being zero.

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = -W_{-(n-1)} + 2W_{-(n-2)} + W_{-(n-3)} - W_{-(n-4)}.$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n . Soykan has conducted a study on this particular sequence, for more details, see [10]

Characteristic equation of $\{W_n\}$ is

$$z^4 - 2z^3 + z^2 - z + 1 = (z^3 - z^2 - 1)(z - 1) = 0.$$

whose roots are

$$\begin{aligned}\alpha &= \frac{1}{3} + \left(\frac{29}{54} + \sqrt{\frac{31}{108}}\right)^{1/3} + \left(\frac{29}{54} - \sqrt{\frac{31}{108}}\right)^{1/3}, \\ \beta &= \frac{1}{3} + \omega \left(\frac{29}{54} + \sqrt{\frac{31}{108}}\right)^{1/3} + \omega^2 \left(\frac{29}{54} - \sqrt{\frac{31}{108}}\right)^{1/3}, \\ \gamma &= \frac{1}{3} + \omega^2 \left(\frac{29}{54} + \sqrt{\frac{31}{108}}\right)^{1/3} + \omega \left(\frac{29}{54} - \sqrt{\frac{31}{108}}\right)^{1/3}, \\ \delta &= 1.\end{aligned}$$

where

$$\omega = \frac{-1 + i\sqrt{3}}{2} = \exp(2\pi i/3).$$

Note that

$$\begin{aligned}\alpha + \beta + \gamma + \delta &= 2, \\ \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta &= 1, \\ \alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta &= 1, \\ \alpha\beta\gamma\delta &= 1.\end{aligned}$$

Note also that

$$\begin{aligned}\alpha + \beta + \gamma &= 1, \\ \alpha\beta + \alpha\gamma + \beta\gamma &= 0, \\ \alpha\beta\gamma &= 1.\end{aligned}$$

For $n = 1, 2, 3, \dots$. Hence, recurrence (1.1) is true for all integer n .

For the fourth-order recurrence relations has been studied by many authors, for more detail see [17, 18, 12, 13, 16, 15, 22, 11, 10, 19].

We now present Binet's formula for the generalized Pandita numbers.

THEOREM 1.1. *[10] Binet formula of generalized Pandita numbers can be presented as follows:*

$$\begin{aligned}W_n &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)\alpha^n}{3\alpha - 2} \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)\beta^n}{3\beta - 2} \\ &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)\gamma^n}{3\gamma - 2} \\ &- W_3 + W_2 + W_0.\end{aligned}$$

Now we define two special cases of the sequence $\{W_n\}$ as follows: The Pandita sequence $\{P_n\}_{n \geq 0}$ and the Pandita-Lucas sequence $\{S_n\}_{n \geq 0}$ are respectively defined by the fourth-order recurrence relations as:

$$P_n = 2P_{n-1} - P_{n-2} + P_{n-3} - P_{n-4}, \quad P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 3, \quad (1.2)$$

$$S_n = 2S_{n-1} - S_{n-2} + S_{n-3} - S_{n-4}, \quad S_0 = 4, S_1 = 2, S_2 = 2, S_3 = 5. \quad (1.3)$$

The sequences $\{P_n\}_{n \geq 0}$, $\{S_n\}_{n \geq 0}$, can be extended to negative subscripts by defining,

$$P_{-n} = P_{-(n-1)} - P_{-(n-2)} + 2P_{-(n-3)} - P_{-(n-4)},$$

$$S_{-n} = S_{-(n-1)} - S_{-(n-2)} + 2S_{-(n-3)} - S_{-(n-4)}.$$

for $n = 1, 2, 3, \dots$ respectively. As a result, recurrences (1.2)-(1.3) hold for all integer n . Binet's formulas of P_n and S_n are given as follows.

COROLLARY 1.2. *For all integers n , Binet's formula of Pandita and Pandita-Lucas numbers are*

$$P_n = \frac{\alpha^{n+3}}{3\alpha - 2} + \frac{\beta^{n+3}}{3\beta - 2} + \frac{\gamma^{n+3}}{3\gamma - 2} - 1,$$

and

$$S_n = \alpha^n + \beta^n + \gamma^n + 1.$$

respectively.

Next, we give the ordinary generating function $\sum_{n=0}^{\infty} W_n z^n$ of the sequence W_n .

LEMMA 1.3. *Suppose that $f_{W_n}(z) = \sum_{n=0}^{\infty} W_n z^n$ is the ordinary generating function of the generalized Pandita sequence $\{W_n\}$. Then, $\sum_{n=0}^{\infty} W_n z^n$ is given by*

$$\sum_{n=0}^{\infty} W_n z^n = \frac{W_0 + (W_1 - 2W_0)z + (W_2 - 2W_1 + W_0)z^2 + (W_3 - 2W_2 + W_1 - W_0)z^3}{1 - 2z + z^2 - z^3 + z^4}.$$

Proof. Take $r = 2, s = -1, t = 1, u = -1$ in Lemma 10. $\square$

Next, we give some information about Gaussian sequences from literature.

We provide some Gaussian numbers that satisfy second-order and third-order recurrence relations.

- Horadam [8] introduced Gaussian Fibonacci numbers and defined by

$$GF_n = F_n + iF_{n-1}.$$

where $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$ (in fact, he defined these numbers as $GF_n = F_n + iF_{n+1}$ and he called them as complex Fibonacci numbers.).

- Pethe and Horadam [9] introduced Gaussian generalized Fibonacci numbers by

$$GF_n = F_n + iF_{n-1}.$$

where $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$.

- Hahcı and Öz [7] studied Gaussian Pell and Pell Lucas numbers by written, respectively,

$$GP_n = P_n + iP_{n-1},$$

$$GQ_n = Q_n + iQ_{n-1}.$$

where $P_n = 2P_{n-1} + P_{n-2}$, $P_0 = 0$, $P_1 = 1$ and $Q_n = 2Q_{n-1} + Q_{n-2}$, $Q_0 = 2$, $Q_1 = 2$.

- Aşcı and Gürel [1] presented Gaussian Jacobsthal and Gaussian Jacobsthal Lucas numbers given by, respectively,

$$GJ_n = J_n + iJ_{n-1},$$

$$Gj_n = j_n + ij_{n-1}.$$

where $J_n = J_{n-1} + 2J_{n-2}$, $J_0 = 0$, $J_1 = 1$ and $j_n = j_{n-1} + 2j_{n-2}$, $j_0 = 2$, $j_1 = 1$.

- Taşçı [23] introduced and studied Gaussian Mersenne numbers defined by

$$GM_n = M_n + iM_{n-1}.$$

where $M_n = 3M_{n-1} - 2M_{n-2}$, $M_0 = 0$, $M_1 = 1$.

- Taşçı [25] introduced and studied Gaussian balancing and Gaussian Lucas Balancing numbers given by, respectively,

$$GB_n = B_n + iB_{n-1},$$

$$GC_n = C_n + iC_{n-1}.$$

where $B_n = 6B_{n-1} - B_{n-2}$, $B_0 = 0$, $B_1 = 1$ and $C_n = 6C_{n-1} - C_{n-2}$, $C_0 = 1$, $C_1 = 3$.

- Ertaş and Yılmaz [5] studied Gaussian Oresme numbers and defined them as

$$GS_n = S_n + iS_{n-1}.$$

where oresme numbers are given by $S_n = S_{n-1} - \frac{1}{4}S_{n-2}$, $S_0 = 0$, $S_1 = \frac{1}{2}$.

Now, we present some Gaussian numbers with third order recurrence relations.

- Soykan et al [20] presented Gaussian generalized Tribonacci numbers given by

$$GW_n = W_n + iW_{n-1}.$$

where $W_n = W_{n-1} + W_{n-2} + W_{n-3}$, with the initial condition W_0, W_1, W_2 .

- Taşcı [24] studied Gaussian Padovan and Gaussian Pell- Padovan numbers by written, respectively,

$$GP_n = P_n + iP_{n-1},$$

$$GR_n = R_n + iR_{n-1}.$$

where $P_n = P_{n-2} + P_{n-3}$, $P_0 = 1$, $P_1 = 1$, $P_2 = 1$, and $R_n = 2R_{n-2} + R_{n-3}$, $R_0 = 1$, $R_1 = 1$, $R_2 = 1$.

- Cerda-Morales [3] defined Gaussian third-order Jacobsthal numbers as

$$GJ_n = J_n + iJ_{n-1}.$$

where $J_n = J_{n-1} + J_{n-2} + 2J_{n-3}$, $J_1 = 0$, $J_2 = 1$, $J_3 = 1$.

- Yılmaz and Soykan [26] presented Gaussian Guglielmo and Guglielmo-Lucas numbers by written respectively,

$$GT_n = T_n + iT_{n-1},$$

$$GH_n = H_n + iH_{n-1}.$$

where $T_n = 3T_{n-1} - 3T_{n-2} + T_{n-3}$, $T_0 = 0$, $T_1 = 1$, $T_2 = 3$, and $H_n = 3H_{n-1} - 3H_{n-2} + H_{n-3}$, $H_0 = 3$, $H_1 = 3$, $H_2 = 3$.

- Dikmen [4] presented Gaussian Leonardo and Leonardo-Lucas numbers by written respectively,

$$Gl_n = l_n + il_{n-1},$$

$$GH_n = H_n + iH_{n-1}.$$

where $l_n = 2l_{n-1} - l_{n-3}$, $l_0 = 1$, $l_1 = 1$, $l_2 = 3$, and $H_n = 2H_{n-1} - H_{n-3}$, $H_0 = 3$, $H_1 = 2$, $H_2 = 4$.

- Ayrılma and Soykan [2] presented Gaussian Edouard and Edouard-Lucas numbers by written respectively,

$$GE_n = E_n + iE_{n-1},$$

$$GK_n = K_n + iK_{n-1}.$$

where $E_n = 7E_{n-1} - 7E_{n-2} + E_{n-3}$, $E_0 = 0$, $E_1 = 1$, $E_2 = 7$, and $K_n = 7K_{n-1} - 7K_{n-2} + K_{n-3}$, $K_0 = 3$, $K_1 = 7$, $K_2 = 35$.

- Soykan et al [21] presented Gaussian Bigollo and Bigollo-Lucas numbers by written respectively,

$$GB_n = B_n + iB_{n-1},$$

$$GC_n = C_n + iC_{n-1}.$$

where $B_n = 4B_{n-1} - 5B_{n-2} + 2B_{n-3}$, $B_0 = 0$, $B_1 = 1$, $B_2 = 4$, and $C_n = 4C_{n-1} - 5C_{n-2} + 2C_{n-3}$, $C_0 = 3$, $C_1 = 4$, $C_2 = 6$.

Next, we give the exponential generating function of $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ of the sequence W_n .

LEMMA 1.4. Suppose that $f_{GW_n}(x) = \sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ is the exponential generating function of the generalized Pandita sequence $\{W_n\}$.

Then $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ is given by

$$\begin{aligned} \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} e^{\alpha x} \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} e^{\beta x} \\ &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} e^{\gamma x} \\ &+ (-W_3 + W_2 + W_0)e^x. \end{aligned}$$

Proof: Using the Binet's formula of generating Pandita numbers we get

$$\begin{aligned} \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} \left(\frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)\alpha^n}{3\alpha - 2} \right. \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)\beta^n}{3\beta - 2} \\ &+ \left. \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)\gamma^n}{3\gamma - 2} - W_3 + W_2 + W_0 \right) \frac{x^n}{n!} \\ &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} \sum_{n=0}^{\infty} \alpha^n \frac{x^n}{n!} \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} \sum_{n=0}^{\infty} \beta^n \frac{x^n}{n!} \\ &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} \sum_{n=0}^{\infty} \gamma^n \frac{x^n}{n!} \\ &+ (-W_3 + W_2 + W_0) \sum_{n=0}^{\infty} \frac{x^n}{n!} \end{aligned}$$

$$\begin{aligned}
 &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} e^{\alpha x} \\
 &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} e^{\beta x} \\
 &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} e^{\gamma x} + (-W_3 + W_2 + W_0)e^x. \quad \square
 \end{aligned}$$

The previous Lemma 1.4 gives the following results as particular examples.

COROLLARY 1.5. *Exponential generating function of Pandita and Pandita-Lucas numbers*

$$\begin{aligned}
 \text{a): } \sum_{n=0}^{\infty} P_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} \left(\frac{\alpha^{n+3}}{3\alpha - 2} + \frac{\beta^{n+3}}{3\beta - 2} + \frac{\gamma^{n+3}}{3\gamma - 2} - 1 \right) \frac{x^n}{n!} = \frac{\alpha^3 e^{\alpha x}}{3\alpha - 2} + \frac{\beta^3 e^{\beta x}}{3\beta - 2} + \frac{\gamma^3 e^{\gamma x}}{3\gamma - 2} - e^x. \\
 \text{b): } \sum_{n=0}^{\infty} S_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} (\alpha^n + \beta^n + \gamma^n + 1) \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^{\gamma x} + e^x.
 \end{aligned}$$

2. Gaussian Generalized Pandita Numbers

This section introduces the Gaussian generalized Pandita numbers and explores key properties, including Binet's formula and their generating function.

Gaussian generalized Pandita numbers $\{GW_n\}_{n \geq 0} = \{GW_n(GW_0, GW_1, GW_2, GW_3)\}_{n \geq 0}$ are defined by

$$GW_n = 2GW_{n-1} - GW_{n-2} + GW_{n-3} - GW_{n-4}. \quad (2.1)$$

with the initial conditions

$$\begin{aligned}
 GW_0 &= W_0 + i(W_0 - W_1 + 2W_2 - W_3), \\
 GW_1 &= W_1 + iW_0, \\
 GW_2 &= W_2 + iW_1, \\
 GW_3 &= W_3 + iW_2.
 \end{aligned}$$

not all being zero. The sequences $\{GW_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$GW_{-n} = GW_{-(n-1)} - GW_{-(n-2)} + 2GW_{-(n-3)} - GW_{-(n-4)}. \quad (2.2)$$

for $n = 1, 2, 3, \dots$. Thus, recurrence (2.1) hold for all integer n . Note that for all integers n , we get

$$GW_n = W_n + iW_{n-1}, \quad (2.3)$$

and

$$GW_{-n} = W_{-n} + iW_{-n-1}. \quad (2.4)$$

The first few generalized Gaussian Pandita numbers with positive subscript and negative subscript are presented in the following table.

Table 1. The first few generalized Gaussian Pandita numbers with positive subscript

n	GW_n
0	$W_0 + i(W_0 - W_1 + 2W_2 - W_3)$
1	$W_1 + iW_0$
2	$W_2 + iW_1$
3	$W_3 + iW_2$
4	$W_1 - W_0 - W_2 + 2W_3 + iW_3$
5	$W_1 - 2W_0 - W_2 + 3W_3 + i(W_1 - W_0 - W_2 + 2W_3)$

and with a negative subscript shown in Table 2

Table 2. First few generalized Gaussian Pandita numbers with negative subscript

n	GW_{-n}
0	$W_0 + i(W_0 - W_1 + 2W_2 - W_3)$
1	$W_0 - W_1 + 2W_2 - W_3 + i(W_1 + W_2 - W_3)$
2	$W_1 + W_2 - W_3 + i(W_0 + W_1 - W_2)$
3	$W_0 + W_1 - W_2 + i(2W_0 - 2W_1 + 2W_2 - W_3)$
4	$2W_0 - 2W_1 + 2W_2 - W_3 + i(3W_2 - 2W_3)$
5	$3W_2 - 2W_3 + i(3W_1 - 2W_2)$

We can define two special cases of GW_n : $GW_n(0, 1, 2 + i, 3 + 2i) = GP_n$ is the sequence of Gaussian Pandita numbers , $GW_n(4 + i, 2 + 4i, 2 + 2i, 5 + 2i) = GS_n$ is the sequence of Gaussian Pandita-Lucas numbers.

So Gaussian Pandita numbers are defined by

$$GP_n = 2P_{n-1} - P_{n-2} + P_{n-3} - P_{n-4}. \quad (2.5)$$

with the initial conditions

$$GP_0 = 0, GP_1 = 1, GP_2 = 2 + i, GP_3 = 3 + 2i.$$

Gaussian Pandita-Lucas numbers are defined by

$$GS_n = 2S_{n-1} - S_{n-2} + S_{n-3} - S_{n-4}. \quad (2.6)$$

with the initial conditions

$$GS_0 = 4 + i, GS_1 = 2 + 4i, GS_2 = 2 + 2i, GS_3 = 5 + 2i.$$

That for all integer we have

$$GP_n = P_n + iP_{n-1},$$

$$GS_n = S_n + iS_{n-1}.$$

The initial values of the Gaussian Pandita and Gaussian Pandita-Lucas numbers, for both positive and negative subscripts, are presented in Table 3.

Table 3. Gaussian Pandita numbers, Gaussian Pandita-Lucas numbers, with positive and negative subscripts, special cases of generalized Pandita numbers.

n	0	1	2	3	4	5	6
GP_n	0	1	$2+i$	$3+2i$	$5+3i$	$8+5i$	$12+8i$
GP_{-n}	0	0	$-i$	$-1-i$	-1	$-i$	$-1-2i$
GS_n	$4+i$	$2+4i$	$2+2i$	$5+2i$	$6+5i$	$7+6i$	$11+7i$
GS_{-n}	$4+i$	$1-i$	$-1+4i$	$4+3i$	$3-4i$	$-4+2i$	$2+8i$

Next, we present the Binet's formula for the Gaussian generalized Pandita numbers.

THEOREM 2.1. *The Binet's formula for the Gaussian generalized Pandita numbers is*

$$\begin{aligned}
 GW_n = & \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)\alpha^n}{3\alpha - 2} \\
 & + \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)\beta^n}{3\beta - 2} \\
 & + \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)\gamma^n}{3\gamma - 2} - W_3 + W_2 + W_0 \\
 & + i \left(\frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)\alpha^{n-1}}{3\alpha - 2} \right. \\
 & + \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)\beta^{n-1}}{3\beta - 2} \\
 & \left. + \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)\gamma^{n-1}}{3\gamma - 2} - W_3 + W_2 + W_0 \right).
 \end{aligned}$$

Proof. The proof follows from (1.1) and (2.3). $\square$

The following results are immediate consequences of the preceding Theorem.

COROLLARY 2.2. *For all integers n , we have following identities:*

$$\begin{aligned}
 \text{(a): } GP_n &= \frac{\alpha^{n+3}}{3\alpha - 2} + \frac{\beta^{n+3}}{3\beta - 2} + \frac{\gamma^{n+3}}{3\gamma - 2} - 1 + i \left(\frac{\alpha^{n+2}}{3\alpha - 2} + \frac{\beta^{n+2}}{3\beta - 2} + \frac{\gamma^{n+2}}{3\gamma - 2} - 1 \right). \\
 \text{(b): } GS_n &= \alpha^n + \beta^n + \gamma^n + 1 + i(\alpha^{n-1} + \beta^{n-1} + \gamma^{n-1} + 1).
 \end{aligned}$$

The next Theorem presents the generating function of Gaussian generalized Pandita numbers.

THEOREM 2.3. *Let $f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n x^n$ denote the generating function of Gaussian generalized Pandita numbers is given as follows:*

$$f_{GW_n}(z) = \sum_{n=0}^{\infty} GW_n x^n = \frac{1}{1-2x+x^2-x^3+x^4} GW_0 + (GW_1 - 2GW_0)x + (GW_2 - 2GW_1 + GW_0)x^2 + (GW_3 - 2GW_2 + GW_1 - GW_0)x^3.$$

Proof. Using the definition of Gaussian Pandita numbers, and subtracting $xf(x)$, $x^2f(x)$ and $x^3f(x)$ from $f(x)$ we obtain $(1 - 2x + x^2 - x^3 + x^4)f_{GW_n}(x)$

$$\begin{aligned}
 & (1 - 2x + x^2 - x^3 + x^4)f_{GW_n}(x) \\
 = & \sum_{n=0}^{\infty} GW_n x^n - 2x \sum_{n=0}^{\infty} GW_n x^n + x^2 \sum_{n=0}^{\infty} GW_n x^n - x^3 \sum_{n=0}^{\infty} GW_n x^n + x^4 \sum_{n=0}^{\infty} GW_n x^n, \\
 = & \sum_{n=0}^{\infty} GW_n x^n - 2 \sum_{n=0}^{\infty} GW_n x^{n+1} + \sum_{n=0}^{\infty} GW_n x^{n+2} - \sum_{n=0}^{\infty} GW_n x^{n+3} + \sum_{n=0}^{\infty} GW_n x^{n+4}, \\
 = & \sum_{n=0}^{\infty} GW_n x^n - 2 \sum_{n=1}^{\infty} GW_{(n-1)} x^n + \sum_{n=2}^{\infty} GW_{(n-2)} x^n - \sum_{n=3}^{\infty} GW_{(n-3)} x^n + \sum_{n=4}^{\infty} GW_{(n-4)} x^n, \\
 = & (GW_0 + GW_1 x + GW_2 x^2 + GW_3 x^3) - 2(GW_0 x + GW_1 x^2 + GW_2 x^3) + (GW_0 x^2 + GW_1 x^3) - GW_0 x^3 \\
 & + \sum_{n=4}^{\infty} (GW_n - 2GW_{n-1} - GW_{n-2} - GW_{n-3} + GW_{n-4}) x^n, \\
 = & GW_0 + (GW_1 - 2GW_0)x + (GW_2 - 2GW_1 + GW_0)x^2 + (GW_3 - 2GW_2 + GW_1 - GW_0)x^3.
 \end{aligned}$$

And rearranging above equation, we get (2.3). $\square$

The following results are immediate consequences of the preceding Theorem.

COROLLARY 2.4. *For all integers n , we have following identities:*

$$\begin{aligned}
 \text{(a): } f_{GP_n}(z) &= \sum_{n=0}^{\infty} GP_n x^n = \frac{ix^2 + x}{x^4 - x^3 + x^2 - 2x + 1}. \\
 \text{(b): } f_{GS_n}(z) &= \sum_{n=0}^{\infty} GS_n x^n = -\frac{(1-i)x^3 - (2-5i)x^2 + (6-2i)x - 4 - i}{x^4 - x^3 + x^2 - 2x + 1}.
 \end{aligned}$$

Theorem (2.3) gives the following results as special cases,

$$\begin{aligned}
 (1 - 2x + x^2 - x^3 + x^4)f_{GP_n}(x) &= GP_0 + (GP_1 - 2GP_0)x + (GP_2 - 2GP_1 + GP_0)x^2 + (GP_3 - 2GP_2 + \\
 GP_1 - GP_0)x^3 &= ix^2 + x, \\
 (1 - 2x + x^2 - x^3 + x^4)f_{GS_n}(x) &= GS_0 + (GS_1 - 2GS_0)x + (GS_2 - 2GS_1 + GS_0)x^2 + (GS_3 - 2GS_2 + \\
 GS_1 - GS_0)x^3 &= -(1-i)x^3 + (2-5i)x^2 - (6-2i)x + 4 + i
 \end{aligned}$$

Next, we give the exponential Gaussian generating function of $\sum_{n=0}^{\infty} GW_n x^n$ of the sequence GW_n .

LEMMA 2.5. *Suppose that $f_{GW_n}(x) = \sum_{n=0}^{\infty} GW_n \frac{x^n}{n!}$ is the exponential Gaussian generating function of the generalized Pandita sequence $\{GW_n\}$.*

Then $\sum_{n=0}^{\infty} GW_n \frac{x^n}{n!}$ is given by

$$\begin{aligned}
\sum_{n=0}^{\infty} GW_n \frac{x^n}{n!} &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} e^{\alpha x} \\
&+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} e^{\beta x} \\
&+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} e^{\gamma x} + (-W_3 + W_2 + W_0)e^x \\
&+ i\left(\frac{1}{\alpha} \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{3\alpha - 2} e^{\alpha x} \right. \\
&+ \frac{1}{\beta} \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{3\beta - 2} e^{\beta x} \\
&\left. + \frac{1}{\gamma} \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{3\gamma - 2} e^{\gamma x} + (-W_3 + W_2 + W_0)e^x\right).
\end{aligned}$$

Proof. The proof follows from the Binet's formula of GW_n and $GW_n = GW_n + iGW_{n-1}$ (Lemma1.4). $\square$

The previous Lemma 2.5 gives the following results as particular examples.

COROLLARY 2.6. *Exponential Gaussian generating function of Pandita and Pandita-Lucas numbers*

$$\begin{aligned}
\mathbf{a):} \quad \sum_{n=0}^{\infty} GP_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} \left(\frac{\alpha^{n+3}}{3\alpha-2} + \frac{\beta^{n+3}}{3\beta-2} + \frac{\gamma^{n+3}}{3\gamma-2} - 1 + i\left(\frac{\alpha^{n+2}}{3\alpha-2} + \frac{\beta^{n+2}}{3\beta-2} + \frac{\gamma^{n+2}}{3\gamma-2} - 1\right) \right) \frac{x^n}{n!} = \frac{\alpha^3 e^{\alpha x}}{3\alpha-2} + \frac{\beta^3 e^{\beta x}}{3\beta-2} + \\
&\frac{\gamma^3 e^{\gamma x}}{3\gamma-2} - e^x + i\left(\frac{\alpha^2 e^{\alpha x}}{3\alpha-2} + \frac{\beta^2 e^{\beta x}}{3\beta-2} + \frac{\gamma^2 e^{\gamma x}}{3\gamma-2} - e^x\right). \\
\mathbf{b):} \quad \sum_{n=0}^{\infty} GS_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} (\alpha^n + \beta^n + \gamma^n + 1 + i(\alpha^{n-1} + \beta^{n-1} + \gamma^{n-1} + 1)) \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^{\gamma x} + e^x + \\
&i\left(\frac{1}{\alpha} e^{\alpha x} + \frac{1}{\beta} e^{\beta x} + \frac{1}{\gamma} e^{\gamma x} + e^x\right).
\end{aligned}$$

3. Obtaining Binet Formula From Generating Function

We next find Binet's formula generalized Gaussian Pandita number $\{GW_n\}$ by the use of generating function for GW_n .

THEOREM 3.1. *Binet's formula of generalized Gaussian Pandita numbers)*

$$GW_n = \frac{q_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{q_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{q_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{q_4 \delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}. \quad (3.1)$$

where

$$\begin{aligned}
q_1 &= W_0 \alpha^3 + (W_1 - 2W_0) \alpha^2 + (W_0 - 2W_1 + W_2) \alpha - W_0 + W_1 - 2W_2 + W_3, \\
q_2 &= W_0 \beta^3 + (W_1 - 2W_0) \beta^2 + (W_0 - 2W_1 + W_2) \beta - W_0 + W_1 - 2W_2 + W_3, \\
q_3 &= W_0 \gamma^3 + (W_1 - 2W_0) \gamma^2 + (W_0 - 2W_1 + W_2) \gamma - W_0 + W_1 - 2W_2 + W_3, \\
q_4 &= W_0 \delta^3 + (W_1 - 2W_0) \delta^2 + (W_0 - 2W_1 + W_2) \delta - W_0 + W_1 - 2W_2 + W_3.
\end{aligned}$$

Proof. Let

$$h(x) = x^4 - x^3 + x^2 - 2x + 1.$$

Then for some α, β, γ and δ we write

$$h(x) = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x).$$

i.e.,

$$x^4 - x^3 + x^2 - 2x + 1 = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x). \quad (3.2)$$

Hence $\frac{1}{\alpha}, \frac{1}{\beta}, \frac{1}{\gamma}$ and $\frac{1}{\delta}$ are the roots of $h(x)$. This gives α, β, γ and δ as the roots of

$$h\left(\frac{1}{x}\right) = \frac{1}{x^2} - \frac{2}{x} - \frac{1}{x^3} + \frac{1}{x^4} + 1 = 0.$$

This implies $x^4 - x^3 + x^2 - 2x + 1 = 0$. Now, by it follows that

$$\sum_{n=0}^{\infty} GW_n x^n = \frac{(GW_1 - GW_0 - 2GW_2 + GW_3)x^3 + (GW_0 - 2GW_1 + GW_2)x^2 + (GW_1 - 2GW_0)x + GW_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)}.$$

Then we write

$$\begin{aligned} \frac{(W_1 - W_0 - 2W_2 + W_3)x^3 + (W_0 - 2W_1 + W_2)x^2 + (W_1 - 2W_0)x + W_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)} &= \frac{B_1}{(1 - \alpha x)} + \frac{B_2}{(1 - \beta x)} \\ &+ \frac{B_3}{(1 - \gamma x)} + \frac{B_4}{(1 - \delta x)}. \end{aligned} \quad (3.3)$$

So

$$\begin{aligned} &(W_1 - W_0 - 2W_2 + W_3)x^3 + (W_0 - 2W_1 + W_2)x^2 + (W_1 - 2W_0)x + W_0 \\ &= B_1(1 - \beta x)(1 - \gamma x)(1 - \delta x) + B_2(1 - \alpha x)(1 - \gamma x)(1 - \delta x) \\ &+ B_3(1 - \alpha x)(1 - \beta x)(1 - \delta x) + B_4(1 - \alpha x)(1 - \beta x)(1 - \gamma x). \end{aligned}$$

If we consider $x = \frac{1}{\alpha}$, we get $W_0 + \frac{1}{\alpha^2}(W_0 - 2W_1 + W_2) - \frac{1}{\alpha^3}(W_0 - W_1 + 2W_2 - W_3) + \frac{1}{\alpha}(W_1 - 2W_0) = -B_1\left(\frac{1}{\alpha}\beta - 1\right)\left(\frac{1}{\alpha}\gamma - 1\right)\left(\frac{1}{\alpha}\delta - 1\right)$.

This gives

$$\begin{aligned} B_1 &= \alpha^3(GW_0 + \frac{1}{\alpha^2}(GW_0 - 2GW_1 + GW_2) + \frac{1}{\alpha^3}(GW_1 - 5GW_0 - 4GW_2 + GW_3) + \frac{1}{\alpha}(GW_1 - 2GW_0)) \\ &= \frac{GW_0\alpha^3 + (GW_1 - 2GW_0)\alpha^2 + (GW_0 - 2GW_1 + GW_2)\alpha - GW_0 + GW_1 - 2GW_2 + GW_3}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} B_2 &= \frac{GW_0\beta^3 + (GW_1 - 2GW_0)\beta^2 + (GW_0 - 2GW_1 + GW_2)\beta - GW_0 + GW_1 - 2GW_2 + GW_3}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)}, \\ B_3 &= \frac{GW_0\gamma^3 + (GW_1 - 2GW_0)\gamma^2 + (GW_0 - 2GW_1 + GW_2)\gamma - GW_0 + GW_1 - 2GW_2 + GW_3}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)}, \\ B_4 &= \frac{GW_0\delta^3 + (GW_1 - 2GW_0)\delta^2 + (GW_0 - 2GW_1 + GW_2)\delta - GW_0 + GW_1 - 2GW_2 + GW_3}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}. \end{aligned}$$

Thus (3.3) can be written as

$$\sum_{n=0}^{\infty} GW_n x^n = B_1(1 - \alpha x)^{-1} + B_2(1 - \beta x)^{-1} + B_3(1 - \gamma x)^{-1} + B_4(1 - \delta x)^{-1}.$$

This gives

$$\sum_{n=0}^{\infty} GW_n x^n = B_1 \sum_{n=0}^{\infty} \alpha^n x^n + B_2 \sum_{n=0}^{\infty} \beta^n x^n + B_3 \sum_{n=0}^{\infty} \gamma^n x^n + B_4 \sum_{n=0}^{\infty} \delta^n x^n = \sum_{n=0}^{\infty} (B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n) x^n.$$

Therefore, comparing coefficients on both sides of the above equality, we obtain

$$GW_n = B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n.$$

and then we get (3.1). $\square$

4. Some Identities About Recurrence Relations of Gaussian Generalized Pandita Numbers

In this section, we present some identities on Gaussian Pandita, Gaussian Pandita-Lucas,

THEOREM 4.1. *The following equations hold for all integer n*

$$\begin{aligned} GP_n &= \frac{54}{31} GS_{n+3} - \frac{41}{31} GS_{n+2} + \frac{6}{31} GS_{n+1} - \frac{50}{31} GS_n, \\ GS_n &= -GP_{n+3} + 3GP_{n+2} + GP_{n+1} - 4GP_n. \end{aligned} \tag{4.1}$$

Proof. To proof identity (4.1), we can write

$$GP_n = aGS_{n+3} + bGS_{n+2} + cGS_{n+1} + dGS_n.$$

Solving the system of equations

$$\begin{aligned} GP_0 &= aGS_3 + bGS_2 + cGS_1 + dGS_0, \\ GP_1 &= aGS_4 + bGS_3 + cGS_2 + dGS_1, \\ GP_2 &= aGS_5 + bGS_4 + cGS_3 + dGS_2, \\ GP_3 &= aGS_6 + bGS_5 + cGS_4 + dGS_3. \end{aligned}$$

we get $a = \frac{54}{31}, b = -\frac{41}{31}, c = \frac{6}{31}, d = -\frac{50}{31}$.

The other identities can be found similarly.

$$GS_n = aGP_{n+3} + bGP_{n+2} + cGP_{n+1} + dGP_n.$$

$$\begin{aligned} GS_0 &= aGP_3 + bGP_2 + cGP_1 + dGP_0, \\ GS_1 &= aGP_4 + bGP_3 + cGP_2 + dGP_1, \\ GS_2 &= aGP_5 + bGP_4 + cGP_3 + dGP_2, \\ GS_3 &= aGP_6 + bGP_5 + cGP_4 + dGP_3. \end{aligned}$$

we get $a = -1, b = 3, c = 1, d = -4$.

LEMMA 4.2. 6, Let's assume that $f(x) = \sum_{n=0}^{\infty} a_n x^n$ is the generating function of the sequence $\{a_n\}_{n \geq 0}$. Then the generating functions of the sequences $\{a_{2n}\}_{n \geq 0}$ and $\{a_{2n+1}\}_{n \geq 0}$ are stated as

$$f_{a_{2n}}(x) = \sum_{n=0}^{\infty} a_{2n} x^n = \frac{f(\sqrt{x}) + f(-\sqrt{x})}{2},$$

and

$$f_{a_{2n+1}}(x) = \sum_{n=0}^{\infty} a_{2n+1} x^n = \frac{f(\sqrt{x}) - f(-\sqrt{x})}{2\sqrt{x}}.$$

respectively.

The generating functions of the even and odd-indexed Gaussian generalized Pandita sequences are provided by the following theorem.

THEOREM 4.3. The generating functions of the sequence GW_{2n} and GW_{2n+1} are provided by

$$f_{GW_{2n}}(x) = \frac{GW_2(x^3 + 3x^2 - x) + GW_0(2x^2 + 2x - 1) - GW_1(x^2 - x^3) - GW_3(x^3 + 2x^2)}{-x^4 - x^3 + x^2 + 2x - 1}, \quad (4.2)$$

$$f_{GW_{2n+1}}(x) = \frac{GW_0(x^3 + 2x^2) - GW_3(x^3 + x^2 + x) - GW_1(x^3 - 2x + 1) + GW_2(2x^3 + x^2)}{-x^4 - x^3 + x^2 + 2x - 1}. \quad (4.3)$$

Proof. We only proof (4.2). From Theorem (2.3) we can obtain following identities.

$$\begin{aligned} f_{GW_n}(\sqrt{x}) &= \frac{1}{\sqrt{x^3 - x^2 + x + 2\sqrt{x} - 1}} ((GW_1 + GW_2 - GW_3)\sqrt{x^3} + (2GW_0 + GW_2 - GW_3)x \\ &\quad + (-GW_2 + 2GW_0)\sqrt{x} - GW_0), \\ f_{GW_n}(-\sqrt{x}) &= -\frac{1}{-\sqrt{x^3 - x^2 + x + 2\sqrt{x} - 1}} ((GW_0 - GW_1 + 2GW_2 - GW_3)\sqrt{x^3} \\ &\quad + (-GW_3 - 3GW_2 + 2GW_1 + 2GW_0)x + (2GW_1 - GW_3)\sqrt{x} - GW_1). \end{aligned}$$

Thus, the result follows from Lemma (4.2). and the other identity can be derived analogously. $\square$

From Theorem (4.3), we get the following Corollary.

$$\begin{aligned} f_{GP_{2n}}(x) &= \frac{ix^3 + (1+i)x^2 + (2+i)x}{x^4 + x^3 - x^2 - 2x + 1}, \\ f_{GP_{2n+1}}(x) &= \frac{(1+i)x^2 + (1+2i)x + 1}{x^4 + x^3 - x^2 - 2x + 1}, \\ f_{GS_{2n}}(x) &= \frac{(1-4i)x^3 - 2x^2 - 6x + 4 + i}{x^4 + x^3 - x^2 - 2x + 1}, \\ f_{GS_{2n+1}}(x) &= \frac{(-1+i)x^3 + (-5-2i)x^2 + (1-6i)x + 2 + 4i}{x^4 + x^3 - x^2 - 2x + 1}. \end{aligned}$$

From Corollary 4 we can obtain the following Corollary which presents the identities on Gaussian Pandita sequences.

COROLLARY 4.4. **a):** $(2+i)GS_{2n-2} + (1+i)GS_{2n-4} + iGS_{2n-6} = (4+i)GP_{2n} + (-6)GP_{2n-2} + (-2)GP_{2n-4} + (1-4i)GP_{2n-6}.$

$$\mathbf{b):} \quad GS_{2n} + (1 + 2i)GS_{2n-2} + (1 + i)GS_{2n-4} = (4 + i)GP_{2n+1} + (-6)GP_{2n-1} + (-2)GP_{2n-3} + (1 - 4i)GP_{2n-5}.$$

$$\mathbf{c):} \quad (2+4i)GS_{2n}+(1-6i)GS_{2n-2}+(-5-2i)GS_{2n-4}+(-1+i)GS_{2n-6} = (4+i)GS_{2n+1}+(-6)GS_{2n-1}+(-2)GS_{2n-3}+(1-4i)GS_{2n-5}.$$

$$\mathbf{d):} \quad GS_{2n+1}+(1+2i)GS_{2n-1}+(1+i)GS_{2n-3} = (2+4i)GP_{2n+1}+(1-6i)GP_{2n-1}+(-5-2i)GP_{2n-3}+(-1+i)GP_{2n-5}.$$

$$\mathbf{e):} \quad (2+i)GS_{2n-1}+(1+i)GS_{2n-3}+iGS_{2n-5} = (2+4i)GP_{2n}+(1-6i)GP_{2n-2}+(-5-2i)GP_{2n-4}+(-1+i)GP_{2n-6}.$$

$$\mathbf{f):} \quad GP_{2n} + (1 + 2i)GP_{2n-2} + (1 + i)GP_{2n-4} = (2 + i)GP_{2n-1} + (1 + i)GP_{2n-3} + iGP_{2n-5}.$$

Proof. From corollary (4) we obtain

$$(ix^3 + (1 + i)x^2 + (2 + i)x)f_{GS_{2n}}(x) = ((1 - 4i)x^3 - 2x^2 - 6x + 4 + i)f_{GP_{2n}}(x).$$

LHS is equal to

$$\begin{aligned} LHS &= (ix^3 + (1 + i)x^2 + (2 + i)x) \sum_{n=0}^{\infty} GS_{2n}x^n, \\ &= (2 + i)x \sum_{n=0}^{\infty} GS_{2n}x^n + (1 + i)x^2 \sum_{n=0}^{\infty} GS_{2n}x^n + ix^3 \sum_{n=0}^{\infty} GS_{2n}x^n \\ &= (2 + i) \sum_{n=0}^{\infty} GS_{2n}x^{n+1} + (1 + i) \sum_{n=0}^{\infty} GS_{2n}x^{n+2} + i \sum_{n=0}^{\infty} GS_{2n}x^{n+3} \\ &= (2 + i) \sum_{n=1}^{\infty} GS_{2n-2}x^n + (1 + i) \sum_{n=2}^{\infty} GS_{2n-4}x^n + i \sum_{n=3}^{\infty} GS_{2n-6}x^n, \\ &= (7 + 6i)x + (5 + 11i)x^2 + \sum_{n=2}^{\infty} ((2 + i)GS_{2n-2} + (1 + i)GS_{2n-4} + iGS_{2n-6})x^n. \end{aligned}$$

Whereas the RHS is equal to

$$\begin{aligned} RHS &= ((1 - 4i)x^3 - 2x^2 - 6x + (4 + i)) \sum_{n=0}^{\infty} GP_{2n}x^n, \\ &= (4 + i) \sum_{n=0}^{\infty} GP_{2n}x^n - 6x \sum_{n=0}^{\infty} GP_{2n}x^n - 2x^2 \sum_{n=0}^{\infty} GP_{2n}x^n + (1 - 4i)x^3 \sum_{n=0}^{\infty} GP_{2n}x^n \\ &= (4 + i) \sum_{n=0}^{\infty} GP_{2n}x^n + (-6) \sum_{n=0}^{\infty} GP_{2n}x^{n+1} + (-2) \sum_{n=0}^{\infty} GP_{2n}x^{n+2} + (1 - 4i) \sum_{n=0}^{\infty} GP_{2n}x^{n+3} \\ &= (4 + i) \sum_{n=0}^{\infty} GP_{2n}x^n + (-6) \sum_{n=1}^{\infty} GP_{2n-2}x^n + (-2) \sum_{n=2}^{\infty} GP_{2n-4}x^n + (1 - 4i) \sum_{n=3}^{\infty} GP_{2n-6}x^n \\ &= (7 + 6i)x + (5 + 11i)x^2 + \sum_{n=3}^{\infty} ((4 + i)GP_{2n} + (-6)GP_{2n-2} + (-2)GP_{2n-4} + (1 - 4i)GP_{2n-6}) \end{aligned}$$

By comparing the coefficients, the proof of the first identity (a) is done. We can prove other identity similarly. $\square$

The following identity establishes a relationship between the Gaussian Pandita numbers and the Pandita–Lucas numbers.

COROLLARY 4.5. *For all integers m, n the following identities holds:*

$$GW_{m+n} = P_{m-2}GW_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})GW_{n+2} + (P_{m-3} - P_{m-4})GW_{n+1} - GW_n P_{m-3}.$$

Proof. First we assume that $m, n \geq 0$. The Theorem (4.5) can be proved by mathematical induction on m . If $m = 0$ we get

$$GW_n = P_{-2}GW_{n+3} + (P_{-4} - P_{-3} - P_{-5})GW_{n+2} + (P_{-3} - P_{-4})GW_{n+1} - GW_n P_{-3}.$$

which is true since $P_{-2} = 0, P_{-1} = -1, P_{-4} = -1, P_{-5} = 0$. Assume that the equality holds for $m \leq k$. For $m = k + 1$, we get

$$\begin{aligned} GW_{k+1+n} &= 2GW_{n+k} - GW_{n+k-1} + GW_{n+k-2} - GW_{n+k-3}, \\ &2(P_{m-2}GW_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})GW_{n+2} + (P_{m-3} - P_{m-4})GW_{n+1} - GW_n P_{m-3}) \\ &- (P_{m-3}GW_{n+3} + (P_{m-5} - P_{m-4} - P_{m-6})GW_{n+2} + (P_{m-4} - P_{m-5})GW_{n+1} - GW_n P_{m-4}) \\ &+ (P_{m-4}GW_{n+3} + (P_{m-6} - P_{m-5} - P_{m-7})GW_{n+2} + (P_{m-5} - P_{m-6})GW_{n+1} - GW_n P_{m-5}) \\ &- (P_{m-5}GW_{n+3} + (P_{m-7} - P_{m-6} - P_{m-8})GW_{n+2} + (P_{m-6} - P_{m-7})GW_{n+1} - GW_n P_{m-6}). \end{aligned}$$

Consequently, by mathematical induction on m , this proves Theorem 4.5.

The other cases of m, n can be proved similarly for all integers m, n . $\square$

Taking $GW_n = GP_n$ or $GW_n = GS_n$ in above Theorem, respectively, we get:

COROLLARY 4.6.

$$GP_{m+n} = P_{m-2}GP_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})GP_{n+2} + (P_{m-3} - P_{m-4})GP_{n+1} - GP_n P_{m-3},$$

$$GS_{m+n} = P_{m-2}GS_{n+3} + (P_{m-4} - P_{m-3} - P_{m-5})GS_{n+2} + (P_{m-3} - P_{m-4})GS_{n+1} - GS_n P_{m-3}.$$

5. SIMSON'S FORMULA

This section is devoted to the presentation of Simson's formula associated with the generalized Gaussian Pandita numbers. This is a special case of [14, Theorem 4.1].

THEOREM 5.1. *For all integers n , we can write the following equality:*

$$\begin{vmatrix} GW_{n+3} & GW_{n+2} & GW_{n+1} & GW_n \\ GW_{n+2} & GW_{n+1} & GW_n & GW_{n-1} \\ GW_{n+1} & GW_n & GW_{n-1} & GW_{n-2} \\ GW_n & GW_{n-1} & GW_{n-2} & GW_{n-3} \end{vmatrix} = \begin{vmatrix} GW_3 & GW_2 & GW_1 & GW_0 \\ GW_2 & GW_1 & GW_0 & GW_{-1} \\ GW_1 & GW_0 & GW_{-1} & GW_{-2} \\ GW_0 & GW_{-1} & GW_{-2} & GW_{-3} \end{vmatrix}$$

$$= (GW_3 - 2GW_2 + GW_0)(GW_3 - 2GW_1 + GW_0)(GW_3^2 - GW_2^2 + GW_1^2 - GW_0^2 - GW_2GW_3 - 2GW_1GW_3 + GW_1GW_2 + GW_0GW_3 + 2GW_0GW_2 - GW_0GW_1).$$

Proof. Using Theorem 2.1 it can be proved by using induction use [14, Theorem 4.1]

From the Theorem 5.1 we get the following Corollary.

COROLLARY 5.2. *For all integers n , the Simson's formulas of Pandita and Pandita Lucas numbers are given as respectively.*

$$\begin{aligned} \text{a): } & \begin{vmatrix} GP_{n+3} & GP_{n+2} & GP_{n+1} & GP_n \\ GP_{n+2} & GP_{n+1} & GP_n & GP_{n-1} \\ GP_{n+1} & GP_n & GP_{n-1} & GP_{n-2} \\ GP_n & GP_{n-1} & GP_{n-2} & GP_{n-3} \end{vmatrix} = 1 - i. \\ \text{b): } & \begin{vmatrix} GS_{n+3} & GS_{n+2} & GS_{n+1} & GS_n \\ GS_{n+2} & GS_{n+1} & GS_n & GS_{n-1} \\ GS_{n+1} & GS_n & GS_{n-1} & GS_{n-2} \\ GS_n & GS_{n-1} & GS_{n-2} & GS_{n-3} \end{vmatrix} = 31 + 31i. \end{aligned}$$

6. SUM FORMULAS

In this section, we identify some sum formulas of generalized Gaussian Pandita numbers.

THEOREM 6.1. *For all integers $n \geq 0$, we get sum formulas below*

$$\begin{aligned} \text{a) } & \sum_{k=0}^n GW_k = -(n+3)GW_{n+3} + (n+4)GW_{n+2} + (n+4)GW_n + 3GW_3 - 4GW_2 - 3GW_0. \\ \text{b) } & \sum_{k=0}^n GW_{2k} = \frac{1}{3}(-3(n+2)GW_{2n+2} + (3n+8)GW_{2n+1} + 2GW_{2n} + (3n+7)GW_{2n-1} + 7GW_3 - 8GW_2 - GW_1 - 6GW_0). \\ \text{c) } & \sum_{k=0}^n GW_{2k+1} = \frac{1}{3}(-(3n+4)GW_{2n+2} + (3n+8)GW_{2n+1} + GW_{2n} + 3(n+2)GW_{2n-1} + 6GW_3 - 8GW_2 + GW_1 - 7GW_0). \end{aligned}$$

Proof. It is given in Soykan [16, Theorem 3.12]. $\square$

As a special case of the theorem 6.1, we present the following Corollary.

COROLLARY 6.2. *For all integers $n \geq 0$, we get sum formulas below:*

$$\begin{aligned} \text{a) } & \sum_{k=0}^n GP_k = -(n+3)GP_{n+3} + (n+4)GP_{n+2} + (n+4)GP_n + 1 + 2i. \\ \text{b) } & \sum_{k=0}^n GP_{2k} = \frac{1}{3}(-3(n+2)GP_{2n+2} + (3n+8)GP_{2n+1} + 2GP_{2n} + (3n+7)GP_{2n-1} + 4 + 6i). \\ \text{c) } & \sum_{k=0}^n GP_{2k+1} = \frac{1}{3}(-(3n+4)GP_{2n+2} + (3n+8)GP_{2n+1} + GP_{2n} + 3(n+2)GP_{2n-1} + 3 + 4i). \end{aligned}$$

As a special case of the theorem 6.1, we present the following Corollary.

COROLLARY 6.3. *For all integers $n \geq 0$, we get sum formulas below:*

$$\begin{aligned} \text{a)} \sum_{k=0}^n GS_k &= -(n+3)GS_{n+3} + (n+4)GS_{n+2} + (n+4)GS_n - 5 - 5i. \\ \text{b)} \sum_{k=0}^n GS_{2k} &= \frac{1}{3}(-3(n+2)GS_{2n+2} + (3n+8)GS_{2n+1} + 2GS_{2n} + (3n+7)GS_{2n-1} - 7 - 12i). \\ \text{c)} \sum_{k=0}^n GS_{2k+1} &= \frac{1}{3}(-(3n+4)GS_{2n+2} + (3n+8)GS_{2n+1} + GS_{2n} + 3(n+2)GS_{2n-1} - 12 - 7i). \end{aligned}$$

Next, we give the ordinary generating functions of some special cases of Gaussian generalized Pandita numbers.

THEOREM 6.4. *The ordinary generating functions of the sequences W_{2n} , W_{2n+1} are given as follows:*

$$\begin{aligned} \text{a)} \sum_{n=0}^{\infty} GW_{2n}x^n &= \frac{GW_2(x^3 + 3x^2 - x) + GW_0(2x^2 + 2x - 1) - GW_1(x^2 - x^3) - GW_3(x^3 + 2x^2)}{-x^4 - x^3 + x^2 + 2x - 1}. \\ \text{b)} \sum_{n=0}^{\infty} GW_{2n+1}x^n &= \frac{GW_0(x^3 + 2x^2) - GW_3(x^3 + x^2 + x) - GW_1(x^3 - 2x + 1) + GW_2(2x^3 + x^2)}{-x^4 - x^3 + x^2 + 2x - 1}. \end{aligned}$$

From the last Theorem, we have the following Corollary which gives sum formula of Gaussian Pandita numbers (Take $W_n = GP_n$ whit $GP_0 = 0, GP_1 = 1, GP_2 = 2 + i, GP_3 = 3 + 2i$).

COROLLARY 6.5. *For $n \geq 0$ Gaussian Pandita numbers have the following properties.*

$$\begin{aligned} \text{a)} \sum_{n=0}^{\infty} GP_{2n}x^n &= \frac{ix^3 + (1+i)x^2 + (2+i)x}{x^4 + x^3 - x^2 - 2x + 1}. \\ \text{b)} \sum_{n=0}^{\infty} GP_{2n+1}x^n &= \frac{(1+i)x^2 + (1+2i)x + 1}{x^4 + x^3 - x^2 - 2x + 1}. \end{aligned}$$

7. Matrix Formulation of GW_n

In this section, we present the matrix representation of generalized Gaussian Pandita numbers

We define the square matrix A of order 4 as

$$A = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

such that $\det A = 1$. Note that

$$A^n = \begin{pmatrix} P_{n+1} & -P_n + P_{n-1} - P_{n-2} & P_n - P_{n-1} & -P_n \\ P_n & -P_{n-1} + P_{n-2} - P_{n-3} & P_{n-1} - P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-2} + P_{n-3} - P_{n-4} & P_{n-2} - P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-3} + P_{n-4} - P_{n-5} & P_{n-3} - P_{n-4} & -P_{n-3} \end{pmatrix}$$

for the proof see [19].

Then we present the following lemma.

For $n \geq 0$ the following identity is true:

$$\begin{pmatrix} GW_{n+3} \\ GW_{n+2} \\ GW_{n+1} \\ GW_n \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}.$$

Proof. The identity (7) can be proved by mathematical induction on n . If $n = 0$ we obtain

$$\begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}$$

which is true. We assume that the identity given holds for $n = k$, we deduce that the following identity is true

$$\begin{pmatrix} GW_{k+3} \\ GW_{k+2} \\ GW_{k+1} \\ GW_k \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix}.$$

For $n = k + 1$, we obtain

$$\begin{aligned} \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} &= \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} GW_3 \\ GW_2 \\ GW_1 \\ GW_0 \end{pmatrix} \\ &= \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} GW_{k+3} \\ GW_{k+2} \\ GW_{k+1} \\ GW_k \end{pmatrix} \\ &= \begin{pmatrix} GW_{k+4} \\ GW_{k+3} \\ GW_{k+2} \\ GW_{k+1} \end{pmatrix}. \end{aligned}$$

Consequently, by applying mathematical induction on n , the proof completed. $\square$

We define

$$N_{Gw} = \begin{pmatrix} GW_3 & GW_2 & GW_1 & GW_0 \\ GW_2 & GW_1 & GW_0 & GW_{-1} \\ GW_1 & GW_0 & GW_{-1} & GW_{-2} \\ GW_0 & GW_{-1} & GW_{-2} & GW_{-3} \end{pmatrix}, \quad (7.1)$$

$$E_{Gw} = \begin{pmatrix} GW_{n+3} & GW_{n+2} & GW_{n+1} & GW_n \\ GW_{n+2} & GW_{n+1} & GW_n & GW_{n-1} \\ GW_{n+1} & GW_n & GW_{n-1} & GW_{n-2} \\ GW_n & GW_{n-1} & GW_{n-2} & GW_{n-3} \end{pmatrix}. \quad (7.2)$$

Now, we have the following theorem with N_{Gw} and E_{Gw}

THEOREM 7.1. *Using N_{Gw} and E_{Gw} , we get*

$$A^n N_{Gw} = E_{Gw}.$$

Proof. Note that we get

$$\begin{aligned} A^n N_{Gw} &= \begin{pmatrix} P_{n+1} & -P_n + P_{n-1} - P_{n-2} & P_n - P_{n-1} & -P_n \\ P_n & -P_{n-1} + P_{n-2} - P_{n-3} & P_{n-1} - P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-2} + P_{n-3} - P_{n-4} & P_{n-2} - P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-3} + P_{n-4} - P_{n-5} & P_{n-3} - P_{n-4} & -P_{n-3} \end{pmatrix} \begin{pmatrix} GW_3 & GW_2 & GW_1 & GW_0 \\ GW_2 & GW_1 & GW_0 & GW_{-1} \\ GW_1 & GW_0 & GW_{-1} & GW_{-2} \\ GW_0 & GW_{-1} & GW_{-2} & GW_{-3} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix} \end{aligned}$$

where

$$\begin{aligned}
 a_{11} &= GW_1(P_n - P_{n-1}) - GW_2(P_n - P_{n-1} + P_{n-2}) - GW_0P_n + GW_3P_{n+1} = GW_3, \\
 a_{12} &= GW_0(P_n - P_{n-1}) - GW_1(P_n - P_{n-1} + P_{n-2}) - GP_nW_{-1} + GW_2P_{n+1} = GW_2, \\
 a_{13} &= GW_{-1}(P_n - P_{n-1}) - GW_0(P_n - P_{n-1} + P_{n-2}) - GP_nW_{-2} + GW_1P_{n+1} = GW_1, \\
 a_{14} &= GW_{-2}(P_n - P_{n-1}) - GW_{-1}(P_n - P_{n-1} + P_{n-2}) - GP_nW_{-3} + GW_0P_{n+1} = GW_0, \\
 a_{21} &= GW_3P_n - GW_2(P_{n-1} - P_{n-2} + P_{n-3}) + GW_1(P_{n-1} - P_{n-2}) - GW_0P_{n-1} = GW_2, \\
 a_{22} &= GW_2P_n - GW_{-1}P_{n-1} - GW_1(P_{n-1} - P_{n-2} + P_{n-3}) + GW_0(P_{n-1} - P_{n-2}) = GW_1, \\
 a_{23} &= G(P_{n-1} - P_{n-2})W_{-1} - GW_{-2}P_{n-1} + GW_1P_n - GW_0(P_{n-1} - P_{n-2} + P_{n-3}) = GW_0, \\
 a_{24} &= G(P_{n-1} - P_{n-2})W_{-2} - GW_{-3}P_{n-1} + GW_0P_n - GW_{-1}(P_{n-1} - P_{n-2} + P_{n-3}) = GW_{-1}, \\
 a_{31} &= GW_1(P_{n-2} - P_{n-3}) - GW_2(P_{n-2} - P_{n-3} + P_{n-4}) - GW_0P_{n-2} + GW_3P_{n-1} = GW_1, \\
 a_{32} &= GW_0(P_{n-2} - P_{n-3}) - GW_1(P_{n-2} - P_{n-3} + P_{n-4}) - GW_{-1}P_{n-2} + GW_2P_{n-1} = GW_0, \\
 a_{33} &= G(P_{n-2} - P_{n-3})W_{-1} - GW_{-2}P_{n-2} - GW_0(P_{n-2} - P_{n-3} + P_{n-4}) + GW_1P_{n-1} = GW_{-1}, \\
 a_{34} &= G(P_{n-2} - P_{n-3})W_{-2} - GW_{-3}P_{n-2} - GW_{-1}(P_{n-2} - P_{n-3} + P_{n-4}) + GW_0P_{n-1} = GW_{-2}, \\
 a_{41} &= GW_1(P_{n-3} - P_{n-4}) - GW_2(P_{n-3} - P_{n-4} + P_{n-5}) - GW_0P_{n-3} + GW_3P_{n-2} = GW_0, \\
 a_{42} &= GW_0(P_{n-3} - P_{n-4}) - GW_1(P_{n-3} - P_{n-4} + P_{n-5}) - GW_{-1}P_{n-3} + GW_2P_{n-2} = GW_{-1}, \\
 a_{43} &= G(P_{n-3} - P_{n-4})W_{-1} - GW_{-2}P_{n-3} - GW_0(P_{n-3} - P_{n-4} + P_{n-5}) + GW_1P_{n-2} = GW_{-2}, \\
 a_{44} &= G(P_{n-3} - P_{n-4})W_{-2} - GW_{-3}P_{n-3} - GW_{-1}(P_{n-3} - P_{n-4} + P_{n-5}) + GW_0P_{n-2} = GW_{-3}.
 \end{aligned}$$

Using the Theorem 4.5 the proof is done. $\square$

By taking $GW_n = GP_n$ with GP_0, GP_1, GP_2, GP_3 in 7.1 and 7.2,

and

$$GW_n = GS_n \text{ with } GS_0, GS_1, GS_2, GS_3 \text{ in 7.1 and 7.2.}$$

respectively, we get:

$$\begin{aligned}
 N_{GP} &= \begin{pmatrix} 3+2i & 2+i & 1 & 0 \\ 2+i & 1 & 0 & 0 \\ 1 & 0 & 0 & -i \\ 0 & 0 & -i & -1-i \end{pmatrix}, \quad E_{GO} = \begin{pmatrix} GP_{n+3} & GP_{n+2} & GP_{n+1} & GP_n \\ GP_{n+2} & GP_{n+1} & GP_n & GP_{n-1} \\ GP_{n+1} & GP_n & GP_{n-1} & GP_{n-2} \\ GP_n & GP_{n-1} & GP_{n-2} & GP_{n-3} \end{pmatrix}, \\
 N_{GS} &= \begin{pmatrix} 5+2i & 2+2i & 2+4i & 4+i \\ 2+2i & 2+4i & 4+i & 1-i \\ 2+4i & 4+i & 1-i & -1+4i \\ 4+i & 1-i & -1+4i & -4+3i \end{pmatrix}, \quad E_{GS} = \begin{pmatrix} GS_{n+3} & GS_{n+2} & GS_{n+1} & GS_n \\ GS_{n+2} & GS_{n+1} & GS_n & GS_{n-1} \\ GS_{n+1} & GS_n & GS_{n-1} & GS_{n-2} \\ GS_n & GS_{n-1} & GS_{n-2} & GS_{n-3} \end{pmatrix}.
 \end{aligned}$$

From Theorem 7.1, we can write the following corollary.

COROLLARY 7.2. *The following identities are hold:*

a): $A^n N_{GP} = E_{GP}.$

b): $A^n N_{GS} = E_{GS}.$

References

- [1] Aşçı M., Gürel E., Gaussian Jacobsthal and Gaussian Jacobsthal Lucas numbers, *Ars Combinatoria*, 111, 2013.
- [2] Ayrıлма, E.E., Soykan, Y., A Study On Gaussian Generalized Edouard Numbers, *Asian Journal of Advanced Research and Reports*, 19(5), 421-438, 2025.
- [3] Cerda-Morales, G., On Gauss third-order Jacobsthal numbers and their applications, , *Annals of the Alexandru Ioan Cuza University-Mathematics*, 67(2), 231-241, 2022.
- [4] Dikmen, C.M., Properties of Gaussian Generalized Leonardo Numbers, *Karaelmas Science and Engineering Journal*, 15(1), 134-145, 2025. DOI: 10.7212/karaelmasfen.1578154
- [5] Ertuş A., Yılmaz F., On Quaternions with Gaussian Oresme Coefficients, *Turk. J. Math. Comput. Sci.*, 15(1), 192-202, 2023.
- [6] Frontczak, R., Convolutions for Generalized Tribonacci Numbers and Related Results, *International Journal of Mathematical Analysis*, Vol. 12, 2018, no. 7, 307 - 324.
- [7] Hacı S., Öz S., On Some Gaussian Pell And Pell-Lucas Numbers, *Ordu Univ. J. Sci. Tech.*, 6(1), 8-18, 2016.
- [8] Horadam, A.F., Complex Fibonacci Numbers and Fibonacci Quaternions, *Amer. Math. Monthly*, 70 , 289-291, 1963.
- [9] Pethe, S., Horadam, A.F., Gaussian Generalized Fibonacci Numbers, *Bull. Austral. Math. Soc.*, 33 , 37-48., 1986.
- [10] Soykan Y., Generalized Pandita Numbers, *International Journal of Mathematics, Statistics and Operations Research*, 3(1), 107-123, 2023.
- [11] Soykan Y., Sum Formulas For Generalized Tetranacci Numbers: Closed Forms of the Sum Formulas $\sum_{k=0}^n x^k W_k$ and $\sum_{k=1}^n x^k W_{-k}$, *Journal of Progressive Research in Mathematics* 18(1), 24-26, 2021
- [12] Soykan, Y., A Study on Generalized Blaise Numbers, *Asian Journal of Advanced Research and Reports*, 17(1), 32-53, 2023. DOI: 10.9734/AJARR/2023/v17i1463
- [13] Soykan, Y., Generalized Friedrich Numbers, *Journal of Advances in Mathematics and Computer Science*, 38(3), 12-31, 2023. DOI: 10.9734/JAMCS/2023/v38i31748
- [14] Soykan, Y., Simson Identity of Generalized m-step Fibonacci Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 45-56, 2019.
- [15] Soykan, Y., Generalized Richard Numbers, *International Journal of Advances in Applied Mathematics and Mechanics*, 10(3), 38-51, 2023.
- [16] Soykan, Y., Sums and Generating Functions of Special Cases of Generalized Tetranacci Polynomials, *International Journal of Advances in Applied Mathematics and Mechanics*, 12(4), 34-101, 2025.
- [17] Soykan, Y., On Generalized 4-primes Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(4), 20-33, 2020, (ISSN: 2347-2529).
- [18] Soykan, Y., Generalized Olivier Numbers, *Asian Research Journal of Mathematics*, 19(1), Page 1-22, 2023. DOI: 10.9734/ARJOM/2023/v19i1634
- [19] Soykan, Y., Properties of Generalized (r,s,t,u)-Numbers, *Earthline Journal of Mathematical Sciences*, 5(2), 297-327, 2021. <https://doi.org/10.34198/ejms.5221.297327>
- [20] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, *Journal of Progressive Research in Mathematics*, 14(2) , 2018.
- [21] Soykan, Y., Okumuş, İ., Bilgin, N.G., On Generalized Bigollo Numbers, *Asian Research Journal of Mathematics*, 19(8), 72-88, 2023. DOI: 10.9734/ARJOM/2023/v19i8690

- [22] Soykan, Y., Generalized Pierre Numbers, *Journal of Progressive Research in Mathematics*, 20(1), 16-38, 2023.
- [23] Taşçı, D., On Mersenne Numbers, *Journal of Science and Arts*, 4(57), 1021-1028, 2021.
- [24] Taşçı, D., Gaussian Padovan and Gaussian Pell- Padovan numbers, *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 67(2), 82-88, 2018.
- [25] Taşçı, D., Gaussian Balancing and Gaussian Lucas Balancing Numbers, *Journal of Science and Arts*, 3(44), 661-666, 2018.
- [26] Yılmaz, B., Soykan, Y., Gaussian Generalized Guglielmo Numbers, *Asian Journal of Advanced Research and Reports*, 17(12), 1-18, 2023. DOI: 10.9734/AJARR/2023/v17i12581