

A Study on $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$: Eulerian Conditions, Kernel, and Domination

Abstract

In this research paper, we undertake a detailed study of a subdigraph of $\vec{\mathcal{G}}(\mathbb{Z}_n)$ for $n \geq 2$, formed by removing the vertex 0. This resulting subdigraph is denoted by $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$. We establish the necessary and sufficient conditions under which $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is Eulerian. Furthermore, we investigate the kernel of this digraph and analyze domination and twin domination properties within the framework of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.

Keywords: Directed Power graph $\vec{\mathcal{G}}(\mathbb{Z}_n)$, $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$, Eulerian Graph, Kernel, Domination, Twin dominating number.

2000 Mathematics Subject Classification: 05C25, 05C78

1 Introduction

In today's context, digraphs corresponding to algebraic structures such as groups and rings are highly significant. The directed power graph $\vec{\mathcal{G}}(G)$ of a group G , introduced by Kelarev et al. in (1), is a digraph with vertex set G and for any $a, b \in G$, there is a directed edge from a to b in $\vec{\mathcal{G}}(G)$ if and only if $a^k = b$, where $k \in \mathbb{N}$. In (6) Manuel et al. defined the concept of a directed power graph associated with the finite cyclic group \mathbb{Z}_n as a simple digraph with a vertex set \mathbb{Z}_n and two distinct vertices in $\vec{\mathcal{G}}(\mathbb{Z}_n)$ are joined by a directed edge or an arc \vec{uv} from u to v if and only if there exists a non-negative integer r such that $v \equiv ru \pmod{n}$. They delve into the $\vec{\mathcal{G}}(\mathbb{Z}_n)$, the directed power graph of the cyclic group \mathbb{Z}_n using the help of congruence and the definition of cyclic subgroups. Through this exploration, we unveil several characteristics of $\vec{\mathcal{G}}(\mathbb{Z}_n)$, leveraging the notions of kernel, domination, and connectedness. In this paper, we examine a vertex-deleted sub-digraph of $\vec{\mathcal{G}}(\mathbb{Z}_n)$ formed by its non-zero elements. ■

Definition 1.1. $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is a subdigraph of $\vec{\mathcal{G}}(\mathbb{Z}_n)$ obtained by removing the vertex 0. That is, $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is obtained from $\vec{\mathcal{G}}(\mathbb{Z}_n)$ by removing the vertex 0 and the $n - 1$ arcs incident with 0.

Figure 1 gives $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ for $n = 3$ and $n = 4$. $\vec{\mathcal{G}}_0(\mathbb{Z}_2)$ is a digraph with only one vertex and no arcs.




Figure 1: $\vec{\mathcal{G}}_0(\mathbb{Z}_3)$ and $\vec{\mathcal{G}}_0(\mathbb{Z}_4)$.

Theorem 1.1. (6) Let $n \in \mathbb{Z}$ and let $1 < m_1 < m_2 < \dots < m_r = n$ be the divisors of n . Then the number of arcs a in $\vec{\mathcal{G}}(\mathbb{Z}_n)$ is

$$a = \sum_{i=1}^r \left[2 \binom{\phi(m_i)}{2} + \phi(m_i)[m_i - \phi(m_i)] \right]. \quad (1.1)$$

From Theorem 1.1, the number of arcs in $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is

$$\sum_{i=1}^r \left[2 \binom{\phi(m_i)}{2} + \phi(m_i)(m_i - \phi(m_i)) \right] - (n - 1).$$

In particular, for a prime number p , the number of arcs in $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is $(p - 1)(p - 2)$. $\vec{\mathcal{G}}(\mathbb{Z}_n)$ can not be Eulerian, since $od(0) = 0$. But $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is Eulerian when n is a prime number. The following Theorem gives a characterization of Eulerian $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.

Theorem 1.2. $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is Eulerian if and only if n is a prime number.

Proof. Suppose that $n = p$ is a prime number. Then for each of its vertices v , $od(v) = id(v)$. Hence $\vec{\mathcal{G}}_0(\mathbb{Z}_p)$ is Eulerian. More precisely, let $\{1, 2, \dots, p - 1\}$ be the vertices of $\vec{\mathcal{G}}_0(\mathbb{Z}_p)$. Also, $\vec{\mathcal{G}}_0(\mathbb{Z}_p)$ has $(p - 1)(p - 2)$ arcs. Then $1 \ 2 \ 3 \ \dots \ (p - 1) \ (p - 2) \ (p - 3) \ \dots \ 2 \ 1 \ 3 \ 1 \ 4 \ \dots \ 1 \ (p - 1) \ 2 \ 4 \ 2 \ 5 \ \dots \ 2 \ (p - 1) \ 3 \ 5 \ 3 \ 6 \ \dots \ 3 \ (p - 1) \ \dots \ i \ (i + 2) \ i \ (i + 3) \ \dots \ i \ (p - 1) \ \dots \ (p - 3) \ (p - 1) \ 1$ is an Euler tour $\vec{\mathcal{G}}_0(\mathbb{Z}_p)$.

Conversely, suppose that n is not a prime number. Let a be a generator of \mathbb{Z}_n , then $id(a) = \phi(n) - 1$ and $od(a) = n - 1$, where $\phi(n)$ is the number of generators of \mathbb{Z}_n . Since n is not a prime number, $od(a) \neq id(a)$ and consequently $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is not Eulerian. \square

Figure 2 gives an Euler tour in $\vec{\mathcal{G}}_0(\mathbb{Z}_5)$. Here $1 \ 2 \ 3 \ 4 \ 3 \ 2 \ 1 \ 3 \ 1 \ 4 \ 2 \ 4 \ 1$ is the Euler tour.

Now we give a characterization of symmetric $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.




Figure 2: An Euler tour of $\vec{\mathcal{G}}_0(\mathbb{Z}_5)$

Theorem 1.3. For $n \geq 2$, $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is a symmetric digraph if and only if n is a prime number.

Proof. Suppose $n = p$, a prime. Then there exist arcs between each distinct vertices $u, v \in V(\vec{\mathcal{G}}_0(\mathbb{Z}_n))$. Therefore $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is symmetric.

Conversely, suppose that $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ are symmetric digraphs. We claim that n is prime. Since 1 is a generator of \mathbb{Z}_n , there exist arcs from 1 to u , for every $u \in V(\vec{\mathcal{G}}_0(\mathbb{Z}_n))$. Since $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is symmetric, there exists an arc from u to 1 also. This is possible only if u is a generator of \mathbb{Z}_n . Thus every non-zero element in \mathbb{Z}_n is a generator and hence n is a prime. \square

Now Let us go through the following results relating kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.

Remark 1.1. If $n = p$, a prime number, the kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_p)$ are the singleton subsets of non-zero elements of \mathbb{Z}_p .

Theorem 1.4. No kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ contains a generator of \mathbb{Z}_n , if $n \neq p$, a prime.

Proof. Let K be a generator of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ and let a be a generator of \mathbb{Z}_n . Since K is an independent set $K = \{a\}$. Since n is not a prime, there exists $u \in V - K$, which is not a generator of \mathbb{Z}_n . Then there exists no arc from u to a which is a contradiction. Hence no kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ contains a generator of \mathbb{Z}_n if n is not a prime. \square

Let $C^0 = \{S_{a_1}, S_{a_2}, \dots, S_{a_r}\}$ be the subset of C explained in notations ???. By the above Theorem 1.4 we can prove the following characterizations on kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.

Theorem 1.5. For $n \neq p$, a prime, K be a kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ if and only if K contains exactly one element from each member of C^0 .

Proof. Suppose that K is a kernel of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$. Since $n \neq p$, by the above Theorem, K contains no generator. Since K is independent there exists at most one element from each member of C^0 . If K contains no member from S_{a_i} , then there exists no arc from a_i to any element in K . Since K is a kernel, this is not possible.

Conversely, suppose that K contains exactly one element from each member of C^0 . Then K is independent. Let $b \in V - K$, if b is a generator of \mathbb{Z}_n , then there exists an arc from b to elements of K . If b is not a generator, then $b \in S_b$. Let $b' \in S_b$ such that $b' \in K$, then there exists an arc bb' . Therefore K is a kernel. \square

Theorem 1.6. $S \subseteq V$ is a twin-dominating set of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ if and only if S contains at least one element from each strong component.

Proof. Let S_1, S_{a_1}, \dots , and S_{a_r} are the strong components of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$. Let $S = \{b, b_1, b_2, \dots, b_r\}$ be a subset of V containing exactly one element from each strong component of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$, where b is a generator of \mathbb{Z}_n . Now since b is a generator of \mathbb{Z}_n , there exist arcs \vec{bv} for every $v \in V$, in particular for every $v \in V - S$. Therefore $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$ is out-dominated by b . Now it is enough to prove that S is an in-dominating set. For that let $v \in V - S$. If v is a generator, then v is adjacent to every vertex in S . If v is not a generator, then $v \in S_{a_i}$, for some $i = 1, 2, \dots, r$. Since $b_i \in S_{a_i}$, there exists an arc from v to b_i . Thus S is a twin-dominating set.

Conversely, suppose S be a twin dominating set. We claim to prove that S contains at least one element from each strong component. If possible suppose that S contains no element from a strong component S_{a_i} , the generators of $\langle a_i \rangle$. Let $v \in S_{a_i}$, then there is no arc from v to any vertex of S . Therefore S cannot be a twin-dominating set. So S must contain at least one element from each strong component. \square

Remark 1.2. From the above Theorem, we can realize that the twin-dominating number, $\gamma^*(\vec{\mathcal{G}}_0(\mathbb{Z}_n))$ is the same as the number of strong components of $\vec{\mathcal{G}}_0(\mathbb{Z}_n)$.

References

- [1] A. V. Kelarev and S. J. Quinn, A combinatorial property and power graphs of groups, Contrib. General Algebra, 12:229–235, 2000.
- [2] John Clark, Derek Allan Holton A First Look at Graph Theory, Allied Publishers Ltd, 1995
- [3] F. Harary, *Graph Theory*, Narosa Publishing House, 2001.
- [4] J. B. Fraleigh, *A First Course in Abstract Algebra*, Seventh edition, Pearson
- [5] D.M. Burton, *Elementary Number Theory*, Mc Graw Hill Education(India) Private Limited, New Delhi, 2012
- [6] Jimly Manuel, Bindhu K Thomas, *Properties of Digraphs Associated with Finite Cyclic Groups*, International Journal of Scientific Research in Mathematical and Statistical Sciences, Vol. 6, Issue.5, pp 52-56, October (2019).
- [7] Joseph A Gallian, *A dynamic survey of graph labeling*, The Electronic Journal of Combinatorics, 25th edn, December 2, 2022
- [8] Narsingh Deo, *Graph Theory with Applications to Engineering and Computer Science*, PHI Learning Private Limited, New Delhi-110001, 2011.

-
- [9] Douglas B. West, Introduction to Graph Theory, Pearson, 2019.
 - [10] Douglas B. West, *On certain valuations of the vertices of a graph*, Theory of Graphs (Int. Symp. Rome 1966), Gordon and Breach, N. Y., and Dunod Paris, 1967, 349-355