

A Study On Dual Hyperbolic Generalized Edouard Numbers

Abstract.In this research, we introduce the generalized dual hyperbolic Edouard numbers, a novel class of number sequences that extends existing recurrence relations into a new mathematical framework. Several special cases of these numbers are examined in detail, including the dual hyperbolic Edouard numbers and the dual hyperbolic Edouard-Lucas numbers, each revealing intriguing combinatorial and algebraic properties.

Explicit expressions for these sequences are derived, such as Binet-type formulas, generating functions, and summation identities, which offer analytical insight into their behavior and structural patterns. In addition, we explore matrix representations associated with these sequences, providing an elegant algebraic tool for further theoretical development and potential applications.

2010 Mathematics Subject Classification. 11B39, 11B83.

Keywords. Edouard numbers, Edouard-Lucas numbers, Dual hyperbolic Edouard numbers, dual hyperbolic Edouard-Lucas numbers.

1. Introduction

1.1. Dual Hyperbolic Numbers. In mathematical and geometric contexts, a hypercomplex system refers to a framework that generalizes the principles of complex numbers. These systems possess rich algebraic structures and are frequently studied for their diverse applications in physics and engineering. Below, we provide a concise overview of the key application areas of hypercomplex number systems in these fields.

In contrast to complex numbers, hypercomplex systems provide a more sophisticated framework for representing transformations and symmetries in higher-dimensional spaces. As noted by Kantor in [38], these systems can be viewed as extensions of the real number line, offering algebraic tools tailored to multidimensional analysis. The principal types of hypercomplex number systems encompass complex numbers, hyperbolic numbers, and dual numbers. Complex numbers, defined by a real and an imaginary component, serve as the foundational structure for more advanced hypercomplex systems. Hyperbolic numbers build

upon the complex number framework and are employed in diverse mathematical models, particularly those involving Lorentz transformations and spacetime geometries. Dual numbers, distinguished by the presence of a dual unit whose square is zero, are instrumental in various algebraic constructions, including automatic differentiation and kinematic analysis.

The following sections offer more detailed insights into the mathematical properties and application areas of these hypercomplex systems.

- Complex numbers are constructed by extending the real number system through the introduction of an imaginary unit, denoted as " i ", which satisfies the identity $i^2 = -1$. A complex number is typically expressed in the form $z = a + bi$, where a and b are real numbers, and i represents the imaginary unit.
- Hyperbolic numbers also referred to as double numbers or split complex numbers extend the real number system by introducing a new unit element j , which satisfies the identity $j^2 = 1$ [44]. These numbers are distinct from real and complex numbers due to their unique algebraic properties. A hyperbolic number is defined as:

$$\mathbb{H} = \{h = a + jb : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\}.$$

where a and b are real numbers and j is the hyperbolic unit. This structure enables the modeling of systems with split-signature metrics and has notable applications in areas such as special relativity and signal processing.

- Dual numbers [20] expand the real number system through the incorporation of a new element ε , which satisfies the identity $\varepsilon^2 = 0$. This infinitesimal unit distinguishes dual numbers from other hypercomplex systems and makes them especially valuable in modeling instantaneous rates of change. A dual number is defined as:

$$\mathbb{D} = \{d = a + \varepsilon b : a, b \in \mathbb{R}, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

where a and b are real numbers, and ε is the nilpotent unit. Dual numbers are commonly used in applications such as automatic differentiation, kinematics, and perturbation analysis, due to their ability to elegantly encode infinitesimal variations.

- Among the non-commutative examples of hypercomplex number systems are quaternions [26]. Quaternions generalize complex numbers by incorporating three distinct imaginary units, typically denoted as i, j , and k . A quaternion has the form $a_0 + ia_1 + ja_2 + ka_3$, where $a_0, a_1, a_2, a_3 \in \mathbb{R}$. These multiplication rules result in a non-commutative structure, meaning the order of multiplication affects the result. The set of quaternion numbers is formally defined as:

$$\mathbb{H}_{\mathbb{Q}} = \{q = a_0 + ia_1 + ja_2 + ka_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\},$$

- Additional hypercomplex systems include octonions and sedenions, which are discussed in [29] and [45]. The algebras \mathbb{C} (complex numbers), $\mathbb{H}_{\mathbb{Q}}$ (quaternions), \mathbb{O} (octonions), and \mathbb{S} (sedenions) are all constructed as real algebras derived from the real numbers \mathbb{R} using a recursive procedure known as the Cayley–Dickson Process. This technique successively doubles the dimension of each algebra and continues beyond sedenions to produce what are collectively referred to as the 2^n -ions. The following table highlights selected publications from the literature that investigate the properties and applications of these extended number systems.

For more information on hypercomplex algebra, see [39,27,41,75]

Table 1. Papers that have been published in the literature related to 2^n -ions.

Authors and Title of the paper↓	Papers↓
Biss, D.K., Dugger, D., Isaksen, D.C., Large annihilators in Cayley-Dickson algebras	[8]
Hamilton, W.R., Elements of Quaternions	[26]
Imaeda, K., Sedenions: algebra and analysis	[28]
Moreno, G., The zero divisors of the Cayley-Dickson algebras over the real numbers	[40]
Göcen, M., Soykan, Y., Horadam 2^k -Ions	[21]
Soykan, Y., Tribonacci and Tribonacci-Lucas Sedenions	[45]
On higher order Fibonacci hyper complex numbers	[37]

A dual hyperbolic number is a type of hypercomplex number, specifically a member of the hyperbolic number system. A dual hyperbolic number is defined as follows

$$q = (a_0 + ja_1) + \varepsilon(a_2 + ja_3) = a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3$$

where $a_0, a_1, a_2, a_3 \in \mathbb{R}$.

$\mathbb{H}_{\mathbb{D}}$, the set of all dual hyperbolic numbers, are generally denoted by

$$\mathbb{H}_{\mathbb{D}} = \{a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, j^2 = 1, j \neq \pm 1, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

The $\{1, j, \varepsilon, \varepsilon j\}$ is linearly independent, and the algebra $\mathbb{H}_{\mathbb{D}}$ is generated by their span, i.e. $\mathbb{H}_{\mathbb{D}} = \text{sp}\{1, j, \varepsilon, \varepsilon j\}$

Therefore, $\{1, j, \varepsilon, \varepsilon j\}$ forms a basis for the dual hyperbolic algebra $\mathbb{H}_{\mathbb{D}}$. For more detail, see [2].

The next properties are holds for the base elements $\{1, j, \varepsilon, \varepsilon j\}$ of dual hyperbolic numbers (commutative multiplications):

$$\begin{aligned} 1.\varepsilon &= \varepsilon, 1.j = j, \varepsilon^2 = \varepsilon.\varepsilon = (j\varepsilon)^2 = 0, j^2 = j.j = 1 \\ \varepsilon.j &= j.\varepsilon, \varepsilon.(\varepsilon j) = (\varepsilon j).\varepsilon = 0, j(\varepsilon j) = (\varepsilon j)j = \varepsilon \end{aligned}$$

where ε denotes the pure dual unit ($\varepsilon^2 = 0, \varepsilon \neq 0$), j denotes the hyperbolic unit ($j^2 = 1$), and εj denotes the dual hyperbolic unit ($(j\varepsilon)^2 = 0$).

We claim that p and q be two dual hyperbolic numbers that $q = a_0 + ja_1 + \varepsilon a_2 + j\varepsilon a_3$ and $p = b_0 + jb_1 + \varepsilon b_2 + j\varepsilon b_3$ and then we can write the product of p and q as

$$qp = a_0b_0 + a_1b_1 + j(a_0b_1 + a_1b_0) + \varepsilon(a_0b_2 + a_2b_0 + a_1b_3 + a_3b_1) + j\varepsilon(a_0b_3 + a_1b_2 + a_2b_1 + b_0a_3)$$

and we can write the sum dual hyperbolic numbers p and q as componentwise.

The dual hyperbolic numbers form a commutative ring, real vector space and an algebra. $\mathbb{H}_{\mathbb{D}}$ is not field since every dual hyperbolic numbers doesn't have an inverse. For more detail about dual hyperbolic numbers, see [2].

1.2. General Linear Recurrence Relations. Linear recurrence relations of second, third, and fourth order are specific instances of the more general k -th order linear recurrence relations. These relations define sequences in which each term is a linear combination of a fixed number of preceding terms. A homogeneous linear recurrence relation of order k is expressed as:

$$a_n = r_1a_{n-1} + r_2a_{n-2} + \cdots + r_ka_{n-k}, \quad \text{for } n \geq k,$$

where r_1, r_2, \dots, r_k are constant coefficients, and the initial terms a_0, a_1, \dots, a_{k-1} are given.

Examples of Specific Orders:

- **Second-order:**

$$a_n = ra_{n-1} + sa_{n-2}$$

Characteristic equation:

$$x^2 - rx - s = 0$$

- **Third-order:**

$$a_n = ra_{n-1} + sa_{n-2} + ta_{n-3}$$

Characteristic equation:

$$x^3 - rx^2 - sx - t = 0$$

- **Fourth-order:**

$$a_n = ra_{n-1} + sa_{n-2} + ta_{n-3} + ua_{n-4}$$

Characteristic equation:

$$x^4 - rx^3 - sx^2 - tx - u = 0$$

The general solution depends on the roots of the characteristic equation. If the polynomial has k distinct roots r_1, r_2, \dots, r_k , the solution is given by:

$$a_n = \alpha_1 r_1^n + \alpha_2 r_2^n + \cdots + \alpha_k r_k^n,$$

where $\alpha_1, \alpha_2, \dots, \alpha_k$ are constants determined by initial conditions.

In the case of repeated roots or complex conjugate pairs, the solution modifies accordingly:

- Repeated roots:

$$a_n = P(n)r^n$$

where $P(n)$ is a polynomial in n .

- Complex conjugate roots:

$$a_n = r^n (\alpha \cos(n\theta) + \beta \sin(n\theta))$$

An *inhomogeneous* linear recurrence relation includes a non-zero forcing term:

$$a_n = r_1 a_{n-1} + \cdots + r_k a_{n-k} + f(n),$$

where $f(n)$ is a known function. The general solution is the sum of the homogeneous solution and a particular solution to the nonhomogeneous recurrence.

Higher-order recurrence relations are foundational in numerous domains:

- **Combinatorics:** Problems involving partitions, tilings, and lattice paths.
- **Computer Science:** Analysis of recursive algorithms and dynamic programming.
- **Mathematical Physics:** Discrete dynamical systems and numerical modeling.
- **Number Theory:** Extensions of sequences such as Fibonacci, Lucas, and others.

1.3. Background on Edouard Numbers. This section presents key foundational results on Edouard numbers, which are governed by a third-order homogeneous linear recurrence relation. These numbers provide a framework for exploring generalized integer sequences and their associated algebraic structures.

A generalized Edouard sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2)\}_{n \geq 0}$ is defined by the third-order recurrence relations

$$W_n = 7W_{n-1} - 7W_{n-2} + W_{n-3}; \quad W_0, W_1, W_2 \quad (n \geq 3) \quad (1.1)$$

with the initial values W_0, W_1, W_2 not all being zero. The sequence $\{W_n\}_{n \geq 0}$ can be given to negative subscripts by defining

$$W_{-n} = 7W_{-(n-1)} - 7W_{-(n-2)} + W_{-(n-3)}$$

for $n = 1, 2, 3, \dots$. Thus, recurrence (1.1) is true for all integer n .

In the Table 2 we give the first few generalized Edouard numbers with positive subscript and negative subscript

Table 2. A few generalized Edouard numbers

n	W_n	W_{-n}
0	W_0	W_0
1	W_1	$7W_0 - 7W_1 + W_2$
2	W_2	$42W_0 - 48W_1 + 7W_2$
3	$W_0 - 7W_1 + 7W_2$	$246W_0 - 287W_1 + 42W_2$
4	$7W_0 - 48W_1 + 42W_2$	$1435W_0 - 1680W_1 + 246W_2$
5	$42W_0 - 287W_1 + 246W_2$	$8365W_0 - 9799W_1 + 1435W_2$
6	$246W_0 - 1680W_1 + 1435W_2$	$48756W_0 - 57120W_1 + 8365W_2$

If we take $W_0 = 0, W_1 = 1, W_2 = 7$ then $\{E_n\}$ is the Edouard sequence, if we take $W_0 = 3, W_1 = 7, W_2 = 35$ then $\{K_n\}$ is the Edouard-Lucas sequence. In other words, Edouard sequence $\{E_n\}_{n \geq 0}$, Edouard-Lucas sequence $\{K_n\}_{n \geq 0}$ are given by the third-order recurrence relations

$$E_n = 7E_{n-1} - 7E_{n-2} + E_{n-3}, \quad E_0 = 0, E_1 = 1, E_2 = 7, \quad (1.2)$$

$$K_n = 7K_{n-1} - 7K_{n-2} + K_{n-3}, \quad K_0 = 3, K_1 = 7, K_2 = 35, \quad (1.3)$$

In addition that the sequences given above can be extended to negative subscripts by defining,

$$E_{-n} = 7E_{-(n-1)} - 7E_{-(n-2)} + E_{-(n-3)},$$

$$K_{-n} = 7K_{-(n-1)} - 7K_{-(n-2)} + K_{-(n-3)},$$

for $n = 1, 2, 3, \dots$ respectively. As a result, recurrences (1.2)-(1.3) are true for all integer n .

We can enumerate several essential properties of generalized Edouard numbers that are required.

Binet formula of generalized Edouard sequence can be calculated using its characteristic equation given as

$$z^3 - 7z^2 + 7z - 1 = (z^2 - 6z + 1)(z - 1) = 0,$$

where the roots of above equation are

$$\alpha = 3 + 2\sqrt{2},$$

$$\beta = 3 - 2\sqrt{2},$$

$$\gamma = 1.$$

Using these roots and the recurrence relation of $\{W_n\}$, we can write the Binet's formula can be written as

$$\begin{aligned} W_n &= \frac{z_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)} + \frac{z_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)} + \frac{z_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)} \\ &= \frac{z_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)} + \frac{z_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)} - \frac{z_3 \gamma^n}{4} \\ &= A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n, \end{aligned} \quad (1.4)$$

where z_1, z_2 and z_3 are given below

$$\begin{aligned} z_1 &= W_2 - (\beta + 1)W_1 + \beta W_0, \\ z_2 &= W_2 - (\alpha + 1)W_1 + \alpha W_0, \\ z_3 &= W_2 - 6W_1 + W_0, \end{aligned}$$

and

$$\begin{aligned} A_1 &= \frac{W_2 - (\beta + 1)W_1 + \beta W_0}{(\alpha - \beta)(\alpha - \gamma)}, \\ A_2 &= \frac{W_2 - (\alpha + 1)W_1 + \alpha W_0}{(\beta - \alpha)(\beta - \gamma)}, \\ A_3 &= \frac{W_2 - 6W_1 + W_0}{(\gamma - \alpha)(\gamma - \beta)}. \end{aligned} \tag{1.5}$$

Binet's formula of Edouard, Edouard-Lucas sequences can be written as

$$\begin{aligned} E_n &= \frac{\alpha^{n+1}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^{n+1}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}, \\ K_n &= \alpha^n + \beta^n + 1. \end{aligned}$$

After then we can write the generating function of generalized Edouard numbers,

$$\sum_{n=0}^{\infty} W_n x^n = \frac{W_0 + (W_1 - 7W_0)x + (W_2 - 7W_1 + 7W_0)x^2}{1 - 7x + 7x^2 - x^3}. \tag{1.6}$$

Next, we give the exponential generating function of $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ of the sequence W_n .

LEMMA 1. [3, Lemma 1.4]. Suppose that $f_{W_n}(x) = \sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$ is the exponential generating function of the generalized Edouard sequence $\{W_n\}$. Then

$$\sum_{n=0}^{\infty} W_n \frac{x^n}{n!} = \frac{(W_2 - (\beta + 1)W_1 + \beta W_0)}{(\alpha - \beta)(\alpha - 1)} e^{\alpha x} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)}{(\beta - \alpha)(\beta - 1)} e^{\beta x} - \frac{(W_2 - 6W_1 + W_0)}{4} e^x.$$

The previous Lemma gives the following results as particular examples.

COROLLARY 2. Exponential generating function of Edouard and Edouard-Lucas numbers are

a):

$$\sum_{n=0}^{\infty} E_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} \left(\frac{\alpha^{n+1}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^{n+1}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4} \right) \frac{x^n}{n!} = \frac{\alpha e^{\alpha x}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta e^{\beta x}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4} e^x.$$

b):

$$\sum_{n=0}^{\infty} K_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} (\alpha^n + \beta^n + 1) \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^x.$$

For more details about generalized Edouard numbers, see [59].

Now, we give some information on published papers related to hyperbolic and dual hyperbolic numbers in literature.

- Cockle [12] explored hyperbolic numbers with complex coefficients, contributing to the early development of hypercomplex algebra.
- Cheng and Thompson [10] introduced dual numbers with complex coefficients, expanding the algebraic versatility of dual number systems for applications in polynomial equations and transformation theory.
- Akar et al [2] introduced the concept of dual hyperbolic numbers, combining characteristics of dual and hyperbolic systems into a unified algebraic structure.

Next, we give some information related to dual hyperbolic sequences presented in literature.

- Soykan et al [56] introduced the concept of dual hyperbolic generalized Pell numbers, extending classical Pell sequences into the framework of hypercomplex algebra. These numbers are defined as:

$$\widehat{V}_n = V_n + jV_{n+1} + \varepsilon V_{n+2} + j\varepsilon V_{n+3}$$

where $j^2 = 1$ and $\varepsilon^2 = 0$, reflecting the underlying dual hyperbolic structure. The scalar components V_n follow the recurrence relation for generalized Pell numbers: $V_n = 2V_{n-1} + V_{n-2}$, $V_0 = a$, $V_1 = b$ ($n \geq 2$) with initial values V_0, V_1 not both zero.

- Cihan et al [1] examined the structure of dual hyperbolic Fibonacci and dual hyperbolic Lucas numbers, which integrate classical sequences into a hypercomplex framework. These numbers are defined as:

$$\begin{aligned} DHF_n &= F_n + jF_{n+1} + \varepsilon F_{n+2} + j\varepsilon F_{n+3}, \\ DHL_n &= L_n + jL_{n+1} + \varepsilon L_{n+2} + j\varepsilon L_{n+3} \end{aligned}$$

where $j^2 = 1$, $\varepsilon^2 = 0$. The scalar components are governed by the classical recurrence relations. Fibonacci sequence: $F_n = F_{n-1} + F_{n-2}$, $F_0 = 0$, $F_1 = 1$, Lucas sequence: $L_n = L_{n-1} + L_{n-2}$, $L_0 = 2$, $L_1 = 1$.

- Soykan et al [[57] investigated dual hyperbolic generalized Jacobsthal numbers, extending classical recurrence relations within a hypercomplex framework. These numbers are expressed as:

$$\widehat{J}_n = J_n + jJ_{n+1} + \varepsilon J_{n+2} + j\varepsilon J_{n+3}$$

where $j^2 = 1$, $\varepsilon^2 = 0$ and the scalar components J_n follow the recurrence relation $J_n = J_{n-1} + 2J_{n-2}$, $J_0 = a$, $J_1 = b$.

- Bród et al [9] examined the structure of dual hyperbolic generalized balancing numbers, integrating classical recurrence sequences with dual and hyperbolic number theory. These numbers are defined as:

$$DHB_n = B_n + jB_{n+1} + \varepsilon B_{n+2} + j\varepsilon B_{n+3}$$

where $j^2 = 1$, $\varepsilon^2 = 0$ and the base sequence B_n follows the recurrence $B_n = 6B_{n-1} - B_{n-2}$, $B_0 = 0$, $B_1 = 1$.

Next section, we present the dual hyperbolic generalized Edouard numbers and give some properties of these numbers.

2. Dual Hyperbolic Generalized Edouard Numbers and their Generating Functions and Binet's Formulas

In this section, we define dual hyperbolic generalized Edouard numbers then using this definition, we present generating functions and Binet's formula of dual hyperbolic generalized Edouard numbers.

We now examine dual hyperbolic generalized Edouard numbers within the algebra \mathbb{H}_D . The n th such number is defined as

$$\widehat{W}_n = W_n + jW_{n+1} + \varepsilon W_{n+2} + j\varepsilon W_{n+3}. \quad (2.1)$$

with the initial values $\widehat{W}_0, \widehat{W}_1, \widehat{W}_2$. (2.1) can be written to negative subscripts by defining,

$$\widehat{W}_{-n} = W_{-n} + jW_{-n+1} + \varepsilon W_{-n+2} + j\varepsilon W_{-n+3}. \quad (2.2)$$

so identity (2.1) holds for all integers n .

Now, we define some special cases of dual hyperbolic generalized Edouard numbers. The n th dual hyperbolic Edouard numbers, the n th dual hyperbolic Edouard-Lucas numbers, respectively, are given as the n th dual hyperbolic Edouard numbers is given $\widehat{E}_n = E_n + jE_{n+1} + \varepsilon E_{n+2} + j\varepsilon E_{n+3}$, with the initial values

$$\begin{aligned} \widehat{E}_0 &= E_0 + jE_1 + \varepsilon E_2 + j\varepsilon E_3, \\ \widehat{E}_1 &= E_1 + jE_2 + \varepsilon E_3 + j\varepsilon E_4, \\ \widehat{E}_2 &= E_2 + jE_3 + \varepsilon E_4 + j\varepsilon E_5, \end{aligned}$$

the n th dual hyperbolic Edouard-Lucas numbers is given $\widehat{K}_n = K_n + jK_{n+1} + \varepsilon K_{n+2} + j\varepsilon K_{n+3}$ with the initial values

$$\begin{aligned} \widehat{K}_0 &= K_0 + jK_1 + \varepsilon K_2 + j\varepsilon K_3, \\ \widehat{K}_1 &= K_1 + jK_2 + \varepsilon K_3 + j\varepsilon K_4, \\ \widehat{K}_2 &= K_2 + jK_3 + \varepsilon K_4 + j\varepsilon K_5. \end{aligned}$$

Note that, for dual hyperbolic Edouard numbers (by using $W_n = E_n$, $E_0 = 0$, $E_1 = 1$, $E_2 = 7$) we get

$$\begin{aligned} \widehat{E}_0 &= j + 7\varepsilon + 42j\varepsilon, \\ \widehat{E}_1 &= 1 + 7j + 42\varepsilon + 246j\varepsilon, \\ \widehat{E}_2 &= 7 + 42j + 246\varepsilon + 1435j\varepsilon, \end{aligned}$$

for dual hyperbolic Edouard-Lucas numbers (by using $W_n = K_n$, $K_0 = 3$, $K_1 = 7$, $K_2 = 35$) we obtain

$$\begin{aligned}\widehat{K}_0 &= 3 + 7j + 35\varepsilon + 199j\varepsilon, \\ \widehat{K}_1 &= 7 + 35j + 199\varepsilon + 1155j\varepsilon, \\ \widehat{K}_2 &= 35 + 199j + 1155\varepsilon + 6727j\varepsilon.\end{aligned}$$

So, using (2.1), we can write the following identity for non negative integers n ,

$$\widehat{W}_n = 7\widehat{W}_{n-1} - 7\widehat{W}_{n-2} + \widehat{W}_{n-3}, \quad (2.3)$$

and the sequence $\{\widehat{W}_n\}_{n \geq 0}$ can be given as

$$\widehat{W}_{-n} = 7\widehat{W}_{-(n-1)} - 7\widehat{W}_{-(n-2)} + \widehat{W}_{-(n-3)},$$

for $n = 1, 2, 3, \dots$ by using (2.2). As a result., recurrence (2.3) holds for all integer n .

Table 3 presents the initial values of the dual hyperbolic generalized Edouard numbers \widehat{W}_n , showcasing terms with both positive and negative subscripts for a comprehensive view of the sequence's symmetric structure.

Table 3. A few dual hyperbolic generalized Edouard numbers

n	\widehat{W}_n	\widehat{W}_{-n}
0	\widehat{W}_0	\widehat{W}_0
1	\widehat{W}_1	$7\widehat{W}_0 - 7\widehat{W}_1 + \widehat{W}_2$
2	\widehat{W}_2	$42\widehat{W}_0 - 48\widehat{W}_1 + 7\widehat{W}_2$
3	$\widehat{W}_0 - 7\widehat{W}_1 + 7\widehat{W}_2$	$246\widehat{W}_0 - 287\widehat{W}_1 + 42\widehat{W}_2$
4	$7\widehat{W}_0 - 48\widehat{W}_1 + 42\widehat{W}_2$	$1435\widehat{W}_0 - 1680\widehat{W}_1 + 246\widehat{W}_2$
5	$42\widehat{W}_0 - 287\widehat{W}_1 + 246\widehat{W}_2$	$8365\widehat{W}_0 - 9799\widehat{W}_1 + 1435\widehat{W}_2$
6	$246\widehat{W}_0 - 1680\widehat{W}_1 + 1435\widehat{W}_2$	$48756\widehat{W}_0 - 57120\widehat{W}_1 + 8365\widehat{W}_2$

Note that

$$\begin{aligned}\widehat{W}_0 &= W_0 + jW_1 + \varepsilon W_2 + j\varepsilon W_3, \\ \widehat{W}_1 &= W_1 + jW_2 + \varepsilon W_3 + j\varepsilon W_4, \\ \widehat{W}_2 &= W_2 + jW_3 + \varepsilon W_4 + j\varepsilon W_5.\end{aligned}$$

A few dual hyperbolic Edouard numbers, dual hyperbolic Edouard-Lucas numbers with positive subscript and negative subscript are given in the following Table 4, Table 5.

Table 4. Dual hyperbolic Edouard numbers

n	\widehat{E}_n	\widehat{E}_{-n}
0	$j + 7\varepsilon + 42j\varepsilon$	
1	$1 + 7j + 42\varepsilon + 246j\varepsilon$	$\varepsilon + 7j\varepsilon$
2	$7 + 42j + 246\varepsilon + 1435j\varepsilon$	$1 + j\varepsilon$
3	$42 + 246j + 1435\varepsilon + 8365j\varepsilon$	$7 + j$
4	$246 + 1435j + 8365\varepsilon + 48756j\varepsilon$	$42 + 7j + \varepsilon$
5	$1435 + 8365j + 48756\varepsilon + 284172j\varepsilon$	$246 + 42j + 7\varepsilon + j\varepsilon$

Table 5. Dual hyperbolic Edouard-Lucas numbers

n	\widehat{K}_n	\widehat{K}_{-n}
0	$3 + 7j + 35\varepsilon + 199j\varepsilon$	
1	$7 + 35j + 199\varepsilon + 1155j\varepsilon$	$7 + 3j + 7\varepsilon + 35j\varepsilon$
2	$35 + 199j + 1155\varepsilon + 6727j\varepsilon$	$35 + 7j + 3\varepsilon + 7j\varepsilon$
3	$199 + 1155j + 6727\varepsilon + 39203j\varepsilon$	$199 + 35j + 7\varepsilon + 3j\varepsilon$
4	$1155 + 6727j + 39203\varepsilon + 228487j\varepsilon$	$1155 + 199j + 35\varepsilon + 7j\varepsilon$
5	$6727 + 39203j + 228487\varepsilon + 1331715j\varepsilon$	$6727 + 1155j + 199\varepsilon + 35j\varepsilon$

Now, we will give some expressions that we will use in the rest of the paper and then we define Binet's formula for the dual hyperbolic generalized Edouard numbers.

$$\widehat{\alpha} = 1 + j\alpha + \varepsilon\alpha^2 + j\varepsilon\alpha^3, \quad (2.4)$$

$$\widehat{\beta} = 1 + j\beta + \varepsilon\beta^2 + j\varepsilon\beta^3, \quad (2.5)$$

$$\widehat{\gamma} = 1 + j + \varepsilon + j\varepsilon. \quad (2.6)$$

Note that using above equalities we can write the following identities:

$$\widehat{\alpha}^2 = 1 + \alpha^2 + 2j\alpha + 2\varepsilon(\alpha^4 + \alpha^2) + 4j\varepsilon\alpha^3,$$

$$\widehat{\beta}^2 = 1 + \beta^2 + 2j\beta + 2\varepsilon(\beta^2 + \beta^4) + 4j\varepsilon\beta^3,$$

$$\widehat{\gamma}^2 = 3 + 2j + 4\varepsilon + 4j\varepsilon,$$

$$\widehat{\alpha}\widehat{\beta} = 1 + \alpha\beta + j(\beta + \alpha) + \varepsilon(\beta^2 + \alpha^2 + \alpha^3\beta + \alpha\beta^3) + \varepsilon j(\alpha^3 + \alpha^2\beta + \alpha\beta^2 + \beta^3),$$

$$\widehat{\alpha}\widehat{\gamma} = 1 + \alpha + j(1 + \alpha) + \varepsilon(1 + \alpha + \alpha^2 + \alpha^3) + j\varepsilon(1 + \alpha + \alpha^2 + \alpha^3),$$

$$\widehat{\beta}\widehat{\gamma} = 1 + \alpha + j(1 + \alpha) + \varepsilon(1 + \alpha + \alpha^2 + \alpha^3) + j\varepsilon(1 + \alpha + \alpha^2 + \alpha^3).$$

THEOREM 3. (*Binet's Formula*) Let n be any integer then the Binet's formula of dual hyperbolic generalized Edouard number is

$$\widehat{W}_n = \widehat{\alpha}A_1\alpha^n + \widehat{\beta}A_2\beta^n + \widehat{\gamma}A_3 \quad (2.7)$$

where $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}$ are given as (2.4)-(2.5)-(2.6).

Proof. Using Binet's formula of the generalized Edouard numbers given below

$$W_n = A_1\alpha^n + A_2\beta^n + A_3$$

where A_1, A_2, A_3 are given (1.5) we get

$$\begin{aligned} \widehat{W}_n &= W_n + jW_{n+1} + \varepsilon W_{n+2} + j\varepsilon W_{n+3}, \\ &= A_1\alpha^n + A_2\beta^n + A_3\gamma^n + (A_1\alpha^{n+1} + A_2\beta^{n+1} + A_3\gamma^{n+1})j + (A_1\alpha^{n+2} + A_2\beta^{n+2} + A_3\gamma^{n+2})\varepsilon \\ &\quad + (A_1\alpha^{n+3} + A_2\beta^{n+3} + A_3)\varepsilon j. \\ &= \widehat{\alpha}A_1\alpha^n + \widehat{\beta}A_2\beta^n + \widehat{\gamma}A_3. \end{aligned}$$

This proves (2.7). \square

In particular, for any integer n , the Binet's Formula of n th dual hyperbolic Edouard number, Edouard-Lucas numbers, respectively, provided by

$$\begin{aligned} \widehat{E}_n &= \frac{\widehat{\alpha}\alpha^{n+1}}{(\alpha - \beta)(\alpha - 1)} + \frac{\widehat{\beta}\beta^{n+1}}{(\beta - \alpha)(\beta - 1)} - \frac{\widehat{\gamma}}{4}, \\ \widehat{K}_n &= \widehat{\alpha}\alpha^n + \widehat{\beta}\beta^n + \widehat{\gamma}, \end{aligned}$$

In the following Theorem, we now derive the generating function for the sequence of dual hyperbolic generalized Edouard numbers, providing a compact analytical representation of their structure and recursive behavior.

THEOREM 4. *The generating function for the dual hyperbolic generalized Edouard numbers is*

$$f_{\widehat{W}_n}(x) = \frac{\widehat{W}_0 + (\widehat{W}_1 - 7\widehat{W}_0)x + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)x^2}{(1 - 7x + 7x^2 - x^3)}. \quad (2.8)$$

Proof. We assume that $f_{\widehat{W}_n}(x)$ is the generating function of the dual hyperbolic generalized Edouard numbers and then we can write

$$f_{\widehat{W}_n}(x) = \sum_{n=0}^{\infty} \widehat{W}_n x^n.$$

Then, in light of the definition of the dual hyperbolic generalized Edouard numbers, and subtracting $7xg(x)$ and $-7x^2g(x)$ from $x^3g(x)$, we get

$$\begin{aligned}
(1 - 7x + 7x^2 - x^3)f_{G\widehat{W}_n}(x) &= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 7x \sum_{n=0}^{\infty} \widehat{W}_n x^n + 7x^2 \sum_{n=0}^{\infty} \widehat{W}_n x^n - x^3 \sum_{n=0}^{\infty} \widehat{W}_n x^n, \\
&= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 7 \sum_{n=0}^{\infty} \widehat{W}_n x^{n+1} + 7 \sum_{n=0}^{\infty} \widehat{W}_n x^{n+2} - \sum_{n=0}^{\infty} \widehat{W}_n x^{n+3}, \\
&= \sum_{n=0}^{\infty} \widehat{W}_n x^n - 7 \sum_{n=1}^{\infty} \widehat{W}_{n-1} x^n + 7 \sum_{n=2}^{\infty} \widehat{W}_{n-2} x^n - \sum_{n=3}^{\infty} \widehat{W}_{n-3} x^n, \\
&= (\widehat{W}_0 + \widehat{W}_1 x + \widehat{W}_2 x^2) - 7(\widehat{W}_0 x + \widehat{W}_1 x^2) + 7G\widehat{W}_0 x^2 \\
&\quad + \sum_{n=3}^{\infty} (\widehat{W}_n - 7\widehat{W}_{n-1} + 7\widehat{W}_{n-2} - \widehat{W}_{n-3}) x^n, \\
&= \widehat{W}_0 + \widehat{W}_1 x + \widehat{W}_2 x^2 - 7\widehat{W}_0 x - 7\widehat{W}_1 x^2 + 7\widehat{W}_0 x^2, \\
&= \widehat{W}_0 + (\widehat{W}_1 - 7\widehat{W}_0)x + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)x^2.
\end{aligned}$$

Note that, using the recurrence relation $\widehat{W}_n = 7\widehat{W}_{n-1} - 7\widehat{W}_{n-2} + \widehat{W}_{n-3}$ and rearranging above equation, the (2.8) has been obtained. \square

Now we can write the generating functions of the dual hyperbolic Edouard, Edouard-Lucas numbers as

$$\begin{aligned}
f_{\widehat{E}_n}(x) &= \frac{(j + 7\varepsilon + 42j\varepsilon) + (1 - 48j\varepsilon - 7\varepsilon)x + (\varepsilon + 7j\varepsilon)x^2}{(1 - 7x + 7x^2 - x^3)}, \\
f_{\widehat{K}_n}(x) &= \frac{(3 + 7j + 35\varepsilon + 199j\varepsilon) + (-14 - 14j - 46\varepsilon - 238j\varepsilon)x + (7 + 3j + 7\varepsilon + 35j\varepsilon)x^2}{(1 - 7x + 7x^2 - x^3)},
\end{aligned}$$

respectively. \square

Next, we give the exponential generating function of $\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ of the sequence \widehat{W}_n .

LEMMA 5. Suppose that $f_{\widehat{W}_n}(x) = \sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ is the exponential generating function of the dual hyperbolic generalized Edouard sequence $\{\widehat{W}_n\}$.

Then $\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!}$ is given by

$$\begin{aligned}
\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} + j \sum_{n=0}^{\infty} W_{n+1} \frac{x^n}{n!} + \varepsilon \sum_{n=0}^{\infty} W_{n+2} \frac{x^n}{n!} + j\varepsilon \sum_{n=0}^{\infty} W_{n+3} \frac{x^n}{n!} \\
&= \frac{(W_2 - (\beta + 1)W_1 + \beta W_0)}{(\alpha - \beta)(\alpha - 1)} e^{\alpha x} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)}{(\beta - \alpha)(\beta - 1)} e^{\beta x} - \frac{(W_2 - 6W_1 + W_0)}{4} e^x \\
&\quad + j \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha}{(\alpha - \beta)(\alpha - 1)} e^{\alpha x} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta}{(\beta - \alpha)(\beta - 1)} e^{\beta x} - \frac{(W_2 - 6W_1 + W_0)}{4} e^x \right) \\
&\quad + \varepsilon \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha^2}{(\alpha - \beta)(\alpha - 1)} e^{\alpha x} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta^2}{(\beta - \alpha)(\beta - 1)} e^{\beta x} - \frac{(W_2 - 6W_1 + W_0)}{4} e^x \right) \\
&\quad + j\varepsilon \left(\frac{(W_2 - (\beta + 1)W_1 + \beta W_0)\alpha^3}{(\alpha - \beta)(\alpha - 1)} e^{\alpha x} + \frac{(W_2 - (\alpha + 1)W_1 + \alpha W_0)\beta^3}{(\beta - \alpha)(\beta - 1)} e^{\beta x} - \frac{(W_2 - 6W_1 + W_0)}{4} e^x \right).
\end{aligned}$$

Proof: Note that we have

$$\sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} (W_n + jW_{n+1} + \varepsilon W_{n+2} + j\varepsilon W_{n+3}) \frac{x^n}{n!}.$$

Then using the Binet's formula of dual hiperbolic generalized Edouard numbers or exponential generating function of the generalized Edouard sequence we get the required identity.

The previous Lemma gives the following results as particular examples.

COROLLARY 6. *Exponential generating function of dual hiperbolic Edouard and dual hyperbolic Edouard-Lucas numbers are*

a):

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{E}_n \frac{x^n}{n!} &= \frac{\alpha e^{\alpha x}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta e^{\beta x}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}e^x + j\left(\frac{\alpha^2 e^{\alpha x}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^2 e^{\beta x}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}e^x\right) \\ &+ \varepsilon\left(\frac{\alpha^3 e^{\alpha x}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^3 e^{\beta x}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}e^x\right) + j\varepsilon\left(\frac{\alpha^4 e^{\alpha x}}{(\alpha - \beta)(\alpha - 1)} + \frac{\beta^4 e^{\beta x}}{(\beta - \alpha)(\beta - 1)} - \frac{1}{4}e^x\right). \end{aligned}$$

b):

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{K}_n \frac{x^n}{n!} &= e^{\alpha x} + e^{\beta x} + e^x + j(\alpha e^{\alpha x} + \beta e^{\beta x} + e^x) \\ &+ \varepsilon(\alpha^2 e^{\alpha x} + \beta^2 e^{\beta x} + e^x) + j\varepsilon(\alpha^3 e^{\alpha x} + \beta^3 e^{\beta x} + e^x). \end{aligned}$$

3. Obtaining Binet Formula From Generating Function

Next, by using generating function $f_{\widehat{W}_n}(x)$, we investigate Binet formula of $\{\widehat{W}_n\}$.

THEOREM 7. *(Binet formula of dual hyperbolic generalized Edouard numbers)*

$$\widehat{W}_n = \widehat{\alpha}A_1\alpha^n + \widehat{\beta}A_2\beta^n + \widehat{\gamma}A_3 \quad (3.1)$$

Proof. Using the $\sum_{n=0}^{\infty} \widehat{W}_n x^n$ we can write

$$\sum_{n=0}^{\infty} \widehat{W}_n x^n = \frac{\widehat{W}_0 + (\widehat{W}_1 - 7\widehat{W}_0)x + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)x^2}{(1 - 7x + 7x^2 - x^3)} = \frac{d_1}{(1 - \alpha x)} + \frac{d_2}{(1 - \beta x)} + \frac{d_3}{(1 - x)}, \quad (3.2)$$

so that

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{W}_n x^n &= \frac{d_1}{(1 - \alpha x)} + \frac{d_2}{(1 - \beta x)} + \frac{d_3}{(1 - x)}, \\ &= \frac{d_1(1 - x)(1 - \beta x) + d_2(1 - \alpha x)(1 - x) + d_3(1 - \alpha x)(1 - \beta x)}{(x^2 - 6x + 1)(1 - x)}, \end{aligned}$$

thus, we obtain

$$\widehat{W}_0 + (\widehat{W}_1 - 7\widehat{W}_0)x + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)x^2 = d_1 + d_2 + d_3 + (-d_2 - \alpha d_2 - \beta d_1 - \alpha d_3 - \beta d_3)x + (\alpha d_2 + \beta d_1 + \alpha \beta d_3)x^2.$$

By equating the coefficients of corresponding powers of x in the above equation, we get

$$\begin{aligned}\widehat{W}_0 &= d_1 + d_2 + d_3, \\ \widehat{W}_1 - 7\widehat{W}_0 &= -d_2 - \alpha d_2 - \beta d_1 - \alpha d_3 - \beta d_3, \\ \widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0 &= \alpha d_2 + \beta d_1 + \alpha \beta d_3.\end{aligned}\tag{3.3}$$

If we solve (3.3) we obtain

$$\begin{aligned}d_1 &= \frac{\widehat{W}_0 \alpha^2 + (\widehat{W}_1 - 7\widehat{W}_0) \alpha + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)}{(\alpha - \beta)(\alpha - \gamma)}, \\ d_2 &= \frac{\widehat{W}_0 \beta^2 + (\widehat{W}_1 - 7\widehat{W}_0) \beta + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)}{(\beta - \alpha)(\beta - \gamma)}, \\ d_3 &= \frac{\widehat{W}_0 \gamma^2 + (\widehat{W}_1 - 7\widehat{W}_0) \gamma + (\widehat{W}_2 - 7\widehat{W}_1 + 7\widehat{W}_0)}{(\gamma - \alpha)(\gamma - \beta)},\end{aligned}$$

Thus (3.2) can be given as

$$\begin{aligned}\sum_{n=0}^{\infty} \widehat{W}_n x^n &= d_1 \sum_{n=0}^{\infty} \alpha^n x^n + d_2 \sum_{n=0}^{\infty} \beta^n x^n + d_3 \sum_{n=0}^{\infty} x^n, \\ &= \sum_{n=0}^{\infty} (d_1 \alpha^n + d_2 \beta^n + d_3) x^n, \\ &= \sum_{n=0}^{\infty} \left(\frac{\widehat{W}_2 - (\beta + 1)\widehat{W}_1 + \beta \widehat{W}_0}{(\alpha - \beta)(\alpha - \gamma)} \alpha^n + \frac{\widehat{W}_2 - (\alpha + 1)\widehat{W}_1 + \alpha \widehat{W}_0}{(\beta - \alpha)(\beta - \gamma)} \beta^n + \frac{\widehat{W}_2 - 6\widehat{W}_1 + \widehat{W}_0}{(\gamma - \alpha)(\gamma - \beta)} \right) x^n.\end{aligned}$$

Hence, we get

$$\widehat{W}_n = \widehat{\alpha} A_1 \alpha^n + \widehat{\beta} A_2 \beta^n + \widehat{\gamma} A_3. \quad \square$$

4. Some Identities

We now introduce distinctive identities pertaining to the sequence $\{\widehat{W}_n\}$ of dual hyperbolic generalized Edouard numbers. The forthcoming theorem establishes a Simpson type formula within this framework, characterizing the structural relationships among consecutive terms of the sequence.

THEOREM 8. (*Simpson's formula for dual hyperbolic generalized Edouard numbers*) For all integers n we have,

$$\begin{vmatrix} \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n \\ \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} \\ \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} \end{vmatrix} = \begin{vmatrix} \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \end{vmatrix}.\tag{4.1}$$

Proof. To proof the above theorem, we can use mathematical induction. First we assume that $n \geq 0$. For $n = 0$ identity (4.1) is true. Let (4.1) is true for $n = k$. Consequently, the identity can be stated as

follows

$$\begin{vmatrix} \widehat{W}_{k+2} & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_{k+1} & \widehat{W}_k & \widehat{W}_{k-1} \\ \widehat{W}_k & \widehat{W}_{k-1} & \widehat{W}_{k-2} \end{vmatrix} = \begin{vmatrix} \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \end{vmatrix}.$$

For $n = k + 1$, and using above equality, we can write

$$\begin{aligned} \begin{vmatrix} \widehat{W}_{k+3} & \widehat{W}_{k+2} & \widehat{W}_{k+1} \\ \widehat{W}_{k+2} & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_{k+1} & \widehat{W}_k & \widehat{W}_{k-1} \end{vmatrix} &= \begin{vmatrix} 7\widehat{W}_{k+2} - 7\widehat{W}_{k+1} + \widehat{W}_k & \widehat{W}_{k+2} & \widehat{W}_{k+1} \\ 7\widehat{W}_{k+1} - 7\widehat{W}_k + \widehat{W}_{k-1} & \widehat{W}_{k+1} & \widehat{W}_k \\ 7\widehat{W}_k - 7\widehat{W}_{k-1} + \widehat{W}_{k-2} & \widehat{W}_k & \widehat{W}_{k-1} \end{vmatrix} \\ &= 7 \begin{vmatrix} \widehat{W}_{k+2} & \widehat{W}_{k+2} & \widehat{W}_{k+1} \\ \widehat{W}_{k+1} & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_k & \widehat{W}_k & \widehat{W}_{k-1} \end{vmatrix} - 7 \begin{vmatrix} \widehat{W}_{k+1} & \widehat{W}_{k+2} & \widehat{W}_{k+1} \\ \widehat{W}_k & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_{k-1} & \widehat{W}_k & \widehat{W}_{k-1} \end{vmatrix} \\ &\quad + \begin{vmatrix} \widehat{W}_k & \widehat{W}_{k+2} & \widehat{W}_{k+1} \\ \widehat{W}_{k-1} & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_{k-2} & \widehat{W}_k & \widehat{W}_{k-1} \end{vmatrix} \\ &= \begin{vmatrix} \widehat{W}_{k+2} & \widehat{W}_{k+1} & \widehat{W}_k \\ \widehat{W}_{k+1} & \widehat{W}_k & \widehat{W}_{k-1} \\ \widehat{W}_k & \widehat{W}_{k-1} & \widehat{W}_{k-2} \end{vmatrix}. \end{aligned}$$

Note that, for the case $n < 0$ the proof can be done similarly. Thus, the proof is completed. \square

From Theorem 4.1 we get the following corollary.

COROLLARY 9.

$$\begin{aligned} \text{(a): } \begin{vmatrix} \widehat{E}_{n+2} & \widehat{E}_{n+1} & \widehat{E}_n \\ \widehat{E}_{n+1} & \widehat{E}_n & \widehat{E}_{n-1} \\ \widehat{E}_n & \widehat{E}_{n-1} & \widehat{E}_{n-2} \end{vmatrix} &= -8j - 280\varepsilon - 280j\varepsilon - 8. \\ \text{(b): } \begin{vmatrix} \widehat{K}_{n+2} & \widehat{K}_{n+1} & \widehat{K}_n \\ \widehat{K}_{n+1} & \widehat{K}_n & \widehat{K}_{n-1} \\ \widehat{K}_n & \widehat{K}_{n-1} & \widehat{K}_{n-2} \end{vmatrix} &= 4096j + 143\,360\varepsilon + 143\,360j\varepsilon + 4096. \end{aligned}$$

THEOREM 10. Suppose that n and m be positive integers, E_n is Edouard numbers, the following equality is valid:

$$\widehat{W}_{m+n} = E_{m-1}\widehat{W}_{n+2} + (E_{m-3} - 7E_{m-2})\widehat{W}_{n+1} + E_{m-2}\widehat{W}_n. \quad (4.2)$$

Proof. First for the proof, we assume that $m \geq 0$. The identity (10) can be proved by mathematical induction on m . Taking $m = 0$, we get

$$\widehat{W}_n = E_{-1}\widehat{W}_{n+2} + (E_{-3} - 7E_{-2})\widehat{W}_{n+1} + E_{-2}\widehat{W}_n$$

which is true by seeing that $E_{-1} = 0, E_{-2} = 1, E_{-3} = 7$. We assume that the identity (4.2) holds for $m = k$. Then for $m = k + 1$, we get

$$\begin{aligned}
\widehat{W}_{(k+1)+n} &= 7\widehat{W}_{n+k} - 7\widehat{W}_{n+k-1} + \widehat{W}_{n+k-2} \\
&= 7(E_{k-1}\widehat{W}_{n+2} + (E_{k-3} - 7E_{k-2})\widehat{W}_{n+1} + E_{k-2}\widehat{W}_n) \\
&\quad - 7(E_{k-2}\widehat{W}_{n+2} + (E_{k-4} - 3E_{k-3})\widehat{W}_{n+1} + E_{k-3}\widehat{W}_n) \\
&\quad + (E_{k-3}\widehat{W}_{n+2} + (E_{k-5} - 3E_{k-4})\widehat{W}_{n+1} + E_{k-4}\widehat{W}_n) \\
&= (7E_{k-1} - 7E_{k-2} + E_{k-3})\widehat{W}_{n+2} + ((7E_{k-3} - 7E_{k-4} + E_{k-5}) \\
&\quad - 7(7E_{k-2} - 7E_{k-3} + E_{k-4}))\widehat{W}_{n+1} + (7E_{k-2} - 7E_{k-3} + E_{k-4})\widehat{W}_n \\
&= E_k\widehat{W}_{n+2} + (E_{k-2} - 7E_{k-1})\widehat{W}_{n+1} + E_{k-1}\widehat{W}_n \\
&= E_{(k+1)-1}\widehat{W}_{n+2} + (E_{(k+1)-3} - 7E_{(k+1)-2})\widehat{W}_{n+1} + E_{(k+1)-2}\widehat{W}_n.
\end{aligned}$$

Consequently, by mathematical induction on m , this proves (10). Note that, for the other cases the proof can be done similarly. \square

5. Linear Sums

In this section, we provide summation formulas for hyperbolic generalized Edouard numbers covering positive subscripts.

PROPOSITION 11. *For the generalized Edouard numbers, we have the following formulas:*

- (a): $\sum_{k=0}^n W_k = \frac{1}{4}(-(n+3)W_n + (n+2)(7W_{n+1} - W_{n+2}) - (n+1)W_{n+1} + 2W_2 - 13W_1 + 7W_0).$
- (b): $\sum_{k=0}^n W_{2k} = \frac{1}{32}(-(n+3)W_{2n} + (n+2)(-7W_{2n+2} + 48W_{2n+1} - 7W_{2n}) - (n+1)W_{2n+2} + 15W_2 - 96W_1 + 49W_0).$
- (c): $\sum_{k=0}^n W_{2k+1} = \frac{1}{32}(-(n+3)W_{2n+1} + (n+2)(-W_{2n+2} + 42W_{2n+1} - 7W_{2n}) - (n+1)(7W_{2n+2} - 7W_{2n+1} + W_{2n}) + 9W_2 - 56W_1 + 15W_0).$

Proof. It is given in Soykan [60, Theorem 3.3]. \square

Next, we present the formulas which give the summation of the dual hyperbolic generalized Edouard numbers.

THEOREM 12. *For $n \geq 0$ then the following sum formulas holds for dual hyperbolic generalized Edouard numbers.*

- (a): $\sum_{k=0}^n \widehat{W}_k = \frac{1}{4}(-(n+3)\widehat{W}_n + (n+2)(7\widehat{W}_{n+1} - \widehat{W}_{n+2}) - (n+1)\widehat{W}_{n+1} + 2\widehat{W}_2 - 13\widehat{W}_1 + 7\widehat{W}_0).$
- (b): $\sum_{k=0}^n \widehat{W}_{2k} = \frac{1}{32}(-(n+3)\widehat{W}_{2n} + (n+2)(-7\widehat{W}_{2n+2} + 48\widehat{W}_{2n+1} - 7\widehat{W}_{2n}) - (n+1)\widehat{W}_{2n+2} + 15\widehat{W}_2 - 96\widehat{W}_1 + 49\widehat{W}_0).$
- (c): $\sum_{k=0}^n \widehat{W}_{2k+1} = \frac{1}{32}(-(n+3)\widehat{W}_{2n+1} + (n+2)(-\widehat{W}_{2n+2} + 42\widehat{W}_{2n+1} - 7\widehat{W}_{2n}) - (n+1)(7\widehat{W}_{2n+2} - 7\widehat{W}_{2n+1} + \widehat{W}_{2n}) + 9\widehat{W}_2 - 56\widehat{W}_1 + 15\widehat{W}_0).$

Proof.

(a): Note that using (2.1), we get

$$\sum_{k=0}^n \widehat{W}_k = \sum_{k=0}^n W_k + j \sum_{k=0}^n W_{k+1} + \varepsilon \sum_{k=0}^n W_{k+2} + j\varepsilon \sum_{k=0}^n W_{k+3}$$

and using Proposition 11 the proof is easily attainable.

(b): Note that using (2.1), we get

$$\sum_{k=0}^n \widehat{W}_{2k} = \sum_{k=0}^n W_{2k} + j \sum_{k=0}^n W_{2k+1} + \varepsilon \sum_{k=0}^n W_{2k+2} + j\varepsilon \sum_{k=0}^n W_{2k+3}$$

and using Proposition 11 the proof is easily attainable.

(c): Note that using (2.1), we get

$$\sum_{k=0}^n \widehat{W}_{2k+1} = \sum_{k=0}^n W_{2k+1} + j \sum_{k=0}^n W_{2k+2} + \varepsilon \sum_{k=0}^n W_{2k+3} + j\varepsilon \sum_{k=0}^n W_{2k+4}$$

and using Proposition 11 the proof is easily attainable. \square

As a particular case of the Theorem 12 (a), we present the following corollary.

COROLLARY 13.

- (a):** $\sum_{k=0}^n \widehat{E}_k = \frac{1}{4}(-(n+3)\widehat{E}_n + (n+2)(7\widehat{E}_{n+1} - \widehat{E}_{n+2}) - (n+1)\widehat{E}_{n+1} + 1 - 34j\varepsilon - 5\varepsilon).$
(b): $\sum_{k=0}^n \widehat{K}_k = \frac{1}{4}(-(n+3)\widehat{K}_n + (n+2)(7\widehat{K}_{n+1} - \widehat{K}_{n+2}) - (n+1)\widehat{K}_{n+1} - 8j - 32\varepsilon - 168j\varepsilon).$

As a particular case of the Theorem 12 (b), we present the following corollary.

COROLLARY 14.

- (a):** $\sum_{k=0}^n \widehat{E}_{2k} = \frac{1}{32}(-(n+3)\widehat{E}_{2n} + (n+2)(-7\widehat{E}_{2n+2} + 48\widehat{E}_{2n+1} - 7\widehat{E}_{2n}) - (n+1)\widehat{E}_{2n+2} + 7j + \varepsilon - 33j\varepsilon + 9).$
(b): $\sum_{k=0}^n \widehat{K}_{2k} = \frac{1}{32}(-(n+3)\widehat{K}_{2n} + (n+2)(-7\widehat{K}_{2n+2} + 48\widehat{K}_{2n+1} - 7\widehat{K}_{2n}) - (n+1)\widehat{K}_{2n+2} - 32j - 64\varepsilon - 224j\varepsilon).$

As a particular case of the Theorem 12 (c), we present the following corollary.

COROLLARY 15.

- (a):** $\sum_{k=0}^n \widehat{E}_{2k+1} = \frac{1}{32}(-(n+3)\widehat{E}_{2n+1} + (n+2)(-\widehat{E}_{2n+2} + 42\widehat{E}_{2n+1} - 7\widehat{E}_{2n}) - (n+1)(7\widehat{E}_{2n+2} - 7\widehat{E}_{2n+1} + \widehat{E}_{2n}) + j - 33\varepsilon - 231j\varepsilon + 7).$
(b): $\sum_{k=0}^n \widehat{K}_{2k+1} = \frac{1}{32}(-(n+3)\widehat{K}_{2n+1} + (n+2)(-\widehat{K}_{2n+2} + 42\widehat{K}_{2n+1} - 7\widehat{K}_{2n}) - (n+1)(7\widehat{K}_{2n+2} - 7\widehat{K}_{2n+1} + \widehat{K}_{2n}) - 64j - 224\varepsilon - 1152j\varepsilon - 32).$

6. Matrices related with Dual Hyperbolic Generalized Edouard Numbers

In this section, using dual hyperbolic Edouard numbers, we give some matrices related to dual hyperbolic Edouard numbers.

We consider the triangular sequence $\{E_n\}$ defined by the third-order recurrence relation as follows

$$E_n = 7E_{n-1} - 7E_{n-2} + E_{n-3}$$

with the initial conditions

$$E_0 = 0, E_1 = 1, E_2 = 7.$$

We present the square matrix A of order 3 as

$$A = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

under the condition that $\det A = 1$. Then, we give the following Lemma.

LEMMA 16. *For any integers n the following identity can be written*

$$\begin{pmatrix} \widehat{W}_{n+2} \\ \widehat{W}_{n+1} \\ \widehat{W}_n \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}. \quad (6.1)$$

Proof. First, we prove the assertion for the case $n \geq 0$. Lemma 16 can be given by mathematical induction on n . If $n = 0$ we get

$$\begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}$$

which is true. We assume that (6.1) is true for $n = k$. Thus the following identity is true.

$$\begin{pmatrix} \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \\ \widehat{W}_k \end{pmatrix} = \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix}.$$

For $n = k + 1$, we get

$$\begin{aligned}
\begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} &= \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} \widehat{W}_2 \\ \widehat{W}_1 \\ \widehat{W}_0 \end{pmatrix} \\
&= \begin{pmatrix} 7 & -7 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \\ \widehat{W}_k \end{pmatrix} \\
&= \begin{pmatrix} 7\widehat{W}_{k+2} - 7\widehat{W}_{k+1} + \widehat{W}_k \\ \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \end{pmatrix} \\
&= \begin{pmatrix} \widehat{W}_{k+3} \\ \widehat{W}_{k+2} \\ \widehat{W}_{k+1} \end{pmatrix}.
\end{aligned}$$

For the other case $n < 0$ the proof is easily attainable. Consequently, using mathematical induction on n , the proof is completed.

Note that, see [58],

$$A^n = \begin{pmatrix} E_{n+1} & -7E_n + E_{n-1} & E_n \\ E_n & -7E_{n-1} + E_{n-2} & E_{n-1} \\ E_{n-1} & -7E_{n-2} + E_{n-3} & E_{n-2} \end{pmatrix}.$$

THEOREM 17. *If we define the matrices $N_{\widehat{W}}$ and $S_{\widehat{W}}$ as follow*

$$\begin{aligned}
N_{\widehat{W}} &= \begin{pmatrix} \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \end{pmatrix}, \\
S_{\widehat{W}} &= \begin{pmatrix} \widehat{W}_{n+2} & \widehat{W}_{n+1} & \widehat{W}_n \\ \widehat{W}_{n+1} & \widehat{W}_n & \widehat{W}_{n-1} \\ \widehat{W}_n & \widehat{W}_{n-1} & \widehat{W}_{n-2} \end{pmatrix}.
\end{aligned}$$

then the following identity is true:

$$A^n N_{\widehat{W}} = S_{\widehat{W}}.$$

Proof. For the proof, we can use the following identities

$$\begin{aligned} A^n N_{\widehat{W}} &= \begin{pmatrix} E_{n+1} & -7E_n + E_{n-1} & E_n \\ E_n & -7E_{n-1} + E_{n-2} & E_{n-1} \\ E_{n-1} & -7E_{n-2} + E_{n-3} & E_{n-2} \end{pmatrix} \begin{pmatrix} \widehat{W}_2 & \widehat{W}_1 & \widehat{W}_0 \\ \widehat{W}_1 & \widehat{W}_0 & \widehat{W}_{-1} \\ \widehat{W}_0 & \widehat{W}_{-1} & \widehat{W}_{-2} \end{pmatrix}, \\ &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \end{aligned}$$

where

$$\begin{aligned} a_{11} &= \widehat{W}_2 E_{n+1} + \widehat{W}_1 (E_{n-1} - 7E_n) + \widehat{W}_0 E_n, \\ a_{12} &= \widehat{W}_1 E_{n+1} + \widehat{W}_0 (E_{n-1} - 7E_n) + \widehat{W}_{-1} E_n, \\ a_{13} &= \widehat{W}_0 E_{n+1} + \widehat{W}_{-1} (E_{n-1} - 7E_n) + \widehat{W}_{-2} E_n, \\ a_{21} &= \widehat{W}_2 E_n + \widehat{W}_1 (E_{n-2} - 7E_{n-1}) + \widehat{W}_0 E_{n-1}, \\ a_{22} &= \widehat{W}_1 E_n + \widehat{W}_0 (E_{n-2} - 7E_{n-1}) + \widehat{W}_{-1} E_{n-1}, \\ a_{23} &= \widehat{W}_0 E_n + \widehat{W}_{-1} (E_{n-2} - 7E_{n-1}) + \widehat{W}_{-2} E_{n-1}, \\ a_{31} &= \widehat{W}_2 E_{n-1} + \widehat{W}_1 (E_{n-3} - 7E_{n-2}) + \widehat{W}_0 E_{n-2}, \\ a_{32} &= \widehat{W}_1 E_{n-1} + \widehat{W}_0 (E_{n-3} - 7E_{n-2}) + \widehat{W}_{-1} E_{n-2}, \\ a_{33} &= \widehat{W}_0 E_{n-1} + \widehat{W}_{-1} (E_{n-3} - 7E_{n-2}) + \widehat{W}_{-2} E_{n-2}. \end{aligned}$$

Using the Theorem 10, the proof is done. \square

From Theorem 17, we can write the following corollary.

COROLLARY 18.

(a): We suppose that the matrices $N_{\widehat{E}}$ and $S_{\widehat{E}}$ are defined as the following

$$\begin{aligned} N_{\widehat{E}} &= \begin{pmatrix} \widehat{E}_2 & \widehat{E}_1 & \widehat{E}_0 \\ \widehat{E}_1 & \widehat{E}_0 & \widehat{E}_{-1} \\ \widehat{E}_0 & \widehat{E}_{-1} & \widehat{E}_{-2} \end{pmatrix}, \\ S_{\widehat{E}} &= \begin{pmatrix} \widehat{E}_{n+2} & \widehat{E}_{n+1} & \widehat{E}_n \\ \widehat{E}_{n+1} & \widehat{E}_n & \widehat{E}_{n-1} \\ \widehat{E}_n & \widehat{E}_{n-1} & \widehat{E}_{n-2} \end{pmatrix}, \end{aligned}$$

so that the following identity is true for A^n , $N_{\widehat{E}}$, $S_{\widehat{E}}$,

$$A^n N_{\widehat{E}} = S_{\widehat{E}},$$

(b): We suppose that the matrices $N_{\hat{K}}$ and $S_{\hat{K}}$ are defined as the following

$$N_{\hat{K}} = \begin{pmatrix} \hat{K}_2 & \hat{K}_1 & \hat{K}_0 \\ \hat{K}_1 & \hat{K}_0 & \hat{K}_{-1} \\ \hat{K}_0 & \hat{K}_{-1} & \hat{K}_{-2} \end{pmatrix},$$

$$S_{\hat{K}} = \begin{pmatrix} \hat{K}_{n+2} & \hat{K}_{n+1} & \hat{K}_n \\ \hat{K}_{n+1} & \hat{K}_n & \hat{K}_{n-1} \\ \hat{K}_n & \hat{K}_{n-1} & \hat{K}_{n-2} \end{pmatrix},$$

so that the following identity is true for A^n , $N_{\hat{K}}$, $S_{\hat{K}}$,

$$A^n N_{\hat{K}} = S_{\hat{K}}.$$

7. Conclusion

In the literature, numerous studies have explored numerical sequences, which play a significant role in a wide range of disciplines, including physics, engineering, architecture, nature, and art. Among these, integer sequences such as Fibonacci, Lucas, Pell, and Jacobsthal are among the most well-known examples of second-order recurrence relations. The Fibonacci sequence, in particular, gained fame through its appearance in the rabbit population problem introduced by Leonardo of Pisa in his 1202 book *Liber Abaci*. Both the Fibonacci and Lucas sequences have inspired a wealth of intriguing and elegant mathematical identities. For rich applications of these second order sequences in science and nature, one can see the citations in [33,34,35].

In this work, we introduce the concept of dual hyperbolic generalized Edouard sequences and focus on four specific cases: dual hyperbolic Edouard numbers, dual hyperbolic Edouard-Lucas numbers.

- In section 1, we present information on the application areas of hypercomplex number systems in physics and engineering fields. Also, we give some properties about generalized Edouard numbers.
- In section 2, we define dual hyperbolic generalized Edouard numbers then using this definition, we present generating functions and Binet's formula of dual hyperbolic generalized Edouard numbers.
- In section 3, we obtain Binet's formula from generating function.
- In section 4, we give certain distinctive identities for the dual hyperbolic generalized Edouard sequence $\{\widehat{W}_n\}$ that named Simpson's formula.
- In section 5, we present summation formulas for dual hyperbolic generalized Edouard numbers.
- In section 6, we give some matrices related to dual hyperbolic Edouard numbers.

Linear recurrence relations (sequences) have many applications. Next, we list applications of sequences which are linear recurrence relations.

First, we present some applications of second order sequences.

- For the applications of Gaussian Fibonacci and Gaussian Lucas numbers to Pauli Fibonacci and Pauli Lucas quaternions, see [4].

- For the application of Pell Numbers to the solutions of three-dimensional difference equation systems, see [6].
- For the application of Jacobsthal numbers to special matrices, see [72].
- For the application of generalized k -order Fibonacci numbers to hybrid quaternions, see [24].
- For the applications of Fibonacci and Lucas numbers to Split Complex Bi-Periodic numbers, see [73].
- For the applications of generalized bivariate Fibonacci and Lucas polynomials to matrix polynomials, see [74].
- For the applications of generalized Fibonacci numbers to binomial sums, see [71].
- For the application of generalized Jacobsthal numbers to hyperbolic numbers, see [68].
- For the application of generalized Fibonacci numbers to dual hyperbolic numbers, see [69].
- For the application of Laplace transform and various matrix operations to the characteristic polynomial of the Fibonacci numbers, see [16].
- For the application of Generalized Fibonacci Matrices to Cryptography, see [43].
- For the application of higher order Jacobsthal numbers to quaternions, see [42].
- For the application of Fibonacci and Lucas Identities to Toeplitz-Hessenberg matrices, see [22].
- For the applications of Fibonacci numbers to lacunary statistical convergence, see [5].
- For the applications of Fibonacci numbers to lacunary statistical convergence in intuitionistic fuzzy normed linear spaces, see [30].
- For the applications of Fibonacci numbers to ideal convergence on intuitionistic fuzzy normed linear spaces, see [31].
- For the applications of k -Fibonacci and k -Lucas numbers to spinors, see [32].
- For the application of dual-generalized complex Fibonacci and Lucas numbers to Quaternions, see [70].
- For the application of special cases of Horadam numbers to Neutrosophic analysis see [23].
- For the application of Hyperbolic Fibonacci numbers to Quaternions, see [15].

We now present some applications of third order sequences.

- For the applications of third order Jacobsthal numbers and Tribonacci numbers to quaternions, see [14] and [13], respectively.
- For the application of Tribonacci numbers to special matrices, see [55].
- For the applications of Padovan numbers and Tribonacci numbers to coding theory, see [46] and [7], respectively.
- For the application of Pell-Padovan numbers to groups, see [19].
- For the application of adjusted Jacobsthal-Padovan numbers to the exact solutions of some difference equations, see [25].

- For the application of Gaussian Tribonacci numbers to various graphs, see [53].
- For the application of third-order Jacobsthal numbers to hyperbolic numbers, see [17]. For the application of Narayan numbers to finite groups see [36].
- For the application of generalized third-order Jacobsthal sequence to binomial transform, see [47].
- For the application of generalized Generalized Padovan numbers to Binomial Transform, see [48].
- For the application of generalized Tribonacci numbers to Gaussian numbers, see [49].
- For the application of generalized Tribonacci numbers to Sedenions, see [50].
- For the application of Tribonacci and Tribonacci-Lucas numbers to matrices, see [51].
- For the application of generalized Tribonacci numbers to circulant matrix, see [52].
- For the application of Tribonacci and Tribonacci-Lucas numbers to hybrinomials, see [54].
- For the application of hyperbolic Leonardo and hyperbolic Francois numbers to quaternions, see [18].

Next, we now list some applications of fourth order sequences.

- For the application of Tetranacci and Tetranacci-Lucas numbers to quaternions, see [63].
- For the application of generalized Tetranacci numbers to Gaussian numbers, see [64].
- For the application of Tetranacci and Tetranacci-Lucas numbers to matrices, see [65].
- For the application of generalized Tetranacci numbers to binomial transform, see [66].

We now present some applications of fifth order sequences.

- For the application of Pentanacci numbers to matrices, see [62].
- For the application of generalized Pentanacci numbers to quaternions, see [61].
- For the application of generalized Pentanacci numbers to binomial transform, see [67].

References

- [1] A. Cihan, A. Z. Azak, M. A. Güngör, M. Tosun, A Study on Dual Hyperbolic Fibonacci and Lucas Numbers, *An. Şt. Univ. Ovidius Constanta*, 27(1), 35–48, 2019.
- [2] Akar, M., Yüce, S., Şahin, Ş., On the Dual Hyperbolic Numbers and the Complex Hyperbolic Numbers, *Journal of Computer Science & Computational Mathematics*, 8(1), 1-6, 2018.
- [3] Ayrilma, E. E., Soykan, Y., A Study On Gaussian Generalized Edouard Numbers, *Asian Journal of Advanced Research and Reports*, 19(5), 421–438, 2025.
- [4] Azak, A.Z., Pauli Gaussian Fibonacci and Pauli Gaussian Lucas Quaternions. *Mathematics*, 2022, 10, 4655. <https://doi.org/10.3390/math10244655>
- [5] Bilgin, N.G., Fibonacci Lacunary Statistical Convergence of Order γ in IFNLS, *International Journal of Advances in Applied Mathematics and Mechanics*, 8(4), 28-36, 2021.
- [6] Büyük, H., Taşkara, N., On The Solutions of Three-Dimensional Difference Equation Systems Via Pell Numbers, *European Journal of Science and Technology, Special Issue* 34, 433-440, 2022.
- [7] Basu, M., Das, M., Tribonacci Matrices and a New Coding Theory, *Discrete Mathematics, Algorithms and Applications*, 6 (1), 1450008, (17 pages), 2014.

- [8] Biss, D.K., Dugger, D., Isaksen, D.C., Large annihilators in Cayley-Dickson algebras, *Communication in Algebra*, 36 (2), 632-664, 2008.
- [9] Bród, D., Liana, A., Włoch, I., Two Generalizations of Dual-Hyperbolic Balancing Numbers, *Symmetry*, 12(11), 1866, 2020
- [10] Cheng, H. H., Thompson, S., Dual Polynomials and Complex Dual Numbers for Analysis of Spatial Mechanisms, *Proc. of ASME 24th Biennial Mechanisms Conference*, Irvine, CA, August, 19-22, 1996.
- [11] Cihan, A., Azak, A. Z., Güngör, M. A., Tosun, M., A Study on Dual Hyperbolic Fibonacci and Lucas Numbers, *An. Şt. Univ. Ovidius Constanta*, 27(1), 35-48, (2019).
- [12] Cockle, J., On a New Imaginary in Algebra, *Philosophical magazine*, London-Dublin-Edinburgh, 3(34), 37-47, 1849.
- [13] Cerda-Morales, G., On a Generalization of Tribonacci Quaternions, *Mediterranean Journal of Mathematics* 14:239, 1-12, 2017.
- [14] Cerda-Morales, G., Identities for Third Order Jacobsthal Quaternions, *Advances in Applied Clifford Algebras* 27(2), 1043-1053, 2017.
- [15] Daşdemir, A., On Recursive Hyperbolic Fibonacci Quaternions, *Communications in Advanced Mathematical Sciences*, 4(4), 198-207, 2021. DOI:10.33434/cams.997824
- [16] Deveci, Ö., Shannon, A.G., On Recurrence Results From Matrix Transforms, *Notes on Number Theory and Discrete Mathematics*, 28(4), 589-592, 2022. DOI: 10.7546/nntdm.2022.28.4.589-592
- [17] Dikmen, C.M., Altınsoy, M., On Third Order Hyperbolic Jacobsthal Numbers, *Konuralp Journal of Mathematics*, 10(1), 118-126, 2022.
- [18] Dişkaya, O., Menken, H., Catarino, P.M.M.C., On the Hyperbolic Leonardo and Hyperbolic Francois Quaternions, *Journal of New Theory*, 42, 74-85, 2023.
- [19] Deveci, Ö., Shannon, A.G., Pell-Padovan-Circulant Sequences and Their Applications, *Notes on Number Theory and Discrete Mathematics*, 23(3), 100-114, 2017.
- [20] Fjelstad, P., Gal, S.G., n-dimensional Hyperbolic Complex Numbers, *Advances in Applied Clifford Algebras*, 8(1), 47-68, 1998, .
- [21] Göcen, M., Soykan, Y., Horadam 2^k -Ions, *Konuralp Journal of Mathematics*, 7(2), 492-501, 2019.
- [22] Goy, T., Shattuck, M., Fibonacci and Lucas Identities from Toeplitz-Hessenberg Matrices, *Appl. Appl. Math.*, 14(2), 699-715, 2019.
- [23] Gökbaşı, H., Topal, S., Smarandache, F., Neutrosophic Number Sequences: An Introductory Study, *International Journal of Neutrosophic Science (IJNS)*, 20(01), 27-48, 2023. <https://doi.org/10.54216/IJNS.200103>
- [24] Gül, K., Generalized k-Order Fibonacci Hybrid Quaternions, *Erzincan University Journal of Science and Technology*, 15(2), 670-683, 2022. DOI: 10.18185/erzifbed.1132164
- [25] Göcen, M., The Exact Solutions of Some Difference Equations Associated with Adjusted Jacobsthal-Padovan Numbers, *Kırklareli University Journal of Engineering and Science* 8(1), 1-14, 2022. DOI: 10.34186/klujes.1078836
- [26] Hamilton, W.R., *Elements of Quaternions*, Chelsea Publishing Company, New York, 1969.
- [27] Hestenes, D., *New Foundations for Classical Mechanics*, Springer, 1999.
- [28] Imaeda, K., Imaeda, M., Sedenions: algebra and analysis, *Applied Mathematics and Computation*, 115, 77-88, 2000.
- [29] J. Baez, The octonions, *Bull. Amer. Math. Soc.* 39(2), 145-205, 2002.
- [30] Kişi, Ö., Tuzcuoglu, I., Fibonacci Lacunary Statistical Convergence in Intuitionistic Fuzzy Normed Linear Spaces, *Journal of Progressive Research in Mathematics* 16(3), 3001-3007, 2020.
- [31] Kişi, Ö., Debnath, P., Fibonacci Ideal Convergence on Intuitionistic Fuzzy Normed Linear Spaces, *Fuzzy Information and Engineering*, 1-13, 2022. <https://doi.org/10.1080/16168658.2022.2160226>
- [32] Kumari, M., Prasad, K., Frontczak, R., On the k -Fibonacci and k -Lucas Spinors, *Notes on Number Theory and Discrete Mathematics*, 29(2), 322-335, 2023. DOI: 10.7546/nntdm.2023.29.2.322-335

- [33] Koshy, T., Fibonacci and Lucas Numbers with Applications, Volume 1 (Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts), Second Edition, John Wiley&Sons, New York, 2018.
- [34] Koshy, T., Fibonacci and Lucas Numbers with Applications, Volume 2 (Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts), John Wiley&Sons, New York, 2019.
- [35] Koshy, T., Pell and Pell-Lucas Numbers with Applications, Springer, New York, 2014.
- [36] Kuloglu, B., Özkan, E., Shannon, A.G., The Narayana Sequence in Finite Groups, Fibonacci Quarterly. 60(5), 212–221, 2022. 2014.
- [37] Kızılateş, C., Kone, T., On higher order Fibonacci hyper complex numbers, Chaos Solitons & Fractals, 148(1), 6 pages (2021).
- [38] Kantor, I., Solodovnikov, A., Hypercomplex Numbers, Springer-Verlag, New York, 1989.
- [39] Lounesto, R., Clifford Algebras and Spinors, Cambridge University Press, 2001.
- [40] Moreno, G., The zero divisors of the Cayley-Dickson algebras over the real numbers, Bol. Soc. Mat. Mexicana 3(4), 13-28, 1998.
- [41] Olver, P.J., Applications of Lie Groups to Differential Equations, Springer, 2012.
- [42] Özkan, E., Uysal, M., On Quaternions with Higher Order Jacobsthal Numbers Components, Gazi University Journal of Science, 36(1), 336-347, 2023. DOI: 10.35378/gujs. 1002454
- [43] Prasad, K., Mahato, H., Cryptography Using Generalized Fibonacci Matrices with Affine-Hill Cipher, Journal of Discrete Mathematical Sciences & Cryptography, 25(8-A), 2341–2352, 2022. DOI : 10.1080/09720529.2020.1838744
- [44] Sobczyk, G., The Hyperbolic Number Plane, The College Mathematics Journal, 26(4), 268-280, 1995.
- [45] Soykan, Y., Tribonacci and Tribonacci-Lucas Sedenions. Mathematics 7(1), 74, 2019.
- [46] Shtayat, J., Al-Kateeb, A., An Encoding-Decoding algorithm based on Padovan numbers, arXiv:1907.02007, 2019.
- [47] Soykan, Y., Taşdemir, E., Göcen, M., Binomial Transform of the Generalized Third-Order Jacobsthal Sequence, Asian-European Journal of Mathematics, 15(12), 2022. <https://doi.org/10.1142/S1793557122502242>.
- [48] Soykan, Y., Taşdemir, E., Okumuş, İ., A Study on Binomial Transform of the Generalized Padovan Sequence, Journal of Science and Arts, 22(1), 63-90, 2022. <https://doi.org/10.46939/J.Sci.Arts-22.1-a06>
- [49] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, Journal of Progressive Research in Mathematics(JPRM), 14 (2), 2373-2387, 2018.
- [50] Soykan, Y., Okumuş, İ., Taşdemir, E., On Generalized Tribonacci Sedenions, Sarajevo Journal of Mathematics, 16(1), 103-122, 2020. ISSN 2233-1964, DOI: 10.5644/SJM.16.01.08
- [51] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, Communications in Mathematics and Applications, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102
- [52] Soykan, Y. Explicit Euclidean Norm, Eigenvalues, Spectral Norm and Determinant of Circulant Matrix with the Generalized Tribonacci Numbers, Earthline Journal of Mathematical Sciences, 6(1), 131-151, 2021. <https://doi.org/10.34198/ejms.6121.131151>
- [53] Sunitha, K., Sheriba. M., Gaussian Tribonacci R-Graceful Labeling of Some Tree Related Graphs, Ratio Mathematica, 44, 188-196, 2022.
- [54] Taşyurdu, Y., Polat, Y.E., Tribonacci and Tribonacci-Lucas Hybrinomials, Journal of Mathematics Research; 13(5), 2021.
- [55] Yilmaz, N., Taskara, N., Tribonacci and Tribonacci-Lucas Numbers via the Determinants of Special Matrices, Applied Mathematical Sciences, 8(39), 1947-1955,
- [56] Soykan, Y., Gümüş, M., Göcen, M., A study on dual hyperbolic generalized Pell numbers, Malaya Journal Of Matematik, 09(03), 99-116, 2021.
- [57] Soykan, Y., Taşdemir, E., Okumuş, İ., On dual hyperbolic numbers with generalized Jacobsthal numbers components, Indian J Pure Appl Math, 54, 824–840, 2023.

- [58] Soykan Y., A Study On Generalized (r,s,t)-Numbers, MathLAB Journal, 7, 101-129, 2020.
- [59] Soykan, Y., Generalized Edouard Numbers, International Journal of Advances in Applied Mathematics and Mechanics, 9(3), 41-52, 2022.
- [60] Soykan, Y., Sums and Generating Functions of Special Cases of Generalized Tribonacci Polynomials, International Journal of Advances in Applied Mathematics and Mechanics, 11(2), 80-173, 2023.
- [61] Soykan, Y., Özmen, N., Göcen, M., On Generalized Pentanacci Quaternions, Tbilisi Mathematical Journal, 13(4), 169-181, 2020.
- [62] Sivakumar, B., James, V., A Notes on Matrix Sequence of Pentanacci Numbers and Pentanacci Cubes, Communications in Mathematics and Applications, 13(2), 603–611, 2022. DOI: 10.26713/cma.v13i2.1725
- [63] Soykan Y, Tetranacci and Tetranacci-Lucas Quaternions, Asian Research Journal of Mathematics, 15(1): 1-24, 2019; Article no.ARJOM.50749.
- [64] Soykan, Y., Gaussian Generalized Tetranacci Numbers, Journal of Advances in Mathematics and Computer Science, 31(3): 1-21, Article no.JAMCS.48063, 2019.
- [65] Soykan, Y., Matrix Sequences of Tetranacci and Tetranacci-Lucas Numbers, Int. J. Adv. Appl. Math. and Mech. 7(2), 57-69, 2019, (ISSN: 2347-2529).
- [66] Soykan, Y., On Binomial Transform of the Generalized Tetranacci Sequence, International Journal of Advances in Applied Mathematics and Mechanics, 9(2), 8-27, 2021.
- [67] Soykan, Y., Binomial Transform of the Generalized Pentanacci Sequence, Asian Research Journal of Current Science, 3(1), 209-231, 2021.
- [68] Soykan, Y., Taşdemir, E., A Study On Hyperbolic Numbers With Generalized Jacobsthal Numbers Components, International Journal of Nonlinear Analysis and Applications, 13(2), 1965–1981, 2022. <http://dx.doi.org/10.22075/ijnaa.2021.22113.2328>
- [69] Soykan, Y., On Dual Hyperbolic Generalized Fibonacci Numbers, Indian J Pure Appl Math, 2021. <https://doi.org/10.1007/s13226-021-00128-2>
- [70] Şentürk, G.Y., Gürses, N., Yüce, S., Construction of Dual-Generalized Complex Fibonacci and Lucas Quaternions, Carpathian Math. Publ. 2022, 14 (2), 406-418, 2022. doi:10.15330/cmp.14.2.406-418
- [71] Ulutaş, Y.T., Toy, D., Some Equalities and Binomial Sums about the Generalized Fibonacci Number u_n , Notes on Number Theory and Discrete Mathematics, 28(2), 252–260, 2022. DOI: 10.7546/nntdm.2022.28.2.252-260
- [72] Vasanthi, S., Sivakumar, B., Jacobsthal Matrices and their Properties. Indian Journal of Science and Technology 15(5): 207-215, 2022, <https://doi.org/10.17485/IJST/v15i5.1948>
- [73] Yılmaz, N., Split Complex Bi-Periodic Fibonacci and Lucas Numbers, Commun.Fac.Sci.Univ.Ank.Ser. A1 Math. Stat. 71(1), 153–164, 2022. DOI:10.31801/cfsuasmas.704435
- [74] Yılmaz, N., The Generalized Bivariate Fibonacci and Lucas Matrix Polynomials, Mathematica Montisnigri, Vol LIII, 33-44, 2022. DOI: 10.20948/mathmontis-2022-53-5
- [75] Yılmaz, B., Soykan, Y. (2024). On Dual Hyperbolic Guglielmo Numbers. Journal of Advances in Mathematics and Computer Science, 39(4), 37–61.