

---

## Exploring Graph Density for Basic and Derived Graphs

### Abstract

This research paper explores the concept of **density of graph**, with a main focus on different graphs and their corona products. We have started with defining a few graphs like; path, cycle, ladder, kite, triangular snake, H, and circular ladder graph along with terms like order, size, and corona product of graph. Then we have investigated the density property of some basic graphs, the corona product of some basic graph with  $K_1$ , and corona product of two or more identical basic graphs. This newly derived findings can contribute to understanding some basic properties of graphs. These results can help in future research on graph theory related to density.

*Keywords: Density, Corona product, Graph theory, Properties of graph.*

## 1 Introduction

Graph theory is a fundamental branch of discrete mathematics that provides a powerful framework for modeling relationships and interactions in complex systems. Graph theory has evolved into a study with profound significance from Euler's work on the Königsberg bridge problem in the 18th century. All graphs considered in this research are assumed to be undirected, simple, and finite. **Density** is one of the essential structural property of a graph. In simple language, density of graph quantifies the ratio of number of edges available in the graph to maximum number of edges. The value of density ranges between 0 and 1. Density of graph serves as an indicator of the structural complexity and connectivity within a network. This makes it highly valuable in various real world applications. Applications includes social networks, computer networks, biological networks, and transportation systems. In general, density of graph provides meaningful insights into the resilience, efficiency, and patterns of complex systems across multiple disciplines.

---

---

---

## 2 Basic Terminology

### 2.1 Order, Size: (Sango and Jr., 2025)

The **size** of a graph  $G$  is the cardinality of its edge set and is denoted by  $|E(G)|$ .  
The **order** of a graph  $G$  is the cardinality of its vertex set and is denoted by  $|V(G)|$ .

### 2.2 Path Graph ( $P_n$ ): (Sango and Jr., 2025; West, 2002)

A **path** graph, denoted by  $P_n$ , is a graph consisting of a sequence of  $n$  distinct vertices in which each vertex(except the first and last) is connected to exactly two others, forming a single, straight-line structure, where  $n \geq 2$ .

### 2.3 Cycle Graph ( $C_n$ ): (Sango and Jr., 2025; West, 2002)

A **cycle** graph, denoted by  $C_n$ , is a graph that consists of a single cycle with  $n$  vertices. In which each vertex is connected to exactly two other vertices, forming a closed loop, where  $n \geq 3$ .

### 2.4 Ladder Graph ( $L_{n,1}$ ): (Gallian, 2000; Joshi and Parmar, 2021)

A **ladder** graph, denoted by  $L_{n,1}$ , is a graph that resembles the structure of a ladder. It is constructed by taking two parallel path graphs  $P_n$  and connecting corresponding pairs of vertices with edges, where  $n \geq 2$ .

### 2.5 Kite Graph ( $K_{n,m}$ ): (Joshi and Parmar, 2021)

A **kite** graph, denoted by  $K_{n,m}$ , is a graph formed by joining a complete graph  $K_n$  to a path graph  $P_m$  by identifying one vertex of  $K_n$  with one endpoint of  $P_m$ , where  $n \geq 3$  and  $m \geq 2$ .

### 2.6 Triangular Snake Graph ( $T_n$ ): (Parmar and Joshi, 2019)

A **triangular snake** graph, denoted by  $T_n$ , is a graph constructed by replacing each edge of a path graph  $P_n$  with a triangle, where  $n \geq 2$ .

### 2.7 H-Graph ( $H_n$ ): (Joshi and Parmar, 2021)

A **H-graph**, denoted by  $H_n$ , is a graph constructed by taking two copies of the path graph  $P_n$  and connecting corresponding pairs of middle vertices with an edge, forming a structure resembling the letter "H", where  $n \geq 3$ .

### 2.8 Circular Ladder Graph ( $CL_n$ ): (M. et al., 2014)

A **circular ladder** graph, denoted by  $CL_n$ , is constructed by connecting corresponding vertices of two  $n$ -vertex cycle graphs  $C_n$  with edges, effectively forming two concentric cycles with additional edges joining corresponding pairs of vertices, where  $n \geq 3$ .

## 2.9 Corona Product: (Gallian, 2000; Sango and Jr., 2025)

A **corona product** of two graphs  $G_1$  and  $G_2$ , denoted by  $G_1 \circ G_2$ , is constructed by taking one copy of  $G_1$  and  $|V(G_1)|$  copies of  $G_2$ , and then joining the  $i^{\text{th}}$  vertex of  $G_1$  to every vertex in the  $i^{\text{th}}$  copy of  $G_2$ , for all  $i = 1, 2, \dots, |V(G_1)|$ .

## 2.10 Density of Graph: (Sango and Jr., 2025)

For a graph  $G(V, E)$  with  $n$  vertices, the **density**  $D$  is mathematically expressed as below, where  $m$  is the total number of edges in the graph:

$$D = \frac{2m}{n(n-1)}$$

## 3 Results on Density of Some Graph

**Theorem 3.1.** (Triangular Snake Graph  $T_n$ ) (Parmar and Joshi, 2019; Sango and Jr., 2025)

For every positive integer  $n \geq 2$ , the density of the triangular snake graph  $T_n$ , denoted by  $D(T_n)$ , is precisely characterized by the below mathematical expression:

$$D(T_n) = \frac{3}{2n-1}$$

*Proof.* From the construction and schematic layout of the  $T_n$  as depicted, it can be deduced that the size of the graph is  $3(n-1)$ , while the order of the graph is  $2n-1$ .

$$\therefore D(T_n) = \frac{2m}{n(n-1)} = \frac{2 \cdot 3(n-1)}{(2n-1)(2n-1-1)} = \frac{3}{2n-1}.$$

□

**Theorem 3.2.** (H-Graph  $H_n$ ) (Joshi and Parmar, 2021; Sango and Jr., 2025)

For every positive integer  $n \geq 3$ , the density of the H-graph  $H_n$ , denoted by  $D(H_n)$ , is precisely characterized by the below mathematical expression:

$$D(H_n) = \frac{1}{n}$$

*Proof.* From the construction and schematic layout of the  $H_n$  as depicted, it can be deduced that the size of the graph is  $2n-1$ , while the order of the graph is  $2n$ .

$$\therefore D(H_n) = \frac{2m}{n(n-1)} = \frac{2 \cdot (2n-1)}{2n(2n-1)} = \frac{1}{n}.$$

□

**Theorem 3.3.** (Kite Graph  $K_{3,n}$ ) (Joshi and Parmar, 2021; Sango and Jr., 2025)

For every positive integer  $n \geq 2$ , the density of the kite graph  $K_{3,n}$ , denoted by  $D(K_{3,n})$ , is precisely characterized by the below mathematical expression:

$$D(K_{3,n}) = \frac{2}{n+2}$$

*Proof.* From the construction and schematic layout of the  $K_{3,n}$  as depicted, it can be deduced that the size of the graph is  $n + 3$ , while the order of the graph is  $n + 3$ .

$$\therefore D(K_{3,n}) = \frac{2m}{n(n-1)} = \frac{2 \cdot (n+3)}{(n+3)(n+3-1)} = \frac{2}{n+2}.$$

□

**Theorem 3.4.** (Circular Ladder Graph  $CL_n$ ) (M. et al., 2014; Sango and Jr., 2025)

For every positive integer  $n \geq 3$ , the density of the circular ladder graph  $CL_n$ , denoted by  $D(CL_n)$ , is precisely characterized by the below mathematical expression:

$$D(CL_n) = \frac{3}{2n-1}$$

*Proof.* From the construction and schematic layout of the  $CL_n$  as depicted, it can be deduced that the size of the graph is  $3n$ , while the order of the graph is  $2n$ .

$$\therefore D(CL_n) = \frac{2m}{n(n-1)} = \frac{2 \cdot 3n}{2n(2n-1)} = \frac{3}{2n-1}.$$

□

**Corollary 3.1.** ( $T_n$  and  $CL_n$ )

Density of the triangular snake graph  $T_n$  and the circular ladder graph  $CL_n$  are the same;  $n \geq 3$ .

$$D(T_n) = D(CL_n) = \frac{3}{2n-1}$$

## 4 Results on Density of Corona Product of Some Graph with $K_1$

**Theorem 4.1.** ( $P_n \circ K_1$ ) (Gallian, 2000; Sango and Jr., 2025; West, 2002)

For every positive integer  $n \geq 2$ , the density of the corona product of  $P_n$  with  $K_1$ , denoted by  $D(P_n \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(P_n \circ K_1) = \frac{1}{n}$$

*Proof.* From the construction and schematic layout of the  $P_n \circ K_1$  as depicted, it can be deduced that the size of the graph is  $2n - 1$ , while the order of the graph is  $2n$ .

$$\therefore D(P_n \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot (2n-1)}{2n(2n-1)} = \frac{1}{n}.$$

□

**Corollary 4.1.** ( $H_n$  and  $P_n \circ K_1$ )

Density of the H-graph  $H_n$  and the corona product of  $P_n$  with  $K_1$  are the same;  $n \geq 3$ .

$$D(H_n) = D(P_n \circ K_1) = \frac{1}{n}$$

**Theorem 4.2.** ( $C_n \circ K_1$ ) (Gallian, 2000; Sango and Jr., 2025; West, 2002)

For every positive integer  $n \geq 3$ , the density of the corona product of  $C_n$  with  $K_1$ , denoted by  $D(C_n \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(C_n \circ K_1) = \frac{2}{2n-1}$$

*Proof.* From the construction and schematic layout of the  $C_n \circ K_1$  as depicted, it can be deduced that the size of the graph is  $2n$ , while the order of the graph is  $2n$ .

$$\therefore D(C_n \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot 2n}{2n(2n-1)} = \frac{2}{2n-1}.$$

□

**Theorem 4.3.** ( $L_{n,1} \circ K_1$ ) (Gallian, 2000; Joshi and Parmar, 2021; Sango and Jr., 2025)  
 For every positive integer  $n \geq 2$ , the density of the corona product of  $L_{n,1}$  with  $K_1$ , denoted by  $D(L_{n,1} \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(L_{n,1} \circ K_1) = \frac{5n-2}{2n(4n-1)}$$

*Proof.* From the construction and schematic layout of the  $L_{n,1} \circ K_1$  as depicted, it can be deduced that the size of the graph is  $5n-2$ , while the order of the graph is  $4n$ .

$$\therefore D(L_{n,1} \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot (5n-2)}{4n(4n-1)} = \frac{5n-2}{2n(4n-1)}.$$

□

**Theorem 4.4.** ( $K_{3,n} \circ K_1$ ) (Gallian, 2000; Joshi and Parmar, 2021; Sango and Jr., 2025)  
 For every positive integer  $n \geq 2$ , the density of the corona product of  $K_{3,n}$  with  $K_1$ , denoted by  $D(K_{3,n} \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(K_{3,n} \circ K_1) = \frac{2}{2n+5}$$

*Proof.* From the construction and schematic layout of the  $K_{3,n} \circ K_1$  as depicted, it can be deduced that the size of the graph is  $2(n+3)$ , while the order of the graph is  $2(n+3)$ .

$$\therefore D(K_{3,n} \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot 2(n+3)}{2(n+3)(2(n+3)-1)} = \frac{2}{2n+5}.$$

□

**Theorem 4.5.** ( $T_n \circ K_1$ ) (Gallian, 2000; Parmar and Joshi, 2019; Sango and Jr., 2025)  
 For every positive integer  $n \geq 2$ , the density of the corona product of  $T_n$  with  $K_1$ , denoted by  $D(T_n \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(T_n \circ K_1) = \frac{5n-4}{(2n-1)(4n-3)}$$

*Proof.* From the construction and schematic layout of the  $T_n \circ K_1$  as depicted, it can be deduced that the size of the graph is  $5n-4$ , while the order of the graph is  $2(2n-1)$ .

$$\therefore D(T_n \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot (5n-4)}{2(2n-1)(2(2n-1)-1)} = \frac{5n-4}{(2n-1)(4n-3)}.$$

□

**Theorem 4.6.** ( $H_n \circ K_1$ ) (Gallian, 2000; Joshi and Parmar, 2021; Sango and Jr., 2025)  
 For every positive integer  $n \geq 3$ , the density of the corona product of  $H_n$  with  $K_1$ , denoted by  $D(H_n \circ K_1)$ , is precisely characterized by the below mathematical expression:

$$D(H_n \circ K_1) = \frac{1}{2n}$$

*Proof.* From the construction and schematic layout of the  $H_n \circ K_1$  as depicted, it can be deduced that the size of the graph is  $4n - 1$ , while the order of the graph is  $4n$ .

$$\therefore D(H_n \circ K_1) = \frac{2m}{n(n-1)} = \frac{2 \cdot (4n-1)}{4n(4n-1)} = \frac{1}{2n}.$$

□

## 5 Results on Density of Corona Product of two same Graph

**Theorem 5.1.** ( $T_n \circ T_n$ ) (Gallian, 2000; Parmar and Joshi, 2019; Sango and Jr., 2025)

For every positive integer  $n \geq 2$ , the density of the corona product of two  $T_n$  graphs, denoted by  $D(T_n \circ T_n)$ , is precisely characterized by the below mathematical expression:

$$D(T_n \circ T_n) = \frac{10n(n-1) + 1}{n(2n-1)(4n^2 - 2n - 1)}$$

*Proof.* From the construction and schematic layout of the  $T_n \circ T_n$  as depicted, it can be deduced that the size of the graph is  $10n^2 - 10n + 1$ , while the order of the graph is  $2n(2n - 1)$ .

$$\therefore D(T_n \circ T_n) = \frac{2m}{n(n-1)} = \frac{2 \cdot (10n^2 - 10n + 1)}{2n(2n-1)(2n(2n-1) - 1)} = \frac{10n(n-1) + 1}{n(2n-1)(4n^2 - 2n - 1)}.$$

□

**Example 5.1.** For the graph  $T_3 \circ T_3$




Figure 1: Graph of  $T_3 \circ T_3$

Using definition,

$$D(T_3 \circ T_3) = \frac{2m}{n(n-1)} = \frac{2 \cdot 61}{30 \cdot 29} = \frac{61}{435}$$

Using theorem,

$$D(T_3 \circ T_3) = \frac{10n(n-1) + 1}{n(2n-1)(4n^2 - 2n - 1)} = \frac{10 \cdot 3(3-1) + 1}{3(2 \cdot 3 - 1)(4 \cdot 3^2 - 2 \cdot 3 - 1)} = \frac{61}{435}$$

**Theorem 5.2.**  $(H_n \circ H_n)$  (Gallian, 2000; Joshi and Parmar, 2021; Sango and Jr., 2025)  
 For every positive integer  $n \geq 3$ , the density of the corona product of two  $H_n$  graphs, denoted by  $D(H_n \circ H_n)$ , is precisely characterized by the below mathematical expression:

$$D(H_n \circ H_n) = \frac{8n^2 - 1}{n(2n + 1)(4n^2 + 2n - 1)}$$

*Proof.* From the construction and schematic layout of the  $H_n \circ H_n$  as depicted, it can be deduced that the size of the graph is  $8n^2 - 1$ , while the order of the graph is  $2n(2n + 1)$ .

$$\therefore D(H_n \circ H_n) = \frac{2m}{n(n - 1)} = \frac{2 \cdot (8n^2 - 1)}{2n(2n + 1)(2n(2n + 1) - 1)} = \frac{8n^2 - 1}{n(2n + 1)(4n^2 + 2n - 1)}.$$

□

**Example 5.2.** For the graph  $H_3 \circ H_3$




Figure 2: Graph of  $H_3 \circ H_3$

Using definition,

$$D(H_3 \circ H_3) = \frac{2m}{n(n - 1)} = \frac{2 \cdot 71}{42 \cdot 41} = \frac{71}{861}$$

Using theorem,

$$D(H_3 \circ H_3) = \frac{8n^2 - 1}{n(2n + 1)(4n^2 + 2n - 1)} = \frac{8 \cdot 3^2 - 1}{3(2 \cdot 3 + 1)(4 \cdot 3^2 + 2 \cdot 3 - 1)} = \frac{71}{861}$$

**Theorem 5.3.**  $(K_{3,n} \circ K_{3,n})$  (Gallian, 2000; Joshi and Parmar, 2021; Sango and Jr., 2025)  
 For every positive integer  $n \geq 2$ , the density of the corona product of two  $K_{3,n}$  graphs, denoted by  $D(K_{3,n} \circ K_{3,n})$ , is precisely characterized by the below mathematical expression:

$$D((K_{3,n} \circ K_{3,n})) = \frac{2(2n + 7)}{(n + 4)(n^2 + 7n + 11)}$$

*Proof.* From the construction and schematic layout of the  $K_{3,n} \circ K_{3,n}$  as depicted, it can be deduced that the size of the graph is  $(n + 3)(2n + 7)$ , while the order of the graph is  $(n + 3)(n + 4)$ .

$$\therefore D(K_{3,n} \circ K_{3,n}) = \frac{2m}{n(n - 1)} = \frac{2(n + 3)(2n + 7)}{(n + 3)(n + 4)((n + 3)(n + 4) - 1)} = \frac{2(2n + 7)}{(n + 4)(n^2 + 7n + 11)}.$$

□

**Example 5.3.** For the graph  $K_{3,2} \circ K_{3,2}$




Figure 3: Graph of  $K_{3,2} \circ K_{3,2}$

Using definition,

$$D(K_{3,2} \circ K_{3,2}) = \frac{2m}{n(n-1)} = \frac{2 \cdot 55}{30 \cdot 29} = \frac{11}{87}$$

Using theorem,

$$D(K_{3,2} \circ K_{3,2}) = \frac{2(2n+7)}{(n+4)(n^2+7n+11)} = \frac{2(2 \cdot 2+7)}{(2+4)(2^2+7 \cdot 2+11)} = \frac{11}{87}$$

**Theorem 5.4.**  $((P_n \circ K_1) \circ (P_n \circ K_1))$  (Gallian, 2000; Sango and Jr., 2025; West, 2002)

For every positive integer  $n \geq 2$ , the density of the corona product of two  $P_n \circ K_1$  graphs, denoted by  $D((P_n \circ K_1) \circ (P_n \circ K_1))$ , is precisely characterized by the below mathematical expression:

$$D((P_n \circ K_1) \circ (P_n \circ K_1)) = \frac{8n^2 - 1}{n(2n+1)(4n^2+2n-1)}$$

*Proof.* From the construction and schematic layout of the  $(P_n \circ K_1) \circ (P_n \circ K_1)$  as depicted, it can be deduced that the size of the graph is  $8n^2 - 1$ , while the order of the graph is  $2n(2n+1)$ .

$$\therefore D((P_n \circ K_1) \circ (P_n \circ K_1)) = \frac{2m}{n(n-1)} = \frac{2 \cdot (8n^2 - 1)}{2n(2n+1)(2n(2n+1) - 1)} = \frac{8n^2 - 1}{n(2n+1)(4n^2+2n-1)}$$

□

**Example 5.4.** For the graph  $(P_3 \circ K_1) \circ (P_3 \circ K_1)$




Figure 4: Graph of  $(P_3 \circ K_1) \circ (P_3 \circ K_1)$

Using definition,

$$D((P_3 \circ K_1) \circ (P_3 \circ K_1)) = \frac{2m}{n(n-1)} = \frac{2 \cdot 71}{42 \cdot 41} = \frac{71}{861}$$

Using theorem,

$$D((P_3 \circ K_1) \circ (P_3 \circ K_1)) = \frac{8n^2 - 1}{n(2n+1)(4n^2 + 2n - 1)} = \frac{8 \cdot 3^2 - 1}{3(2 \cdot 3 + 1)(4 \cdot 3^2 + 2 \cdot 3 - 1)} = \frac{71}{861}$$

**Corollary 5.1.**  $(H_n \circ H_n \text{ and } (P_n \circ K_1) \circ (P_n \circ K_1))$

Density of the two corona product of  $H_n$  and the two corona product of  $P_n \circ K_1$  are the same;  $n \geq 3$ .

$$D(H_n \circ H_n) = D((P_n \circ K_1) \circ (P_n \circ K_1)) = \frac{8n^2 - 1}{n(2n+1)(4n^2 + 2n - 1)}$$

## 6 CONCLUSION

This study presents the density analysis of the corona product of various standard graphs with the complete graph  $K_1$  as well as the corona product of two identical graphs. Furthermore, the investigation of the corona product of two similar graphs highlights the significant structural changes that occur in terms of connectivity and compactness. These findings may also be helpful for researchers working in areas where graph density plays a key role, such as network analysis, computer science, and structural modeling. Future work may involve extending this approach to other graph classes or analyzing the practical relevance of these density changes in real-world networks.

## References

Joseph A. Gallian. Dynamic survey of graph labeling. The Electronic Journal of Combinatorics, 2000. doi: 10.37236/11668.

---

Jayshree Ratilal Joshi and Dharamvirsinh Parmar. Hk-cordial labeling of some graph and its corona graph. International Journal of Aquatic Science, 12(02):1519–1534, 2021. ISSN 2008-8019.

Joice Punitha M., S. Rajakumari, and Indra Rajasingh. Skew chromatic index of circular ladder graphs. Annals of Pure and Applied Mathematics, 8(2):1–7, 2014. ISSN 2279-087X(P), 2279-0888(online).

Dharamvirsinh Parmar and Jayshree Ratilal Joshi. Hk-cordial labeling of triangular snake graph. Journal of Applied Science and Computations, VI(III):2118–2123, 2019. ISSN 1076-5131.

Racma L. Sango and Isagani S. Cabahug Jr. On the density of some graphs and corona graphs. Asian Research Journal of Mathematics, 21(6):1–6, 2025. ISSN 456-477X.

Douglas B. West. Introduction to graph theory. Pearson Education (Singapore) Pte. Ltd., 2002. ISBN 81-7808-830-4.