

A Novel Formula for the nth Power of a 2x2 Matrix with Applications to Linear Recurrences and the Birth-Death

**Original Research
Article**

Abstract

After exploring a robust method for calculating the nth power of a 2x2 matrix A^n without resorting to diagonalization techniques, we have provided a unique formula for the nth power of any type of 2x2 matrix, diagonalizable or not, with or without two distinct eigenvalues, and whether these eigenvalues are real or not.

In this article, we will discuss applications of the above, such as:

- 1) Application of the nth power of 2x2 matrices to solve second-order linear recurrence problems;
- 2) The (unsolved) problem of the birth and death process (BDP) with constant coefficients and an infinite number of states. That is to say we will apply our matrix power formula to determine the eigenvectors of the generating matrix A of this process.

Keywords: Power of a matrix; Recurrent linear sequences of order 2; The birth and death process

2010 Mathematics Subject Classification: 15A24; 60J27; 39A05

1 A New Approach to Matrix Powers with Wide Applications

(Introduction):

In 1992 *Kenneth S. Williams* (see (8)) gives us a general formula by discussing the equality or not of the eigenvalues and working with the Matrices $X = \frac{A-\beta I}{\alpha-\beta}$, $Y = \frac{A-\alpha I}{\beta-\alpha}$ et $Z = A - \alpha I$, which is

$$A^n = \begin{cases} \alpha^n \left(\frac{A-\beta I}{\alpha-\beta} \right) + \beta^n \left(\frac{A-\alpha I}{\beta-\alpha} \right) & \text{if } \alpha \neq \beta \\ \alpha^{n-1} (nA - (n-1)\alpha I) & \text{if } \alpha = \beta \end{cases}$$

In the article (9), they study and determine a general formula for the nth power of 2x2 matrices using the eigenvalues of A, and give some Excel programming examples.

On the other hand, in paper (13) the authors seek to determine other combinatorial identities deriving from the power n of a 2×2 matrix.

But the real problem for mathematicians and even computer scientists is determining the exponential of a matrix (see (10), (11), (12), (17)) since this problem is often found especially in systems of differential equations of the type $X(t)' = AX(t)$ (see (14), (15), (16))

This paper presents a unique and efficient method for calculating the n th power of a 2×2 matrix (A^n) without resorting to the traditional diagonalization method. Although diagonalization is a widely used and effective technique for certain classes of matrices, its dependence on distinct and real eigenvalues can be limiting. Our approach provides a single, unified formula applicable to "any" 2×2 matrix, regardless of its characteristics. This includes non-diagonalizable matrices, matrices with repeated eigenvalues, and those with complex eigenvalues. By providing a general and universally applicable solution, we eliminate the need for case-by-case analysis, thus simplifying and streamlining a fundamental operation in linear algebra.

This will surely help solve many important problems, and to begin with, we will apply it to the following two famous problems:

1. Solving Second-Order Linear Recurrence Sequences: We show how the shorthand solution of the n th term of a second-order linear recurrence relation, such as the Fibonacci sequence, can be directly derived using our general formula for A^n . By representing the recurrence relation as a matrix equation, we transform a recursive problem into a direct and explicit computation. This method offers a more elegant and computationally efficient alternative to traditional characteristic equation methods, especially for large values of n .

2. Treating the Birth-Death Process (BDP) with an Infinite Number of States: We apply our methodology to the notoriously complex problem of the birth-death process (BDP) with constant coefficients and an infinite number of states. Although the BDP is a cornerstone of stochastic modeling in fields such as queueing theory, population dynamics, and epidemiology, the analytical solution of its state probabilities in the case of an infinite number of states remains an open problem. Our approach to the computation of matrix powers, a key element in solving the BDP system of differential equations (as is the case for the exponential matrix), offers a new perspective and a potential path toward an analytical solution. We believe this work could open new avenues for research on this long-unsolved problem. The use of spectral methods (using eigenvalues and eigenvectors) to study birth and death processes, and especially to determine their eigenvectors, was pioneered by S. Karlin and J. McGregor (see (6) and (7)). They defined a sequence of polynomials $Q_k(x)$ such that $Q_0(x) = 1$ and $xQ = AQ$. These polynomials are orthogonal with respect to a probability measure ψ on the interval $[0, +\infty[$ and

$$P_{i,j}(t) = \frac{\int_0^{\infty} e^{-xt} Q_i(x) Q_j(x) d\psi(x)}{\int_0^{\infty} Q_j(x)^2 d\psi(x)}$$

Which is a nice formula, but complicated in practice. And so it is for this purpose that we apply our formula to search for these eigenvectors.

2 The power of any (diagonalizable or non-diagonalizable) 2x2 matrix:

Theorem 2.1. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ a 2x2 matrix, so $\forall n \geq 2$ ($n \in \mathbb{N}$)

$$A^n = \begin{pmatrix} a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} & b \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ c \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} & \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \end{pmatrix} \quad (i)$$

with α_1 and α_2 solutions of $|A - \alpha I| = 0$.

Proof. We will demonstrate our formula using recurrence.

1) For $n = 2$, we will show that we actually have

$$A^2 = \begin{pmatrix} a^2 + bc & b(a+d) \\ c(a+d) & d^2 + bc \end{pmatrix}$$

$$A^2 = \begin{pmatrix} a \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} - \alpha_1 \alpha_2 & b \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} \\ c \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} & \sum_{k=0}^2 \alpha_1^k \alpha_2^{2-k} - a \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} \end{pmatrix}$$

we must therefore show the following 4 equalities

$$\begin{cases} a^2 + bc = a \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} - \alpha_1 \alpha_2 \\ d^2 + bc = \sum_{k=0}^2 \alpha_1^k \alpha_2^{2-k} - a \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} \\ b(a+d) = b \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} \\ c(a+d) = c \sum_{k=0}^1 \alpha_1^k \alpha_2^{1-k} \end{cases}$$

what does it mean

$$\begin{cases} a^2 + bc = a(\alpha_2 + \alpha_1) - \alpha_1 \alpha_2 \\ d^2 + bc = \alpha_2^2 + \alpha_1 \alpha_2 + \alpha_1^2 - a(\alpha_2 + \alpha_1) \\ b(a+d) = b(\alpha_2 + \alpha_1) \\ c(a+d) = c(\alpha_2 + \alpha_1) \end{cases}$$

α_1 and α_2 solutions of $|A - \alpha I| = 0$, so

$$\begin{cases} \alpha_1 + \alpha_2 = \text{Tr}(A) = a + d \\ \alpha_1 \alpha_2 = |A| = ad - bc \end{cases}$$

So

$$a(\alpha_2 + \alpha_1) - \alpha_1 \alpha_2 = a(a+d) - (ad - bc) = a^2 + ad - ad + bc = a^2 + bc$$

$$\alpha_2^2 + \alpha_1 \alpha_2 + \alpha_1^2 - a(\alpha_2 + \alpha_1) = (\alpha_1 + \alpha_2)^2 - \alpha_1 \alpha_2 - a(\alpha_2 + \alpha_1) \implies$$

$$\alpha_2^2 + \alpha_1 \alpha_2 + \alpha_1^2 - a(\alpha_2 + \alpha_1) = (a+d)^2 - (ad - bc) - a(a+d) = d^2 + bc$$

$$b(\alpha_2 + \alpha_1) = b(a+d)$$

$$c(\alpha_2 + \alpha_1) = c(a+d)$$

hence the result is true for $n = 2$.

2) Let's suppose

$$A^n = \begin{pmatrix} a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} & b \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ c \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} & \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \end{pmatrix}$$

3) Let us show that

$$A^{n+1} = A^n \cdot A$$

$$A^{n+1} = \begin{pmatrix} a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - \sum_{k=0}^{n-1} \alpha_1^{k+1} \alpha_2^{n-k} & b \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \\ c \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} & \sum_{k=0}^{n+1} \alpha_1^k \alpha_2^{n+1-k} - a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \end{pmatrix}$$

We have $A^{n+1} = A^n \cdot A =$

$$= \begin{pmatrix} a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} & b \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ c \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} & \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \end{pmatrix} A$$

$$A^{n+1} = \begin{pmatrix} x & y \\ z & t \end{pmatrix}$$

$$x = a^2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} + cb \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

$$y = ba \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - b \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} + db \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

$$z = ac \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + c \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - ca \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

$$t = bc \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + d \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - da \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

$$\Rightarrow \begin{cases} x = (a^2 + cb) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \\ y = b \left[(a + d) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \right] \\ z = c \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \\ t = (bc - da) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + d \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \end{cases}$$

\Rightarrow

$$x = [a(\alpha_1 + \alpha_2) - \alpha_1 \alpha_2] \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k}$$

$$y = b \left[(\alpha_1 + \alpha_2) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \right]$$

$$z = c \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k}$$

$$t = -\alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + d \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k}$$

because $\alpha_1 + \alpha_2 = a + d$, $\alpha_1 \alpha_2 = ad - bc$, $a(\alpha_1 + \alpha_2) - \alpha_1 \alpha_2 = a^2 + cb$

$$A^{n+1} = \begin{pmatrix} a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - \alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} & b \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \\ c \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} & \sum_{k=0}^{n+1} \alpha_1^k \alpha_2^{n+1-k} - a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \end{pmatrix}$$

because

$$x = [a(\alpha_1 + \alpha_2) - \alpha_1 \alpha_2] \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k}$$

$$= a(\alpha_1 + \alpha_2) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k}$$

$$= a\alpha_1 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + a\alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - a \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k}$$

$$= a\alpha_1^n + a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-k} - \alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

$$\begin{aligned}
x &= a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - \alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} ; \\
y &= b \left[(\alpha_1 + \alpha_2) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \right] \\
y &= b \left[\sum_{k=0}^{n-1} \alpha_1^{k+1} \alpha_2^{n-1-k} + \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \right] \\
y &= b \left[\alpha_1^n + \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-k} \right] = b \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k}
\end{aligned}$$

and

$$\begin{aligned}
t &= -\alpha_1 \alpha_2 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + d \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \\
t &= -\sum_{k=0}^{n-1} \alpha_1^{k+1} \alpha_2^{n-k} + (\alpha_1 + \alpha_2 - a) \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \text{ (as } \alpha_1 + \alpha_2 = a + d) \\
t &= -\sum_{k=0}^{n-1} \alpha_1^{k+1} \alpha_2^{n-k} - a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} + \sum_{k=0}^n \alpha_1^{k+1} \alpha_2^{n-k} + \sum_{k=0}^n \alpha_1^k \alpha_2^{n+1-k} \\
t &= -a \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} + \sum_{k=0}^{n+1} \alpha_1^k \alpha_2^{n+1-k}
\end{aligned}$$

hence the result. \square

rem 2.1. Let $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ a 2×2 matrix and α_1 and α_2 solutions of $|A - \alpha I| = 0$.

1) If $\alpha_1 = \alpha_2 = \alpha$, so

$$A^n = \left(\frac{a+d}{2} \right)^{n-1} \begin{pmatrix} \frac{(n+1)a - (n-1)d}{2} & nb \\ nc & \frac{(n+1)d - (n-1)a}{2} \end{pmatrix}, \forall n \geq 2 \ (n \in \mathbb{N})$$

with $\alpha = \frac{a+d}{2}$ and $(a+d)^2 = 4(ad-bc)$ ($(a-d)^2 = -4bc$)

And

$$A^n = n\alpha^{n-1}A - (n-1)\alpha^n I$$

2) If $\alpha_1 \neq \alpha_2$ so $\forall n \geq 2 \ (n \in \mathbb{N})$

$$\begin{aligned}
A^n &= \frac{1}{\alpha_1 - \alpha_2} \begin{pmatrix} (a - \alpha_2)\alpha_1^n - (a - \alpha_1)\alpha_2^n & b(\alpha_1^n - \alpha_2^n) \\ c(\alpha_1^n - \alpha_2^n) & (a - \alpha_2)\alpha_2^n - (a - \alpha_1)\alpha_1^n \end{pmatrix} \\
A^n - \alpha_2^n I &= \frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} (A - \alpha_2 I) \text{ and } A^n - \alpha_1^n I = \frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} (A - \alpha_1 I) \\
A^n &= \left(\frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} \right) A - \alpha_1 \alpha_2 \left(\frac{\alpha_1^{n-1} - \alpha_2^{n-1}}{\alpha_1 - \alpha_2} \right) I
\end{aligned}$$

Proof. 1) $A^n = \alpha^{n-1} \begin{pmatrix} na - (n-1)\alpha & nb \\ nc & (n+1)\alpha - na \end{pmatrix}$

And

$$\begin{aligned}
A^n &= \alpha^{n-1} \begin{pmatrix} na - (n-1)\alpha & nb \\ nc & nd + (n+1)\alpha - n(a+d) \end{pmatrix} \\
A^n &= \alpha^{n-1} \begin{pmatrix} na - (n-1)\alpha & nb \\ nc & nd + (n+1)\alpha - 2n\alpha \end{pmatrix} \\
A^n &= \alpha^{n-1} \begin{pmatrix} na - (n-1)\alpha & nb \\ nc & nd + (1-n)\alpha \end{pmatrix} \\
A^n &= \alpha^{n-1} \begin{pmatrix} na & nb \\ nc & nd \end{pmatrix} + \alpha^{n-1} \begin{pmatrix} -(n-1)\alpha & 0 \\ 0 & -(n-1)\alpha \end{pmatrix}
\end{aligned}$$

$$A^n = n\alpha^{n-1}A - (n-1)\alpha^n I$$

2)

$$A^n = \frac{1}{\alpha_1 - \alpha_2} \begin{pmatrix} a(\alpha_1^n - \alpha_2^n) - \alpha_1\alpha_2(\alpha_1^{n-1} - \alpha_2^{n-1}) & b(\alpha_1^n - \alpha_2^n) \\ c(\alpha_1^n - \alpha_2^n) & (\alpha_1^{n+1} - \alpha_2^{n+1}) - a(\alpha_1^n - \alpha_2^n) \end{pmatrix}$$

□

3 Applications:

3.1 Linear recurrent sequence order 2:

A second-order linear recurrence sequence is a sequence in which each term is defined as a linear combination of the two preceding terms. In simpler terms, to find the next number in the sequence, simply multiply the two preceding numbers by constants and add the results. A classic example is the Fibonacci sequence, where each number is the sum of the two preceding numbers.

$$u_{n+1} = au_n + bu_{n-1} \text{ with } (u_1, u_0) \in IK^2 \text{ and } (a, b) \in IK^2$$

$IK = IR$ or \mathbb{C}

$$\Leftrightarrow \begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u_n \\ u_{n-1} \end{pmatrix} \text{ and } (u_1, u_0) \in IK^2$$

$$\Leftrightarrow \begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}^n \begin{pmatrix} u_1 \\ u_0 \end{pmatrix} \text{ and } (u_1, u_0) \in IK^2$$

$$\Leftrightarrow \begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = \begin{pmatrix} x_n & y_n \\ z_n & t_n \end{pmatrix} \begin{pmatrix} u_1 \\ u_0 \end{pmatrix} \text{ and } (u_1, u_0) \in IK^2$$

such

$$\begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}^n = \begin{pmatrix} x_n & y_n \\ z_n & t_n \end{pmatrix}$$

So, according to our formula (i) "the n-th Power of 2x2 Matrix" in the first theorem

$$\begin{cases} x_n = a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \\ y_n = b \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ z_n = \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ t_n = \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \end{cases}$$

with α_1 and α_2 are solutions of $\alpha^2 - a\alpha - b = 0$

rem 3.1. $\alpha_1 + \alpha_2 = a$ and $\alpha_1\alpha_2 = -b$

Theorem 3.1. Let $u_{n+1} = au_n + bu_{n-1}$ with $(u_1, u_0) \in IK^2$, so

$$u_n = (u_1 - au_0) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + u_0 \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k}$$

such α_1 and α_2 are solutions of $\alpha^2 - a\alpha - b = 0$.

Proof. $\begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = \begin{pmatrix} a & b \\ 1 & 0 \end{pmatrix}^n \begin{pmatrix} u_1 \\ u_0 \end{pmatrix} = \begin{pmatrix} x_n & y_n \\ z_n & t_n \end{pmatrix} \begin{pmatrix} u_1 \\ u_0 \end{pmatrix}$

$$\implies \begin{pmatrix} u_{n+1} \\ u_n \end{pmatrix} = \begin{pmatrix} x_n u_1 + y_n u_0 \\ z_n u_1 + t_n u_0 \end{pmatrix}$$

$$\text{such } \begin{cases} x_n = a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} - \sum_{k=0}^{n-2} \alpha_1^{k+1} \alpha_2^{n-1-k} \\ y_n = b \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ z_n = \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \\ t_n = \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \end{cases}$$

$$\text{so, } u_n = z_n u_1 + t_n u_0 = u_1 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + u_0 \left(\sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} - a \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} \right)$$

$$\implies u_n = (u_1 - a u_0) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + u_0 \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k}. \quad \square$$

rem 3.2. 1st case: If $a^2 + 4b = 0$, so $\alpha_1 = \alpha_2 = \frac{a}{2}$ and

$$\begin{aligned} u_n &= [(u_1 + (1-a)u_0)n + u_0] \alpha^{n-1} \\ u_n &= [n(u_0 + u_1) + (1-2n)\alpha u_0] \alpha^{n-1} \end{aligned}$$

2nd case: If $a^2 + 4b \neq 0$, so $\alpha_1 \neq \alpha_2 = \frac{a}{2}$ and

$$u_n = \frac{1}{\alpha_1 - \alpha_2} [(u_1 - a u_0)(\alpha_1^n - \alpha_2^n) + u_0(\alpha_1^{n+1} - \alpha_2^{n+1})]$$

Proof. 1)

$$u_n = (u_1 - a u_0) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + u_0 \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \implies$$

$$u_n = (u_1 - a u_0) n \alpha^{n-1} + u_0 (n+1) \alpha^n$$

$$\text{since } \alpha_1 = \alpha_2 \implies \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} = \sum_{k=0}^{n-1} \alpha_2^{n-1} = \alpha_2^{n-1} \sum_{k=0}^{n-1} 1 = n \alpha_2^{n-1} \text{ and } \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} = (n+1) \alpha_2^n$$

2)

$$\begin{aligned} u_n &= (u_1 - a u_0) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + u_0 \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \\ &= (u_1 - a u_0) \left(\frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} \right) + u_0 \left(\frac{\alpha_1^{n+1} - \alpha_2^{n+1}}{\alpha_1 - \alpha_2} \right) \\ &= \frac{1}{\alpha_1 - \alpha_2} [(u_1 - a u_0)(\alpha_1^n - \alpha_2^n) + u_0(\alpha_1^{n+1} - \alpha_2^{n+1})] \end{aligned}$$

□

3.2 The birth-death process (BDP):

The birth-death process (BDP) is a classic example of a continuous-time, discrete-state Markov process that is used to model population growth over time. It was introduced by Feller (1938) (5) and has been used in various fields, such as biology, physics, ecology, queuing theory, ...

Mathematically, birth and death processes are often modeled using systems of differential equations, called direct Kolmogorov equations. These equations describe the evolution of the probability distribution of the number of individuals in the system over time. However, these equations can be notoriously

difficult to solve analytically or even impossible, especially for complex systems with non-constant birth and death rates.

As a result, researchers often resort to approximation methods or numerical simulations to study birth-death processes (see (3)). These methods can provide valuable insights into the behavior of these systems, but they also have limitations. Therefore, there is a constant search for new methods to resolve these equations.

The use of spectral methods to study birth and death processes was pioneered by S. Karlin and J. McGregor (see (6) and (7)). They defined a sequence of polynomials $Q_k(x)$ such that $Q_0(x) = 1$ and $xQ = AQ$.

This article focuses on general birth and death processes with an infinite number of states:

Postulates. The system changes only through transitions from states to their nearest neighbors (from E_n to E_{n+1} or E_{n-1} if $n \geq 1$, but from E_0 to E_1 only). If at epoch t the system is in state E_n , the probability that between t and $t+h$ the transition $E_n \rightarrow E_{n+1}$ occurs equals $\lambda_n h + o(h)$, and the probability of $E_n \rightarrow E_{n-1}$ (if $n \geq 1$) equals $\mu_n h + o(h)$. The probability that during $(t, t+h)$ more than one change occurs is $o(h)$. ((4) page 454).

This paper builds upon our previous work (1) and (2) which laid the foundations for this approach. However, the current study extends and refines these methods, making significant new contributions, especially by applying our formula that gives the n th power of any 2×2 matrix.

Let $(X_t)_{t \in \mathbb{R}^+}$ be the discrete and homogeneous "Birth and Death" stochastic process such $\forall (i, j) \in \mathbb{N}^2$

$$P_i(t) = P(X_t = i), P(t) = \begin{pmatrix} P_1(t) \\ P_2(t) \\ \vdots \\ P_n(t) \\ \vdots \end{pmatrix}$$

$$\text{with } \sum_{i \geq 1} P(X_t = i) = 1, P(0) = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ \vdots \end{pmatrix} \text{ and}$$

$$\forall (i, j) \in \mathbb{N}^2$$

$$P_{ij}(\Delta t) = P(X_{t+\Delta t} = j / X_t = i) = \begin{cases} \lambda_i \Delta t + o(\Delta t); & \text{if } j = i + 1 \\ \mu_i \Delta t + o(\Delta t); & \text{if } j = i - 1 \\ 1 - (\lambda_i + \mu_i) \Delta t + o(\Delta t); & \text{if } j = i \\ o(\Delta t); & \text{if } |j - i| \geq 2 \end{cases}$$

λ_i is the birth rate and μ_i is the death rate.

Proposition 3.1. *Let $(X_t)_{t \in \mathbb{R}^+}$ be the discrete and homogeneous "Birth and Death" stochastic process with conditions (1), so $P(t) = (P_0(t), P_1(t), \dots)^t = (P_i(t))_{i \geq 1}$ is solution of the following linear differential equations system.*

$$\begin{cases} P_1'(t) = -(\lambda_1 + \mu_1) P_1(t) + \mu_2 P_2(t) \\ P_j'(t) = \lambda_{j-1} P_{j-1}(t) - (\lambda_j + \mu_j) P_j(t) + \mu_{j+1} P_{j+1}(t) \quad \forall j \geq 2 \\ \dots \end{cases}$$

and

$$P'(t) = AP(t)$$

called the backward equation.

with

$$A = \begin{pmatrix} -(\lambda_1 + \mu_1) & \mu_2 & 0 & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ \lambda_1 & \ddots & \ddots & \ddots & & & & & \\ 0 & \ddots & \ddots & \ddots & \ddots & & & & \\ \vdots & \ddots & \ddots & \ddots & \ddots & & & & \\ \vdots & & 0 & \lambda_{k-1} & -(\lambda_k + \mu_k) & \mu_{k+1} & 0 & & \\ \vdots & & & \ddots & \ddots & \ddots & \ddots & \ddots & \\ \vdots & & & & \ddots & \ddots & \ddots & \ddots & \\ \vdots & & & & & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

$$\& P(0) = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ \vdots \\ \vdots \end{pmatrix}$$

Proof.

$$\begin{aligned} P(X_{t+\Delta t} = j) &= \sum_{i \in \mathbb{N}} P(X_{t+\Delta t} = j / X_t = i) P(X_t = i) \\ &= P(X_{t+\Delta t} = j / X_t = j-1) P(X_t = j-1) + \\ &\quad + P(X_{t+\Delta t} = j / X_t = j+1) P(X_t = j+1) + \\ &\quad + P(X_{t+\Delta t} = j / X_t = j) P(X_t = j) + \\ &\quad + \sum_{\substack{i \in \mathbb{N} \\ i \neq j \\ i \neq j \pm 1}} P(X_{t+\Delta t} = j / X_t = i) P(X_t = i) \\ &= \lambda_{j-1} \Delta t P(X_t = j-1) + \mu_{j+1} \Delta t P(X_t = j+1) + \\ &\quad + [1 - (\lambda_j + \mu_j) \Delta t] P(X_t = j) + o(\Delta t) \end{aligned}$$

$$\begin{aligned} P(X_{t+\Delta t} = j) - P(X_t = j) &= \lambda_{j-1} P(X_t = j-1) \Delta t + \\ &\quad + \mu_{j+1} P(X_t = j+1) \Delta t + \\ &\quad - (\lambda_j + \mu_j) P(X_t = j) \Delta t + o(\Delta t) \end{aligned}$$

$$\begin{aligned} \frac{P(X_{t+\Delta t} = j) - P(X_t = j)}{\Delta t} &= \lambda_{j-1} P(X_t = j-1) + \mu_{j+1} P(X_t = j+1) + \\ &\quad - (\lambda_j + \mu_j) P(X_t = j) + \\ &\quad + \frac{o(\Delta t)}{\Delta t} \end{aligned}$$

$$P'_j(t) = \lambda_{j-1} P_{j-1}(t) - (\lambda_j + \mu_j) P_j(t) + \mu_{j+1} P_{j+1}(t), \forall j \geq 2$$

(When $\Delta t \rightarrow 0$)

...

□

let β be an eigenvalue of the matrix A and $x = (x_k)_{k \in \mathbb{N}^*}$ and an associated right eigenvector, so $Ax = \beta x$

$$\left\{ \begin{array}{l} -(\lambda_1 + \mu_1)x_1 + \mu_2x_2 = \beta x_1 \\ \dots \\ \lambda_{k-1}x_{k-1} - (\lambda_k + \mu_k)x_k + \mu_{k+1}x_{k+1} = \beta x_k \quad 2 \leq k \\ \dots \end{array} \right.$$

$$\mu_{k+1}x_{k+1} - (\lambda_k + \mu_k + \beta)x_k + \lambda_{k-1}x_{k-1} = 0, \quad \forall 1 \leq k$$

with $x_0 = 0$ or $\lambda_0 = 0$.

3.2.1 The case of constant birth and death rates: $\lambda_k = \lambda$ and $\mu_k = \mu$

The associated right eigenvectors ($Ax = \beta x$, $x = (x_k)_{k \in \mathbb{N}^*}$) will therefore verify the following system

$$(E_k): \quad \mu x_{k+1} - (\lambda + \mu + \beta)x_k + \lambda x_{k-1} = 0, \quad \forall 1 \leq k$$

with $x_0 = 0$ and $x_1 \in IK$.

If $\mu \neq 0$, (E_k) is a second-order linear recurrence sequence with constant coefficients.

$$(E_k) \iff x_{k+1} = \left(\frac{\lambda + \mu + \beta}{\mu} \right) x_k - \frac{\lambda}{\mu} x_{k-1}, \quad \forall 1 \leq k$$

Applying the results of the last subsection concerning second-order linear recurrence sequences with constant coefficients with $a = \frac{\lambda + \mu + \beta}{\mu}$ and $b = -\frac{\lambda}{\mu}$, we will have

$$x_n = u_n = (x_1 - ax_0) \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k} + x_0 \sum_{k=0}^n \alpha_1^k \alpha_2^{n-k} \implies \forall n \geq 1$$

$$x_n = x_1 \sum_{k=0}^{n-1} \alpha_1^k \alpha_2^{n-1-k}$$

such α_1 and α_2 are solutions of $\mu\alpha^2 - (\lambda + \mu + \beta)\alpha + \lambda = 0$ (and $x_0 = 0$).

Let $\Delta = (\lambda + \mu + \beta)^2 - 4\lambda\mu$

1st case: If $(\lambda + \mu + \beta)^2 = 4\lambda\mu$, so $\alpha_1 = \alpha_2 = \frac{\lambda + \mu + \beta}{2\mu} = \pm \sqrt{\frac{\lambda}{\mu}}$, and $\forall n \geq 1$

$$x_n = n \left(\frac{\lambda + \mu + \beta}{2\mu} \right)^{n-1} x_1$$

2nd case: If $(\lambda + \mu + \beta)^2 \neq 4\lambda\mu$, so $\alpha_1 \neq \alpha_2$, and $\forall n \geq 1$

$$x_n = \left(\frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} \right) x_1$$

such $\alpha_1 = \frac{(\lambda + \mu + \beta) + \Delta^{\frac{1}{2}}}{2\mu} \in IK$ and $\alpha_2 = \frac{(\lambda + \mu + \beta) - \Delta^{\frac{1}{2}}}{2\mu} \in IK$ ($IK = IR$ or \mathbb{C}).

$$\frac{\alpha_1^n - \alpha_2^n}{\alpha_1 - \alpha_2} \in IR$$

since, $\alpha_1 = \frac{(\lambda + \mu + \beta) + \Delta^{\frac{1}{2}}}{2\mu} \in \mathbb{C} \setminus IR \implies \alpha_2 = \frac{(\lambda + \mu + \beta) - \Delta^{\frac{1}{2}}}{2\mu} = \overline{\alpha_1}$ (as $(\lambda, \mu, \beta) \in IR^3$)

Conclusion 3.2. By applying our matrix power formula to the resolution of the second-order linear differential equation deduced from the infinite system of equations $Ax = \beta x$, we obtained the eigenvectors of the generating matrix A of the BDP process in the case of constant birth and death rates.

Competing Interests

As the sole author of this work, I declare that I have no conflict of interest with anyone.

Authors' Contributions

I am the sole author and contributor to the writing of this article. I have read and approved the final manuscript.

References

- [1] Arbai A., Benslimane M., ElOussati M. Birth and Death Processes with Finished Number of State. *International Journal of Mathematics and Statistics Invention (IJMSI)*. <https://www.ijmsi.org/Papers/Volume.5.Issue.7/A05070110.pdf>, Vol. 5 (7) . 2017
- [2] Arbai A., El Merouani M., Bellekbir A. Exact Solutions for Finite-State Birth-Death Processes: A Matrix-Theoretic Approach and Generalizations. *Franco-Moroccan Mathematics Days (F2MDays'25)*. Abdelmalek Essaadi University, Faculty of Sciences Tetouan, Morocco. Conference, <https://www.f2mdays.uae.ma/assets/img/BookofAbstractsFinalF2MDays'25.pdf>. page 40. 2025.
- [3] Crawford, F.W., Suchard, M.A. Transition probabilities for general birth–death processes with applications in ecology, genetics, and evolution. *J. Math. Biol.* 65, 553–580. <https://doi.org/10.1007/s00285-011-0471-z>. 2012
- [4] Feller W. *An introduction to probability theory and its applications*. John Wiley and Sons, (<https://bitcoinwords.github.io/assets/papers/an-introduction-to-probability-theory-and-its-applications.pdf>). 1950
- [5] Feller, W. Die Grundlagen der Volterraschen Theorie des Kampfes ums Dasein in wahrscheinlichkeitstheoretischer Behandlung. *Acta Biotheor* 5, 11–40. <https://doi.org/10.1007/BF01602932>. 1939
- [6] Karlin S., McGregor J. Representation of a class of stochastic processes. *Proc. Natl. Acad. Sci. U.S.A.* 41 (6) 387-391, <https://doi.org/10.1073/pnas.41.6.387>. 1955
- [7] Samuel Karlin, James McGregor "Random walks," *Illinois Journal of Mathematics*, *Illinois J. Math.* 3(1), 66-81. DOI: 10.1215/ijm/1255454999. 1959
- [8] Williams, K. S. The n th power of a 2×2 matrix (in Notes). *Math. Magazine*, Vol. 65(5), 336, 1992
- [9] Jung Hyeon Ryu and Won Kyu Kim. (2011). A formula on power n of 2×2 Matrices. *Far East Journal of Mathematical Sciences (FJMS)*. <http://dx.doi.org/10.17654/FJMSFeb2015.365.377>. Vol. 96(3). 365–377.2015
- [10] Sidje, Roger B., Stewart, William J. A numerical study of large sparse matrix exponentials arising in Markov chains. *Computational Statistics and Data Analysis*, Elsevier. <https://www.sciencedirect.com/science/article/abs/pii/S0167947398000620>. Vol. 29(3). 345–368. 1999
- [11] Respondek, J. Another Dubious Way to Compute the Exponential of a Matrix. In: Gervasi, O., et al. *Computational Science and Its Applications*. Lecture Notes in Computer Science. Springer, Cham. https://doi.org/10.1007/978-3-030-86653-2_34. 2021
- [12] Junghenn, H.D. Matrix Algebra. In: *Symbolic Mathematics with Python*. Synthesis Lectures on Mathematics and Statistics. Springer, Cham. https://doi.org/10.1007/978-3-031-90522-3_10. 2025
- [13] James Mc Laughlin, Nancy J. Wyshinski. Further combinatorial identities deriving from the n th power of a 2×2 matrix. *Discrete Applied Mathematics*. <https://doi.org/10.1016/j.dam.2006.01.003>. Vol. 154(8), 1301–1308, 2008
- [14] Cleve Moler and Charles Van Loan. Nineteen Dubious Ways to Compute the Exponential of a Matrix. *SIAM Review*. DOI: 10.1137/1020098. Vol. 20 (4). 801–836. 1978

-
- [15] Hall, B. The Matrix Exponential. In: Lie Groups, Lie Algebras, and Representations. Graduate Texts in Mathematics, Springer, Cham. Vol. 222. https://doi.org/10.1007/978-3-319-13467-3_2. 2015
- [16] Leonard I. E. The Matrix Exponential. SIAM Review. Vol. 38 (3). Pages: 507 - 512. DOI: 10.1137/S0036144595286488. 1996
- [17] Asadi, F. Equations and Linear Algebra. In: Engineering Mathematics with MATLAB and Simulink. Springer, Cham. doi.org/10.1007/978-3-031-85244-2_3. 2025