

## The Impact of Integrated Weed Management Practices on Nutrient Uptake in Rice (*Oryza sativa* L.)

### Abstract

The impact of integrated weed management practices on nutrient uptake in rice (*Oryza sativa* L.) and weeds at 90 days after transplanting (DAT) over the 2022 and 2023 growing seasons. Various weed control strategies, including pre-emergence and post-emergence herbicides combined with manual weeding, were assessed for their effectiveness in enhancing nitrogen (N), phosphorus (P), and potassium (K) uptake by rice while minimizing nutrient loss to weeds. The combination of Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> (pre-emergence) and Penoxsulam 22 g a.i. ha<sup>-1</sup> (post-emergence), along with one hand weeding at 40 DAT, was the most effective, resulting in the highest nutrient uptake by rice and the lowest nutrient uptake by weeds. Nitrogen uptake in grains reached 62.7 kg ha<sup>-1</sup> in 2022 and 67.4 kg ha<sup>-1</sup> in 2023, significantly outperforming other treatments. The weed-free treatment also demonstrated excellent nutrient absorption, emphasizing the importance of complete weed control for optimal rice growth. Conversely, the weedy check treatment, where no weed control was applied, led to the highest nutrient uptake by weeds, severely limiting nutrient availability for the rice crop. Manual weeding alone showed moderate effectiveness but was less efficient than integrated methods.

**Keywords:** *Nutrient uptake, Pyrazosulfuron, Penoxsulam, Nutrient Uptake, Rice*

### 1. Introduction

Rice (*Oryza sativa* L.) holds an indispensable position as the most important staple food crop worldwide, serving as the primary food source for nearly 60% of the global population. In India, rice accounts for approximately 43% of total food grain production and 46% of total cereal production, making it a cornerstone of food security. Rice cultivation is deeply rooted in the agricultural landscape, providing income and employment for over 50 million households, particularly in rural areas [1]. With one-third of Asia's rice production consumed in China and one-fifth in India, the global demand for rice continues to rise, reaching a cultivation area of 158 million hectares and an annual production of over 527 million tonnes. India, being the second-largest producer and consumer of rice after China, has seen its rice cultivated on 43.57 million hectares, yielding an annual production of 104.32 million tonnes with a productivity rate of 2.98 tonnes per hectare. Rice production faces numerous challenges, particularly in India, where factors such as declining natural resources, labor shortages, and increasing weed infestations pose significant threats to productivity [2]. Weeds are a major hindrance to rice cultivation, especially in the critical growth stages when they compete with rice for essential resources like nutrients, light, and space. Uncontrolled weed growth can lead to yield losses ranging from 12 to 51%, depending on the severity of the infestation and the control methods employed. In transplanted rice, heavy weed infestations are especially problematic, as weeds emerge simultaneously with rice seedlings, exacerbating

the competition for nutrients and other growth factors. Weeds contribute to 45% of the total annual agricultural production losses in India, surpassing other agricultural pests like insects and diseases [3]. Their rapid growth and adaptability enable them to dominate the crop habitat, significantly reducing rice yields. The critical period for rice-weed competition typically occurs during the vegetative phase of rice, and effective weed management during this time is crucial for maximizing rice productivity. Traditional mechanical methods, such as hand weeding, while effective, are labor-intensive and costly, making them less feasible for large-scale operations. Chemical weed control, particularly the use of herbicides, offers a more practical and cost-effective solution for managing weed infestations, especially in transplanted rice systems. Despite the benefits of herbicides, the continuous use of the same chemical agents can lead to shifts in weed flora and the development of herbicide-resistant weed species. This has led to the growing recognition of the need for integrated weed management (IWM) strategies that combine cultural, mechanical, and chemical methods to achieve sustainable weed control. IWM not only reduces dependency on herbicides but also promotes eco-friendly practices, ensuring long-term productivity and environmental health [4]. In India, the adoption of IWM in rice cultivation has been driven by labor shortages, the rising cost of herbicides, and the need for more efficient weed management systems. Weed control in rice is essential not only for improving yield but also for enhancing nutrient uptake by the crop. Weeds, by competing with rice for nutrients, can significantly reduce the availability of essential nutrients like nitrogen, phosphorus, and potassium, which are critical for rice growth and development [5]. Effective weed management practices, including the use of herbicides and timely mechanical interventions, can mitigate these losses and promote better nutrient absorption by rice plants. In this context, understanding the impact of integrated weed management practices on nutrient uptake in rice is crucial for optimizing rice productivity and ensuring food security in India and other rice-producing countries. This study aims to evaluate the effectiveness of integrated weed management practices in transplanted rice ecosystems and their impact on nutrient uptake by both rice and weeds. By analyzing various weed control methods, including chemical and mechanical interventions, this research seeks to identify sustainable and cost-effective strategies for enhancing rice yield while minimizing nutrient loss to weed competition. The findings will contribute to the ongoing efforts to improve rice production systems in India and other rice-dependent regions, ultimately supporting the goal of achieving food self-sufficiency and meeting future food demands.

## **2. Material and Method**

The field experiment was conducted over two consecutive years, 2022 and 2023, at the Crop Research Centre (CRC) farm of Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, located in the Indo-Gangetic plains of Western Uttar Pradesh, India. The geographical coordinates of the study area are 29°40' N latitude and 77°42' E longitude, with an elevation of 237 meters above mean sea level. The site experiences a semi-arid and subtropical climate, characterized by extreme temperature fluctuations, with hot summers and cold winters. During both years of the study, the temperature patterns showed a gradual decrease from the third week of November, reaching the lowest in December and January, followed by a steady increase from early February. The weekly mean maximum temperature during the crop-growing period ranged from 37.6°C to 15.4°C, and the mean minimum

temperature ranged from 5.9°C to 20.7°C, ensuring stable climatic conditions for the experiment.

The experiment was laid out in a randomized block design (RBD) with 12 treatment combinations and three replications. The treatments were designed to evaluate different weed management practices, including a weedy check (no weed control), weed-free conditions (complete removal of weeds), farmer's practices (one hand weeding at 40 days after transplanting [DAT]), and the application of herbicides such as Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> (pre-emergence), Bispyribac Sodium 25 g a.i. ha<sup>-1</sup> (post-emergence at 15 DAT), Ethoxy sulfuron 20 g a.i. ha<sup>-1</sup> (pre-emergence), and Penoxsulam 22 g a.i. ha<sup>-1</sup> (post-emergence). The herbicides were applied either alone or in combination with one hand weeding at 40 DAT. The plot sizes were maintained with a gross plot size of 5 x 4 m<sup>2</sup> and a net plot size of 4 x 3 m<sup>2</sup>, with a spacing of 20 x 10 cm between the rice plants. The rice variety PB-1637 was used for planting in both 2022 and 2023. Nutrient uptake (N, P, and K) by the rice crop was measured at harvest in both grain and straw. Weed nutrient uptake (N, P, and K) was also assessed at 90 DAT to evaluate the competition between weeds and rice for nutrients. Statistical analysis was performed using the analysis of variance (ANOVA) method in SPSS software. Treatment means were compared using the least significant difference (LSD) test at a 5% probability level to assess the significance of the results [6]. The comparison of the treatments across both years allowed for a comprehensive understanding of the efficacy of the weed management practices in improving rice nutrient uptake and yield while minimizing competition from weeds.

### 3. Result and Discussion

#### Nutrient Uptake of Weeds

The data presented in (Table 1) clearly shows the significant impact of different weed management practices on nutrient uptake (kg ha<sup>-1</sup>) by weeds at 90 days after transplanting (DAT) during the 2022 and 2023 growing seasons. The weedy check treatment (T1: No weed control) exhibited the highest nutrient uptake, with nitrogen levels reaching 29.3 kg ha<sup>-1</sup> in 2022 and 36.8 kg ha<sup>-1</sup> in 2023, phosphorus uptake at 8.0 kg ha<sup>-1</sup> and 10.4 kg ha<sup>-1</sup> in the respective years, and potassium uptake at 30.5 kg ha<sup>-1</sup> in 2022 and 37.9 kg ha<sup>-1</sup> in 2023. This indicates the significant nutrient depletion caused by uncontrolled weeds, emphasizing their aggressive competition for resources, which reduces nutrient availability for rice. In stark contrast, the weed-free treatment (T2: Complete weed removal) recorded zero nutrient uptake in both years, illustrating the efficacy of thorough weed control in preventing nutrient loss to weeds. Farmer's practices involving one hand weeding at 40 DAT (T3) moderately reduced nutrient uptake, with nitrogen uptake of 13.1 kg ha<sup>-1</sup> in 2022 and 16.5 kg ha<sup>-1</sup> in 2023, along with reduced phosphorus and potassium uptake, indicating that manual weeding effectively controls weed growth but does not fully eliminate competition. The application of Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> as a pre-emergence herbicide (T4) showed a further reduction in nutrient uptake, with nitrogen values dropping to 8.4 kg ha<sup>-1</sup> in 2022 and 10.6 kg ha<sup>-1</sup> in 2023. Combining Pyrazosulfuron with one hand weeding at 40 DAT (T5) resulted in even lower nutrient uptake, with nitrogen levels at 7.4 kg ha<sup>-1</sup> in 2022 and 9.3 kg ha<sup>-1</sup> in 2023. This combination highlights the added benefit of integrated methods, where both chemical and manual controls suppress weed competition more effectively than using a single

method. Post-emergence application of Bispyribac Sodium 25 g a.i. ha<sup>-1</sup> at 15 DAT (T6) also led to significant reductions in nutrient uptake, though slightly less effective than pre-emergence applications. Nutrient uptake further decreased when Bispyribac Sodium was combined with one hand weeding at 40 DAT (T7), with nitrogen uptake of 7.9 kg ha<sup>-1</sup> in 2022 and 10.0 kg ha<sup>-1</sup> in 2023. Similar trends were observed with Ethoxy sulfuron 20 g a.i. ha<sup>-1</sup> (T8), and its combination with hand weeding (T9), as well as with Penoxsulam 22 g a.i. ha<sup>-1</sup> (T10), and its combination with hand weeding (T11), where nutrient uptake consistently dropped, demonstrating the synergistic effect of integrating chemical and manual weed control practices. The most effective treatment in reducing nutrient uptake was the combination of Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> (pre-emergence) and Penoxsulam 22 g a.i. ha<sup>-1</sup> (post-emergence) with one hand weeding at 40 DAT (T12), which recorded the lowest nutrient uptake across both years, with nitrogen levels at 6.0 kg ha<sup>-1</sup> in 2022 and 7.6 kg ha<sup>-1</sup> in 2023. This treatment exemplifies the effectiveness of integrated weed management (IWM) in minimizing nutrient competition from weeds and ensuring optimal nutrient availability for the rice crop, as supported [7], [8], [9]. IWM strategies, which combine herbicide application with manual interventions, provide a sustainable and cost-effective approach to weed control, improving crop nutrient uptake and overall productivity.

### **Nutrient Uptake of Rice**

The data presented in (Table 2) reveals the significant effects of various weed management practices on nitrogen (N), phosphorus (P), and potassium (K) uptake in rice grain and straw at harvest in 2022 and 2023. The weedy check treatment (T1: No weed control) exhibited the lowest nutrient uptake, with nitrogen uptake in grains at 37.8 kg ha<sup>-1</sup> in 2022 and 41.1 kg ha<sup>-1</sup> in 2023, and in straw at 21.3 kg ha<sup>-1</sup> and 23.2 kg ha<sup>-1</sup> in 2022 and 2023, respectively. This treatment also had minimal phosphorus and potassium uptake, highlighting the detrimental impact of unchecked weed growth on nutrient availability for rice. In contrast, the weed-free treatment (T2: Complete weed removal) showed the highest nutrient uptake, with nitrogen uptake in grains at 65.1 kg ha<sup>-1</sup> in 2022 and 70.5 kg ha<sup>-1</sup> in 2023, and phosphorus and potassium levels similarly high, emphasizing the importance of eliminating weeds to maximize nutrient absorption. The farmer's practice (T3: One hand weeding at 40 DAT) moderately improved nutrient uptake compared to the weedy check, with nitrogen uptake in grains at 46.0 kg ha<sup>-1</sup> in 2022 and 50.0 kg ha<sup>-1</sup> in 2023. The application of Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> as pre-emergence (T4) further increased nutrient uptake, and when combined with one hand weeding at 40 DAT (T5), nutrient uptake improved even more, reaching nitrogen uptake levels of 58.1 kg ha<sup>-1</sup> in 2022 and 62.4 kg ha<sup>-1</sup> in 2023. Similarly, the post-emergence application of Bispyribac Sodium 25 g a.i. ha<sup>-1</sup> at 15 DAT (T6) enhanced nutrient uptake, which was further boosted when combined with hand weeding (T7). Treatments involving Ethoxy sulfuron 20 g a.i. ha<sup>-1</sup> as pre-emergence (T8) and its combination with one hand weeding (T9), as well as Penoxsulam 22 g a.i. ha<sup>-1</sup> as post-emergence (T10) and its combination with hand weeding (T11), also demonstrated improved nutrient uptake compared to the weedy check, though the results were less pronounced than the Pyrazosulfuron and Bispyribac sodium treatments. The highest nutrient uptake was recorded in the treatment combining Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> (pre-emergence) + Penoxsulam 22 g a.i. ha<sup>-1</sup> (post-emergence) + one hand weeding (T12), with nitrogen uptake in grains at 62.7 kg ha<sup>-1</sup> in 2022 and 67.4 kg ha<sup>-1</sup> in 2023. This integrated treatment demonstrated superior weed control, thereby enhancing nutrient availability for rice, supporting previous

research findings that combining chemical and manual weed control methods is the most effective strategy for optimizing nutrient uptake and rice yields [9], [10],[11].

UNDER PEER REVIEW

**Table :1 Effect of weed management practices on nutrient uptake (kg ha<sup>-1</sup>) of weeds at 90 Days**

	Treatments	Nutrient uptake (kg ha <sup>-1</sup> )					
		Nitrogen		Phosphorus		Potassium	
		2022	2023	2022	2023	2022	2023
<b>T<sub>1</sub></b>	Weedy check	29.3	36.8	8.0	10.4	30.5	37.9
<b>T<sub>2</sub></b>	Weed free	0.0	0.0	0.0	0.0	0.0	0.0
<b>T<sub>3</sub></b>	Farmers practices (One hand weeding 40 DAT)	13.1	16.5	3.5	4.5	14.4	17.9
<b>T<sub>4</sub></b>	Pyrazosulfuron 150 g a.i. ha <sup>-1</sup> (Pre emergence)	8.4	10.6	1.9	2.5	9.4	11.7
<b>T<sub>5</sub></b>	Pyrazosulfuron 150 g a.i. ha <sup>-1</sup> (Pre emergence) + one hand weeding 40 DAT	7.4	9.3	1.7	2.2	8.6	10.7
<b>T<sub>6</sub></b>	Bispyribac Sodium @25 g a.i. ha <sup>-1</sup> 15 DAT ( Post- emergence)	11.6	14.6	2.8	3.7	12.9	16.0
<b>T<sub>7</sub></b>	Bispyribac Sodium @25 g a.i. ha <sup>-1</sup> 15 DAT ( Post- emergence) + one hand weeding 40 DAT	7.9	10.0	1.8	2.4	9.2	11.5
<b>T<sub>8</sub></b>	Ethoxy Sulfuron @20g a.i. ha <sup>-1</sup> (pre-emergence)	12.4	15.6	3.2	4.2	13.6	16.9
<b>T<sub>9</sub></b>	Ethoxy Sulfuron @20g a.i. ha <sup>-1</sup> (pre-emergence) + + one hand weeding 40 DAT	13.6	17.1	3.8	4.9	14.9	18.5
<b>T<sub>10</sub></b>	Penoxsulam 22 g a.i. ha <sup>-1</sup> (post emergence)	11.3	14.1	2.7	3.5	12.5	15.6
<b>T<sub>11</sub></b>	Penoxsulam 22 g a.i. ha <sup>-1</sup> (post emergence) + One hand weeding	7.1	8.9	1.5	2.0	8.0	9.9
<b>T<sub>12</sub></b>	Pyrazosulfuron 150 g ai.ha <sup>-1</sup> (pre emergence) + Penoxsulam 22 g a.i. ha <sup>-1</sup> post emergence + One hand weeding	6.0	7.6	1.3	1.7	6.9	8.6
	<b>SEm+</b>	<b>0.42</b>	<b>0.53</b>	<b>0.11</b>	<b>0.14</b>	<b>0.46</b>	<b>0.57</b>
	<b>C.D.(P=0.05)</b>	<b>1.21</b>	<b>1.52</b>	<b>0.31</b>	<b>0.41</b>	<b>1.32</b>	<b>1.64</b>

**Table :2 Effect of weed management practices on N, P & K uptake (kg ha<sup>-1</sup>) in rice grain & straw at harvest**

	Treatments	Nutrient uptake (kg ha <sup>-1</sup> )											
		Nitrogen				Phosphorus				Potassium			
		Grains		Straw		Grains		Straw		Grains		Straw	
		2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
T <sub>1</sub>	Weedy check	37.8	41.1	21.3	23.2	7.7	8.4	5.0	5.8	9.0	10.0	64.4	68.4
T <sub>2</sub>	Weed free	65.1	70.5	32.7	35.0	17.5	19.3	10.7	11.7	17.5	19.3	93.7	99.1
T <sub>3</sub>	Farmers practices (One hand weeding 40 DAT)	46.0	50.0	25.0	26.7	10.1	11.0	6.1	6.8	10.9	12.1	76.1	80.1
T <sub>4</sub>	Pyrazosulfuron 150 g a.i. ha <sup>-1</sup> (Pre emergence)	53.1	58.1	27.9	29.6	12.2	13.7	7.8	8.5	13.0	14.5	82.9	86.7
T <sub>5</sub>	Pyrazosulfuron 150 g a.i. ha <sup>-1</sup> (Pre emergence) + one hand weeding 40 DAT	58.1	62.4	29.6	31.8	13.5	14.9	8.7	9.7	14.4	15.8	86.0	91.2
T <sub>6</sub>	Bispyribac Sodium @25 g a.i. ha <sup>-1</sup> 15 DAT ( Post- emergence)	49.3	52.9	26.7	28.6	11.2	11.9	7.0	7.8	11.9	13.1	80.6	85.2
T <sub>7</sub>	Bispyribac Sodium @25 g a.i. ha <sup>-1</sup> 15 DAT ( Post- emergence) + one hand weeding 40 DAT	54.4	59.2	28.9	31.0	12.8	14.3	7.9	8.7	13.6	15.1	84.8	89.5
T <sub>8</sub>	Ethoxy Sulfuron @20g a.i. ha <sup>-1</sup> (pre-emergence)	46.8	50.9	26.5	28.4	10.2	11.1	7.0	7.8	11.0	12.3	79.6	84.0
T <sub>9</sub>	Ethoxy Sulfuron @20g a.i. ha <sup>-1</sup> (pre-emergence) + one hand weeding 40 DAT	43.8	47.7	24.1	26.1	9.4	10.2	5.9	6.7	10.4	11.6	73.0	77.7
T <sub>10</sub>	Penoxsulam 22 g a.i. ha <sup>-1</sup> (post emergence)	50.7	54.8	27.3	29.3	11.6	12.9	7.0	7.8	12.4	13.7	80.8	85.4
T <sub>11</sub>	Penoxsulam 22 g a.i. ha <sup>-1</sup> (post emergence) + One hand weeding	59.2	64.6	31.6	34.0	14.2	15.9	9.1	10.1	15.1	16.8	91.4	97.0
T <sub>12</sub>	Pyrazosulfuron 150 g ai.ha <sup>-1</sup> (pre emergence) + Penoxsulam 22 g a.i. ha <sup>-1</sup> post emergence + One hand weeding	62.7	67.4	31.9	34.4	15.0	16.5	9.9	11.0	15.9	17.5	92.7	98.7
	<b>SEm+</b>	<b>1.86</b>	<b>2.02</b>	<b>0.99</b>	<b>1.06</b>	<b>0.43</b>	<b>0.48</b>	<b>0.27</b>	<b>0.30</b>	<b>0.46</b>	<b>0.51</b>	<b>2.92</b>	<b>3.09</b>
	<b>C.D.(P=0.05)</b>	<b>5.34</b>	<b>5.79</b>	<b>2.83</b>	<b>3.04</b>	<b>1.24</b>	<b>1.37</b>	<b>0.79</b>	<b>0.87</b>	<b>1.32</b>	<b>1.46</b>	<b>8.38</b>	<b>8.86</b>

## Conclusion

The study demonstrates that integrated weed management practices significantly improve nutrient uptake in rice by reducing competition from weeds. The combination of Pyrazosulfuron 150 g a.i. ha<sup>-1</sup> (pre-emergence) and Penoxsulam 22 g a.i. ha<sup>-1</sup> (post-emergence), along with one hand weeding at 40 DAT proved to be the most effective resulting in the highest nutrient uptake and improved crop productivity. The weed-free treatment also showed excellent results, underscoring the importance of complete weed control. These findings suggest that combining chemical and manual weed control methods offers a sustainable and efficient approach to optimize nutrient availability, thereby enhancing rice yield and productivity. This integrated approach provides a balanced, cost-effective solution to managing weed competition in rice cultivation.

## References

1. Bouman, B. A., Barker, R., Humphreys, E., Tuong, T. P., Atlin, G., Bennett, J., ... & Wassman, R. (2007). Rice: feeding the billions.
2. Fahad, S., Adnan, M., Noor, M., Arif, M., Alam, M., Khan, I. A., ... & Wang, D. (2019). Major constraints for global rice production. In *Advances in rice research for abiotic stress tolerance* (pp. 1-22). Woodhead Publishing.
3. Junaid, M. D., & Gokce, A. F. (2024). Global agricultural losses and their causes. *Bulletin of Biological and Allied Sciences Research*, 2024(1), 66-66.
4. Bahadur, S., Verma, S. K., Prasad, S. K., Madane, A. J., Maurya, S. P., Gaurav, V. K. V., & Sihag, S. K. (2015). Eco-friendly weed management for sustainable crop production-A review.
5. Ghosh, D., Brahmachari, K., Skalický, M., Roy, D., Das, A., Sarkar, S., ... & Hossain, A. (2022). The combination of organic and inorganic fertilizers influence the weed growth, productivity and soil fertility of monsoon rice. *PloS one*, 17(1), e0262586.
6. Gomez KA, Gomez AA (1976). *Statistical procedure for agricultural research with emphasis on rice*. 2nd Ed. IRRI, Los Banos, Philippine.
7. Onwuchekwa-Henry, C. B., Van Ogtrop, F., Roche, R., & Tan, D. K. (2023). Evaluation of pre-emergence herbicides for weed management and rice yield in direct-seeded rice in Cambodian lowland ecosystems. *Farming System*, 1(2), 100018.
8. Ahmed, M. T., Sarker, U. K., Rahman, M. R., & Uddin, M. R. (2024). Assessing Herbicide-Resistance and Yield Optimization in Boro Rice (cv. BRRI dhan29) through Post-emergence Herbicides Applications. *Asian Journal of Research in Crop Science*, 9(3), 22-36.
9. Tiwari, S., Mondal, T., Soni, R., Naj, I., Tomar, A., & Upadhyay, V. B. (2023). Integrative Weed Management Tactics for Rice (*Oryza sativa* L.) under Direct-Seeding in Kymore Plateau and Satpura Hills. *Int. J. Plant Soil Sci*, 35(20), 848-862.

10. Tiwari, S., Mondal, T., Soni, R., Naj, I., Tomar, A., & Upadhyay, V. B. (2023). Integrative Weed Management Tactics for Rice (*Oryza sativa* L.) under Direct-Seeding in Kymore Plateau and Satpura Hills. *Int. J. Plant Soil Sci*, 35(20), 848-862.
11. Meena, S. N., Sharma, S. K., Singh, P., Jadon, C. K., Jat, M. L., & Meena, R. L. (2022). Crop-management practices influence weed dynamics, yield and economics of soybean (*Glycine max*). *Indian Journal of Agronomy*, 67(3), 282-286.

UNDER PEER REVIEW

UNDER PEER REVIEW