

## A Note on Hyperbolic Generalized Pierre Numbers

**Abstract.** In this paper, we introduce the generalized hyperbolic Pierre numbers defined over the bidimensional Clifford algebra of hyperbolic numbers. As special cases, we examine the hyperbolic Pierre and hyperbolic Pierre Lucas numbers. We derive Binet formulas, construct generating functions, and establish summation identities for these sequences. Furthermore, we present matrix representations associated with the proposed number sequences.

**Keywords.** Pierre numbers, Pierre-Lucas numbers, hyperbolic numbers, hyperbolic Pierre numbers.

**2010 Mathematics Subject Classification.** 11B37, 11B39, 11B83.

### 1. Introduction

Hypercomplex number systems, as introduced in [14], constitute algebraic extensions of the real numbers. Among the commutative examples are:

- **Complex numbers**

$$\mathbb{C} = \{z = a + ib : a, b \in \mathbb{R}, i^2 = -1\},$$

- **Hyperbolic numbers** (also known as double or split-complex numbers)[24]

$$\mathbb{H} = \{h = a + jb : a, b \in \mathbb{R}, j^2 = 1, j \neq \pm 1\},$$

- **Dual numbers** [10]

$$\mathbb{D} = \{d = a + \varepsilon b : a, b \in \mathbb{R}, \varepsilon^2 = 0, \varepsilon \neq 0\},$$

In contrast, non-commutative hypercomplex systems include:

- **Quaternions** [11]

$$\mathbb{H}_{\mathbb{Q}} = \{q = a_0 + ia_1 + ja_2 + ka_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, i^2 = j^2 = k^2 = ijk = -1\},$$

- **Octonions** [3]
- **Sedenions** [26].

The algebras  $\mathbb{C}$  (complex numbers),  $\mathbb{H}_{\mathbb{Q}}$  (quaternions),  $\mathbb{O}$  (octonions), and  $\mathbb{S}$  (sedenions) are real algebras constructed from the real numbers  $\mathbb{R}$  via a recursive doubling procedure known as the Cayley–Dickson process. This process can be extended beyond the sedenions to generate higher-dimensional algebras referred to as  $2^n$ -ions (see, for instance, [4], [12], [19]).

Quaternions were introduced by the Irish mathematician W. R. Hamilton (1805–1865) [11] as a four-dimensional extension of complex numbers. Hyperbolic numbers with complex coefficients were first proposed by J. Cockle in 1848 [6]. Later, H. H. Cheng and S. Thompson[5] extended the concept of dual numbers to the complex domain, referring to them as complex dual numbers. More recently, Akar, Yüce, and Şahin introduced the notion of [1] dual hyperbolic numbers, further enriching the landscape of hypercomplex number systems.

A **dual hyperbolic** number is a type of hypercomplex number defined as

$$q = (a_0 + ja_1) + \varepsilon(a_2 + ja_3) = a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3,$$

where  $a_0, a_1, a_2$  and  $a_3$  are real numbers.

The set of all dual hyperbolic numbers is denoted by

$$\mathbb{H}_{\mathbb{D}} = \{a_0 + ja_1 + \varepsilon a_2 + \varepsilon ja_3 : a_0, a_1, a_2, a_3 \in \mathbb{R}, j^2 = 1, j \neq \pm 1, \varepsilon^2 = 0, \varepsilon \neq 0\}.$$

The basis elements  $\{1, j, \varepsilon, \varepsilon j\}$  satisfy the following commutative multiplication rules:

$$\begin{aligned} 1.\varepsilon &= \varepsilon, 1.j = j, \varepsilon^2 = \varepsilon.\varepsilon = (j\varepsilon)^2 = 0, j^2 = j.j = 1, \\ \varepsilon.j &= j.\varepsilon, \varepsilon.(\varepsilon j) = (\varepsilon j).\varepsilon = 0, j(\varepsilon j) = (\varepsilon j)j = \varepsilon, \end{aligned}$$

Here,  $\varepsilon$  denotes the pure dual unit ( $\varepsilon^2 = 0, \varepsilon \neq 0$ ),  $j$  is the hyperbolic unit ( $j^2 = 1$ ), and  $\varepsilon j$  is the dual hyperbolic unit

(with  $(j\varepsilon)^2 = 0$ ).

The product of two dual hyperbolic numbers  $q = a_0 + ja_1 + \varepsilon a_2 + j\varepsilon a_3$  and  $p = b_0 + jb_1 + \varepsilon b_2 + j\varepsilon b_3$  is given by

$$qp = a_0b_0 + a_1b_1 + j(a_0b_1 + a_1b_0) + \varepsilon(a_0b_2 + a_2b_0 + a_1b_3 + a_3b_1) + j\varepsilon(a_0b_3 + a_1b_2 + a_2b_1 + b_0a_3).$$

Addition of dual hyperbolic numbers is defined componentwise.

The dual hyperbolic numbers form a commutative ring, real vector space and an algebra. But  $\mathbb{H}_{\mathbb{D}}$  is not field because every dual hyperbolic numbers doesn't have an inverse. For more information on the dual hyperbolic numbers, see [1].

Here we use the set of hyperbolic numbers. The set of hyperbolic numbers  $\mathbb{H}$  can be described as

$$\mathbb{H} = \{z = x + hy \mid h \notin \mathbb{R}, h^2 = 1, x, y \in \mathbb{R}\}.$$

Here,  $h$  is the hyperbolic unit satisfying  $h^2 = 1$ , and  $x, y$  are real components.

The hyperbolic ring  $\mathbb{H}$  constitutes a bidimensional Clifford algebra; for further details, see [17]. In the mathematical literature, hyperbolic numbers have appeared under various names, including Lorentz numbers, double numbers, duplex numbers, split-complex numbers, and perplex numbers. These numbers are particularly useful in modeling distances within the Lorentz space-time plane (see Sobczyk [24]). For additional insights into the structure and applications of

hyperbolic numbers, refer to [13,16,20,25].

Addition, subtraction and multiplication of any two hyperbolic numbers  $z_1$  and  $z_2$  are defined by

$$\begin{aligned} z_1 \pm z_2 &= (x_1 + hy_1) \pm (x_2 + hy_2) = (x_1 \pm x_2) + h(y_1 \pm y_2), \\ z_1 \times z_2 &= (x_1 + hy_1) \times (x_2 + hy_2) = x_1x_2 + y_1y_2 + h(x_1y_2 + y_1x_2) \end{aligned}$$

and the division of two hyperbolic numbers are given by

$$\frac{z_1}{z_2} = \frac{x_1 + hy_1}{x_2 + hy_2} = \frac{(x_1 + hy_1)(x_2 - hy_2)}{(x_2 + hy_2)(x_2 - hy_2)} = \frac{x_1x_2 + y_1y_2}{x_2^2 - y_2^2} + h \frac{x_1y_2 + y_1x_2}{x_2^2 - y_2^2}.$$

It is easy to see that this algebra of hyperbolic numbers is commutative and contains zero divisors. The hyperbolic conjugation of  $z = x + hy$  is defined by

$$\bar{z} = z^\dagger = x - hy.$$

Note that  $\bar{\bar{z}} = z$ . Note also that for any hyperbolic numbers  $z_1, z_2, z$  we have

$$\begin{aligned} \overline{z_1 + z_2} &= \bar{z}_1 + \bar{z}_2, \\ \overline{z_1 \times z_2} &= \bar{z}_1 \times \bar{z}_2, \\ \|z\|^2 &= z \times \bar{z} = x^2 - y^2. \end{aligned}$$

Let us now revisit the definition of generalized Pierre numbers.

A generalized Pierre sequence  $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, W_3)\}_{n \geq 0}$  is defined by the fourth-order recurrence relations

$$W_n = 2W_{n-1} - W_{n-4}, \tag{1.1}$$

with the initial values  $W_0, W_1, W_2, W_3$  not all being zero. The sequence  $\{W_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$W_{-n} = 2W_{-(n-3)} - W_{-(n-4)},$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (1.1) holds for all integer  $n$ .

The initial values of the generalized Pierre numbers for both positive and negative subscripts are presented in Table 1.

Table 1. A few generalized Pierre numbers

$n$	$W_n$	$W_{-n}$
0	$W_0$	$W_0$
1	$W_1$	$2W_2 - W_3$
2	$W_2$	$2W_1 - W_2$
3	$W_3$	$2W_0 - W_1$
4	$2W_3 - W_0$	$4W_2 - W_0 - 2W_3$
5	$4W_3 - W_1 - 2W_0$	$4W_1 - 4W_2 + W_3$
6	$8W_3 - 2W_1 - W_2 - 4W_0$	$4W_0 - 4W_1 + W_2$
7	$15W_3 - 4W_1 - 2W_2 - 8W_0$	$W_1 - 4W_0 + 8W_2 - 4W_3$
8	$28W_3 - 8W_1 - 4W_2 - 15W_0$	$W_0 + 8W_1 - 12W_2 + 4W_3$
9	$52W_3 - 15W_1 - 8W_2 - 28W_0$	$8W_0 - 12W_1 + 6W_2 - W_3$
10	$96W_3 - 28W_1 - 15W_2 - 52W_0$	$6W_1 - 12W_0 + 15W_2 - 8W_3$
11	$177W_3 - 52W_1 - 28W_2 - 96W_0$	$6W_0 + 15W_1 - 32W_2 + 12W_3$
12	$326W_3 - 96W_1 - 52W_2 - 177W_0$	$15W_0 - 32W_1 + 24W_2 - 6W_3$
13	$600W_3 - 177W_1 - 96W_2 - 326W_0$	$24W_1 - 32W_0 + 24W_2 - 15W_3$

If we set  $W_0 = 0, W_1 = 1, W_2 = 2, W_3 = 4$  then  $\{W_n\}$  is the well-known Pierre sequence and if we set  $W_0 = 4, W_1 = 2, W_2 = 4, W_3 = 8$  then  $\{W_n\}$  is the well-known Pierre -Lucas sequence. In other words, Pierre sequence  $\{P_n\}_{n \geq 0}$  and Pierre -Lucas sequence  $\{C_n\}_{n \geq 0}$  are defined by the second-order recurrence relations

$$P_n = 2P_{n-1} - P_{n-4}, \quad P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 4, \quad n \geq 4, \tag{1.2}$$

and

$$C_n = 2C_{n-1} - C_{n-4}, \quad C_0 = 4, C_1 = 2, C_2 = 4, C_3 = 8, \quad n \geq 4. \tag{1.3}$$

The sequences  $\{P_n\}_{n \geq 0}$  and  $\{C_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$P_{-n} = 2P_{-(n-3)} - P_{-(n-4)},$$

and

$$C_{-n} = 2C_{-(n-3)} - C_{-(n-4)},$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrences (1.2) and (1.3) hold for all integer  $n$ .

We can list some important properties of generalized Pierre numbers that are needed.

- Binet formula of generalized Pierre sequence can be calculated using its characteristic equation which is given as

$$z^4 - 2z^3 + 1 = (z^3 - z^2 - z - 1)(z - 1) = 0.$$

The roots of characteristic equation are

$$\begin{aligned} \alpha &= \frac{1 + \sqrt[3]{19 + 3\sqrt{33}} + \sqrt[3]{19 - 3\sqrt{33}}}{3}, \\ \beta &= \frac{1 + \omega \sqrt[3]{19 + 3\sqrt{33}} + \omega^2 \sqrt[3]{19 - 3\sqrt{33}}}{3}, \\ \gamma &= \frac{1 + \omega^2 \sqrt[3]{19 + 3\sqrt{33}} + \omega \sqrt[3]{19 - 3\sqrt{33}}}{3}, \\ \delta &= 1, \end{aligned}$$

where

$$\omega = \frac{-1 + i\sqrt{3}}{2} = \exp(2\pi i/3).$$

Using these roots and the recurrence relation, Binet formula can be given as

$$\begin{aligned} W_n &= \frac{p_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{p_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{p_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{p_4 \delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)} \quad (1.4) \\ &= \frac{p_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - 1)} + \frac{p_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - 1)} + \frac{p_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - 1)} + \frac{p_4}{(1 - \alpha)(1 - \beta)(1 - \gamma)} \\ &= A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4 \end{aligned}$$

where  $p_1, p_2, p_3$  and  $p_4$  are given below

$$\begin{aligned} p_1 &= W_3 - (\beta + \gamma + \delta)W_2 + (\beta\gamma + \beta\delta + \gamma\delta)W_1 - \beta\gamma\delta W_0, \\ p_2 &= W_3 - (\alpha + \gamma + \delta)W_2 + (\alpha\gamma + \alpha\delta + \gamma\delta)W_1 - \alpha\gamma\delta W_0, \\ p_3 &= W_3 - (\alpha + \beta + \delta)W_2 + (\alpha\beta + \alpha\delta + \beta\delta)W_1 - \alpha\beta\delta W_0, \\ p_4 &= W_3 - W_2 - W_1 - W_0 \end{aligned} \quad (1.5)$$

and

$$\begin{aligned} A_1 &= \frac{W_3 - (\beta + \gamma + \delta)W_2 + (\beta\gamma + \beta\delta + \gamma\delta)W_1 - \beta\gamma\delta W_0}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)}, \\ A_2 &= \frac{W_3 - (\alpha + \gamma + \delta)W_2 + (\alpha\gamma + \alpha\delta + \gamma\delta)W_1 - \alpha\gamma\delta W_0}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)}, \\ A_3 &= \frac{W_3 - (\alpha + \beta + \delta)W_2 + (\alpha\beta + \alpha\delta + \beta\delta)W_1 - \alpha\beta\delta W_0}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)}, \\ A_4 &= \frac{W_3 - W_2 - W_1 - W_0}{-2}. \end{aligned} \quad (1.6)$$

Binet formula of Pierre and Pierre Lucas sequences are

$$P_n = \frac{\alpha^{n+2}}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{\beta^{n+2}}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{\gamma^{n+2}}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} - \frac{1}{2},$$

and

$$C_n = \alpha^n + \beta^n + \gamma^n + 1,$$

respectively.

The generating function for generalized Pierre numbers is

$$\sum_{n=0}^{\infty} W_n x^n = \frac{W_0 + (W_1 - 2W_0)x + (W_2 - 2W_1)x^2 + (W_3 - 2W_2)x^3}{1 - 2x + x^4}. \tag{1.7}$$

In the following section, we introduce the dual hyperbolic generalized Pierre numbers and investigate several of their fundamental properties.

Next, we give the exponential generating function of  $\sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$  of the sequence  $W_n$ .

LEMMA 1. [21, Lemma 1.4]. Suppose that  $f_{GW_n}(x) = \sum_{n=0}^{\infty} W_n \frac{x^n}{n!}$  is the exponential generating function of the generalized Pierre sequence  $\{W_n\}$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} &= \frac{(\alpha W_3 - \alpha(2 - \alpha)W_2 + (-\alpha^2 + \alpha + 1)W_1 - W_0)}{2\alpha^2 + 2\alpha - 2} e^{\alpha x} \\ &+ \frac{(\beta W_3 - \beta(2 - \beta)W_2 + (-\beta^2 + \beta + 1)W_1 - W_0)}{2\beta^2 + 2\beta - 2} e^{\beta x} \\ &+ \frac{(\gamma W_3 - \gamma(2 - \gamma)W_2 + (-\gamma^2 + \gamma + 1)W_1 - W_0)}{2\gamma^2 + 2\gamma - 2} e^{\gamma x} + \left(\frac{W_3 - W_2 + W_1 - W_0}{-2}\right) e^x. \end{aligned}$$

The previous Lemma gives the following results as particular examples.

COROLLARY 2. Exponential generating function of Pierre and Pierre-Lucas numbers are

$$\begin{aligned} \text{a): } \sum_{n=0}^{\infty} P_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} \left( \frac{(\alpha^2 + \alpha + 1)\alpha^n}{2(\alpha^2 + \alpha - 1)} + \frac{(\beta^2 + \beta + 1)\beta^n}{2(\beta^2 + \beta - 1)} + \frac{(\gamma^2 + \gamma + 1)\gamma^n}{2(\gamma^2 + \gamma - 1)} - \frac{1}{2} \right) \frac{x^n}{n!} \\ &= \frac{(\alpha^2 + \alpha + 1)}{2(\alpha^2 + \alpha - 1)} e^{\alpha x} + \frac{(\beta^2 + \beta + 1)}{2(\beta^2 + \beta - 1)} e^{\beta x} + \frac{(\gamma^2 + \gamma + 1)\gamma^n}{2(\gamma^2 + \gamma - 1)} e^{\gamma x} - \frac{1}{2} e^x. \\ \text{b): } \sum_{n=0}^{\infty} C_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} (\alpha^n + \beta^n + \gamma^n + 1) \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^{\gamma x} + e^x. \end{aligned}$$

Next, we provide an overview of selected publications in the literature that pertain to hyperbolic numbers.

- Cockle [6] presented the hyperbolic numbers with complex coefficients.
- Akar et al [1] introduced the dual hyperbolic numbers.
- Aydin [2] introduced the concept of hyperbolic Fibonacci numbers, defined by the following expression:

$$\tilde{F}_n = F_n + hF_{n+1},$$

where Fibonacci numbers are given by  $F_{n+2} = F_{n+1} + F_n$ , with the initial condition  $F_0 = 0, F_1 = 1$ .

- Soykan and Taşdemir [23] studied hyperbolic generalized Jacobsthal numbers given by

$$\tilde{V}_n = V_n + hV_{n+1}$$

where generalized Jacobsthal numbers are  $V_{n+2} = V_{n+1} + 2V_n$  with the initial conditation  $V_0 = a, V_1 = b$ .

- Dikmen and Altınsoy, [8] introduced On Third Order Hyperbolic Jacobsthal Numbers are

$$\begin{aligned} \hat{J}_n^{(3)} &= J_n^{(3)} + hJ_{n+1}^{(3)}, \\ \hat{j}_n^{(3)} &= j_n^{(3)} + hj_{n+1}^{(3)} \end{aligned}$$

where Jacobsthal numbers, respectively, given by  $J_n^{(3)} = J_{n-1}^{(3)} + J_{n-2}^{(3)} + 2J_{n-3}^{(3)}, J_0^{(3)} = 0, J_1^{(3)} = 1, J_2^{(3)} = 1, j_n^{(3)} = j_{n-1}^{(3)} + j_{n-2}^{(3)} + 2j_{n-3}^{(3)}, j_0^{(3)} = 2, j_1^{(3)} = 1, j_2^{(3)} = 5$ .

- Yılmaz and Soykan , [22] studied hyperbolic generalized Guglielmo numbers given by

$$HW_n = W_n + jW_{n+1}$$

where generalized Guglielmo numbers are  $W_n = 3W_{n-1} - 3W_{n-2} + W_{n-3}$  with the initial conditation  $W_0, W_1, W_2 \quad (n \geq 2)$ .

- Ayrılma and Soykan , [9] introduced On Hyperbolic Edouard Numbers are

$$HE_n = 7HE_{n-1} - 7HE_{n-2} + HE_{n-3}$$

where generalized Edouard numbers are  $E_n = 7E_{n-1} - 7E_{n-2} + E_{n-3}$  with the initial conditation  $E_0 = 0, E_1 = 1, E_2 = 7$

Following this, we provide details on dual hyperbolic sequences as they are presented in literature.

- Demirci and Soykan, [7] studied dual hyperbolic generalized Adrien numbers given by

$$\hat{A}_n = 3\hat{A}_{n-1} - \hat{A}_{n-2} - \hat{A}_{n-4}$$

where generalized Adrien numbers are  $A_n = 3A_{n-1} - A_{n-2} + A_{n-4}$  with the initial conditation  $A_0 = 0, A_1 = 1, A_2 = 3, A_3 = 8, \quad n \geq 4$ .

- Kalca and Soykan , [15] studied dual hyperbolic generalized Pandita numbers given by

$$\hat{P}_n = 2\hat{P}_{n-1} - \hat{P}_{n-2} + \hat{P}_{n-3} - \hat{P}_{n-4}$$

where generalized Pandita numbers are  $P_n = 2P_{n-1} - P_{n-2} + P_{n-3} - P_{n-4}$  with the initial conditation  $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 3, \quad n \geq 4$ .

In this paper, we define the hyperbolic generalized Pierre numbers in the next section and give some properties of them.

## 2. Hyperbolic Generalized Pierre Numbers and their Generating Functions and Binet's Formulas

In this section, we introduce the hyperbolic generalized Pierre numbers and derive their corresponding generating functions and Binet formulas. We now define the hyperbolic generalized Pierre numbers over the algebra  $\mathbb{H}_{\mathbb{D}}$  of dual hyperbolic numbers. The  $n$ th hyperbolic generalized Pierre number is

$$HW_n = W_n + jW_{n+1}. \quad (2.1)$$

The sequence  $\{HW_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$HW_{-n} = W_{-n} + jW_{-n+1},$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrence (2.2) holds for all integer  $n$ .

Note that

$$\begin{aligned} HW_0 &= W_0 + jW_1 = W_0 + jW_1, \\ HW_1 &= W_1 + jW_2 = W_1 + jHW_2, \\ HW_2 &= W_2 + jW_3 = W_2 + jHW_3, \\ HW_3 &= W_3 + jW_4 = W_3 + j(2HW_3 - HW_0). \end{aligned}$$

It can be easily shown that

$$HW_n = 2HW_{n-1} - HW_{n-4} \quad (2.2)$$

and

$$HW_{-n} = 2HW_{-(n-3)} - HW_{-(n-4)}.$$

The initial values of the hyperbolic generalized Pierre numbers for both positive and negative subscripts are listed in Table 2.

Table 2. A few hyperbolic generalized Pierre numbers

$n$	$HW_n$	$HW_{-n}$
0	$HW_0$	$HW_0$
1	$HW_1$	$2HW_2 - HW_3$
2	$HW_2$	$2HW_1 - HW_3$
3	$HW_3$	$2W_0 - W_1$
4	$2HW_3 - HW_0$	$4HW_2 - HW_0 - 2HW_3$
5	$4HW_3 - HW_1 - 2HW_0$	$4HW_1 - 4HW_2 + HW_3$
6	$8HW_3 - HW_2 - 2HW_1 - 4HW_0$	$4HW_0 - 4HW_1 + HW_2$
7	$15HW_3 - 2HW_2 - 4HW_1 - 8HW_0$	$HW_1 - 4HW_0 + 8HW_2 - 4HW_3$
8	$28HW_3 - 4HW_2 - 8HW_1 - 15HW_0$	$HW_0 + 8HW_1 - 12HW_2 + 4HW_3$
9	$52HW_3 - 8HW_2 - 15HW_1 - 28HW_0$	$8HW_0 - 12HW_1 + 6HW_2 - HW_3$
10	$96HW_3 - 15HW_2 - 28HW_1 - 52HW_0$	$6HW_1 - 12HW_0 + 15HW_2 - 8HW_3$
11	$177HW_3 - 15HW_2 - 28HW_1 - 96HW_0$	$6HW_0 + 15HW_1 - 32HW_2 + 12HW_3$
12	$326HW_3 - 52HW_2 - 96HW_1 - 177HW_0$	$15HW_0 - 32HW_1 + 24HW_2 - 6HW_3$
13	$600HW_3 - 96HW_2 - 177HW_1 - 326HW_0$	$24HW_1 - 32HW_0 + 24HW_2 - 15HW_3$

As special cases, the  $n$ th dual hyperbolic Pierre numbers and the  $n$ th dual hyperbolic Pierre Lucas numbers are given as

$$HP_n = P_n + jP_{n+1} \tag{2.3}$$

and

$$HC_n = C_n + jC_{n+1} \tag{2.4}$$

respectively. The sequences  $\{HP_n\}_{n \geq 0}$  and  $\{HC_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$HP_{-n} = 2P_{-(n-3)} - P_{-(n-4)},$$

and

$$HC_{-n} = 2C_{-(n-3)} - C_{-(n-4)},$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrence (2.3) and (2.4) holds for all integer  $n$ .

For hyperbolic Pierre numbers (taking  $W_n = P_n$ ,  $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 4$ .) we get

$$\begin{aligned} HP_0 &= j, \\ HP_1 &= 2j + 1, \\ HP_2 &= 4j + 2, \end{aligned}$$

and for hyperbolic Pierre Lucas numbers (taking  $W_n = C_n$ ,  $C_0 = 4, C_1 = 2, C_2 = 4, C_3 = 8$ ,) we get

$$HC_0 = 2j + 4,$$

$$HC_1 = 4j + 2,$$

$$HC_2 = 8j + 4.$$

Selected values of the hyperbolic Pierre numbers and hyperbolic Pierre Lucas numbers for both positive and negative subscripts are presented in Table 3 and Table 4, respectively.

Table 3. Hyperbolic Pierre numbers

$n$	$HP_n$	$HP_{-n}$
0	$j$	$j$
1	$2j + 1$	0
2	$4j + 2$	0
3	$8j + 4$	-1
4	$15j + 8$	$-j$
5	$28j + 15$	0

Table 4. Hyperbolic Pierre- Lucas numbers

$n$	$HC_n$	$HC_{-n}$
0	$2j + 4$	$2j + 4$
1	$4j + 2$	$4j$
2	$8j + 4$	0
3	$12j + 8$	6
4	$22j + 12$	$-4 + 6j$
5	$40j + 22$	$-4j$

We now present the Binet formula for the hyperbolic generalized Pierre numbers, and for the remainder of the paper, we adopt the following notational conventions.

$$\hat{\alpha} = 1 + j\alpha, \tag{2.5}$$

$$\hat{\beta} = 1 + j\beta, \tag{2.6}$$

$$\hat{\gamma} = 1 + j\gamma \tag{2.7}$$

$$\hat{\delta} = \hat{1} = 1 + j, \tag{2.8}$$

Note that we have the following identities:

$$\begin{aligned}\widehat{\alpha}^2 &= 1 + \alpha^2 + 2\alpha j, \\ \widehat{\beta}^2 &= 1 + \beta^2 + 2j\beta, \\ \widehat{\alpha}\widehat{\beta} &= 1 + \alpha\beta + (\alpha + \beta)j, \\ \widehat{\gamma}^2 &= 1 + \gamma^2 + 2j\gamma, \\ \widehat{\delta}^2 &= \widehat{1}^2 = 2 + 2j, \\ \widehat{\gamma}\widehat{\delta} &= 1 + \gamma + j + j\gamma.\end{aligned}$$

**THEOREM 3.** (*Binet's Formula*) For any integer  $n$ , the  $n$ th hyperbolic generalized Pierre number is

$$HW_n = \widehat{\alpha}A_1\alpha^n + \widehat{\beta}A_2\beta^n + \widehat{\gamma}A_3\gamma^n + \widehat{\delta}A_4 \tag{2.9}$$

where  $\widehat{\alpha}, \widehat{\beta}, \widehat{\gamma}, \widehat{\delta}$  are given as (2.5)-(2.8)

*Proof.* Using Binet's formula of the generalized Pierre numbers given below

$$W_n = A_1\alpha^n + A_2\beta^n + A_3\gamma^n + A_4$$

where  $A_1, A_2, A_3, A_4$  are given (1.6) we get

$$\begin{aligned}HW_n &= W_n + jW_{n+1}, \\ &= A_1\alpha^n + A_2\beta^n + A_3\gamma^n + A_4 + (A_1\alpha^{n+1} + A_2\beta^{n+1} + A_3\gamma^{n+1} + A_4)j \\ &= \alpha A_1\alpha^n + \widehat{\beta}A_2\beta^n + \widehat{\gamma}A_3\gamma^n + \widehat{\delta}A_4.\end{aligned}$$

This proves (2.9).

As special cases, for any integer  $n$ , the Binet's Formula of  $n$ th hyperbolic Pierre number is

$$HP_n = \frac{(\alpha^2 + \alpha + 1)\alpha^n\widehat{\alpha}}{2(\alpha^2 + \alpha - 1)} + \frac{(\beta^2 + \beta + 1)\beta^n\widehat{\beta}}{2(\beta^2 + \beta - 1)} + \frac{(\gamma^2 + \gamma + 1)\gamma^n\widehat{\gamma}}{2(\gamma^2 + \gamma - 1)} - \frac{\widehat{1}}{2} \tag{2.10}$$

and the Binet's Formula of  $n$ th hyperbolic Pierre Lucas number is

$$HC_n = \widehat{\alpha}\alpha^n + \widehat{\beta}\beta^n + \widehat{\gamma}\gamma^n + \widehat{1}. \tag{2.11}$$

Next, we present generating function.

**THEOREM 4.** *The generating function for the hyperbolic generalized Pierre numbers is*

$$f_{HW_n}(x) = \sum_{n=0}^{\infty} HW_n x^n = \frac{HW_0 + (HW_1 - 2HW_0)x + (HW_2 - 2HW_1)x^2 + (HW_3 - 2HW_2)x^3}{1 - 2x + x^4}.$$

*Proof.* We assume that  $f_{HW_n}(x)$  is the generating function of the hyperbolic generalized Pierre numbers and then we can write

$$f_{HW_n}(x) = \sum_{n=0}^{\infty} HW_n x^n.$$

Then, using the definition of the hyperbolic generalized Pierre numbers, and subtracting  $xf(x)$  and  $x^2f(x)$  from  $f(x)$ , we obtain (note the shift in the index  $n$  in the third line)

$$\begin{aligned}
 (1 - 2x + x^4)f_{HW_n}(x) &= \sum_{n=0}^{\infty} HW_n x^n - 2x \sum_{n=0}^{\infty} HW_n x^n + x^4 \sum_{n=0}^{\infty} HW_n x^n \\
 &= \sum_{n=0}^{\infty} HW_n x^n - 2 \sum_{n=0}^{\infty} HW_n x^{n+1} + \sum_{n=0}^{\infty} HW_n x^{n+4} \\
 &= \sum_{n=0}^{\infty} HW_n x^n - 2 \sum_{n=1}^{\infty} HW_{(n-1)} x^n + \sum_{n=4}^{\infty} HW_{(n-4)} x^n \\
 &= (HW_0 + HW_1 x + HW_2 x^2 + HW_3 x^3) - 2(HW_0 x + HW_1 x^2 + HW_2 x^3) \\
 &\quad + \sum_{n=4}^{\infty} (HW_n - 2HW_{n-1} + HW_{n-4}) x^n \\
 &= HW_0 + (HW_1 - 2HW_0)x + (HW_2 - 2HW_1)x^2 + (HW_3 - 2HW_2)x^3.
 \end{aligned}$$

As special cases, the generating functions for the hyperbolic Pierre and hyperbolic Pierre Lucas numbers are

$$\sum_{n=0}^{\infty} HP_n x^n = \frac{j + x}{1 - 2x + x^4}$$

and

$$\sum_{n=0}^{\infty} HC_n x^n = \frac{2j + 4 - 6x - 4jx^3}{1 - 2x + x^4}$$

respectively.

Next, we give the exponential generating function of  $\sum_{n=0}^{\infty} HW_n \frac{x^n}{n!}$  of the sequence  $HW_n$ .

LEMMA 5. Suppose that  $f_{HW_n}(x) = \sum_{n=0}^{\infty} HW_n \frac{x^n}{n!}$  is the exponential hyperbolic generating function of the generalized Pierre sequence  $\{HW_n\}$ .

Then  $\sum_{n=0}^{\infty} HW_n \frac{x^n}{n!}$  is given by

$$\sum_{n=0}^{\infty} HW_n \frac{x^n}{n!} = A_1 e^{\alpha x} \hat{\alpha} + A_2 e^{\beta x} \hat{\beta} + A_3 e^{\gamma x} \hat{\gamma} + A_4 e^{x} \hat{1}.$$

where  $\hat{\alpha}, \hat{\beta}, \hat{\gamma}, \hat{\delta}$  are given as (2.5)-(2.8)

*Proof.* Using Binet's formula

$$W_n = A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4.$$

where  $A_1, A_2, A_3, A_4$  are given as in (1.6) we get

$$\begin{aligned} \sum_{n=0}^{\infty} \widehat{W}_n \frac{x^n}{n!} &= \sum_{n=0}^{\infty} W_n \frac{x^n}{n!} + j \sum_{n=0}^{\infty} W_{n+1} \frac{x^n}{n!} \\ &= \sum_{n=0}^{\infty} (A_1 \alpha^n + A_2 \beta^n + A_3 \gamma^n + A_4) \frac{x^n}{n!} + j \sum_{n=0}^{\infty} (A_1 \alpha^{n+1} + A_2 \beta^{n+1} + A_3 \gamma^{n+1} + A_4) \frac{x^n}{n!} \\ &= (A_1 e^{\alpha x} + A_2 e^{\beta x} + A_3 e^{\gamma x} + A_4 e^x) + j(A_1 \alpha e^{\alpha x} + A_2 \beta e^{\beta x} + A_3 \gamma e^{\gamma x} + A_4 e^x) \\ &= A_1 e^{\alpha x} (1 + j\alpha) + A_2 e^{\beta x} (1 + j\beta) + A_3 e^{\gamma x} (1 + j\gamma) + A_4 e^x (1 + j) \\ &= A_1 e^{\alpha x} \widehat{\alpha} + A_2 e^{\beta x} \widehat{\beta} + A_3 e^{\gamma x} \widehat{\gamma} + A_4 e^x \widehat{1} \end{aligned}$$

This proves (5).  $\square$

The previous Lemma gives the following results as particular examples.

**COROLLARY 6.** *Exponential generating function of hiperbolic Pierre and hiperbolic Pierre-Lucas numbers are*

a):

$$\begin{aligned} \sum_{n=0}^{\infty} HP_n \frac{x^n}{n!} &= \left( \frac{(\alpha^2 + \alpha + 1)}{2(\alpha^2 + \alpha - 1)} e^{\alpha x} + \frac{(\beta^2 + \beta + 1)}{2(\beta^2 + \beta - 1)} e^{\beta x} + \frac{(\gamma^2 + \gamma + 1)}{2(\gamma^2 + \gamma - 1)} e^{\gamma x} - \frac{1}{2} e^x \right) \\ &\quad + j \left( \frac{\alpha(\alpha^2 + \alpha + 1)}{2(\alpha^2 + \alpha - 1)} e^{\alpha x} + \frac{\beta(\beta^2 + \beta + 1)}{2(\beta^2 + \beta - 1)} e^{\beta x} + \frac{\gamma(\gamma^2 + \gamma + 1)}{2(\gamma^2 + \gamma - 1)} e^{\gamma x} - \frac{1}{2} e^x \right) \end{aligned}$$

b):

$$\sum_{n=0}^{\infty} HC_n \frac{x^n}{n!} = e^{\alpha x} + e^{\beta x} + e^{\gamma x} + e^x + j(\alpha e^{\alpha x} + \beta e^{\beta x} + \gamma e^{\gamma x} + e^x).$$

### 3. Obtaining Binet Formula From Generating Function

Next, we derive the Binet formula for the generalized hyperbolic Pierre numbers  $\{HW_n\}$  by utilizing their corresponding generating function.

**THEOREM 7.** *Binet's formula of generalized hyperbolic Pierre numbers:*

$$HW_n = \frac{q_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{q_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{q_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{q_4 \delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}, \tag{3.1}$$

where

$$\begin{aligned} q_1 &= HW_0 \alpha^3 + (HW_1 - 2HW_0) \alpha^2 + (HW_2 - 2HW_1) \alpha + HW_3 - 2HW_2, \\ q_2 &= HW_0 \beta^3 + (HW_1 - 2HW_0) \beta^2 + (HW_2 - 2HW_1) \beta + HW_3 - 2HW_2, \\ q_3 &= HW_0 \gamma^3 + (HW_1 - 2HW_0) \gamma^2 + (HW_2 - 2HW_1) \gamma + HW_3 - 2HW_2, \\ q_4 &= HW_0 \delta^3 + (HW_1 - 2HW_0) \delta^2 + (HW_2 - 2HW_1) \delta + HW_3 - 2HW_2. \end{aligned}$$

*Proof.* Let

$$h(x) = x^4 - 2x + 1.$$

Then for some  $\alpha, \beta, \gamma$  and  $\delta$  we write

$$h(x) = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x).$$

i.e.,

$$x^4 - 2x + 1 = (1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x). \tag{3.2}$$

Hence  $\frac{1}{\alpha}, \frac{1}{\beta}, \frac{1}{\gamma}$  and  $\frac{1}{\delta}$  are the roots of  $h(x)$ . This gives  $\alpha, \beta, \gamma$  and  $\delta$  as the roots of

$$h\left(\frac{1}{x}\right) = h\left(\frac{1}{x}\right) = 1 - \frac{2}{x} + \frac{1}{x^4} = 0.$$

This implies  $x^4 - 2x + 1 = 0$ . Now, by it follows that

$$\sum_{n=0}^{\infty} HW_n x^n = \frac{(HW_3 - 2HW_2)x^3 + (HW_2 - 2HW_1)x^2 + (HW_1 - 2HW_0)x + HW_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)}.$$

Then we write

$$\frac{(HW_3 - 2HW_2)x^3 + (HW_2 - 2HW_1)x^2 + (HW_1 - 2HW_0)x + HW_0}{(1 - \alpha x)(1 - \beta x)(1 - \gamma x)(1 - \delta x)} = \frac{B_1}{(1 - \alpha x)} + \frac{B_2}{(1 - \beta x)} + \frac{B_3}{(1 - \gamma x)} + \frac{B_4}{(1 - \delta x)}. \tag{3.3}$$

So

$$\begin{aligned} & (HW_3 - 2HW_2)x^3 + (HW_2 - 2HW_1)x^2 + (HW_1 - 2HW_0)x + HW_0 \\ &= B_1(1 - \beta x)(1 - \gamma x)(1 - \delta x) + B_2(1 - \alpha x)(1 - \gamma x)(1 - \delta x) \\ & \quad + B_3(1 - \alpha x)(1 - \beta x)(1 - \delta x) + B_4(1 - \alpha x)(1 - \beta x)(1 - \gamma x). \end{aligned}$$

If we consider  $x = \frac{1}{\alpha}$ , we get  $HW_0 + \frac{1}{\alpha}(HW_1 - 2HW_0) + \frac{1}{\alpha^2}(HW_2 - 2HW_1) + \frac{1}{\alpha^3}(HW_3 - 2HW_2) = -B_1\left(\frac{1}{\alpha}\beta - 1\right)\left(\frac{1}{\alpha}\gamma - 1\right)\left(\frac{1}{\alpha}\delta - 1\right)$ .

This gives

$$\begin{aligned} B_1 &= \alpha^3(HW_0 + \frac{1}{\alpha^2}(HW_2 - 2HW_1) + \frac{1}{\alpha^3}(HW_3 - 2HW_2) + \frac{1}{\alpha}(HW_1 - 2HW_0)) \\ &= \frac{HW_0\alpha^3 + (HW_1 - 2HW_0)\alpha^2 + (HW_2 - 2HW_1)\alpha + HW_3 - 2HW_2}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} B_2 &= \frac{HW_0\beta^3 + (HW_1 - 2HW_0)\beta^2 + (HW_2 - 2HW_1)\beta + HW_3 - 2HW_2}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)}, \\ B_3 &= \frac{HW_0\gamma^3 + (HW_1 - 2HW_0)\gamma^2 + (HW_2 - 2HW_1)\gamma + HW_3 - 2HW_2}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)}, \\ B_4 &= \frac{HW_0\delta^3 + (HW_1 - 2HW_0)\delta^2 + (HW_2 - 2HW_1)\delta + HW_3 - 2HW_2}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}. \end{aligned}$$

Thus (3.3) can be written as

$$\sum_{n=0}^{\infty} HW_n x^n = B_1(1 - \alpha x)^{-1} + B_2(1 - \beta x)^{-1} + B_3(1 - \gamma x)^{-1} + B_4(1 - \delta x)^{-1}.$$

This gives

$$\sum_{n=0}^{\infty} HW_n x^n = B_1 \sum_{n=0}^{\infty} \alpha^n x^n + B_2 \sum_{n=0}^{\infty} \beta^n x^n + B_3 \sum_{n=0}^{\infty} \gamma^n x^n + B_4 \sum_{n=0}^{\infty} \delta^n x^n = \sum_{n=0}^{\infty} (B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n) x^n.$$

Therefore, comparing coefficients on both sides of the above equality, we obtain

$$HW_n = B_1 \alpha^n + B_2 \beta^n + B_3 \gamma^n + B_4 \delta^n.$$

**THEOREM 8.** *For all integers  $m, n$  the following identities holds:*

$$HW_{m+n} = P_{m-2} HW_{n+3} - P_{m-5} HW_{n+2} - P_{m-4} HW_{n+1} - P_{m-3} HW_n.$$

*Proof.* First we assume that  $m, n \geq 0$  theorem 8 can be proved by mathematical induction on  $m$ . If  $m = 0$  we get

$$HW_n = P_{-2} HW_{n+3} - P_{-5} HW_{n+2} - P_{-4} HW_{n+1} - P_{-3} HW_n.$$

which is true since  $P_{-2} = 0, P_{-3} = -1, P_{-4} = 0, P_{-5} = 0$ . Suppose that the equality holds for  $m \leq k$ . For  $m = k + 1$ , we obtain

$$\begin{aligned} HW_{k+1+n} &= 2HW_{n+k} + -HW_{n+k-3}, \\ &= 2(P_{k-2} HW_{n+3} - P_{k-5} HW_{n+2} - P_{k-4} HW_{n+1} - P_{k-3} HW_n) \\ &\quad - (2P_{k-5} HW_{n+3} - P_{k-8} HW_{n+2} - P_{k-6} HW_{n+1} - P_{k-6} HW_n) \end{aligned}$$

by mathematical induction on  $m$ , this proves Theorem 8.

The other cases of  $m, n$  can be proved similarly for all integers  $m, n$ .  $\square$

Taking  $HW_n = HP_n$  or  $HW_n = HC_n$  in above Theorem, respectively, we obtain:

**COROLLARY 9.**

$$\begin{aligned} HP_{m+n} &= P_{m-2} HP_{n+3} - P_{m-5} HP_{n+2} - P_{m-4} HP_{n+1} - P_{m-3} HP_n, \\ HC_{m+n} &= P_{m-2} HC_{n+3} - P_{m-5} HC_{n+2} - P_{m-4} HC_{n+1} - P_{m-3} HC_n. \end{aligned}$$

#### 4. SIMSON'S FORMULA

In this section, we present Simpson's formula for the hyperbolic generalized Pierre numbers, which constitutes a special case of [29, Theorem 4.1].

**THEOREM 10.** *(Simpson's formula for hyperbolic generalized Pierre numbers) For all integers  $n$  we have,*

$$\begin{aligned} & \begin{vmatrix} HW_{n+3} & HW_{n+2} & HW_{n+1} & HW_n \\ HW_{n+2} & HW_{n+1} & HW_n & HW_{n-1} \\ HW_{n+1} & HW_n & HW_{n-1} & HW_{n-2} \\ HW_n & HW_{n-1} & HW_{n-2} & HW_{n-3} \end{vmatrix} = \begin{vmatrix} HW_3 & HW_2 & HW_1 & HW_0 \\ HW_2 & HW_1 & HW_0 & HW_{-1} \\ HW_1 & HW_0 & HW_{-1} & HW_{-2} \\ HW_0 & HW_{-1} & HW_{-2} & HW_{-3} \end{vmatrix} \\ & = (HW_3 - HW_2 - HW_1 - HW_0)(HW_3^3 - HW_2^3 - HW_1^3 - HW_0^3 + (-5HW_2 + HW_1 + HW_0)HW_3^2 + \\ & (7HW_3 - 3HW_0 - HW_1)HW_2^2 \\ & + (3HW_3 + HW_2 - HW_0)HW_1^2 + (HW_3 + HW_2 + HW_1)HW_0^2 + 4(-HW_2HW_3 - HW_0HW_3 + HW_0HW_2)HW_1). \end{aligned}$$

Proof. Using Theorem 10 it can be proved by using induction or use [29, Theorem 4.1]

From the Theorem 10 we get the following Corollary.

**COROLLARY 11.** *For all integers  $n$ , the Simson's formulas of dual hyperbolic Pierre numbers and dual hyperbolic Pierre Lucas numbers are given as,*

$$\begin{aligned} \text{a): } & \begin{vmatrix} HP_{n+3} & HP_{n+2} & HP_{n+1} & HP_n \\ HP_{n+2} & HP_{n+1} & HP_n & HP_{n-1} \\ HP_{n+1} & HP_n & HP_{n-1} & HP_{n-2} \\ HP_n & HP_{n-1} & HP_{n-2} & HP_{n-3} \end{vmatrix} = 2 + 2j, \\ \text{b): } & \begin{vmatrix} HC_{n+3} & HC_{n+2} & HC_{n+1} & HC_n \\ HC_{n+2} & HC_{n+1} & HC_n & HC_{n-1} \\ HC_{n+1} & HC_n & HC_{n-1} & HC_{n-2} \\ HC_n & HC_{n-1} & HC_{n-2} & HC_{n-3} \end{vmatrix} = -352 - 352j, \end{aligned}$$

respectively.

### 5. Linear Sums

In this section, we present the summation formulas for the hyperbolic generalized Pierre numbers corresponding to both positive and negative subscripts.

We now present the summation formulas for the generalized Pierre numbers.

**THEOREM 12.** *For the dual hyperbolic Pierre numbers, we have the following formulas:*

$$\begin{aligned} \text{(a): } & \sum_{k=0}^n W_k = \frac{1}{2}(-(n+3)W_{n+3} + (n+4)W_{n+2} + (n+3)W_{n+1} + (n+4)W_n + 3W_3 - 4W_2 - 3W_1 - 2W_0). \\ \text{(b): } & \sum_{k=0}^n W_{2k} = \frac{1}{2}(-(n+2)W_{2n+2} + (n+3)W_{2n+1} + (n+3)W_{2n} + (n+2)W_{2n-1} + 2W_3 - 2W_2 - 3W_1 - W_0). \\ \text{(c): } & \sum_{k=0}^n W_{2k+1} = \frac{1}{2}(-(n+1)W_{2n+2} + (n+3)W_{2n+1} + (n+2)W_{2n} + (n+2)W_{2n-1} + 2W_3 - 3W_2 - \\ & W_1 - 2W_0). \end{aligned}$$

Proof. For the proof, see Soykan [27, Theorem 3.10].  $\square$

**THEOREM 13.** *For the hyperbolic Pierre numbers, we have the following formulas:*

- (a):  $\sum_{k=0}^n HW_k = \frac{1}{2}(- (n + 3))HW_{n+3} + (n + 4)HW_{n+2} + (n + 3)HW_{n+1} + (n + 4)HW_n + 3HW_3 - 4HW_2 - 3HW_1 - 2HW_0$ .
- (b):  $\sum_{k=0}^n HW_{2k} = \frac{1}{2}(- (n + 2))HW_{2n+2} + (n + 3)HW_{2n+1} + (n + 3)HW_{2n} + (n + 2)HW_{2n-1} + 2HW_3 - 2HW_2 - 3HW_1 - HW_0$ .
- (c):  $\sum_{k=0}^n HW_{2k+1} = \frac{1}{2}(- (n + 1))HW_{2n+2} + (n + 3)HW_{2n+1} + (n + 2)HW_{2n} + (n + 2)HW_{2n-1} + 2HW_3 - 3HW_2 - HW_1 - 2HW_0$ .

Proof. Use Theorem 12 and the definition of  $HW_n$ .  $\square$

As a special case of Theorem 13, we state the following Corollary.

COROLLARY 14. For  $n \geq 0$ , dual hyperbolic Pierre numbers have the following properties:

- (a):  $\sum_{k=0}^n HP_k = \frac{1}{2}(- (n + 3))HP_{n+3} + (n + 4)HP_{n+2} + (n + 3)HP_{n+1} + (n + 4)HP_n + 1$ .
- (b):  $\sum_{k=0}^n HP_{2k} = \frac{1}{2}(- (n + 2))HP_{2n+2} + (n + 3)HP_{2n+1} + (n + 3)HP_{2n} + (n + 2)HP_{2n-1} + j + 1$ .
- (c):  $\sum_{k=0}^n HP_{2k+1} = \frac{1}{2}(- (n + 1))HP_{2n+2} + (n + 3)HP_{2n+1} + (n + 2)HP_{2n} + (n + 2)HP_{2n-1} + 1$ .

As a second special case of the above theorem, we obtain the following summation formulas for the hyperbolic Pierre Lucas numbers:

COROLLARY 15. For  $n \geq 0$ , the hyperbolic Pierre Lucas numbers satisfy the following properties.

- (a):  $\sum_{k=0}^n HC_k = \frac{1}{2}(- (n + 3))HC_{n+3} + (n + 4)HC_{n+2} + (n + 3)HC_{n+1} + (n + 4)HC_n - 12j - 6$ .
- (b):  $\sum_{k=0}^n HC_{2k} = \frac{1}{2}(- (n + 2))HC_{2n+2} + (n + 3)HC_{2n+1} + (n + 3)HC_{2n} + (n + 2)HC_{2n-1} - 6j - 2$ .
- (c):  $\sum_{k=0}^n HC_{2k+1} = \frac{1}{2}(- (n + 1))HC_{2n+2} + (n + 3)HC_{2n+1} + (n + 2)HC_{2n} + (n + 2)HC_{2n-1} - 8j - 6$ .

Next, we present the ordinary generating functions corresponding to selected special cases of the hyperbolic generalized Pierre numbers.

THEOREM 16. The ordinary generating functions of the sequences  $HW_{2n}$ ,  $HW_{2n+1}$  are given as follows:

- (a):  $\sum_{n=0}^{\infty} HW_{2n}x^n = \frac{HW_3(2x^2)+HW_2(x^3-4x^2+x)-HW_1(2x^3)+HW_0(x^2-4x+1)}{x^4+2x^2-4x+1}$ .
- (b):  $\sum_{n=0}^{\infty} HW_{2n+1}x^n = \frac{HW_3(x^3+x)-HW_2(2x^3)-HW_1(x^2-4x+1)-HW_0(2x^2)}{x^4+2x^2-4x+1}$ .

Proof. Similarly, the proof can be constructed as in [4]

From the preceding theorem, we derive the following Corollary, which provides a summation formula for the hyperbolic Pierre numbers. (Take  $HW_n = HP_n$  with  $HP_0 = j, HP_1 = 2j + 1, HP_2 = 4j + 2, HP_3 = 8j + 4$ )

COROLLARY 17.  $n \geq 0$  the hyperbolic Pierre numbers exhibit the following properties.

- (a):  $\sum_{n=0}^{\infty} HP_{2n}x^n = \frac{j+2x+jx^2}{x^4+2x^2-4x+1}$ .
- (b):  $\sum_{n=0}^{\infty} HP_{2n+1}x^n = \frac{(-2j-1)+(8+16j)x+(-4j-1)x^2}{x^4+2x^2-4x+1}$ .

### 6. Matrices related with Hyperbolic Generalized Pierre Numbers

We define the square matrix  $A$  of order 4 as

$$A = \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

such that  $\det A = 1$ . Note that

$$A^n = \begin{pmatrix} P_{n+1} & -P_{n-2} & -P_{n-1} & -P_n \\ P_n & -P_{n-3} & -P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-4} & -P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-5} & -P_{n-4} & -P_{n-3} \end{pmatrix}$$

for the proof see [28].

Then we give the following lemma.

LEMMA 18. For  $n \geq 0$  the following identity is true:

$$\begin{pmatrix} HW_{n+3} \\ HW_{n+2} \\ HW_{n+1} \\ HW_n \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n \begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix}.$$

*Proof.* The identity(18) can be proved by mathematical induction on  $n$ . If  $n = 0$  we obtain

$$\begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix} = \begin{pmatrix} 2 & -1 & 1 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^0 \begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix}$$

which is true. Assuming that the given identity holds for  $n = k$ , the following identity is consequently valid.

$$\begin{pmatrix} HW_{k+3} \\ HW_{k+2} \\ HW_{k+1} \\ HW_k \end{pmatrix} = \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix}.$$

For  $n = k + 1$ , we get

$$\begin{aligned}
 \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^{k+1} \begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix} &= \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^k \begin{pmatrix} HW_3 \\ HW_2 \\ HW_1 \\ HW_0 \end{pmatrix} \\
 &= \begin{pmatrix} 2 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} HW_{k+3} \\ HW_{k+2} \\ HW_{k+1} \\ HW_k \end{pmatrix} \\
 &= \begin{pmatrix} HW_{k+4} \\ HW_{k+3} \\ HW_{k+2} \\ HW_{k+1} \end{pmatrix}.
 \end{aligned}$$

Thus, the proof is completed.  $\square$

We define

$$N_{HW} = \begin{pmatrix} HW_3 & HW_2 & HW_1 & HW_0 \\ HW_2 & HW_1 & HW_0 & HW_{-1} \\ HW_1 & HW_0 & HW_{-1} & HW_{-2} \\ HW_0 & HW_{-1} & HW_{-2} & HW_{-3} \end{pmatrix}, \tag{6.1}$$

$$E_{HW} = \begin{pmatrix} HW_{n+3} & HW_{n+2} & HW_{n+1} & HW_n \\ HW_{n+2} & HW_{n+1} & HW_n & HW_{n-1} \\ HW_{n+1} & HW_n & HW_{n-1} & HW_{n-2} \\ HW_n & HW_{n-1} & HW_{n-2} & HW_{n-3} \end{pmatrix}. \tag{6.2}$$

Now, we have the following theorem with  $N_{HW}$  and  $E_{HW}$ ,

**THEOREM 19.** *Using  $N_{HW}$  and  $E_{HW}$ , we get*

$$A^n N_{HW} = E_{HW}.$$

*Proof.* Note that we get

$$\begin{aligned}
 A^n N_{HW} &= \begin{pmatrix} P_{n+1} & -P_{n-2} & -P_{n-1} & -P_n \\ P_n & -P_{n-3} & -P_{n-2} & -P_{n-1} \\ P_{n-1} & -P_{n-4} & -P_{n-3} & -P_{n-2} \\ P_{n-2} & -P_{n-5} & -P_{n-4} & -P_{n-3} \end{pmatrix} \begin{pmatrix} HW_3 & HW_2 & HW_1 & HW_0 \\ HW_2 & HW_1 & HW_0 & HW_{-1} \\ HW_1 & HW_0 & HW_{-1} & HW_{-2} \\ HW_0 & HW_{-1} & HW_{-2} & HW_{-3} \end{pmatrix} \\
 &= \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{pmatrix}
 \end{aligned}$$

where

$$\begin{aligned}
 a_{11} &= P_{n+1}HW_3 - P_{n-2}HW_2 - P_{n-1}HW_1 - P_nHW_0 = HW_{n+3}, \\
 a_{12} &= P_{n+1}HW_2 - P_{n-2}HW_1 - P_{n-1}HW_0 - P_nHW_{-1} = HW_{n+2}, \\
 a_{13} &= P_{n+1}HW_1 - P_{n-2}HW_0 - P_{n-1}HW_{-1} - P_nHW_{-2} = HW_{n+1}, \\
 a_{14} &= P_{n+1}HW_0 - P_{n-2}HW_{-1} - P_{n-1}HW_{-2} - P_nHW_{-3} = HW_n, \\
 a_{21} &= P_nHW_3 - P_{n-3}HW_2 - P_{n-2}HW_1 - P_{n-1}HW_0 = HW_{n+2}, \\
 a_{22} &= P_nHW_2 - P_{n-3}HW_1 - P_{n-2}HW_0 - P_{n-1}HW_{-1} = HW_{n+1}, \\
 a_{23} &= P_nHW_1 - P_{n-3}HW_0 - P_{n-2}HW_{-1} - P_{n-1}HW_{-2} = HW_n, \\
 a_{24} &= P_nHW_0 - P_{n-3}HW_{-1} - P_{n-2}HW_{-2} - P_{n-1}HW_{-3} = HW_{n-1}, \\
 a_{31} &= P_{n-1}HW_3 - P_{n-4}HW_2 - P_{n-3}HW_1 - P_{n-2}HW_0 = HW_{n+1}, \\
 a_{32} &= P_{n-1}HW_2 - P_{n-4}HW_1 - P_{n-3}HW_0 - P_{n-2}HW_{-1} = HW_n, \\
 a_{33} &= P_{n-1}HW_1 - P_{n-4}HW_0 - P_{n-3}HW_{-1} - P_{n-2}HW_{-2} = HW_{n-1}, \\
 a_{34} &= P_{n-1}HW_0 - P_{n-4}HW_{-1} - P_{n-3}HW_{-2} - P_{n-2}HW_{-3} = HW_{n-2}, \\
 a_{41} &= P_{n-2}HW_3 - P_{n-5}HW_2 - P_{n-4}HW_1 - P_{n-3}HW_0 = HW_n, \\
 a_{42} &= P_{n-2}HW_2 - P_{n-5}HW_1 - P_{n-4}HW_0 - P_{n-3}HW_{-1} = HW_{n-1}, \\
 a_{43} &= P_{n-2}HW_1 - P_{n-5}HW_0 - P_{n-4}HW_{-1} - P_{n-3}HW_{-2} = HW_{n-2}, \\
 a_{44} &= P_{n-2}HW_0 - P_{n-5}HW_{-1} - P_{n-4}HW_{-2} - P_{n-3}HW_{-3} = HW_{n-3}.
 \end{aligned}$$

Using the theorem (8) the proof is done.  $\square$

By taking  $HW_n = HP_n$  with  $HP_0, HP_1, HP_2, HP_3$  in (6.1) and (6.2)

$HW_n = C_n$  with  $HC_0, HC_1, HC_2, HC_3$  in (6.1) and (6.2)

respectively, we get:

$$\begin{aligned}
 N_{HP} &= \begin{pmatrix} 8j+4 & 4j+2 & 2j+1 & j \\ 4j+2 & 2j+1 & j & 0 \\ 2j+1 & j & 0 & 0 \\ j & 0 & 0 & -1 \end{pmatrix}, \\
 E_{HP} &= \begin{pmatrix} HP_{n+3} & HP_{n+2} & HP_{n+1} & HP_n \\ HP_{n+2} & HP_{n+1} & HP_n & HP_{n-1} \\ HP_{n+1} & HP_n & HP_{n-1} & HP_{n-2} \\ HP_n & HP_{n-1} & HP_{n-2} & HP_{n-3} \end{pmatrix}, \\
 N_{HC} &= \begin{pmatrix} 12j+8 & 8j+4 & 4j+2 & 2j+4 \\ 8j+4 & 4j+2 & 2j+4 & 4j \\ 4j+2 & 2j+4 & 4j & 0 \\ 2j+4 & 4j & 0 & 6 \end{pmatrix}, \\
 E_{HC} &= \begin{pmatrix} HC_{n+3} & HC_{n+2} & HC_{n+1} & HC_n \\ HC_{n+2} & HC_{n+1} & HC_n & HC_{n-1} \\ HC_{n+1} & C_n & HC_{n-1} & HC_{n-2} \\ HC_n & HC_{n-1} & HC_{n-2} & HC_{n-3} \end{pmatrix}.
 \end{aligned}$$

From Theorem [19], we can write the following corollary.

**COROLLARY 20.** *The following identities are hold:*

**a):**  $A^n N_{HP} = E_{HP}$ .

**b):**  $A^n N_{HC} = E_{HC}$ .

### References

- [1] M. Akar, S. Yüce, Ş. Şahin, On the Dual Hyperbolic Numbers and the Complex Hyperbolic Numbers, Journal of Computer Science & Computational Mathematics, 8(1) (2018), 1-6.
- [2] Aydın, F. T., Hyperbolic Fibonacci Sequence, Universal Journal of Mathematics and Applications, 2 (2), 59-64, 2019.
- [3] J. Baez, The octonions, Bull. Amer. Math. Soc. 39(2) (2002), 145-205.
- [4] D.K. Biss, D. Dugger, D.C. Isaksen, Large annihilators in Cayley-Dickson algebras, Communication in Algebra, 36 (2), 632-664, 2008.
- [5] H. H. Cheng, S. Thompson, Dual Polynomials and Complex Dual Numbers for Analysis of Spatial Mechanisms, Proc. of ASME 24th Biennial Mechanisms Conference, Irvine, CA, August, 1996, 19-22.
- [6] J. Cockle, On a New Imaginary in Algebra, Philosophical magazine, London-Dublin-Edinburgh, 3(34) (1849), 37-47.
- [7] Soykan, Y., Demirci, F., Dual Hyperbolic Generalized Adrien Numbers, Asian Research Journal of Mathematics, 21(8), (2025), 63-91.

- [8] Dikmen, C. M., Altınsoy, M., On Third Order Hyperbolic Jacobsthal Numbers, *Konuralp Journal of Mathematics*, 10 (1), 118-126, 2022.
- [9] Soykan, Y. Ayrılmaz, E. A Study On Hyperbolic Generalized Edouard Numbers, *Asian Journal of Probability and Statistics*, 27 (8), 52-79, 2025.
- [10] P. Fjelstad, S.G. Gal, n-dimensional Hyperbolic Complex Numbers, *Advances in Applied Clifford Algebras*, 8(1) (1998), 47-68, .
- [11] W.R. Hamilton, *Elements of Quaternions*, Chelsea Publishing Company, New York (1969).
- [12] K. Imaeda, M. Imaeda, Sedenions: algebra and analysis, *Applied Mathematics and Computation*, 115 (2000), 77-88.
- [13] B. Jancewicz, The extended Grassmann algebra of  $R^3$ , in *Clifford (Geometric) Algebras with Applications and Engineering*, Birkhauser, Boston, 389-421, 1996. [https://doi.org/10.1007/978-1-4612-4104-1\\_28](https://doi.org/10.1007/978-1-4612-4104-1_28)
- [14] I. Kantor, A. Solodovnikov, *Hypercomplex Numbers*, Springer-Verlag, New York (1989).
- [15] Kalca, F. Soykan, Y. A Study on Dual Hyperbolic Generalized Pandita Numbers. *Archives of Current Research International*, 25(8), 412-436, 2025.
- [16] V. V. Kravchenko, *Hyperbolic Numbers and Analytic Functions*. In: *Applied Pseudoanalytic Function Theory. Frontiers in Mathematics*. Birkhäuser Basel, 2009. DOI: [https://doi.org/10.1007/978-3-0346-0004-0\\_11](https://doi.org/10.1007/978-3-0346-0004-0_11).
- [17] A. Khrennikov, and Segre. G, *An Introduction to Hyperbolic Analysis*, <http://arxiv.org/abs/math-ph/0507053v2>, 2005.
- [18] Taş, S., On Hyperbolic Jacobsthal-Lucas Sequence, *Fundamental Journal of Mathematics and Applications*, 5(1), 16-20, 2022.52v3, *MathNT*, 2021.
- [19] G. Moreno, The zero divisors of the Cayley-Dickson algebras over the real numbers, *Bol. Soc. Mat. Mexicana* 3(4) (1998), 13-28.
- [20] A. E. Motter, and A. F. Rosa, Hyperbolic calculus, *Adv. Appl. Clifford Algebr.*, 8(1) (1998), 109-128. <https://doi.org/10.1007/BF03041929>
- [21] Soykan, Y. Dogan, S., Gaussian Generalized Pierre Numbers, *Asian Research Journal of Mathematics*, 21(7), (2025), 99-119.
- [22] Soykan, Y. Yılmaz, B. A Study on Hyperbolic Generalized Guglielmo Numbers, *Asian Research Journal of Mathematics*, 19(12), (2023), 120-147.
- [23] Soykan, Y., Taşdemir, E., Okumuş, İ., On dual hyperbolic numbers with generalized Jacobsthal numbers components, *Indian J Pure Appl Math*, 54, 824–840, 2023.
- [24] G. Sobczyk, The Hyperbolic Number Plane, *The College Mathematics Journal*, 26(4) (1995), 268-280.
- [25] G. Sobczyk, *Complex and Hyperbolic Numbers*. In: *New Foundations in Mathematics*. Birkhäuser, Boston, 2013. DOI: [https://doi.org/10.1007/978-0-8176-8385-6\\_2](https://doi.org/10.1007/978-0-8176-8385-6_2).
- [26] Y. Soykan, Tribonacci and Tribonacci-Lucas Sedenions. *Mathematics* 7(1) (2019), 74.
- [27] Soykan, Y., Sums and Generating Functions of Special Cases of Generalized Tetranacci Polynomials, *International Journal of Advances in Applied Mathematics and Mechanics*, 12(4), 34-101, 2025.
- [28] Soykan, Y., Properties of Generalized (r,s,t,u)-Numbers, *Earthline Journal of Mathematical Sciences*, 5(2), 297-327, 2021. <https://doi.org/10.34198/ejms.5221.297327>.
- [29] Soykan, Y., Simson Identity of Generalized m-step Fibonacci Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 45-56, 2019.