

## **Effect of Biofertilizer and Phosphorus on Yield and Nutrient Uptake by Black gram (*Vigna mungo* L.)**

### **Abstract**

A field experiment was conducted at experimental farm, Department of Agronomy, A field Repetition  
experiment was conducted at experimental farm, Department of Agronomy, Faculty of  
 Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan) during  
 Rabi season of 2023-24 to effect missing word...study... of Biofertilizer and Phosphorus yield and nutrient uptake by  
 of black gram variety “Pratap Urd-1” was used in this study. The result revealed that the  
 maximum yield parameter such as number of pods per plant (29.75), number of seed per pod  
 (7.78), grain yield (12.12 q/ha), straw yield (27.02 q/ha) with application of T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>  
 + PSB + VAM. The maximum nitrogen content and uptake in grain and straw (3.64, 1.52 % and  
 44.12, 41.07 kg/ha), phosphorus content and uptake (0.32, 0.30% and 3.38, 8.11 kg/ha) and  
 potassium content and uptake (1.92, 1.74% and 23.47, 47.01 kg/ha) were recorded with T<sub>10</sub>-50  
 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM.

**Key words: -Biofertilizer; Phosphorus; Nutrient content & uptake**

### **1. Introduction**

Black gram [*Vigna mungo* (L.)] also known as urad, mash bean or black gram, belongs to family Fabaceae (Leguminaceae) is an important pulse crop. It is a self-pollinated leguminous crop. The production is not sufficient to meet the per capita requirement. Pulses are the main source of dietary protein particularly for vegetarians and contribute about 14 per cent of the total protein of average Indian diet. Black gram contains about 24 percent protein, 60 percent carbohydrate, 10.9 percent moisture, 1.4 percent fat, 0.9 percent fiber, 3.2 percent minerals and vitamin viz. calcium -154 mg, phosphorus 385mg, iron 9.1 mg and small amount of vitamin B complex (Gajera *et al.* 2014).

Phosphate Solubilizing Bacteria (PSB) plays an important role in solubilization of soil P through secretion of various organic acids (formic, acetic, butyric, propionic, citric, gluconic,

succinic, oxalic, malic, maleic and lactic acids) and make it available to plant. Many fungi, bacteria and actinomycetes are potential solubilizers of bound phosphates in soil. Application of P along with PSB improved phosphorus uptake by plants and yields indicating that the PSB were able to solubilize phosphates and mobilize phosphorus in crop plants. Phosphorus solubilizing bacteria play important role in phosphorus nutrition by enhancing its availability to plants through release from inorganic and organic soil P pools by solubilization and mineralization. Greater efficiency of phosphorus solubilizing bacteria has been shown through co-inoculation with other beneficial bacteria and mycorrhiza (Khan *et al.* 2009).

PSB & VAM inoculum, include some heterotrophic bacteria and fungi, have the ability to solubilize inorganic phosphorus from insoluble sources, such as, tricalcium phosphate, ferric, aluminum and magnesium phosphate, rock phosphate and bone meal. Vesicular Arbuscular Mycorrhiza (VAM) plays an important role in phosphorus cycling and its uptake by plants (Biswas *et al.* 2006). These symbiotic microorganisms have extensive mycelial network and can increase the transport of other mineral elements such as zinc and copper. VAM fungi can play an important role in enhancing P availability to plants in deficient soils and can save P-fertilizer by 25-30% (Somani *et al.* 2008). It is well known that VAM fungi improve plant growth through increased uptake of relatively immobile nutrients such as P, Zn, Cu). VAM fungi infect the plants, spread inside the root and produce highly branched hyphal structure, known as vesicles and arbuscules, within the host cell (Patel *et al.* 2017).

The present study was conducted to evaluate the beneficial effect of phosphorus and in combination with PSB and VAM inoculums to determine the nitrogen and phosphorus concentration in roots, shoots and grains and their uptake by black gram plant. In most of the pulses, number of branches, number of pods, seed weight and seeds per pod are the major yield components.

## 2. Materials and Methods

A field experiment was conducted during Rabi season of 2023-24 at experimental farm, Department of Agronomy, Faculty of Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan). Soil of the experimental field was sandy loam intexture, saline in reaction with a pH value of 7.6, poor in organic carbon (0.32%), deficient inavailable zinc (0.48 ppm) and iron (1.2 ppm) low in available nitrogen (176 kg/ha) and phosphorus (20.2 kg/ha) but medium in available potassium (320 kg/ha). The experiment was laid out in randomized block design with three replications consisting of ten treatments *viz.* The experiment was laid out in randomized block design with three replications and ten treatments *i.e.* T<sub>1</sub>-Control, T<sub>2</sub>-PSB, T<sub>3</sub>-VAM, T<sub>4</sub>-PSB + VAM, T<sub>5</sub>-25 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB, T<sub>7</sub>-25 kg P<sub>2</sub>O<sub>5</sub>

ha<sup>-1</sup> + PSB + VAM, T<sub>8</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB, T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM and T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB + VAM. The required quantities of fertilizers as per treatments were applied. The doses of NPK were applied in the form of urea, diammonium phosphate, murate of potash respectively. The half dose of nitrogen gives basal dose and remain two split doses after irrigation and full dose of phosphorus and potassium at basal dose.

### 3. Results and Discussion

#### 3.1 Yield attributes and yield

Data presented in Table 1.0 ~~that~~ <sup>reveals</sup> that the maximum number of pods per plant recorded with T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (29.75), it was found at par with T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM and T<sub>8</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB (28.78 and 28.15), respectively. The minimum number of pods per plant recorded with control treatment (28.15). Data reveals <sup>s</sup> that the maximum number of seed per pod recorded with T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (7.78), ~~it was found~~ at par with T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM and T<sub>8</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB (7.65 and 7.25), respectively. The minimum number of seed per pod recorded with control treatment (5.25). Data reveals <sup>s</sup> that the maximum grain yield recorded with T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (12.12 q/ha), ~~it was found~~ at par with T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM and T<sub>8</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB (11.65 and 11.15 q/ha), respectively. The minimum grain yield recorded with control treatment (8.15 q/ha). Data reveals <sup>s</sup> that the maximum straw yield recorded with T<sub>10</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (27.02 q/ha), ~~it was found~~ at par with T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM and T<sub>8</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + PSB (26.45 and 25.75 q/ha), respectively. The minimum straw yield recorded with control treatment (20.33 q/ha). Similar concluded also observed by Hakeem *et al.* (2008), Mehta *et al.* (2010), Mir *et al.* (2013), Singh and Singh (2013), Singh *et al.* (2016), Yadav *et al.* (2017) and Kachave *et al.* (2018).

keep he reporting tense to past tense

#### 3.2 Nutrient content and uptake

##### 3.2.1 Nutrient content (%)

Data revealed (Table 2.0) the maximum nitrogen content in grain was recorded with treatment was recorded T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (3.64 %). The minimum nitrogen content in grain was recorded with control treatment (3.28 %). Data revealed the maximum nitrogen content in straw was recorded with treatment was recorded T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (1.52 %). The minimum nitrogen content in straw was recorded with control treatment (1.18 %). Data revealed the maximum phosphorus content in grain was recorded with treatment ~~was recorded~~ <sup>remove</sup> T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (0.32 %). The minimum phosphorus content in grain was recorded with control treatment (0.20 %). Data revealed the maximum phosphorus content in straw was recorded with treatment was recorded T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup>

+ PSB + VAM (0.30 %). The minimum phosphorus content in straw was recorded with control treatment (0.14 %). Data revealed the maximum potassium content in grain was recorded with treatment was recorded T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (1.92 %). The minimum potassium content in grain was recorded with control treatment (1.70 %). Data revealed the maximum potassium content in straw was recorded with treatment was recorded T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (1.74 %). The minimum potassium content in straw was recorded with control treatment (1.42 %). Similar result also presented by Khan *et al.* (2009), Ali *et al.* (2012), Das *et al.* (2013), Yadav (2013), Rathour *et al.* (2015), Rani *et al.* (2016), Chaudhary *et al.* (2017) and Dhewa *et al.* (2017).

### 3.2.2 Nutrient uptake (kg/ha)

Data revealed (Table 3.0 and Figure 1.0) the maximum nitrogen uptake in grain was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (44.12 kg/ha). The minimum nitrogen uptake in grain was recorded with control treatment (26.73 kg/ha). Data revealed the maximum nitrogen uptake in straw was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (41.07 kg/ha). The minimum nitrogen uptake in straw was recorded with control treatment (23.99 kg/ha). Data revealed the maximum phosphorus uptake in grain was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (3.88 kg/ha). The minimum phosphorus uptake in grain was recorded with control treatment (1.63 kg/ha). Data revealed the maximum phosphorus uptake in straw was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (8.11 kg/ha). The minimum phosphorus uptake in straw was recorded with control treatment (2.85 kg/ha). Data revealed the maximum potassium uptake in grain was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (23.47 kg/ha). The minimum potassium uptake in grain was recorded with control treatment (13.86 kg/ha). Data revealed the maximum potassium uptake in straw was recorded with treatment **was recorded** T<sub>10</sub>-50 kg P<sub>2</sub>O T<sub>9</sub>-50 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> + VAM<sub>5</sub> ha<sup>-1</sup> + PSB + VAM (47.01 kg/ha). The minimum potassium uptake in straw was recorded with control treatment (28.87 kg/ha). These findings also supported by Basu *et al.* (2006), Mohammad *et al.* (2017) and Singh *et al.* (2017).

### Conclusion

It is concluded from the present investigation that the application of phosphorus @ 50 kg/ha with co-inoculation of Bio-fertilizer (PSB + VAM) to get maximum yield and nutrients uptake of Black gram. However, the treatment 50 kg/ha and inoculation with PSB and VAM superior among all treatments.

**Table 1.0 Effect of biofertilizer and phosphorus yield attributes and yield of black gram**


<b>Treatments</b>	<b>Number of pods per plant</b>	<b>Number of seed per pod</b>	<b>Grain yield (q/ha)</b>	<b>Straw yield (q/ha)</b>
T <sub>1</sub> -Control	22.80	5.25	8.15	20.33
T <sub>2</sub> -PSB	23.10	6.52	9.10	22.36
T <sub>3</sub> -VAM	23.96	6.65	9.40	22.85
T <sub>4</sub> -PSB + VAM	24.58	6.78	9.58	23.15
T <sub>5</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	25.25	6.96	9.85	23.96
T <sub>6</sub> -25 P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	26.75	7.10	10.36	24.85
T <sub>7</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	27.63	7.18	10.78	25.25
T <sub>8</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	28.15	7.25	11.15	25.75
T <sub>9</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	28.78	7.65	11.65	26.45
T <sub>10</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	29.75	7.78	12.12	27.02
S. Em. ±	0.55	0.18	0.33	0.60
CD (0.05%)	1.62	0.55	1.00	1.80

**Table 2.0 Effect of biofertilizer and phosphorus on nitrogen, phosphorus and potassium content in grain and straw**

Treatments	Nitrogen content (%)		Phosphorus content (%)		Potassium content (%)	
	Grain	Straw	Grain	Straw	Grain	Straw
T <sub>1</sub> -Control	3.28	1.18	0.20	0.14	1.70	1.42
T <sub>2</sub> -PSB	3.32	1.22	0.23	0.16	1.75	1.48
T <sub>3</sub> -VAM	3.36	1.26	0.22	0.15	1.77	1.55
T <sub>4</sub> -PSB + VAM	3.40	1.28	0.24	0.20	1.80	1.60
T <sub>5</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	3.45	1.32	0.27	0.23	1.82	1.65
T <sub>6</sub> -25 P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	3.48	1.36	0.26	0.22	1.85	1.68
T <sub>7</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	3.52	1.40	0.29	0.25	1.88	1.69
T <sub>8</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	3.56	1.45	0.31	0.28	1.89	1.70
T <sub>9</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	3.60	1.48	0.30	0.26	1.90	1.72
T <sub>10</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	3.64	1.52	0.32	0.30	1.92	1.74
S. Em. ±	0.03	0.02	0.01	0.01	0.01	0.01
CD (0.05%)	0.08	0.07	0.02	0.04	0.03	0.04

**Table 3.0 Effect of biofertilizer and phosphorus on nitrogen, phosphorus and potassium uptake in grain and straw**

Treatments	Nitrogen uptake (kg/ha)		Phosphorus uptake (kg/ha)		Potassium uptake (kg/ha)	
	Grain	Straw	Grain	Straw	Grain	Straw
T <sub>1</sub> -Control	26.73	23.99	1.63	2.85	13.86	28.87
T <sub>2</sub> -PSB	30.21	27.28	2.09	3.58	15.93	33.09
T <sub>3</sub> -VAM	31.58	28.79	2.07	3.43	16.64	35.42
T <sub>4</sub> -PSB + VAM	32.57	29.63	2.30	4.63	17.24	37.04
T <sub>5</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	33.98	31.63	2.66	5.51	17.93	39.53
T <sub>6</sub> -25 P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	36.05	33.80	2.69	5.47	19.17	41.75
T <sub>7</sub> -25 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	37.95	35.35	3.13	6.31	20.27	42.67
T <sub>8</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB	39.69	37.34	3.46	7.21	21.07	43.78
T <sub>9</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + VAM	41.94	39.15	3.50	6.88	22.14	45.49
T <sub>10</sub> -50 kg P <sub>2</sub> O <sub>5</sub> ha <sup>-1</sup> + PSB + VAM	44.12	41.07	3.88	8.11	23.27	47.01
S. Em. ±	1.50	1.26	0.15	0.43	0.75	1.08
CD (0.05%)	4.52	3.75	0.45	1.30	2.26	3.25



**Figure 1.0 Effect of biofertilizer and phosphorus on nitrogen, phosphorus and potassium uptake in grain and straw**

## References

- Ali, M., Hossain, M., Rahman, M. and Islam, M. 2012. Effect of vermicompost on growth, yield, chemical composition and Oil content of rapeseed. *Journal of Experimental Biology Science*, **3**(2): 91-96.
- Basu, Manisha; Mondhal, Pinaki; Basak, R.K.; Basu, T.K. and Mahapatra, S.C. (2006). Effect of cobalt, Rhizobium and phosphobacterium inoculations on yield and nutrient uptake in summer groundnut (*Archis hypogaea* L.) on Alluvial soils. *Journal of the Indian Society of Soil Science*, **54**: 60-64.
- Biswas, B.C., Das, S. and Subhash, K.P. (2006). Crop response to bio-fertilizers. *Fertilizers News*, **46**: 15-24.
- Choudhary, S., Sharma, O. P., Choudhary, G. L. and Jat, L. (2017). Response of urdbean [*Vigna mungo* (L.) Hepper] to phosphorus fertilization and thiourea on yield, quality, nutrient content and uptake. *Int. J. Curr. Microbiol. App. Sci*, **6** (10): 2841- 2847.
- Das, S., Pareek, B. L., Kumawat, A. and Dhikwal, S. R. (2013) Effect of phosphorus and bio-fertilizers on production of chickpea (*Cicer arietinum* L.) in north western Rajasthan, India. *Legume Res.* **36** (6): 511- 514.
- Dhewa, J. S., Daniel, S. and Sulochana, (2017). Effect of different levels of phosphorus and sulphur on growth nutrient uptake of green gram (*Vigna radiate* L.) under teak (*Tectona grandis* L.) based agroforestry system. *Int. J. Curr. Microbiol, App. Sci.*, **6** (2): 520-534.
- Gajera, R. J., Khafi, H. R., Raj, A.D., Yadav, V. and Lad, A. N. (2014). Effect of phosphorus and bio-fertilizers on growth yield and economics of summer green gram [*Vigna radiate* (L.) Wilczek]. *Agric. Update*, **9** (1): 98-102.
- Hakeem, S.A., Thomas, T. and Shagufta, Wani (2008). Effect of different levels of neem cake and biofertilizers on growth, yield attributes and nutrient uptake in black gram (*Vigna mungo* L.) Type-9. *Environment and Ecology*. **26** (1A): 369-371.
- Kachave (2018) *International Journal of Current Microbiology and Applied Sciences* ISSN: 2319-7706 Volume 7 Number 07 (2018)
- Khan, A.A., Jilani, G., Akhtar, M.S. Naqvi, S.M.S. and Rasheed, M. (2009). Phosphorus solubilizing bacteria: Occurrence, mechanisms and their role in crop production. *Journal of Agriculture and Biological Science*, **1**: 48-58.

- Mehta, R.S., Patel, B.S., Meena,S.S. and Meena, R.S. (2010). Influence of nitrogen, phosphorus and biofertilizers on growth characters and yield of fenugreek (*Trigonella foenum greacum* L.). *Journal of Spices and Aromatic Crops*. 19: 23- 28.
- Mir, A. H., Lal, S. B., Salmani, M., Abid, M. and Khan, I. (2013). Growth, yield and nutrient content of black gram (*vigna mungo*) as influenced by levels of phosphorus, sulphur and phosphorus solubilizing bacteria. *SAARC J. Agric.* ,**11** (1): 1-6.
- Mohammad, I., Yadav, B. L. and Ahamad, A. (2017). Effect of phosphorus and bio-organics on yield and soil fertility status of mung bean [*Vigna radiate* (L.) Wilczek under semi-arid condition of Rajasthan, India. *Int. J. Curr. Microbiol. App. Sci* **6** (3): 1545- 1553.
- Patel, H. F., Maheriya, V. D., Attar, S. K. and Patel, H. R. (2017). Nutrient uptake and yield of *kharif* green gram as influenced by levels of sulphur, phosphorus and PSB inoculation. *Legume Research*.
- Rani, M., Prakash, V. and Khan, Khalil. (2016). Response of mung bean [*Vignaradiate* (L.) wilczek] to phosphorus, sulphur and PSB during summer season. *Agric. Sci. Digest.*,**36** (2) 2016: 146-148.
- Rathor, D. K., Gupta, A. K., Chaudhary, R. R. and Sadhu, A. C. (2015). Effect of integrated phosphorus management on growth, yield attributes and yield of summer green gram (*Vigna radiata* L.). *The Bioscan.*, **10** (1): 05-07.
- Singh, A. K., Singh, C. K., Singh, R. K., Sarvjeet and Lavanya, G. R. (2016). Effect of phosphorus and bio- fertilizer on growth and yield of green gram (*Vigna radiiata* L.). *Res. Environ. Life Sci.* **9** (2) 223- 225.
- Singh, A. Kumar, A. and Pandey, A. K. (2017). Response of phosphorus, PSB and pressmud on dry matter accumulation, phosphorus content and uptake of urd bean. *Int. J. of Chemical studies*, **5**(5): 2351- 2353.
- Singh, S. K. and Singh, G. C. (2013). Effect of phosphorus, sulphur and zinc on nutrient composition in black gram. *J. Rural and Agril. Res. Vol.***13** No.2, 63-64.
- Somani, L.L., Bhandari, S.C., Saxena, S.N. and Vyas, K.K. (Ed.) (2008). *Biofertilizers. Scientific Publication*, Jodhpur 142-145.
- Yadav, K. R. (2013). Effect of phosphorus sources and Phosphorus Solubilizing Micro-organisms on performance of mung bean [*Vigna radiate* (L.) Wilczek] grown in Entisols.