

Yield attributes and yield of maize as influenced by different paddy straw management options and fertilizer Nitrogen levels in rice-maize cropping systems.

ABSTRACT

A field experiment was conducted during 2021-22 and 2022-23 at Regional Agricultural Research Station, Arepally, Warangal district. It is situated in the Central Telangana Zone. A total of 18 treatments replicated thrice in factorial RBD design. The study is conducted to know the yield attributes and yield of maize as influenced by different paddy straw management options and fertilizer Nitrogen levels in rice-maize cropping systems. The results of the experiment concluded that highest grain yield, straw yield, cob length (cm), cob girth (cm), number of cobs per plant, number of rows per kernal, 100 seed weight (gms) was obtained in incorporation and least was obtained in cutting and removal. With the different nitrogen levels highest was obtained in T₆130% RDN (20% at basal, 42% 15 days after sowing, 42% knee high stage (30 DAS) and 26 % tasselling stage (60 DAS) and the least was obtained in T₁ 100 % RDN (33.3 % at basal, 33.3% knee high stage (30 DAS) and 33.3% tasselling stage (60 DAS). There is no significant effect of crop residue management and fertilizer N levels on yield attributes of maize and there is no significant interaction between the two factors.

Keywords: Grain yield, cob length, cob girth, incorporation, Nitrogen.

Introduction: Maize is one of the most versatile emerging crops having wider adaptability and compatibility under diverse soil and climatic conditions and hence is cultivated in sequence with different crops under various agro-ecologies of the country (Maurya *et al.*, 2020). It is the third most important food crop after rice and wheat in India (Murdia *et al.*, 2016). Globally, this crop is known as queen of cereals because of its highest genetic yield potential among the cereals (Vishuddha *et al.*, 2018). It is cultivated on nearly 150 m ha in about 160 countries having wider diversity of soil, climate, biodiversity and management practices that contributes 36 % (782 m t) in the global grain production. In India, average area of cultivation is 8.7 m ha mainly during *Kharif* season which covers 80% of the area. The average productivity in India is 2.43 t ha⁻¹ lower than the global average (4.92 t ha⁻¹) (Parihar *et al.*, 2011). USA is the largest producer of maize contributes nearly 35% of the total production in world and highest productivity (>9.6 t ha⁻¹) which is double than the global average (4.92 t ha⁻¹). India contributes nearly 9% in the national food basket. About 25% of N and P, 50% of S and 75% of K taken up by cereal crops are retained in the crop residue, making them viable nutrient sources. Also, residues return carbon (C) to the soil, which improves soil structure, the ability of the soil to hold nutrients, and water holding capacity. The use of organic inputs such as crop residue and manures has great potential for improving soil productivity and crop yield through improvement of

the soil **Physico-chemical** properties and nutrient supply (Abbasi *et al.*, 2009). Incorporation of crop residues can contribute to sustainability mainly through improvement of soil fertility as judged by organic carbon, available P and potassium (K) content. Straw incorporation returns valuable nutrients back into the soil, particularly P and K, leading to potential economic savings through reduced additions of organic and inorganic **fertilizers** (HGCA, 2009).

Material and methods:

A field experiment was conducted during 2021-22 and 2022-23 at Regional Agricultural Research Station, Arepally, Warangal district. It is situated in the Central Telangana Zone. The experiment was conducted in **Factorial Randomized Block** design. The soil was sandy clay loam in texture. The variety choosed for maize is NK-7884 syngenta hybrid corn. Spacing adopted for maize was 60 X 20 cm (Maize) with plot size of 6 X 4 =24 Sq. Meters.

Treatmental details:

Factor 1 (paddy straw management options) Factor 2 (fertilizer N levels). Among paddy straw management options it has burning (one week before land preparation), incorporation (a week before land preparation consortium is sprayed, week after its spray paddy straw will be incorporated) ,Cutting and removal of straw (one week before land preparation paddy straw removed and consortium will be sprayed, week after spray paddy straw is incorporated) .

Among fertilizer N levels T₁100 % RDN (33.3 % at basal, 33.3% knee high stage (30 DAS) and 33.3% tasselling stage (60 DAS), T₂ 100% RDN (40 % at basal, 30% knee-high stage (30 DAS), 30% tasselling stage (60 DAS), T₃ 100% RDN (10% at basal, 30% 15 days after sowing, 30% knee high stage (30 DAS) and 30% tasselling stage (60 DAS), T₄ 100% RDN (15% at basal, 30% 15 days after sowing, 30% knee high stage (30 DAS) and 25% tasselling stage (60 DAS), T₅ 115% RDN (18 % at basal, 36% 15 days after sowing, 36% knee high stage (30 DAS) and 25% tasselling stage (60 DAS), T₆ 130% RDN (20% at basal, 42% 15 days after sowing, 42% knee high stage (30 DAS) and 26 % tasselling stage (60 DAS).

Microbial consortium: Consists of decomposers belonging to *genera Phanerochaeta, Asperigillus, Trichoderma*. RDF for *Rabi* maize - 240-80-80 kg N-P-K kg ha⁻¹. Recommended doses of phosphorus, potassium and FYM (10 t ha⁻¹) will be applied in equal quantities in all treatments.

Five randomly selected plants from each plot were taken and measured for cob length, cob girth, number of cobs per plant, number of grains per cob, 100 seed weight and average values were calculated.

Results and discussion:

Yield Attributes:

There is no significant effect of paddy straw and nitrogen management on yield attributes of maize. There is no significant interaction between the two factors.

Cob length (cm)

~~With factor 1~~ The highest Cob length was obtained in incorporation 23.15 cm followed by burning 21.04 cm and cutting and removal 20.16 cm.

Among the different nitrogen management options levels the highest Cob length was reported in 21.95 cm T₃ followed by T₂ 21.58 cm and the least was observed in T₁ 20.94 cm.

Cob girth (cm)

~~With paddy straw options~~ The highest Cob girth was obtained in paddy straw incorporation 17.19cm followed by burning 16.50cm and cutting and removal 16.20cm.

~~Factor 2~~ Regarding response of Cob girth to different N levels the highest was observed (17.96 cm) in 130% RDN (20% at basal, 42% 15 days after sowing, 42% knee high stage (30 DAS) and 26 % tasselling stage (60 DAS) i.e.T₆ followed by T₅ (17.37 cm) and the least was observed (15.18 cm) in T₁.

Number of cobs per plant

~~With factor 1~~ The highest number of cobs per plants was obtained in incorporation (1.05) followed by burning (1.04) and cutting and removal (1.03).

Among the N levels T₆ recorded the highest number of cobs per plants (1.05) but found at par with T₅. The least number of cobs per plants (1.03) was observed in T₁.

Number of rows per kernel

Regarding effect of different paddy straw options the highest (30.20) was obtained in treatment incorporation followed by burning (28.69) and cutting and removal (28.31).

Similarly, Fertilizer N levels affect number of rows per kernel and the highest (30.31) was observed in T₆ followed by T₅ (29.76) and the least was observed in T₁(27.86)

100 seed weight (gms)

The highest 100 seeds weight was obtained in incorporation treatment i.e. 29.43 gms followed by burning (29.13 gms) and cutting and removal (29.02 gms).

With effect of different N levels the highest (29.92 gms) was observed in T₆ followed by T₅ (29.61gms) and the least was reported in T₁ (28.46 gms)

Singh *et al.*, 2019 reported that plant height, number of effective tillers, no of grain per spike and test weight remained significantly higher under residue incorporation treatment over residue burning and residue removal during 2019 and 2020, respectively. This **was** might be due to presence of residue under incorporation and retention treatment improved the physical, chemical and biological properties of soil by increasing organic matter content over the burning and removal practice. **Under** present study the yield attributes was lower under removal treatments. This was mainly due to **scarcity** of nutrient because there was not any other source of nutrient addition except the fertilizer.

Anil *et al.* (2018) **results** revealed that application of nitrogen 180 kg ha⁻¹ with 4 splits given the better plant height, tiller number m⁻²; dry matter production, root volume and root dry weight that contributed for higher grain yield.

Grain and stover yield (kg ha⁻¹)

Different treatments significantly affect the grain yield of Maize and the highest grain (8266.6 kg ha⁻¹) yield was obtained in incorporation treatments, but found at par with burning (8089.5 kg ha⁻¹) while cutting and removal showed significantly less grain yield 7639.5 kg ha⁻¹.

Different Nitrogen levels and its time of application significantly affect grain yield of Maize and treatment T₆ recorded the highest (8529.7 kg ha⁻¹) grain yield of maize which was at par with T₅ (8362.1 kg ha⁻¹) and T₄ (8138.7 kg ha⁻¹) lowest was recorded in T₁ (7399.5 kg ha⁻¹).

Burning of paddy residue recorded higher stover yield of 10098.2 kg ha⁻¹ which is on par with incorporation of rice residue 10513.6 kg ha⁻¹, while cutting and removal showed lower stover yield of 9400.3 kg ha⁻¹.

Among different levels of nitrogen and time of application stover yield was highest in T₆ 10578.0 kg ha⁻¹ followed by T₃, T₄ and T₅ but found at par with each other, where nitrogen was applied in 4 splits. The lowest stover yield (9281.1 kg) was recorded in T₁ ha⁻¹ in which 100 % RDN was applied in 3 equal splits.

Similar results were observed by Han *et al.* (2020), with an increase in crop yields due to increased biomass production. However, there was a decrease in crop yields where straw burning was done, possibly due to loss of NPK with burning and thereby reducing the fertility of the soil (El-Sobky, El-Sayed., 2017).

Due to the incorporation of straw, there was an increase in grain and straw yield, possibly due to the rise in the availability of nutrients in the soil, which will increase plant biomass production (Zhang *et al.*,2021b). There was an increase in grain yield by

incorporating straw and fertilizer N due to the availability of mineral N, which plays an important role in grain yield (Sui *et.al.*,2020).

UNDER PEER REVIEW

Table 1 : Grain yield and straw yield (kg ha⁻¹) of maize as influenced by paddy straw management options and fertilizer N levels.

Treatments	Grain Yield			Straw Yield			Harvest Index (%)		
	2021-22	2022-23	Pooled data	2021-22	2022-23	Pooled data	2021-22	2022-23	Pooled data
Burning	8052.7	8126.3	8089.5	10038.2	10158.2	10098.2	44.48	44.42	44.45
Incorporation	8230.5	8302.7	8266.6	10483.7	10543.6	10513.6	44.00	44.11	44.06
Cutting and removal	7576.3	7702.7	7639.5	9260.3	9540.3	9400.3	45.02	44.76	44.89
SE(m)±	172.31	160.74	103.01	206.94	198.21	189.21	0.45	0.75	0.40
CD	495.23	461.98	296.07	594.75	569.67	543.80	1.30	2.17	1.16
Nitrogen levels and time of application									
T1	7354.5	7444.6	7399.5	9204.5	9357.8	9281.1	44.45	44.34	44.40
T2	7619.2	7709.3	7664.2	9624.1	9777.4	9700.8	44.19	44.12	44.15
T3	7851.9	7941.9	7896.9	9804.7	9958.0	9881.3	44.42	44.30	44.36
T4	8091.6	8185.8	8138.7	10109.5	10262.8	10186.1	44.47	44.43	44.45
T5	8317.2	8407.1	8362.1	10320.3	10473.6	10396.9	44.70	44.69	44.70
T6	8484.7	8574.6	8529.7	10501.3	10654.7	10578.0	44.76	44.70	44.73
SE(m)±	243.69	227.32	145.69	292.66	280.31	267.58	0.64	1.07	0.57
CD	700.36	653.34	418.70	841.11	805.63	769.05	1.84	3.07	1.64

T1:100 % RDN (33.3 % at basal, 33.3% knee high stage (30 DAS) and 33.3% tasselling stage (60 DAS)

T2:100% RDN (40 % at basal, 30% knee- high stage (30 DAS), 30% tasselling stage (60 DAS)

T3:100% RDN (10% at basal, 30% 15 days after sowing, 30% knee high stage (30 DAS) and 30% tasselling stage (60 DAS)

T4:100% RDN (15% at basal, 30% 15 days after sowing, 30% knee high stage (30 DAS) and 25% tasselling stage (60 DAS)

T5:115% RDN (18 % at basal, 36% 15 days after sowing, 36% knee high stage (30 DAS) and 25% tasselling stage (60 DAS)

T6:130% RDN (20% at basal, 42% 15 days after sowing, 42% knee high stage (30 DAS) and 26 % tasselling stage (60 DAS)

UNDER PEER REVIEW

Conclusions: Paddy straw and different nitrogen levels significantly showed highest grain yield, stover yield and yield attributes in maize. The highest was found in incorporation treatments followed by burning and cutting and removal. With respect to different fertilizer N levels and its application the highest grain yield, stover yield and yield attributes was observed in T₆ (115% RDN) and T₅ (115% RDN) where, higher doses of nitrogen was applied in four different splits doses.

References:

- Han, Y., Ma, W., Zhou, B., Yang, X., Salah, A., Li, C. and Zhao, M. 2020. Effects of Straw-Return Method for the Maize–Rice Rotation System on Soil Properties and Crop Yields. *Agronomy*, 10(4), 461.
- El-Sobky, El-Sayed E.A. 2017. Effect of burned rice straw, phosphorus and nitrogen fertilization on wheat (*Triticum aestivum* L.). *Annals of Agricultural Sciences*, 62(1),113–120.
- Sui, Pengxiang., Tian, Ping., Lian, Hongli., Wang, Zhengyu., Ma, Ziqi., Qi, Hua., Mei, Nan., Sun, Yue., Wang, Yingya., Su, Yehan., Meng, Guangxin., Jiang, Ying 2020. Straw Incorporation Management Affects Maize Grain Yield through Regulating Nitrogen Uptake, Water Use Efficiency, and Root Distribution. *Agronomy*, 10(3), 324–328.
- Zhang, J., Li, W., Zhou, Y., Ding, Y., Xu, L., Jiang, Y., and Li, G. 2021b. Long term straw incorporation increases rice yield stability under high fertilization level conditions in the rice–wheat system. *The Crop Journal*.2214-2220.
- Anil, K., Yakadri, M and Jayasree, G. 2018. Influence of Nitrogen Levels and Times of Application on Growth Parameters of Aerobic Rice. *Int.J.Curr.Microbiol.App. Sci.* 7(5): 1525-1529.
- Singh, R.K., Sharma, G.K., Kumar, P., Singh, S.K and Singh, R. 2019. Effect of crop residues management on soil properties and crop productivity of rice-wheat system in inceptisols of Seemanchal region of Bihar. *Current Journal of Applied Science and Technology*. 37(6): 1-6.
- HGCA. Information Sheet 05: Assessing the nutrient content of cereal straw. HGCA, London, 2009.
- Abbasi, M.K., Mushtaq, A and Tahir, M.M. 2009. Cumulative effects of white clover residues on the changes in soil properties, nutrient uptake, growth and yield of maize crop in the sub-humid hilly region of Azad Jammu and Kashmir, Pakistan. *African Journal of Biotechnol.* 8(10): 2184-2194.
- Parihar, C.M., Jat, S.L., Singh, A.K., Kumar, R.S., Hooda, K.S., Chikkappa, G.K and Singh, D.K. 2011. Maize production technologies in India.
- Murdia, L.K., Wadhvani, R., Wadhawan, N., Bajpai, P and Shekhawat, S. 2016. Maize utilization in India: an overview. *American Journal of Food and Nutrition*. 4(6): 169- 176.
- Maurya, R.N., Verma, V.K., Singh, U.P and Kumar, R. 2020. Response of

mulching and nutrient management practices yield under different cropping system. *Journal of Pharmacognosy and Phytochemistry*. 9(6): 1589-1591.

Vishuddha, N., Rajesh, K.,Doharey, R.K and Vema, S.K. 2018. Maize (*Zea mays* L.) Crop at Various Stages of Growth in Rabi Season as Influenced by Varieties, Plant Geometry and Nutrient Management. *Int.J.Curr.Microbiol.App.Sci*. 7: 989-997.

UNDER PEER REVIEW