

**Intra-familial Child Sexual Abuse among Adolescents in Rivers State, Nigeria:
Analyzing contributing factors and victim-perpetrator characteristics.**

Abstract

Background: The global prevalence of intra familial child sexual abuse (IF CSA) is uncertain because only a fraction of the incidence is ever reported. This study investigated the prevalence and characteristics of IF CSA among adolescents in Rivers State, Nigeria.

Comment [EAMS1]: intra-familial

Methods: This cross sectional retrospective study was done in 2013 using a modified self-administered questionnaire adapted from a previous study. Participants were adolescents attending secondary schools in Obio Akpor Local Government Area in Rivers State, Nigeria. A multistage sampling technique was employed to identify 573 victims of child sexual abuse aged 10-19 years. Descriptive and inferential statistics was carried out using SPSS version 25 and the level of significance was set at ≤ 0.05 .

Comment [EAMS2]: cross-sectional

Results: Results showed that 572 adolescents reported they had experienced sexual abuse. Of this, 408 (71.3%) had reported being victims of IF CSA with majority (55.4%) of them aged between 10-13 years. This number comprised 183 (32.0%) males and 389 (68.0%) females giving a male to female ratio of IF CSA of 1:2. However, there was no significant sex-based difference ($\chi^2 0.12, p 0.73$). The mean age of the victims at time of occurrence was 13.3 ± 1.2 years with younger adolescents more likely to experience IF CSA. The males were slightly younger than the females at the time of abuse ($p=0.04$). Majority of perpetrators were females (75.0%) and most being students. Rate of disclosure of IF CSA was low, reasons for non-disclosure includes fear, shame and the relationship with their perpetrators. Risk factors for IF CSA included low socio-economic class, living status of the participant and younger age.

Comment [EAMS3]: were

Comment [EAMS4]: ,

Comment [EAMS5]: ,

Comment [EAMS6]: the

Comment [EAMS7]: ,

Comment [EAMS8]: male-to-female

Comment [EAMS9]: ,

Comment [EAMS10]: The majority

Comment [EAMS11]: were

Comment [EAMS12]: The rate

Comment [EAMS13]: include

Conclusion: The prevalence of IF CSA is high, risk factors include lower social class, younger age and living status of the child. Females are the major perpetrators and level of disclosure is low. There is a need for targeted prevention and intervention programmes to alert parents and guardians of these crimes.

Comment [EAMS14]: :

Comment [EAMS15]: ,

Comment [EAMS16]: the

Keywords: Intra-familial Abuse, Child Sexual Abuse, Adolescents, Socio-demographic Factors, Perpetrators, Disclosure

Introduction:

Sexual abuse is defined as sexual contact of any kind that takes place without the consent of the individual involved¹. It may also be viewed as a violation of one’s sexual rights infringed on a vulnerable person by someone of a higher power or authority². Child sexual abuse (CSA) is sexual activity involving an individual less than the age of 18 years who does not fully comprehend such an act and is therefore not able to give informed consent to it or may not be developmentally prepared for it and still cannot consent to it. The abuse may be from adults, including family members and close relatives (incest) or older adolescents, and could be either the same or of the opposite sex as the victim^{3,4}.

CSA could involve physical or non-physical contact. Physical contact may include fondling of the adolescent’s genitals or breasts, mutual masturbation, intimate kissing, oral-genital contact, penetration with the **digits**, having anal and/or vaginal intercourse with the victim, non-physical contact may include observing the **victim** bath, and undress, exposing the adolescent to pornography, exhibitionism, the adult masturbates and makes the adolescent watch, or simply gain sexual pleasure from watching the victim naked or engaged in sexual activity^{5,6}.

The effects of CSA are far-reaching, affecting the survivor adversely with psychosocial and behavioural consequences⁷. Research has shown that young women who had experienced sexual abuse between the ages of 12-16 years had higher rates of teenage pregnancies, sexually transmitted infections, unprotected sex, and early sexual intercourse⁸. In addition, survivors of CSA face a triple chance of experiencing future physical/sexual abuse or becoming potential **perpetrator**⁹.

When the sexual abuse of children involves a close family member (siblings, relatives, parents, step-parents, or grandparents), any other person who lives in the same house (a maid

Comment [EAMS17]: and
Comment [EAMS18]: :
Comment [EAMS19]: victims

Comment [EAMS20]: perpetrators

or a caregiver), or someone close to the family (intimate friends) and is perceived as family by the child, it is referred to as Intrafamilial Child Sexual Abuse¹. This is emphasized in a study by Magalhães et al¹⁰ where it was stated that intrafamilial sexual abuse cases most often involved younger victims who had a close relationship with their perpetrators.

The prevalence globally for Intrafamilial Child Sexual Abuse (IF CSA) is uncertain because only a fraction of the incidence has been reported to the authorities¹¹. However, a study by Valle et al.¹ (2018) in Peru estimated that out of a prevalence of 22% of rape cases among adolescents, 73% were intra-familial assaults among adolescents aged 14-17 years. A study in Turkey reported a prevalence of 23.6%¹² and another in Canada reported a prevalence of 44%¹³.

IF CSA has become a widespread issue that significantly affects the lives of the victims, the families and society at large. Despite its untoward impact this type of abuse is underestimated, under-reported, hidden, unaddressed, and not getting the deserved attention. This is especially common in low- and middle-income countries (LMIC) such as Nigeria¹⁴. The victims are usually made to promise or sometimes threatened to secrecy by their perpetrators and the difficulty associated with detection where there was no physical contact makes it a huge challenge to detect^{15,16}. Various studies have shown that adolescents who experience IF CSA hardly disclose such abuse until adulthood¹⁷. Therefore, the true prevalence may not be readily apparent.

In a study by Fayaz¹⁸, participants who had been survivors disclosed that several reasons were responsible for the non-disclosure of such abuse, including the fear of the perpetrator, the fear of not being doubted, and others. Due to the social stigma and shame that the victim and the family are likely to experience, perpetrators of such acts are hardly brought to book as family members would rather have it settled quietly within the household when discovered^{5,19}. This

Comment [EAMS21]: of

Comment [EAMS22]: their

Comment [EAMS23]: ,

Comment [EAMS24]: .

is also expressed in a study by Gencer et al¹² where delays in identifying and addressing inter-familial abuses were noticed as well as withdrawals of claims if ever reported.

Some factors contributing to IF CSA as outlined by Black et al.²⁰ include proximity as well as access to the victim as a result of a familial relationship, a dysfunctional home; usually, the perpetrator will have the advantage of authority and power over the victim, and there will be a level of trust that the victim has for the perpetrator. However, the personality displayed by the perpetrator may include manipulative and deceitful traits exhibited to gain the confidence and compliance of the victim, awkward views on relationships, excessive closeness to the victim, and lack of empathy. From varying literature, most perpetrators were males^{13,21,22}.

In most of the literature, familial dynamics have shown that the victims of intra-familial CSA were mainly younger adolescents who were females²³ (Hinds et al., 2020) and of a lower socioeconomic class¹⁶. However, studies have also shown that boys were more averse to disclosing IF CSA than girls²⁴, which could explain why girls had a greater prevalence than boys and not necessarily because they were more affected.

A community-based study by David et al²⁵ on CSA and disclosure in South-Western Nigeria reported 7.5% of intra-familial cases with 2.1% of them being incest. A literature search indicated that there may be a paucity of studies on IF CSA in the south-south region of the country. Hence, there is a need for a study to aid the understanding of the contributory factors, as well as the characteristics of both the victims and perpetrators, to provide effective preventive and intervention strategies in the region. Therefore, the present study aimed to determine the risk factors associated with IF CSA among adolescents, the socio-demographic characteristics of perpetrators and victims, and the relationship between perpetrators and victims in Rivers State.

Material and Method

Comment [EAMS25]: This is also expressed in a study by Gencer et al. 12, where delays in identifying and addressing inter-familial abuses were noticed, and claims were withdrawn if ever reported. Some factors contributing to IF CSA, as outlined by Black et al.20 include proximity as well as access to the victim as a result of a familial

Comment [EAMS26]: However, the personality displayed by the perpetrator may include:

- Manipulative and deceitful traits are exhibited to gain the confidence and compliance of the victim.
- Awkward views on relationships.
- Excessive closeness to the victim.
- Lack of empathy.

Comment [EAMS27]: might

Comment [EAMS28]: and

Study design: This is a cross-sectional retrospective school-based study conducted among adolescents aged 10–19 years in 2013.

Setting: This study was carried out in Obio Akpor Local Government Area, in Rivers State, Nigeria. Most of the companies and industries are located in the LGA and is also a residential area for their staff as well as civil servants. The LGA is also a home for a diverse population of residents from different cultures, religions, socioeconomic backgrounds and family structures. There are over 100 secondary schools in the LGA, thirteen co-educational schools were selected for this study.

Sample Size Calculation: The following factors were taken into account when determining the minimum sample size, which was determined using the Cochran formula for sample size for proportion.²⁶ Prevalence of 36.0% was used gotten from a previous Nigerian study on sexual abuse²⁷ while a 95% Confidence Interval (CI), 2.5% error margin, and 7.5% non-response allowance were used. Consequently, the study's minimum sample size of 1558 adolescents was determined.

Selection of participants: Using a multi-staged sampling technique, thirteen co-educational secondary schools were selected and from the selected schools, one thousand five hundred and fifty-eight adolescents aged 10-19 years were recruited for the study. However, only participants who responded that they had experienced sexual abuse irrespective of any form were selected for the final study. All participants who were 18 years and above gave a written consent while those that were less than 18 years in addition to assenting to the study, a written consent was also obtained from their Parents before participating in the study.

Study instrument: A self-administered questionnaire modified from the Halperin et al. questionnaire on child sexual abuse²⁸ was used. Socio-demographic information, family

Comment [EAMS29]: the

Comment [EAMS30]: of

Comment [EAMS31]: which

Comment [EAMS32]: to

Comment [EAMS33]: 13

Comment [EAMS34]: considered

Comment [EAMS35]:

Comment [EAMS36]: ,

Comment [EAMS37]: ,

Comment [EAMS38]: ,

Comment [EAMS39]: old

Comment [EAMS40]:

structure, and history of sexual assault were gathered through the questionnaire. A total of one thousand five hundred and fifty-eight questionnaires were distributed. Only participants who responded to YES to the question have you ever experienced sexual abuse? were selected for the final study. Oyedeji's socioeconomic classification was utilized to determine each participant's socioeconomic class²⁹.

Data Analysis: Data were analyzed using the Statistical Package for Social Sciences, version 25. Descriptive statistics were performed, with the results presented as graphs, frequency tables in simple proportions, and charts. The student's t-test was used to compare means, whereas the test for association between two groups was performed using the chi-square test. A p-value of less than or equal to 0.05 was considered significant in all cases.

RESULTS

Five hundred and seventy-two victims of CSA participated in this study. Their ages ranged from 10-19 years with a mean age of 13.28 ± 2.14 years. The mean age for the males was 14.01 ± 2.08 years while that of the female was 13.75 ± 2.18 . There were no significant differences in mean age according to sex. Three hundred and thirteen of them (55.4%) were between 10-13 years when they experienced sexual abuse.

Comment [EAMS41]: ,

Comment [EAMS42]: ,

One hundred and eighty-four (32.0%) of them were male and 388 (68.0%) were female, giving a male-to-female ratio of approximately 1:2.

Comment [EAMS43]:

Comment [EAMS44]: ,

Concerning the social and living status of the participants, 263 (46.0%) and 215 (37.6%) were from the lower and middle social classes, respectively. Fifty three percent of the participants lived with both parents, while 93 (16.3%) lived with their mother only.

Comment [EAMS45]: Fifty-three

Prevalence of intra-familial sexual abuse among the victims: Of the 572 adolescents who experienced sexual abuse, 408 (71.3%) experienced intra-familial sexual abuse. One hundred and thirty three (72.3%) of the 187 adolescent male victims reported intra-familial sexual abuse compared to 275 (70.9%) of the 388 female victims. This sex-based difference was not statistically significant. $\chi^2 0.12$, $p 0.73$ (OR 1.07, CI 0.73-1.58) Table II.

Comment [EAMS46]: thirty-three

The youngest age of the victims of IF CSA at the time of occurrence was 7 years, while the oldest was 16 years. The mean age at occurrence was 13.3 ± 1.2 years. There was a significant sex difference in mean age at occurrence. The mean age for males was 12.3 ± 2.2 years while that of the females was 11.2 ± 1.2 years. ($t = 2.15$, $p = 0.04$)

Comment [EAMS47]: ,

Comment [EAMS48]: for

Table III lists the characteristics of the perpetrators. Their ages ranged from 15 to 46 years, with a mean age of 22.0 ± 4.4 years. The mean age for the males was 21.8 ± 5.5 , while that of the females was 18.7 ± 3.4 years. This sex difference was statistically significant ($t = 1.96$, $p = 0.01$).

Comment [EAMS49]: .

The majority (75.0%) of the intra-familial perpetrators of sexual abuse were females and most of them were students.

Comment [EAMS50]: ,

A total of 134 (23.4%) victims disclosed the abuse; however, 16.2% of those who experienced intra-familial sexual abuse disclosed their abuse compared to 41.5% of those who experienced extra-familial sexual abuse. These differences were statistically significant. $P < 0.0001$, $\chi^2 41.63$ (Fig.1)

Reasons for non-disclosure include fear, no one will believe them, shame, relationship with the perpetrators- if related by blood or not, and self-blame.

Comment [EAMS51]: that

Concerning the victim perpetrators' relationships, 23% were cousins, 16.2% were uncles/aunts, and 14% were family friends. (Fig 2)

Table IV shows the relationship between intra-familial sexual abuse and socio-demographic variables. More adolescents (81.8%) within the younger age range (10-14 years) had experienced intra-familial sexual abuse compared to 120 (54.6%) of the 220 adolescents aged (15-19 years). Adolescents within the younger age group were three times more likely to have experienced intra-familial abuse compared to the older ones. This was significant $p < 0.001$, OR 3.75, 95% CI 2.57-5.47. There was no significant difference in the age at which the abuse occurred. The socioeconomic class of the victims and their living status significantly predicted the odds of experiencing intra-familial sexual abuse.

Table I: Characteristics of participants

Variables	Frequency	Percentages
Age(years)		
10-14	352	61.5
15-19	220	38.5
Sex		
Males	184	32.2
Females	388	67.8
Social class		
Upper	94	16.4
Middle	215	37.6
Lower	263	46.0
Living status		
Lives with mother	93	16.3
Lives with father	74	12.9
Lives with both parents	301	52.6
Lives with relatives	80	14.0
Non-relatives	24	4.2
Intra familial abuse		
Yes	408	71.3
No	164	28.7
Age at the time of abuse		
< 10 years	90	15.7
10-13 years	317	55.4
≥14 years	165	28.9
Total	572	100.0


Table II: Prevalence of intra-familial abuse among study participants

Sex	Intra-familial Sexual Abuse		Total N (%)
	Yes N (%)	No N (%)	
Males	133(72.3)	51(27.7)	187(100.0)
Females	275(70.9)	113(29.1)	388(100.0)
Total	408 (71.3)	164 (28.7)	572(100.0)

χ^2 0.12, p 0.73

Table III: Characteristics of perpetrators of intra-familial child sexual abuse

Variables	Frequency	Percentages
Age		
15-19	108	26.5
20-24	193	47.3
25-29	97	23.8
≥ 30	18	4.4
Unknown	20	4.9
Sex		
Females	306	75.0
Males	102	25.0
Married		
Yes	98	24.0
No	310	76.0
Occupation		
Students	295	72.3
Civil servants	56	13.7
Farmer	3	0.7
Trader	24	5.9
Artisan	9	2.2
Unemployed	21	5.1
Total	408	100.0



$P < 0.001, \chi^2 41.63$

Fig. 1: Disclosure of sexual abuse.




Figure 2: perpetrators of intra-familial child sexual abuse

Table IV: Relationship between intra-family sexual abuse and some socio-demographic variables

Variables	Intra family abuse		Total N (%)	χ^2	p-value	OR 95% CI
	Yes N (%)	No N (%)				
Age range						
10-14	288(81.8)	64(18.2)	352(100.0)	49.2	< 0.001	3.75
15-19	120(54.6)	100(45.5)	220(100.0)			2.57-5.47
Age at abuse						
7-14	290(71.3)	117(28.7)	407(100.0)	0.004	0.95	0.99
15-19	118(71.5)	47(28.5)	165(100.0)			0.65-1.47
Sex						
Males	133(72.3)	51 (27.7)	184(100.0)	0.12	0.73	1.07
Females	275(70.9)	113(29.1)	388(100.0)			0.73-1.58
Social class of subjects						
Upper	52(55.3)	42(44.7)	94(100.0)	14.07	<0.001	0.42
Lower	356(70.7)	122(25.5)	478(100.0)			0.27-0.67
Living status						
Both parents	281(93.4)	20 (6.6)	301(100.0)	150.5	< 0.001	15.9
Not with both parents	127(46.9)	144(53.1)	271(100.0)			9.5-26.6
Total	408(71.3)	164(28.7)	572(100.0)			

Discussion:

Intra-familial child sexual abuse (IF CSA) can cause lifetime trauma and untoward effects on the victims. This study aimed to analyze the contributing factors as well as the victim-perpetrator characteristics and **relationship** among adolescent victims of intra-familial child sexual abuse in Rivers State.

The present study showed that 71.3% of the abused adolescents had experienced IF CSA, a finding that was higher than the 5.4% reported by Valle et al.¹, 7.5% from the study by²⁵ David et al²⁵, and 44.8% reported by Loinaz et al¹⁶. Several factors may have been responsible for the higher prevalence observed in this study. For example, the present study was a school-based **one** which may have ensured an increased awareness and education regarding **CSA** thus aiding the respondents to answer accordingly; the school environment may have also encouraged a willingness to disclose abuse; the data collection methods in the studies were different; the present study had a larger sample size compared to the other studies; in addition, the variations in social structures, cultural norms, and family dynamics across the

Comment [EAMS52]: relationships

Comment [EAMS53]: ,

Comment [EAMS54]: ,

different study regions may have influenced the prevalence and reporting of IF CA significantly. Furthermore, a very low prevalence of 4% was obtained from Turkey by Demirçi et al³⁰. This difference could be because the present study was a school-based study where a self-administered questionnaire was used and confidentiality of obtained information was assured before the study was commenced while the turkey study was the prevalence obtained from forensic cases as only cases that reported to the authority were involved, and due to shame and possible stigmatization, victims may not report to the authority. In addition, the victims may not have preserved the evidence needed for forensic investigation to come out positive.

This study reported a nearly equal prevalence of IF CSA between the males and females and that the likelihood of both males and females experiencing IF CSA is almost the same. It is, therefore, important that male children be given the same kind of attention and care as females to reduce the possibility of sexual abuse. However, this finding differs from that of Gekoski, et al³¹, 2016, who reported that females were more likely to be victims of IF CSA than the males. The methodology and the study population used in the two studies could be the reason for this difference. While the present study was a cross-sectional descriptive study among adolescent students, Gekoski, et al³¹ study employed a rapid evidence assessment method, where previous studies were searched in a given time frame.

In contrast to several previous studies^{32,33}, our study revealed that female victims were exposed early to IF CSA compared to males, as the mean age of the female victims was significantly lower than that of the male victims. A possible reason may be that girls commence puberty earlier with an increased growth spurt compared to boys. In addition, previous studies have focused only on incest, which was not the case in the present study.

Our study revealed that the age range of the perpetrators was 15–46 years, with a mean age of 22.0 ± 4.4 years. This result indicates that most of the perpetrators were either older adolescents or young adults. The results also indicated that there was a noteworthy age difference between male and female perpetrators. The average age of the males was 21.8 ± 5.5 years, while for the females it was 18.7 ± 3.4 years. The disparity in age shows that the female perpetrators were younger. This outcome was in agreement with Bourke et al³⁴ who noted that female perpetrators were likely to be younger than male perpetrators, and their victims were usually male children or young adolescents (9 – 17 years).

Our study also revealed that these female perpetrators constituted a notable number in comparison to male perpetrators. This finding is significant as it contradicts the common

Comment [EAMS55]:

Comment [EAMS56]:

Comment [EAMS57]:

Comment [EAMS58]: than

Comment [EAMS59]:

Comment [EAMS60]: showed

Comment [EAMS61]: ,

Comment [EAMS62]: Compared

perception that males were the predominant perpetrators of IF CSA as seen in other studies^{21,22,35}. However, collaborates with the findings from an investigative report by Gekoski et al³¹. (2016). This high percentage of female perpetrators from our study, who were majorly students, may suggest opportunistic abuse within the family as a result of caregiving responsibilities or co-housing. This setting, therefore, signifies a family in which young females are given the role or responsibility of watching over or caring for young children, thereby leading to potential abuse. This is further supported by the results we obtained regarding the victim-perpetrator relationship, showing that 23% were cousins, 16.2% were uncles/aunts, and 14% were family friends.

The findings of our study showed that the disclosure of abuse by victims of IF CSA was minimal. Fewer victims of IF CSA (16.2%) disclosed abuse than those who experienced extra-familial CSA (41.5%). This finding agrees with Lahtinen et al³⁶ (2018), who reported low rates of non-disclosure. However, another twist to it, as described by Lahtinen et al³⁶ (2018), was that the non-disclosure by these children was towards other adults for fear of being doubted, but they had disclosed their experiences to their friends, which could be a possible reason for the high prevalence of IF CSA noted in this present school-based study.

The various reasons for non-disclosure, as reported in this study, included fear, disbelief, shame, relationship with the perpetrator, and self-blame. These reasons are in line with those outlined in the literature^{18,37}. Studies have shown that an important means of dealing with an experience, such as IF CSA, is to talk about it. Disclosure is therefore a key aspect of healing. Unfortunately, the rate of disclosure remains low in many regions and some of the victims who eventually decide to open up do so after a lengthy period, usually in adulthood³⁸. Non-disclosure fosters the continuity of the act by the perpetrator.^{13,16}

The findings of this study showed that younger age, lower socioeconomic class, and living status of the child determined the risk of experiencing IF CSA.

In terms of the age of IF CSA, we noticed that younger adolescents (10-14 years) had a higher prevalence of IF CSA than older adolescents (15-19 years). In addition, we observed that the younger age group was found to be 3.75 times more likely to experience IF CSA. This is consistent with the outcomes of other studies^{12,13,39}, which noted that younger children were more vulnerable. Similarly, a previous study showed that, in contrast to abuse that occurs outside the family, CSA within the family typically begins earlier, involves more severe forms over an extended period of time, and almost always takes place in the setting of more

Comment [EAMS63]: it

Comment [EAMS64]: conclusions of

Comment [EAMS65]: As reported in this study, the various reasons for non-disclosure included fear, disbelief, shame, relationship with the perpetrator, and self-blame.

Comment [EAMS66]: Regarding

Comment [EAMS67]:

complex relationships involving power and control as well as practical and emotional dependency⁴⁰.

It may be expected that children living with their parents, compared to those from a single-parent household or living with non-relatives, would be less at risk for IF CSA. This is because children from a single-parent household are likely to be perceived as more vulnerable due to reduced supervision from a parent, whereas those from a home with both parents are shielded from such risks. However, we noted that IF CSA occurred more frequently among children who lived with both parents compared to those who were not living with both parents. This finding is in contrast with the results of Valle et al¹. The high prevalence of IF CSA in our study among children living with both parents suggests that perpetrators are often family members or close friends who are trusted by parents. This result, therefore, implies that living arrangements alone may not be regarded as a strong predictor of risk for IF CSA, but rather to consider the characteristics and behaviour of the members of such a family.

Comment [EAMS68]: parents trust.

Our study showed that children from lower socioeconomic classes experienced more IF CSA than those from higher social classes. This agrees with the findings of Muratoğlu et al³² who reported that low levels of education and income are significant factors affecting the prevalence of IF CSA, particularly in cases of incest.

However, the findings of this study showed no significant sex differences in the prevalence of IF CSA, as males and females were equally affected. This finding is in agreement with previous studies that have reported a higher prevalence among females. They however asserted that this lower prevalence was due to lower rates of reporting the abuse among males.^{32,41}

Comment [EAMS69]: ,

Comment [EAMS70]: ,

Conclusion: IF CSA was common, and female family members were the main perpetrators. Low socioeconomic class, living status of the child and young age were factors that increased the risk for IF CSA. These findings, therefore, call for awareness programmes, educational initiatives, as well as protective measures specifically formulated to address these issues as a means of mitigating the risk of such abuse in this age group. There is also a need for targeted prevention and intervention programmes focused on alerting parents to these crimes.

Comment [EAMS71]: of

Comment [EAMS72]: to alert

Disclaimer: Part of this study have been previously published⁴²: Prevalence of child sexual abuse among secondary school adolescents in Obio/Akpor Local Government Area of Rivers State, Nigeria. *Nigerian Journal of Paediatrics*, 46(4), 156–162.

Ethical considerations Ethical clearance was obtained from the Ethics and Research Committee of the University of Port Harcourt Teaching Hospital. Consent and (or) assent were obtained from the parents/guardians and participants. Discretion and anonymity was ensured throughout the study.

References

1. Valle, R., Bernabé-Ortiz, A., Gálvez-Buccollini, J. A., Gutiérrez, C., & Martins, S. S. Intrafamilial and extrafamilial sexual assaults and their association with alcohol consumption. *Rev Saude Publica*, 2018; 52: 86. <https://doi.org/10.11606/s1518-8787.2018052000539>
2. Stewart, L., Sebastiani, A., Delgado, G., & López, G. Consequences of sexual abuse of adolescents. *Reproductive Health Matters* 1996; 4(7):129-134.
3. Ali, S., & Ali, J. Sexual Abuse of Women and Its Impact: Psychosexual Perspectives. *J. Appl. Environ. Biol. Sci*, 2018; 8(5): 1-1.
4. World Health Organization. *Responding to Children and Adolescents who have been sexually abused* 2017; <https://doi.org/https://iris.who.int/bitstream/handle/10665/259270/9789241550147-eng.pdf>
5. Assabu, G., Tefera, M., Abebe, A., & Busse, H. The magnitude of sexual abuse and its physical effects among female children seen at Gandhi Memorial Hospital in Addis Ababa, Ethiopia. *Ethiopian Journal of Health Development*, 2019; 33(3).
6. Gabriel-Job, N. Pattern of Sexual Abuse of Minors in Rivers State, Nigeria: A School-based Study. *West Afr J Med*, 2019; 36(3), 274-279.
7. Papalia, N. L., Luebbers, S., Ogloff, J. R., Cutajar, M., & Mullen, P. E. The long-term co-occurrence of psychiatric illness and behavioral problems following child sexual abuse. *Australian & New Zealand Journal of Psychiatry*, 2017; 51(6), 604-613. <https://doi.org/10.1177/0004867416667232>
8. Fergusson, D. M., McLeod, G. F., & Horwood, L. J. Childhood sexual abuse and adult developmental outcomes: Findings from a 30-year longitudinal study in New Zealand. *Child abuse & neglect*, 2013; 37(9): 664-674.
9. Banvard-Fox, C., Linger, M., Paulson, D. J., Cottrell, L., & Davidov, D. M. Sexual Assault in Adolescents. *Prim Care*, 2020; 47(2): 331-349. <https://doi.org/10.1016/j.pop.2020.02.010>

10. Magalhães, T., Taveira, F., Jardim, P., Santos, L., Matos, E., & Santos, A. Sexual abuse of children. A comparative study of intra and extra-familial cases. *Journal of forensic and legal medicine*, 2009; 16(8):455-459.
11. Valle, R., Bernabé-Ortiz, A., Gálvez-Buccollini, J. A., Gutiérrez, C., & Martins, S. S. Intrafamilial and extrafamilial sexual assault and its association with alcohol consumption. *Rev Saude Publica*, 2018; 52, 86. <https://doi.org/10.11606/s1518-8787.2018052000539>
12. Gencer, Ozlem & Ozbek, Aylin & Ozyurt, Gonca & Kavurma, Canem. A comparison of extra and intra-familial sexual abuse of children and adolescents. *Anatolian Journal of Psychiatry*. 2014; 17. 1. 10.5455/apd.1419190512.
13. Fischer, D. G., & McDonald, W. L. (1998). Characteristics of intrafamilial and extrafamilial child sexual abuse. *Child abuse & neglect*, 22(9), 915-929.
14. Maitanmi, B., Awojobi, Z., Adesuyi, E., Maitanmi Phd, J., Chukwuere, P., & Akingbade, O. Knowledge and Perceived Effect of Sexual Abuse Among Adolescents in Selected Secondary Schools in Mushin Local Government. *African Journal of Health, Nursing and Midwifery*, 2021; 4. <https://doi.org/10.52589/AJHNM-54GMEBUD>
15. Gupta, S., & Garg, D. Causes and Effects of Child Sexual Abuse. *International Journal of Innovative Science and Research Technology*, 2020; 5:1867-1870. <https://doi.org/10.38124/IJISRT20MAY650>
16. Loinaz, I., Bigas, N., & de Sousa, A. M. (2019). Comparing intra and extra-familial child sexual abuse in a forensic context. *Psicothema*, 31(3), 271.
17. McElvaney, R. Disclosure of child sexual abuse: Delays, non-disclosure and partial disclosure. What the research tells us and implications for practice. *Child Abuse Review*, 2015; 24(3), 159-169.
18. Fayaz, I. What stops children from disclosing sexual abuse: perspectives from adults. *Journal of loss and trauma*, 2023; 28(1): 96-99.
19. Thukral S, Rodriguez TD. Child Sexual Abuse. *Advances in psychology, mental health, and behavioral studies (APMHBS) book series*, 2018, p. 229–58. <https://doi.org/10.4018/978-1-5225-3958-2.ch017>.
20. Black, D. A., Heyman, R. E., & Slep, A. M. S. Risk factors for child sexual abuse. *Aggression and violent behavior*, 2001; 6(2-3): 203-229.

21. Craissati, J., McClurg, G., & Browne, K. Characteristics of perpetrators of child sexual abuse who have been sexually victimized as children. *Sexual Abuse: A Journal of Research and Treatment*, 2002; 14: 221-235.
22. Koçtürk, N., & Yüksel, F. Characteristics of victims and perpetrators of intrafamilial sexual abuse. *Child abuse & neglect*, 2019; 96: 104122.
23. Hinds, T. S., Giardino, A. P., Hinds, T. S., & Giardino, A. P. (2020). Incidence and prevalence of child sexual abuse. *Child Sexual Abuse: Current Evidence, Clinical Practice, and Policy Directions*, 1-10.
24. Ungar, M., Barter, K., McConnell, S. M., Tutty, L. M., & Fairholm, J. Patterns of abuse disclosure among youth. *Qualitative Social Work*, 2009; 8(3), 341-356.
25. David, N., Ezechi, O., Wapmuk, A., Gbajabiamila, T., Ohihoin, A., Herbertson, E., & Odeyemi, K. Child sexual abuse and disclosure in South Western Nigeria: a community based study. *African health sciences*, 2018; 18(2): 199-208.
26. Cochran WG. Sampling Technique. 2nded. New York, USA: Wiley and Sons Inc., 1963
27. Mezie-Okoye M.MMI and Folusho F, Alamina FF. Sexual Violence among Female Under-graduates in a Tertiary Institution in Port Harcourt: Prevalence, Pattern, Determinants and Health Consequences. *African Journal of Reproductive Health* 2014; 18(4):79
28. Halperin DS, Bouvier P, Jaffe PD ,Mounoud R, Pawlak CH, Laederach J. et al. Prevalence of child sexual abuse among adolescents in Geneva: results of a cross sectional survey. *BMJ* 1996; 312:1326
29. Oyedeji GA. Socio-economic and cultural background of hospitalized children in Ilesha. *Nig J Paed.* 1985; 12: 111-7
30. DEMİRCİ Ş, DOĞAN KH, ERKOL Z, DENİZ İ. Konya'da cinsel istismar yönünden muayenesi yapılan çocuk olguların değerlendirilmesi. *Türkiye Klinikleri Journal of Forensic Medicine and Forensic Sciences.* 2008;5(2):43-9.
31. Gekoski, A., Davidson, J. C., & Horvath, M. A. The prevalence, nature, and impact of intrafamilial child sexual abuse: Findings from a rapid evidence assessment. *Journal of Criminological Research, Policy and Practice*, 2016; 2(4), 231-243.
32. Muratoglu G, Celikel A, Karbeyaz K. Evaluation of the incest cases in Hatay. *Medicine Science International Medical Journal* 2018:1. <https://doi.org/10.5455/medscience.2017.06.8736>.

33. Gunduz T, Karbeyaz K, Ayrancı U. Evaluation of the adjudicated incest cases in turkey: difficulties in notification of incestuous relationships. *J Forensic Sci.* 2011; 56(2):438-43.
34. Bourke, A, Doherty S, McBride O, Morgan K, & McGee H. Female perpetrators of child sexual abuse: Characteristics of the offender and victim. *Psychology, crime & law*, 2014; 20(8): 769-780.
35. Crawford, S. L. Intrafamilial sexual abuse: What we think we know about mothers, and implications for intervention. *Journal of Child Sexual Abuse*, 1999; 7(3): 55-72.
36. Lahtinen, H.-M., Laitila, A., Korkman, J., & Ellonen, N. Children's disclosures of sexual abuse in a population-based sample. *Child abuse & neglect* 2018; 76:84-94
37. Elliott, S. A., Goodman, K. L., Bardwell, E. S., & Mullin, T. M. Reactions to the disclosure of intrafamilial childhood sexual abuse: Findings from the National Sexual Assault Online Hotline. *Child abuse & neglect*, 2022; 127: 105567.
38. Fuller-Thomson E, Lacombe-Duncan A, Goodman D, Fallon B, Brennenstuhl S. From surviving to thriving: Factors associated with complete mental health among childhood sexual abuse survivors. *Social psychiatry and psychiatric epidemiology*, 2020; 55: 735-744.
39. Stoltenborgh M, Van Ijzendoorn M. H, Euser E. M, & Bakermans-Kranenburg M. J. A global perspective on child sexual abuse: Meta-analysis of prevalence around the world. *Child maltreatment*, 2011; 16(2): 79-101.
40. Intra-familial child sexual abuse | CSA Centre. CSA Centre 2024. Retrieved on 16th of August 2024 from <https://www.csacentre.org.uk/research-resources/key-messages/intra-familial-csa/>.
41. Fis N.P, Arman A.R, Kalaca S, & Berkem M. Psychiatric evaluation of sexual abuse cases: A clinical representative sample from Turkey. *Children and Youth Services Review*, 2010; 32: 1285-1290.
42. Gabriel-Job, N., Alikor, E., & Akani, N. Prevalence of child sexual abuse among secondary school adolescents in Obio/Akpor Local Government Area of Rivers State, Nigeria. *Nigerian Journal of Paediatrics*, 2019; 46(4): 156-162.