

Ethno-Botanical Study of Water Yam (*Dioscorea alata* L.) Tubers in Storage in the South-Eastern Nigeria

ABSTRACT

Water yam tubers are staple food consumed in the South-Eastern Nigeria. This investigation focused on indigenous knowledge and usage of water yam tubers in the South-Eastern Nigeria. A well-structured questionnaire was designed and administered to obtain crucial information from the respondents within the study area. Five South-Eastern States of Nigeria namely; Abia, Anambra, Ebonyi, Enugu and Imo States were randomly surveyed. Descriptive statistics were adopted and relationships among some variables were checked with percentage error bars in MS EXCEL version 2007. Six hundred questionnaires successfully retrieved were responded to. The results were analyzed by gender, age groups and locations. The study revealed that 62% of the respondents were aware of water yam cultivation, 41% grew water yam tubers as food because of its nutritional importance while 94% believed that water yam had medicinal value. The findings from the study will be used as basis for further research on emerging indigenous practices and sustainability impact.

Keywords: ethno-botanical study, Nigeria, South East, storage, water yam,

Introduction

Yam (*Dioscorea* spp) belongs to the genus *Dioscorea* (Family *Dioscoreaceae*) and is the second most important tropical root crop in West Africa after cassava (Adisa *et al.*, 2015) and form an important food source in other tropical countries including East Africa, the Caribbean, South America, India and South East Asia (Coursey, 1967; Okigbo, 2002).

Okigbo (2002) estimated that the world production of yams is around 20 million tonnes per year. The greatest part of the world yam production (over 90%) is derived from West Africa (Coursey, 1967; Okigbo, 2002; Sahel, 2014) and Nigeria alone produces three-quarters of the world total output of yam (Modu *et al.*, 2015; Okigbo, 2002). “Yams are the fifth most harvested crops in Nigeria, following after cassava, maize, guinea corn and beans/cowpeas. More so, after cassava, yams are the most commonly harvested tuber crops in the country (NBS, 2012). Yams do not only serve as the main source of earnings and food consumption, but also as a major employer of labour in Nigeria” (Falola *et al.*, 2017).

Despite the importance of yams to people, the attention to its production is still questionable (Verter and Becvarova, 2014). Studies by Zaknayiba and Tanko (2013) reveal that lack of access to inputs, finance, poor producer prices, inadequate storage facilities, incidence of pests and diseases have negatively affected yam production. Similarly, (Maikasuwa and Ala, 2013) examined some determinants of yam production in particular regions (North Central zone, North West zone, South East zone and South West Zone) in Nigeria. They found that the factors of production such as labour, finance and material inputs like fertilizer have influenced yam production in the region.

Among the wide species reported, only about ten species are estimated to have been domesticated across Africa, Asia and Latin America for food and income generation (Scarcelli *et al.*, 2017). Of the ten cultivated species of yam, the six most important in Nigeria are: *Dioscorea rotundata* Poir (white yam), *Dioscorea alata* L. (water yam), *Dioscorea cayenensis* Lam. (yellow yam), *Dioscorea dumetorum* (Kunth) Pax. (cluster or bitter yam), *Dioscorea bulbifera* L. (aerial yam) and *Dioscorea esculenta* (Loir) Bark (Chinese yam) (Adeniji, 1970a; Okigbo, 2004).

Water yam (*Dioscorea alata* L.) is the most economically important yam species which serve as a staple food for millions of people in tropical and subtropical (South China, India, Sri Lanka, Brazil, South Africa) countries (Coursey, 1967; Hahn, 1995). *Dioscorea alata* is a crop with potential for increased consumer demand due to its low sugar content necessary for diabetic patients (Udensi *et al.*, 2010). According to Scott *et al.* (2000), water yam (*Dioscorea alata* L.) is the most widely distributed species of yam, though the total quantity produced is less than that of white yam. Water yam (*Dioscorea alata* L.) is grown widely in tropical and subtropical regions of the world. They are plants yielding tubers and contain starch between 70% and 80% of dry matter (Zhang and Oates, 1999; Shang *et al.*, 2007).

The tuber is the only economically important part of the crop and according to Sangoyomi (2004), it is consumed roasted, fried, boiled, pounded or as flour which can be reconstituted with hot water. Yam tubers are of a very high value, as in food, where it is a major source of carbohydrate, minerals of calcium, phosphorus, iron and vitamin B and C (Coursey, 1967; Okigbo and Ogbonnaya, 2006). It provides other nutritional benefits such as proteins, lipids, vitamins and minerals (Lasztity *et al.*, 1998).

Across different ethnic communities and geographic regions, diverse species of *Dioscorea* have been adapted within different habitation as a food source due to the high nutritional benefits and therapeutic values toward treatment and cure of certain health problems (Dutta, 2015; Trimanto and Hapsari, 2015). While yam is one of the most important staple root and tuber crops worldwide, it is still classified as an orphan crop because it is highly underutilized and receives little investment or research attention toward crop improvement. Yam plays a significant role in food security, medicine and economy in the developing countries. Its importance places it as the fourth most essential and utilized root and tuber crop globally after potatoes (*Solanum* spp.), cassava (*Manihot esculenta*) and sweet potatoes (*Ipomoea* spp.) and the second in West Africa after cassava (Dansie *et al.*, 2013). This is evident in annual global production, especially in West Africa.

In addition, the significance of yam in the cultural, social and religious environment of West Africa cannot be overemphasized (Obidiegwu and Akpabio, 2017). Its symbolism as king of crops is manifested in its use in ceremonies such as those for fertility and marriages, as well as an annual festival held to celebrate its harvest. Importantly, the cultural and linguistic diversity that cuts across West Africa has no influence on the beliefs, social values and religious practices attached to the yam crop. In spite of the importance of yams as major staple food and its socio-cultural value in the lives of the people of the West and Central Africa sub-region, research and documentation on this important staple food crop is very limited (Sangoyomi, 2004). This present investigation focused on the indigenous knowledge of water yam in storage in the South-Eastern Nigeria.

Materials and Methods

Ethno-botanical study of Water Yam

The method of Akpaja *et al.* (2003) was used in the study area which involved the use of a well-structured questionnaire to obtain vital information from the respondents.

Study Area

The study was conducted in the South-Eastern Nigeria (Figure 1). Geographically, South-Eastern Nigeria extends from latitudes $4^{\circ} 40^I$ to $7^{\circ} 20^I$ North and $6^{\circ} 00^I$ to $8^{\circ} 20^I$ East longitude. The study area occupies about 50,000 km² of Nigeria's total area of 923, 768km². A 2006 Census population estimate by the National Population Commission (NPC) reported that the South-Eastern Nigeria has a population of 30 million people (NPC, 2006). The crops commonly grown in this region include but not limited to yam, cassava, cocoyam, and various vegetables and fruits. The natural vegetation in many parts of South-Eastern states is mainly grassland and woodland as well as tropical rainforest. Also, the annual rainfall is between 1400mm in the north to 2500mm in the south with soil pH in some parts ranging from 3.5 to 6.4 (Igwilo *et al.*, 2006).

Sampling Frame

The respondents to the questionnaires were the people from each of the five South Eastern States of Nigeria. The reason for selecting the area was because they were yam-producing and Igbo-speaking area. Besides, they share similar cultural and traditional beliefs.

Questionnaire Administration

A well-structured questionnaire was designed to aid in obtaining crucial information from the population of the study area. One hundred and twenty (120) questionnaires were distributed to the study population of different age groups in each of the five South Eastern States, making a

total of six hundred (600) questionnaires used in the study. The questionnaires were constructed to get vital information as follows:

- a. Demographic characteristics of the respondents.
- b. Factors responsible for water yam cultivation.
- c. Predisposing factors that deterred the production and storage of water yam.

Data Analysis

Data were analyzed using MS EXCEL version 2007 and a descriptive statistics were used. The relationship between some variables were obtained.

Results

The results of the study is presented in a frequency distribution table as shown in (Figure 1). A total of 600 respondents comprising 192 (32%) from Anambra, 162 (27%) from Ebonyi, 126 (21%) from Enugu, 66 (11%) from Abia and 54 (9%) from Imo States responded to the structured questionnaire administered (Figure 2). Of these, 468 (78%) were males and 132 (22%) were females (Figure 3). The results revealed that water yam (Plate 1) production in the study area was dominated by male (78%). This is probably because yam production is energy demanding thus female farmers prefer to go into the production of other crops. The socio-cultural background of the study area is diverse with four ethnic groups encountered among which the, Igbo, Hausa, Igala and Yoruba were the most prominent (Figure 4). The diverse ethnic composition provides an in-depth understanding of the ethno-botany of the crop plant especially among neighboring ethnic groups. Various tribes emphasize different knowledge about the crop and this would assist in flora documentation and classification. The results revealed that 49% of the respondents were within the ages of 31 to 40 years (Figure 5). This age group is usually believed to be the active age group. On the educational level of the respondents,

168 (28.8%) had no formal education, 72 (12%) had primary education, 306 (51%) had secondary education and 54 (9%) had tertiary education (Figure 6). This shows that yam farmers had a significant level of expertise in yam production. The 47% of the respondents were farmers in the study area, 21% were public servants, and 19% were traders while 13% engaged in other occupations (Figure 7).

Out of the 600 respondents interviewed, 62% of the respondents were aware of water yam cultivation in their area and also involved in its cultivation while 38% were not aware of water yam cultivation in their area (Figure 8). Out of the 600 respondents interviewed, 41% of the respondents grew their water yam tubers as food for consumption, 37% grew the tubers as source of income, 3% as source of employment while 19% grew tubers for land security (Figure 9). The data confirmed previous report that yam tubers are mainly cultivated as source of food. From the data obtained, 36 (6%) of the respondents showed no interest in acquiring the knowledge of water yam cultivation while 564 (94%) indicated interest in water yam cultivation (Figure 10).

The results revealed that 43% of the respondents were within the ages of 41 to 50 years, 30% were between 51-60 years while 20% were between 31-40 years (Figure 11). The age range of 35-45 years is usually believed to be the active age group. Out of the 600 respondents interviewed, 62% grew white yam, 26% grew water yam, 2% grew yellow yam while 10% grew aerial yam respectively (Figure 12). On the method for water yam cultivation in the study area, 12% used mound making, 50% used ridge making, 33% used hole making while only 5% used bagging method. The report confirmed that majority of the farmers use ridge making method for water yam cultivation (Figure 13).

The results revealed that, 53% of the respondents were yam farmers in the chain of water yam production, 13% were distributors and whole sellers, 32% were yam sellers while 2% performed

other role not listed here in the water yam production chain (Figure 14). The 64% of the respondents chose April-June as the most suitable season for water yam cultivation, 22% chose January-March, 10% chose October-December while 4% chose July-September (Figure 15). Majority of the respondents (98%) in the study area confirmed the incidence of pest attack on the water yam both in the field and storage while 2% indicated there was no presence of pest attack (Figure 16). The study showed that yam barn was the storage techniques that were popularly known to most of the respondents (70%). This is followed by straw and stick (17%) which ranked second in the storage method (Figure 17). It is worthy to note, however, that only 2% of the respondents were aware of the raffia bag storage which is modern and improved method of yam storage.

The data obtained confirmed that 63% of the respondents adopt cultural method of preservation of water yam, 16% used biological method (curing, fungicide treatment), 9% used chemical while 12% employed other methods (biocontrol agents, yeasts) for water yam preservation (Figure 18). The study indicated that natural wounds (38%), field disease (35%) (yam anthracnose, yam leaf spot) and field pests (27%) (insect pests, nematode pests) were seen as the major predisposal factors causing water yam rot during storage (Figure 19). On the duration in which water yams are stored after harvest, the study revealed that 57% respondents stored their water yams between 4-6 months before they sell them, 40% stored their water yams between 7-9 months while 2% stored their tubers above 10 months before sending them to the market (Figure 20).

In this study, 94% of the respondents believed that water yam has medicinal value. They revealed that *Dioscorea alata* is used to treat post-partum abdominal pain in women. They further revealed its use in the treatment of the fungal skin infection, *Tinea nigra*, which attacks the uppermost layers of skin. However, 6% of the respondents did not report any medicinal use

of the plant in this study (Figure 21). The 97% of the respondents believed that water yams have a socio-cultural value while 3% responded otherwise (Figure 22). “The socio-cultural value of water yam includes information on nature, mythical beliefs and medicine. Some respondents in the study area used it during marriage and fertility ceremonies. All the tribes studied recognized dry/wet/harvest seasons. Other knowledge on non-food uses is tribe dependent (cultural practices, traditional knowledge and uses of water yam vary among different ethnic groups or tribes in the region). This documented indigenous knowledge can be transferred to other communities that cultivate the crop even though the knowledge recorded in this study is tribe dependent” [5].

The 65.67% of the respondents responded that water yam could be cooked/boiled, 18.83% stated that it could be roasted while 7.50% said that it could be fried as chips (Figure 23). However, this can vary depending on the variation in the eating habits of individuals, some of whom may prefer to consume yams alone, while others may eat them with additional foods. These crops can be consumed to fill the gaps of any meal types especially at breakfast time. This indicates the variation in individual preferences between different parts of the world.




Figure 1: Map of South East Nigeria (Onoja *et al.*, 2015)



Plate 1: Yam barn (*Dioscorea alata* L.)

Source: NRCRI, Umudike, Abia State




Figure 2: Percentage response by respondents from the five South-Eastern States




Figure 3: Percentage response by gender




Figure 4: Percentage response by respondents from various ethnic groups




Figure 5: Age of the respondents




Figure 6: Educational qualification of the respondents




Figure 7: Occupation of respondents




Figure 8: Awareness of water yam cultivation by the respondents



Figure 9: Reason for water yam cultivation




Figure 10: Interest in acquiring the knowledge of water yam cultivation




Figure 11: Age group involved in water yam cultivation




Figure 12: Yam species mostly cultivated




Figure 13: Method of water yam cultivation




Figure 14: Role play in yam production




Figure 15: Most suitable season for water yam production




Figure 16: Pest attack on water yam




Figure 17: Water yam storage method




Figure 18: Water yam preservation method




Figure 19: Causal factors of water yam rot in storage




Figure 20: Storage period of water yam




Figure 21: Medicinal value of water yam




Figure 22: Socio-cultural value of water yam




Figure 23: Water yam consumption

Discussion

A total of 600 respondents comprising 32% from Anambra, 27% from Ebonyi, 21% from Enugu, 11% from Abia and 9% from Imo State responded to the structured questionnaire administered. Of these, 78% were males and 22% were females. The results revealed that water yam production in the study area was dominated by male (78%). This is probably because yam production is energy demanding thus female farmers prefer to go into the production of other crops. This result is in line with the findings of Suleiman (2014) and Ekunwe and Osewa (2007) who noted that 85% of yam farmers in Kaiama Local Government, Kwara State were males. The socio-cultural background of the study area is diverse with four ethnic groups encountered among which the, Igbo, Hausa, Igala and Yoruba were the most prominent. The diverse ethnic composition provides an in-depth understanding of the ethno-botany of the crop plant especially among neighboring ethnic groups. Various tribes emphasize different knowledge about the crop and this would assist in flora documentation and classification. The results revealed that 49% of the respondents were within the ages of 31 to 40 years. This age group is usually believed to be the active age group. These findings are closely supported by that of Oluwatosin (2011) who reported a mean age of 35yrs for yam farmers in Ondo State and Osun State respectively. On the educational level of the respondents, 28% had no formal education, 12% had primary education, 51% had secondary education and 9% had tertiary education. This shows that yam

farmers had a significant level of expertise in yam production. This study supports the findings of Oluwatosin (2011) who suggested a secondary school qualification for farmers in Osun State. “The 47% of the respondents were farmers in the study area, 21% were public servants, and 19% were traders while 13% engaged in other occupations. Out of the 600 respondents interviewed, 62% of the respondents were aware of water yam cultivation in their area and also involved in its cultivation while 38% were not aware of water yam cultivation in their area. This study supports the findings that Nigeria alone produces three-quarters of the world total output of yam” (Modu *et al.*, 2015; Okigbo, 2002). Yams are the fifth most harvested crops in Nigeria, following after cassava, maize, guinea corn and beans/cowpeas. More so, after cassava, yams are the most commonly harvested tuber crops in the country (NBS, 2012). “Out of the 600 respondents interviewed, 41% of the respondents grew their water yam tubers as food for consumption, 37% grew the tubers as source of income, 3% as source of employment while 19% grew tubers for land security. The data confirmed previous report that yam tubers are mainly cultivated as source of food. Water yam (*Dioscorea alata* L.) is the most economically important yam species which serve as a staple food for millions of people in tropical and subtropical countries” (Oben *et al.*, 2016). Studies conducted in Eastern Nigeria showed that yams constituted an average of 32% of the farmer’s gross income derived from arable crops (Orkwor, 2000). From the data obtained, 6% of the respondents showed no interest in acquiring the knowledge of water yam cultivation while 94% indicated interest in water yam cultivation. The results revealed that 43% of the respondents were within the ages of 41 to 50 years, 30% were between 51-60 years while 20% were between 31-40 years. The age range of 35-45 years is usually believed to be the active age group. This finding is closely supported by that of Oluwatosin (2011), who reported a mean age of 35 years for yam farmers in Ondo State and Osun State respectively. Out of the 600

respondents interviewed, 62% grew white yam, 26% grew water yam, 2% grew yellow yam while 10% grew aerial yam respectively. “This study supports the previous findings that *Dioscorea alata* is considered to be the most widely distributed species, but the highest levels of yam production worldwide can be attributed to *Dioscorea rotundata*” (Darkwa *et al.*, 2020).

On the method for water yam cultivation in the study area, 12% used mound making, 50% used ridge making, 33% used hole making while only 5% used bagging method. The report confirmed that majority of the farmers use ridge making method for water yam cultivation. The results revealed that, 53% of the respondents were yam farmers in the chain of water yam production, 13% were distributors and whole sellers, 32% were yam sellers while 2% performed other role not listed in the water yam production chain. The 64% of the respondents chose April-June as the most suitable season for water yam cultivation, 22% chose January-March, 10% chose October-December while 4% chose July-September. Majority of the respondents (98%) in the study area confirmed the incidence of pest attack on the water yam both in the field and storage while 2% indicated no to pest attack. This study is in line with the finding of Opera (1999) who reported that the attack by yam beetle and microorganisms such as nematodes and yam virus are the most devastating.

The study showed that yam barn was the storage techniques that were popularly known to most of the respondents (70%). This is followed by straw and stick (17%) which ranked second in the storage method awareness. This finding is similar to those of (Suleiman, 2014; Okoedo-Okojie and Onemolease, 2009), whose studies revealed that yam barn ranked first, followed by heap on the floor. It is worthy to note, however, that only 2% of the respondents were aware of the raffia bag storage which is modern and improved. The data obtained confirmed that 63% of the respondents adopt cultural method of preservation of water yam, 16% used biological method,

9% used chemical while 12% employed other methods for water yam preservation. The study indicated that natural wounds (38%), field disease (35%) and field pests (27%) were seen as the major predisposal factors causing water yam rot during storage. On the period in which water yams are stored after harvest, the study revealed that 57% respondents stored their water yams between 4-6 months before they sell them, 40% stored their water yams between 7-9 months while 2% stored their tubers above 10 months before sending them to the market. In this study, 94% of the respondents believed that water yam has medicinal value. They revealed that *Dioscorea alata* is used to treat post-partum abdominal pain in women. They further revealed its use in the treatment of the fungal skin infection, *Tinea nigra*, which attacks the uppermost layers of skin. Similar to the current findings, Dutta (2015) has reported the application of *D. alata* paste in the treatment of skin diseases. The local tribal communities of Enugu in Nigeria used *D. alata* against fever and the tubers of *D. cayenensis* are used to treat diarrhea (Aiyeloja and Bello, 2006).

However, 6% of the respondents did not report any medicinal use of the plant in this study. The 97% of the respondents believed that water yams have a socio-cultural value while 3% responded otherwise. The socio-cultural value of water yam includes information on nature, mythical beliefs and medicine. Some respondents in the study area used it during marriage and fertility ceremonies. All the tribes studied can recognize dry/wet/harvest seasons. Other knowledge on non-food uses is tribe dependent. This documented indigenous knowledge can be transferred to other communities that cultivate the crop even though the knowledge recorded in this study is tribe dependent. The 65.67% of the respondents responded that water yam could be cooked/boiled, 18.83% stated that it could be roasted while 7.50% said that it could be fried as chips. In line with these finding, Dutta (2015) and Zinash (2008) have reported the consumption

of *Dioscorea* species by mixing them with mushrooms, meat and other vegetables. According to Sangoyomi (2004), it is consumed roasted, fried, boiled, pounded or as flour which can be reconstituted with hot water. However, this can vary depending on the variation in the eating habits of individuals, some of whom may prefer to consume yams alone, while others may eat them with additional foods. These crops can be consumed to fill the gaps of any meal types especially at breakfast time. This indicates the variation in ethno-medicinal knowledge between different parts of the world.

Conclusion

In the light of the foregoing, water yam tubers are grown in the South-Eastern Nigeria and the people from the study area have indigenous knowledge of water yams and their uses. This documented indigenous knowledge can be transferred to other communities that cultivate the crop.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the

name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

- Adeniji, M.O. (1970 a). Fungi associated with storage decay of yam in Nigeria. *Phytopathology*, 60: 590-592.
- Adisa, R.S., Adefalu, L.L., Olatinwo, L.K., Balogun, K.S. and Ogunmadeko, O.O. (2015). Determinants of postharvest losses of yam among yam farmers in Ekiti State, Nigeria. *Bulletin of Institute of Tropical Agriculture Kyushu University*, 38: 73-78.
- Aiyelaja, A.A. and Bello, O.A. (2006). Ethnobotanical potentials of common herbs in Nigeria: A case study of Enugu State. *Educational Research Review*, 1: 16-22.
- Akpaja, E.O., Isikhuemhen, O.S. and Okhuoya, J.A. (2003). Ethnomycology and usage of edible mushrooms among the Igbo people of Nigeria. *International Journal of Medicinal Mushrooms*. 5: 313-319.
- Coursey, D.G. (1967). *Yams*. Longmans Green London. 230p.
- Dansi, A., Dantsey-Barry, H., Dossou-Aminon, I., N'kpenu, E.K., Agre, A.P., Sunu, Y.D., Kombate, K., Loko, Y.L., Dansi, M. and Assogba, P. (2013). Varietal diversity and genetic erosion of cultivated yams (*Dioscorea cayenensis* Poir- *D. rotundata* Lam complex and *D. alata* L.) in Togo. *International Journal of Biodiversity and Conservation*, 5: 223-239.

- Darkwa, K., Olasanmi, B., Asiedu, R. and Asfaw, A. (2020). Review of empirical and emerging breeding methods and tools for yam (*Dioscorea* spp.) improvement: status and prospects. *Zeitschrift Pflanzenzuchtung*, 139: 474-497.
- Dutta, B. (2015). Food and medicinal values of certain species of *Dioscorea* with reference to Assam. *Journal of Pharmacognosy and Phytochemistry*, 3: 15-18.
- Ekunwe and Osewa (2007). Technical efficiency and productivity of yam in Kogi State, Nigeria. *Journal of Applied Science*, 7 (13): 1818-1820.
- Hahn, S.K. (1995). Yams: *Dioscorea* spp. (*Dioscoreaceae*) In: J. Smaartt and Simmonds, N.W. (Eds). *Evolution of Crop Plants*. Longman Scientific and Technical, UK. Pp. 112-120.
- Igwilo, I.O., Afonne, O.J., Maduabuchi, U.J. and Orisakwe, O.E. (2006). Toxicology Study of the Anam River in Otuocha, Anambra State, Nigeria. *Archives of Environmental and Occupational Health*, 61: 205-208.
- Lasztity, R., Hidvegi, M. and Bata, A. (1998). Saponins in Food. *Food Reviews International*, 14: 371-390.
- Maikasuwa, M.A. and Ala, A.L. (2013). Determination of profitability and resource-use efficiency of yam production by women in Bosso local government area of Niger State, Nigeria. *European Scientific Journal*, 9 (16): 196-205.
- Modu, S., Balkis, J., Hajjagana, L., Falmata, A.S., Bintu, B.P. and Yagana, S. (2015). Nutritional qualities in amala produced from *Dioscorea rotundata* (Poir). *Wudpecker Journal of Food Technology*, 3: 012-018.
- National Bureau of Statistics (2012). LSMS-integrated surveys on agriculture: general household survey panel 2010/2011. www.nigerianstat.gov.ng/pages/download/194
- National Population Commission (2006). *Nigeria National Census: Population Distribution by Sex, State, LGAs and Senatorial District*. 2006 Census Priority Tables. Vol. 3.

- Obidiegwu, J.E. and Akpabio, E.M. (2017). The geography of yam cultivation in Southern Nigeria: Exploring its social meanings and cultural functions. *Journal of Ethnic Foods*, 4: 28-35.
- Okigbo, R.N. (2002). Mycoflora of tuber surface of white yam (*Dioscorea rotundata* Poir) and postharvest control of pathogens with *Bacillus subtilis*. *Mycopathologia*, 156: 81-85.
- Okigbo, R.N. (2004). A review of biological control methods for postharvest yams (*Dioscorea* spp.) in storage in South Eastern Nigeria. *King Mongkut Technology Ladkraba Science Journal*, 4 (1): 207-215.
- Okigbo, R.N. and Ogbonnaya, U.O. (2006). Antifungal effects of two tropical plant leaf extracts (*Ocimum gratissimum* and *Aframomum meleguata*) on postharvest yam (*Dioscorea* spp.) rot. *African Journal of Biotechnology*, 5(9): 727-731.
- Okoedo-Okojie, D. U and Onemolease, E. A. (2009). Factors Affecting the Adoption of Yam Storage Technologies in the Northern Ecological Zone of Edo State. *Nigeria Journal of Human Ecology*, 27(2): 155-160.
- Oluwatosin F.M. (2011). Measuring Technical Efficiency of Yam Farmers in Nigeria: A Stochastic Parametric Approach. *Agricultural Journal*, 6 (2): 40-46.
- Onoja, S.U., Nweze, E.I., Mattias, A. and Ejere, V.C. (2015). Evaluation of bacteriological quality and essential elements in commercially bottled/package water produced and marketed in Southeastern Nigeria. *African Journal of Microbiology Research*, 9(8): 1728-1737.
- Opera, L.U. (1999). Yam storage. In: Bakker, A. (Ed.) *CIGR Handbook of Agricultural Engineering Volume IV Agro Processing*. Pp. 182-214.
- Orkwor, G.C. (2000). The importance of yams In: Orkwor, G.C., Asiedu, R. and Ekanayake, I.J. (Eds.) *Food Yams Advances in Research*. International Institute of Tropical Agriculture (IITA), Ibadan, Oyo State, Nigeria. Pp. 1-2.
- Sahel, S. (2014). Unlocking Opportunities in the Yam Value Chain. *Newsletter*, Volume 3.

- Sangoyomi, T.E. (2004). *Post-harvest Fungal Deterioration of Yam (Dioscorea spp) and its Control*. Ph.D. Thesis University of Ibadan, Nigeria. 179p.
- Scarcelli, N., Chair, H., Causse, S., Vesta, R., Couvreur, T.L. and Vigouroux, Y. (2017). Crop wild relative conservation: wild yams are not that wild. *Biological Conservation*, 210: 325-333.
- Scott, G.J.R., Rosegrant, M. and Bokanga, M. (2000). *Roots and Tubers in the Global Food Vision Statement of Year 2020*. International Potato Centre, Lima Peru. Pp. 67-70.
- Shang, H.F., Cheng, H.C., Liang, H.Y., Liu, S.Y. and Hou, W.C. (2007). Immunostimulatory activities of yam tuber mucilages. *Botanical Studies*, 48 (1): 63-70.
- Suleiman .M. (2014). Profitability and Technical Efficiency of Yam Production in Kaiama Local Government Area of Kwara State, Nigeria: *A Bachelor of Agriculture Project*, Department of Agricultural Economics and Farm Management, University of Ilorin, Ilorin, Kwara State. Pp. 124-128.
- Trimanto, T. and Hapsari, L. (2015). Diversity and utilization of *Dioscorea* spp. tuber as alternative food source in Nganjuk Regency, East Java. *Agrivita*, 37: 97-107.
- Udensi, E.A., Oselebe, H.O. and Onuoha, A.U. (2010). Anti-nutritional assessment of *D. alata* varieties. *Pakistan Journal of Nutrition*, 9(2): 179-181.
- Vertex, N. and Becvarova, V. (2014). Yam production as pillar of food security in Logo Local Government Area of Benue State, Nigeria. *European Scientific Journal*, 10 (31): 27-42.
- Zaknayiba, D.B. and Tanko, L. (2013). Costs and returns analysis of yam production among small scale farmers in Karu local government area, Nasarawa State, Nigeria. *Process Analytical Technology*, 9 (1): 73-80.
- Zhang, T. and Oates, C.G. (1999). Relationship between amylase degradation and physicochemical properties of sweet potato starches. *Food Chemistry*, 65: 157-163.
- Zinash O.O. (2008). Determinants of postharvest losses of yam among yam farmers in Ekiti State, Nigeria. *Bulletin of the Institute of Tropical Agriculture, Kyushu University*, 38: 73-78.

Falola A, Salami MF, Bello AA, Olaoye TA. Effect of yam storage techniques usage on farm income in Kwara State, Nigeria. *Agrosearch*. 2017 Jul 19;17(1):54-65.