

Characterizations of Bounds and Tree Structures for the Hyperbolic Sombor Index

**Original Research
Article**

Received: XX December 20XX

Accepted: XX December 20XX

Online Ready: XX December 20XX

Abstract

For a simple graph G with order n and size m . We study the Hyperbolic Sombor index $HSO(G) = \sum_{uv \in E} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}}$, where d_u and d_v are the degrees of vertices u and v respectively, a recently introduced degree-based topological index. We first establish several new bounds for $HSO(G)$ in terms of fundamental graph parameters, including the maximum and minimum degrees, the number of pendent vertices, vertex deletion processes and the relationship between a graph and its complement. In addition, we compute exact values of $HSO(G)$ for several well-known families, including Kragujevac trees, perfect binary trees, binary caterpillar trees and dendrimer trees.

Keywords: Hyperbolic Sombor index; Kragujevac trees; Binary caterpillar trees; dendrimer trees.

2010 Mathematics Subject Classification: 05C50

1 Introduction

Chemical graph theory has become an essential tool for understanding the structural features of chemical compounds through graph theoretic descriptors known as topological indices. These indices translate molecular architecture into numerical values that correlate with physicochemical properties, biological activity or reactivity patterns. Among degree based indices, the recently introduced Sombor index have gained significant attention due to their strong discriminating power and simple analytical structure.

The *Sombor index* was introduced by Gutman (1) as a geometric degree based topological index defined for a simple graph G as

$$SO(G) = \sum_{uv \in E(G)} \sqrt{d_u^2 + d_v^2},$$

where d_u and d_v denote the degrees of the adjacent vertices u and v . Since its introduction, the Sombor index has inspired various modifications and analytical investigations due to its strong correlation with molecular properties. For further literature on the Sombor index, the reader is referred to papers (2; 3; 4).

To incorporate nonlinear growth more effectively, Barman et al. (5) introduced the *Hyperbolic Sombor index*, which is defined for a graph G as

$$HSO(G) = \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}}.$$

In which it established its basic properties, derived initial bounds and demonstrated potential chemical applicability. Subsequent studies have strengthened this foundation through deeper analytical investigation. Albalahi et al. revisited the index by identifying and correcting earlier inaccuracies and introducing a complementary Diminished Sombor Index (6). In parallel, Das and Ahmad further corrected and refined earlier results while providing sharper and more general bounds for trees, unicyclic and bicyclic graphs (7). Collectively, these works underscore the mathematical richness of the Hyperbolic Sombor index and highlight the need for additional structural results, bounds and closed analytical formulas for broader graph families.

Throughout this work, we adopt the standard terminology, notation and structural conventions of graph theory as formulated in (8; 9).

2 Bounds of Hyperbolic Sombor index

In this section, we derive basic bounds for the Hyperbolic Sombor index $HSO(G)$. The derived results include bounds expressed in terms of the maximum and minimum degrees, the number of pendent vertices and the effect of vertex deletion.

Theorem 2.1. *Let $G(V, E)$ be a simple connected graph of order n , size m , minimum degree δ and maximum degree Δ . Then*

$$HSO(G) \leq m\sqrt{1 + \frac{\Delta^2}{\delta^2}}.$$

Proof. Let $uv \in E(G)$ be any edge and assume without loss of generality that $d_u \leq d_v$, so $\min\{d_u, d_v\} = d_u$.

Define

$$a_i = 1 \quad \text{and} \quad b_i = \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} = \frac{\sqrt{d_u^2 + d_v^2}}{d_u}.$$

By the Cauchy Schwarz inequality, we have

$$\left(\sum_{uv \in E} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \right)^2 \leq \left(\sum_{uv \in E} 1^2 \right) \left(\sum_{uv \in E} \frac{d_u^2 + d_v^2}{d_u^2} \right) = m \sum_{uv \in E} \left(1 + \frac{d_v^2}{d_u^2} \right),$$

Since $\delta \leq d_u \leq d_v \leq \Delta$ for every edge $uv \in E$, it follows that

$$\frac{d_v^2}{d_u^2} \leq \frac{\Delta^2}{\delta^2}.$$

Hence,

$$\sum_{uv \in E} \left(1 + \frac{d_v^2}{d_u^2} \right) \leq \sum_{uv \in E} \left(1 + \frac{\Delta^2}{\delta^2} \right) = m \left(1 + \frac{\Delta^2}{\delta^2} \right).$$

Combining the above inequalities yields

$$HSO(G) = \sum_{uv \in E} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \leq m \sqrt{1 + \frac{\Delta^2}{\delta^2}}.$$

Hence, the theorem is proved. □

Theorem 2.2. Let $G(V, E)$ be a simple connected graph of order n , size m and l pendant vertices. Then

$$HSO(G) \geq \sqrt{2}m + (\sqrt{5} - \sqrt{2})l.$$

Proof. Let $G = (V, E)$ be simple connected graph with l pendant vertices. The contribution of edges to the Hyperbolic Sombor index $HSO(G)$ can be classified into two categories:

- Edges incident to a pendant vertex.
- Edges whose both endpoints have degree at least two.

Case 1: Let $uv \in E(G)$ be an edge such that $d_u = 1$ and $d_v \geq 2$. Then the corresponding term in $HSO(G)$ is

$$\frac{\sqrt{1^2 + d_v^2}}{\min\{1, d_v\}} = \sqrt{1 + d_v^2} \geq \sqrt{1 + 2^2} = \sqrt{5}.$$

Since there are exactly l pendant vertices, so l such edges. Hence their total contribution to $HSO(G)$ is at least $l\sqrt{5}$.

Case 2: Let $uv \in E(G)$ be an edge such that $d_u, d_v \geq 2$. Let $m' = \min\{d_u, d_v\}$, so that $d_u, d_v \geq m'$. Then

$$\sqrt{d_u^2 + d_v^2} \geq m' \sqrt{2}.$$

Thus, for each such edge,

$$\frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \geq \frac{m' \sqrt{2}}{m'} = \sqrt{2}.$$

Since there are $m - l$ edges of this type, their total contribution to $HSO(G)$ is at least $(m - l)\sqrt{2}$. Combining the contributions from both cases, we obtain

$$HSO(G) = \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \geq l\sqrt{5} + (m - l)\sqrt{2} = \sqrt{2}m + (\sqrt{5} - \sqrt{2})l.$$

Thus, we arrive at the required result. □

This provides a stronger lower bound than the classical bound $m\sqrt{2}$, expressed in terms of the number of pendant vertices.

Theorem 2.3. Let G be a simple connected graph and let $v \in V(G)$ be a vertex of degree $d_v = k$. Then the Hyperbolic Sombor index satisfies

$$k\sqrt{2} \leq HSO(G) - HSO(G - v) \leq k \cdot \frac{\sqrt{\Delta^2 + k^2}}{\min\{\Delta, k\}},$$

Proof. From the definition of the Hyperbolic Sombor index,

$$HSO(G) = \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}}.$$

Deleting the vertex v removes precisely the k edges incident with v . Hence,

$$HSO(G) - HSO(G - v) = \sum_{u \in N_G(v)} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}}. \tag{1}$$

Lower bound. For every neighbor $u \in N_G(v)$,

$$d_u \geq 1, \quad d_v = k \geq 1.$$

Therefore

$$\sqrt{d_u^2 + k^2} \geq \sqrt{1^2 + 1^2} = \sqrt{2}.$$

Also,

$$\min\{d_u, k\} \leq k.$$

The expression

$$\frac{\sqrt{d_u^2 + k^2}}{\min\{d_u, k\}}$$

attains its minimum value when $d_u = k = 1$, yielding $\sqrt{2}$. Thus each removed edge contributes at least $\sqrt{2}$, and since there are k such edges,

$$HSO(G) - HSO(G - v) \geq k\sqrt{2}.$$

Upper bound. Since $d_u \leq \Delta$ for all $u \in N_G(v)$,

$$\sqrt{d_u^2 + k^2} \leq \sqrt{\Delta^2 + k^2}.$$

Also,

$$\min\{d_u, k\} \geq \min\{\Delta, k\}.$$

Hence, for each term in (1),

$$\frac{\sqrt{d_u^2 + k^2}}{\min\{d_u, k\}} \leq \frac{\sqrt{\Delta^2 + k^2}}{\min\{\Delta, k\}}.$$

Summing over all k neighbors of v gives the desired upper bound:

$$HSO(G) - HSO(G - v) \leq k \cdot \frac{\sqrt{\Delta^2 + k^2}}{\min\{\Delta, k\}}.$$

This completes the proof. □

Theorem 2.4. Let G be a simple graph on n vertices with degree sequence d_1, d_2, \dots, d_n , and let \overline{G} denote its complement. Then the sum of the Hyperbolic Sombor indices of G and \overline{G} satisfies

$$HSO(G) + HSO(\overline{G}) \geq \binom{n}{2} \sqrt{2},$$

Proof. The Hyperbolic Sombor index of a graph G is defined as

$$HSO(G) = \sum_{uv \in E(G)} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}}.$$

For every edge $uv \in E(G)$,

$$\frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \geq \frac{\sqrt{1^2 + 1^2}}{1} = \sqrt{2}.$$

Thus,

$$HSO(G) \geq m\sqrt{2}. \tag{1}$$

Now consider the complement graph \overline{G} . For each vertex v ,

$$d_{\overline{G}}(v) = n - 1 - d_v.$$

Applying the definition,

$$HSO(\overline{G}) = \sum_{uv \in E(\overline{G})} \frac{\sqrt{d_{\overline{G}}(u)^2 + d_{\overline{G}}(v)^2}}{\min\{d_{\overline{G}}(u), d_{\overline{G}}(v)\}}.$$

Since every vertex in a simple graph has degree at least 0 and at most $n - 1$, for each edge of the complement we still have:

$$\frac{\sqrt{d_{\overline{G}}(u)^2 + d_{\overline{G}}(v)^2}}{\min\{d_{\overline{G}}(u), d_{\overline{G}}(v)\}} \geq \sqrt{1^2 + 1^2} = \sqrt{2},$$

because \overline{G} has no isolated edges unless G has universal vertices. Hence,

$$HSO(\overline{G}) \geq m_{\overline{G}}\sqrt{2}. \tag{2}$$

Adding inequalities (1) and (2) gives

$$HSO(G) + HSO(\overline{G}) \geq m\sqrt{2} + m_{\overline{G}}\sqrt{2}.$$

Since

$$m + m_{\overline{G}} = \binom{n}{2},$$

we obtain

$$HSO(G) + HSO(\overline{G}) \geq \binom{n}{2}\sqrt{2}.$$

This completes the proof. □

3 Hyperbolic Sombor index of Trees

In this section, we investigate the explicit computation of the Hyperbolic Sombor index for several classical families of trees. In particular, we establish closed formulas for Kragujevac trees, perfect binary trees, binary caterpillar trees, dendrimer trees and related structures. Throughout this section, we adopt the notation (u, v) -edge to denote an edge whose end vertices have degree u and v , respectively.

Definition 3.1. (10) Let k_1, k_2, \dots, k_n be integers satisfying $0 \leq k_1 \leq k_2 \leq \dots \leq k_n$. The *Kragujevac tree*, denoted by $Kg(k_1, k_2, \dots, k_n)$, is defined as follows:

1. For each k_i , construct the rooted tree B_{k_i} with $2k_i + 1$ vertices, obtained by attaching k_i copies of a two-vertex branch to the root.
2. Introduce a new central vertex and join it with the root of each B_{k_i} .

The resulting graph is a rooted tree with the new central vertex as the main root.

Theorem 3.1. Let $Kg(k_1, k_2, \dots, k_n)$ is the Kragujevac tree. Then

$$HSO(Kg(k_1, k_2, \dots, k_n)) = \sqrt{5}k + \sum_{i=1}^n \left(\frac{k_i}{2} \sqrt{(k_i + 1)^2 + 4} + \frac{\sqrt{(k_i + 1)^2 + n^2}}{\min\{k_i + 1, n\}} \right),$$

where $k = k_1 + k_2 + \dots + k_n$.

Proof. Let $Kg = Kg(k_1, k_2, \dots, k_n)$ denote the Kragujevac tree. If $k = k_1 + k_2 + \dots + k_n$, then the graph Kg contains $2k + n + 1$ vertices. Furthermore, it has k edges of type $(1, 2)$, exactly k_i edges of type $(k_i + 1, 2)$ for each $i = 1, 2, \dots, n$, and one edge of type $(k_i + 1, n)$ for every $i = 1, 2, \dots, n$. Hence, the Hyperbolic Sombor index of $Kg(k_1, k_2, \dots, k_n)$ is

$$\begin{aligned} HSO(Kg) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\ &= \sum_{(1,2)\text{-edge}} \sqrt{1^2 + 2^2} + \sum_{i=1}^n \sum_{(k_i+1,2)\text{-edges}} \frac{\sqrt{(k_i+1)^2 + 2^2}}{2} \\ &\quad + \sum_{i=1}^n \sum_{(k_i+1,n)\text{-edges}} \frac{\sqrt{(k_i+1)^2 + n^2}}{\min\{k_i+1, n\}} \\ &= \sqrt{5}k + \sum_{i=1}^n \frac{k_i}{2} \sqrt{(k_i+1)^2 + 4} + \sum_{i=1}^n \frac{\sqrt{(k_i+1)^2 + n^2}}{\min\{k_i+1, n\}} \\ &= \sqrt{5}k + \sum_{i=1}^n \left(\frac{k_i}{2} \sqrt{(k_i+1)^2 + 4} + \frac{\sqrt{(k_i+1)^2 + n^2}}{\min\{k_i+1, n\}} \right). \end{aligned}$$

This completes the proof. □

Definition 3.2. (8) A *binary tree* is a rooted tree with at least three vertices in which exactly one vertex, called the root, has degree two, while all other vertices have degree either one (leaves) or three (internal nodes). The root is considered to be at level 0. A vertex v_i is said to be at level k_i if the distance between v_i and the root is k_i .

Definition 3.3. (8) A *perfect binary tree* T_k of height k is a binary tree in which all vertices occur up to level k , and every level contains the maximum possible number of vertices. In particular, all internal vertices have exactly two children and all leaf vertices are at the same level k .

Theorem 3.2. Let T_k be a perfect binary tree of height k . Then

$$HSO(T_k) = \sqrt{13} - 4\sqrt{2} + 2^k(\sqrt{2} + \sqrt{10}).$$

Proof. Let T_k be a perfect binary tree of height k with root vertex r . The tree T_k contains $2^{k+1} - 1$ vertices and consequently, $2^{k+1} - 2$ edges. It has 2 edges of type $(2, 3)$, $2^k - 4$ edges of type $(3, 3)$ and 2^k edges of type $(3, 1)$. Hence, the Hyperbolic Sombor index of T_k is

$$\begin{aligned} HSO(T_k) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\ &= \sum_{(2,3)\text{-edges}} \frac{\sqrt{2^2 + 3^2}}{2} + \sum_{(3,3)\text{-edges}} \frac{\sqrt{3^2 + 3^2}}{3} \\ &\quad + \sum_{(3,1)\text{-edges}} \sqrt{3^2 + 1} \\ &= \sqrt{13} + (2^k - 4)\sqrt{2} + 2^k\sqrt{10} \\ &= \sqrt{13} - 4\sqrt{2} + 2^k(\sqrt{2} + \sqrt{10}). \end{aligned}$$

Thus, the desired result is established. □

Definition 3.4. (11) A *binary caterpillar tree* T is a specialized form of a caterpillar tree. In a caterpillar tree, all vertices are either on a central path, called the *spine* or are directly connected to a vertex on

this spine. In a binary caterpillar tree, every internal vertex on the spine has at most two children, so the tree satisfies the properties of a binary tree. Consequently, each internal vertex is either a leaf or has exactly two children.

Theorem 3.3. *Let T be a binary caterpillar tree on n vertices with height k . Then*

$$HSO(T) = \sqrt{5} + \frac{\sqrt{13}}{2} - 2\sqrt{2} + k(\sqrt{2} + \sqrt{10}).$$

Proof. Let T be a binary caterpillar tree of n vertices and height k . Then T has $2k + 1$ vertices and $2k$ edges, consisting of one edge of type $(1, 2)$, one edge of type $(2, 3)$, k edges of type $(1, 3)$ and $k - 2$ edges of type $(3, 3)$.

Hence, the Hyperbolic Sombor index of T is

$$\begin{aligned} HSO(T) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\ &= \sum_{(1,2)\text{-edge}} \sqrt{1 + 2^2} + \sum_{(2,3)\text{-edge}} \frac{\sqrt{2^2 + 3^2}}{2} \\ &\quad + \sum_{(3,1)\text{-edges}} \sqrt{3^2 + 1} + \sum_{(3,3)\text{-edges}} \frac{\sqrt{3^2 + 3^2}}{3} \\ &= \sqrt{5} + \frac{\sqrt{13}}{2} + k\sqrt{10} + (k - 2)\sqrt{2} \\ &= \sqrt{5} + \frac{\sqrt{13}}{2} - 2\sqrt{2} + k(\sqrt{2} + \sqrt{10}). \end{aligned}$$

Hence, the desired result follows. □

Definition 3.5. (12) For a positive integer k , two special families of trees, denoted by A_{2k} and A_{2k+1} , are constructed recursively. The tree A_{2k+1} is derived from A_{2k-1} , while A_{2k} is derived from A_{2k-2} , according to the following rule: three pendant vertices (one in the horizontal direction and two in the vertical direction) are attached to the extreme leftmost and rightmost pendant vertices of the preceding tree. Furthermore, each of the remaining pendant vertices of A_{2k-1} (or A_{2k-2}) receives one additional pendant vertex attached vertically.

Note that:

1. In A_{2k} , there are $2(2k - 1)$ pendant vertices and $(2k - 2)$ vertices of degree 4.
2. In A_{2k+1} , there are $4k$ pendant vertices and $(2k - 1)$ vertices of degree 4.

Theorem 3.4. *Let A_{2k} be the tree. Then*

$$HSO(A_{2k}) = 6\sqrt{17} + (8k - 16)\sqrt{5} + (2k^2 - 8k + 9)\sqrt{2}.$$

Proof. Let k be any positive integer then, the tree A_{2k} contains 6 edges of type $(1, 4)$, $4k - 8$ edges of type $(1, 2)$, $2k - 3$ edges of type $(4, 4)$, $2(2k - 4)$ edges of type $(2, 4)$ and $2(k - 2)(k - 3)$ edges of type $(2, 2)$.

Hence, the Hyperbolic Sombor index of tree A_{2k} is

$$\begin{aligned}
 HSO(A_{2k}) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\
 &= \sum_{(1,4)\text{-edges}} \sqrt{1 + 4^2} + \sum_{(1,2)\text{-edges}} \sqrt{1 + 2^2} \\
 &\quad + \sum_{(4,4)\text{-edges}} \frac{\sqrt{4^2 + 4^2}}{4} + \sum_{(2,4)\text{-edges}} \frac{\sqrt{2^2 + 4^2}}{2} \\
 &\quad + \sum_{(2,2)\text{-edges}} \frac{\sqrt{2^2 + 2^2}}{2} \\
 &= 6\sqrt{17} + (4k - 8)\sqrt{5} + (2k - 3)\sqrt{2} + (2k - 4)\sqrt{20} \\
 &\quad + (k - 2)(k - 3)\sqrt{8} \\
 &= 6\sqrt{17} + (8k - 16)\sqrt{5} + (2k^2 - 8k + 9)\sqrt{2}.
 \end{aligned}$$

This concludes the proof. □

Theorem 3.5. Let A_{2k+1} be the tree. Then

$$HSO(A_{2k+1}) = 6\sqrt{17} + (8k - 12)\sqrt{5} + (2k^2 - 6k + 6)\sqrt{2}.$$

Proof. Let k be any positive integer then, the tree A_{2k+1} contains 6 edges of type (1, 4), $4k - 6$ edges of type (1, 2), $2k - 2$ edges of type (4, 4), $2(2k - 3)$ edges of type (2, 4) and $2(k - 2)^2$ edges of type (2, 2).

Hence, the Hyperbolic Sombor index of tree A_{2k+1} is

$$\begin{aligned}
 HSO(A_{2k+1}) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\
 &= \sum_{(1,4)\text{-edges}} \sqrt{1 + 4^2} + \sum_{(1,2)\text{-edges}} \sqrt{1 + 2^2} \\
 &\quad + \sum_{(4,4)\text{-edges}} \frac{\sqrt{4^2 + 4^2}}{4} + \sum_{(2,4)\text{-edges}} \frac{\sqrt{2^2 + 4^2}}{2} \\
 &\quad + \sum_{(2,2)\text{-edges}} \frac{\sqrt{2^2 + 2^2}}{2} \\
 &= 6\sqrt{17} + (4k - 6)\sqrt{5} + (2k - 2)\sqrt{2} + (2k - 3)\sqrt{20} \\
 &\quad + 2(k - 2)^2\sqrt{2} \\
 &= 6\sqrt{17} + (8k - 12)\sqrt{5} + (2k^2 - 6k + 6)\sqrt{2}.
 \end{aligned}$$

Thus, we arrive at the required result. □

Definition 3.6. (13) A *dendrimer tree* $T_{k,d}$ is a highly regular branched structure with parameters $k \geq 0$ and $d \geq 3$. For any $d \geq 3$, $T_{0,d}$ is the one-vertex tree and $T_{1,d}$ is the star graph S_d . For $k = 2, 3, \dots$, the tree $T_{k,d}$ is obtained from $T_{k-1,d}$ by attaching $d - 1$ new vertices of degree one to each pendant vertex of $T_{k-1,d}$.

Theorem 3.6. Let $T_{k,d}$ be a dendrimer tree. Then

$$HSO(T_{k,d}) = d(d - 1)^k \left(\frac{\sqrt{2}}{d - 2} + \sqrt{d^2 + 1} \right) - \frac{\sqrt{2}d}{d - 2}.$$

Proof. Let $T_{k,d}$ be a dendrimer tree with $k \geq 0$ and $d \geq 3$. The number of vertices in $T_{k,d}$ is

$$1 + d + d(d-1) + d(d-1)^2 + \dots + d(d-1)^{k-1} = 1 + \frac{d}{d-2}((d-1)^k - 1)$$

and the number of edges is $\frac{d}{d-2}((d-1)^k - 1)$.

In $T_{k,d}$, there are $\frac{d}{d-2}((d-1)^{k-1} - 1)$ edges of type (d, d) and $d(d-1)^{k-1}$ edges of type $(1, d)$. Hence, the Hyperbolic Sombor index of tree $T_{k,d}$ is

$$\begin{aligned} HSO(T_{k,d}) &= \sum_{(u,v)\text{-edge}} \frac{\sqrt{d_u^2 + d_v^2}}{\min\{d_u, d_v\}} \\ &= \sum_{(d,d)\text{-edges}} \frac{\sqrt{d^2 + d^2}}{d} + \sum_{(1,d)\text{-edges}} \sqrt{1 + d^2} \\ &= \frac{d}{d-2}((d-1)^{k-1} - 1)\sqrt{2} + d(d-1)^{k-1}\sqrt{1 + d^2} \\ &= d(d-1)^{k-1} \left(\frac{\sqrt{2}}{d-2} + \sqrt{1 + d^2} \right) - \frac{\sqrt{2}d}{d-2}. \end{aligned}$$

Therefore, the obtained result confirms the statement. □

4 CONCLUSIONS

In this paper, we investigated several important bounds for the Hyperbolic Sombor index $HSO(G)$ that describe its behavior under different graph parameters. Bounds based on the maximum and minimum degrees, the number of pendent vertices, vertex deletion from graph and the sum of HSO of graph with its complement. Furthermore, explicit formulas for $HSO(G)$ were obtained for several fundamental families of trees, including Kragujevac trees, perfect binary trees, binary caterpillar trees and dendrimer trees.

Acknowledgment

The second author Kajal S. Baldaniya, sincerely acknowledges the financial support received from the Knowledge Consortium of Gujarat (KCG), Education Department, Government of Gujarat, India, under the SODH Scholarship Scheme (Reference No. 2023015012).

References

- [1] Gutman, I. (2021). Geometric approach to degree-based topological indices: Sombor indices. *MATCH Communications in Mathematical and in Computer Chemistry*, **86**(1), 11–16.
- [2] V. S. Shigehalli, S. A. Chougale. (2025). Sombor index and its applications. *Open J. Discret. Appl. Math.*, **8**(1), 50-63.
- [3] M. Khan, M. Y. Khan, G. Ali, I. L. Popa. (2025). Investigation of General Sombor Index for Optimal Values in Bicyclic Graphs, Trees, and Unicyclic Graphs Using Well-Known Transformations. *Symmetry*, **17**, 968.

- [4] S. Pirzada, B. A. Rather, K. C. Das, Y. Shang, I. Gutman. (2025). On spectrum of Sombor matrix and Sombor energy of graphs. *Georgian Mathematical Journal*, **32**(4), 649-660.
- [5] J. Barman, S. Das. (2023). Geometric Approach to Degree-Based Topological Index: Hyperbolic Sombor Index, *MATCH Commun. Math. Comput. Chem.*, **90**, 77–92.
- [6] Albalahi, A. M., Das, S., Ali, A., Barman, J., Hamza, A. E. (2025). On the hyperbolic Sombor index and its counterpart, arXiv preprint, October.
- [7] Das, K. C., Ahmad, S. (2025). On hyperbolic Sombor index of graphs, arXiv preprint, October.
- [8] J. A. Bondy, U. S. R. Murty. (1976). *Graph Theory with Applications*, Macmillan, London.
- [9] N. Deo. (1974). *Graph theory with applications to engineering and computer science*, Prentice-Hall.
- [10] S. A. Hosseini, M. B. Ahmadi, I. Gutman. (2014). Kragujevac trees with minimal atom–bond connectivity index, *MATCH Commun. Math. Comput. Chem.*, **71**, 5–20.
- [11] L. A. Székely, H. Wang. (2004). On subtrees of trees, *Ind. Math. Inst. Preprint Ser.*, Univ. South Carolina.
- [12] K. V. Iyer, R. Balakrishnan, K. T. Raghavendra. (2007). Wiener index of two special trees, *MATCH Commun. Math. Comput. Chem.*, **57**, 385–392.
- [13] I. Gutman, Y. N. Yeh, S. L. Lee, J. C. Chen. (1994). Wiener numbers of dendrimers, *MATCH Commun. Math. Comput. Chem.*, **30**, 103–115.

©2011 Author1 & Author2; This is an Open Access article distributed under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/2.0>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.