

1
2 **Optimizing Seed Germination in fresh and aged**
3 **Guava (*Psidium guajava* L.) seeds using**
4 **different Chemicals Scarification Treatments**
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 **ABSTRACT**
31

The present investigation entitled “Optimizing Seed Germination in fresh and aged Guava (*Psidium guajava* L.) seeds using different Chemicals Scarification Treatments” was conducted during 2024–2025 at the Horticulture Laboratory of Shree Guru Gobind Singh Tricentenary University, Gurugram (Haryana). The study evaluated the influence of various pre-sowing treatments, including water soaking, hot water soaking (70° C and 80° C) , sulphuric acid (10% and 20%) , gibberellic acid (GA₃) @200 ppm and 400 ppm , and thiourea (@1000 ppm and 2000 ppm) , on seed germination and seedling growth parameters in both fresh and one-year-old guava seed lots. The data were recorded on standard germination percentage, days to emergence of seedling, mean germination time, survival percentage, seedling length, seedling dry weight, vigour indices- I and II and speed

of germination. The results revealed that GA₃ at 400 ppm for 24 hours consistently outperformed other treatments, recording the highest germination percentage (57.61% in fresh seeds and 50.56% in one-year-old seeds), minimum days to emergence (17 and 20.33 days, respectively), maximum survival percentage (59.67% and 54.00%), longest seedling length (5.60 cm and 4.20 cm), and greatest seedling dry weight (328.00 mg and 281.00 mg). This treatment also enhanced vigour indices- I and II and speed of germination. GA₃ at 200 ppm and thiourea at 1000 ppm also showed promising results, though inferior to GA₃ at 400 ppm. The findings demonstrate that pre-sowing seed treatment with GA₃ (400 ppm) is highly effective in improving germination and seedling growth of guava seeds, thereby offering a practical strategy for enhancing seedling establishment in nurseries.

Keywords: Guava, seed germination, pre-sowing treatment, gibberellic acid, seed vigour, seedling growth.

1. INTRODUCTION

Guava (*Psidium guajava* L.), a member of the family Myrtaceae, is a small to medium-sized fruit tree native to tropical America, particularly Mexico and Peru. In India, it ranks fifth among fruit crops after mango, banana, papaya, and citrus, and is widely known as the “Apple of the Tropics” due to its affordability and year-round availability (Singh, 2011). India is the world’s largest producer, contributing nearly 45% of global guava output, with major cultivation in Uttar Pradesh, Bihar, Madhya Pradesh, and Haryana (NHB, 2024). In 2023–24, Haryana produced about 183,643 metric tonnes of guava from 16,750 hectares, with Allahabad Safeda, Hisar Safeda, Hisar Surkha, and Sardar as prominent cultivars (NHB, 2023). The crop plays a significant role in the state’s fruit economy, generating employment and supporting processing industries.

Guava is valued for its high vitamin C content, dietary fiber, pectin, minerals, and antioxidants, making it important for both nutrition and processing into products such as jelly, jam, and nectar (Naseer *et al.*, 2018). Although commercial propagation relies mainly on vegetative methods like air layering and grafting, seed propagation remains essential for raising rootstocks. However, guava seeds exhibit physical dormancy due to a hard seed coat and have short viability, which results in delayed germination and poor seedling establishment (Nonogaki, 2014). Since raising healthy rootstocks is vital for vegetative propagation in Haryana’s expanding guava orchards, pre-sowing treatments such as soaking, scarification, and growth regulators are needed to improve germination and seedling vigor.

Pre-sowing treatments, including soaking, acid scarification, growth regulators, and chemicals, have been applied to improve seed germination and seedling vigor. The present study was therefore undertaken to assess the effects of different pre-sowing seed treatments on germination, seedling growth, and seed quality of guava seeds of varying ages.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The laboratory experiments were conducted at the Horticulture Laboratory, Faculty of Agricultural Sciences, Shree Guru Gobind Singh Tricentenary University, Gurugram, Haryana, India (28°27' N, 77°00' E) situated at 217 m above mean sea level during 2024–25.

70 Seeds of wild guava (*Psidium guajava* L.) of two age groups, viz. fresh and one-year-old,
71 were used for the study. Healthy seeds were extracted from ripe fruits, separated from fleshy
72 mesocarp, thoroughly washed, shade dried to about 12–15% moisture, and stored in cloth
73 bags under room conditions. Before treatment, the seeds were immersed in 1% NaOCl
74 solution for 10 min, rinsed thoroughly in running water for 3 min, and surface dried. The
75 experiment was laid out in Completely Randomized Design (CRD) with ten treatments
76 replicated thrice. Each treatment consisted of 30 seeds per replication. The treatments were:
77 untreated control (T_1), water soaking for 48 h (T_2), hot water soaking at 70°C (1 min, T_3),
78 hot water soaking at 80°C (1 min, T_4), sulphuric acid quick dip at 10% (T_5) and 20% (T_6),
79 GA₃ soaking @ 200 ppm (24 h, T_7), GA₃ @ 400 ppm (24 h, T_8), thiourea @ 1000 ppm
80 (24 h, T_9) and thiourea @ 2000 ppm (24 h, T_{10}). After treatments, seeds were thoroughly
81 washed with tap water to remove residues.

82 For standard germination test, 30 seeds of each treatment in three replicates were placed in
83 moist sand medium and kept at $27 \pm 1^\circ\text{C}$ with 80–85% RH in a seed germinator.
84 Germination count was recorded at regular intervals up to 30 days, and normal seedlings
85 were considered as per ISTA (2001) rules. Mean germination time (MGT) was calculated
86 using the formula of Ranal (2006). Seedling length was measured from 10 randomly
87 selected normal seedlings, and dry weight was recorded after oven drying at $65 \pm 2^\circ\text{C}$ for 48
88 h.

91 3. RESULTS AND DISCUSSION

92
93 **Standard germination (%):** Different pre-sowing seed treatments significantly influenced
94 the germination percentage of guava seeds of both ages (fresh and one-year-old).
95 Germination ranged from 38.6% to 57.6% in fresh seeds and 30.7% to 50.6% in one-year-
96 old seeds. The maximum germination (57.6%) was observed with soaking in GA₃ @ 400
97 ppm for 24 h, which was statistically at par with GA₃ @ 200 ppm, whereas the minimum
98 was recorded in control (38.6%). Similar trends were observed in aged seeds where GA₃
99 @400ppm recorded 50.6% germination as shown in the table no. 1 given below. The
100 improvement under GA₃ treatments may be attributed to enhanced enzymatic activity,
101 particularly the activation of α -amylase, leading to efficient mobilization of stored food
102 reserves (Dinesh & Padmapriya 2022). These findings are in agreement with Barche *et al.*
103 (2010) and Hassan *et al.*, 2014, who reported improved germination of guava and papaya
104 seeds following GA₃ treatments.

105
106 **Days to emergence of seedlings:** The number of days required for seedling emergence
107 was reduced significantly under GA₃ treatments. The earliest emergence (17.0 days for
108 fresh seeds and 20.3 days for aged seeds) was recorded in GA₃ @400ppm (table no.1),
109 while the maximum days (25.0 and 30.0, respectively) were recorded in control. Faster
110 emergence in GA₃-treated seeds might be due to its role in promoting cell elongation and
111 early radicle protrusion (Taiz *et al.*, 2015).

112
113 **Mean germination time (MGT) :** Mean germination time decreased with pre-sowing
114 treatments. The lowest Mean germination time (14.0 and 16.0 days for fresh and aged
115 seeds, respectively) was recorded in GA₃ @ 400 ppm, while the highest (24.0 and 25.0
116 days) was observed in the control as given in the table no. 1. The reduced Mean germination
117 time under GA₃ treatments suggests accelerated germination due to improved water uptake
118 and enzymatic activity (Finch- Savage, 2006; Weitbrecht *et al.* 2011).

119 Table no. 1 Effect of treatments on standard germination %, days to emergence of seedling
120 and mean germination time .

Treatment Details	Germination (%)		Days to emergence		Mean germination time	
	Fresh	1yr old	Fresh	1yr old	Fresh	1yr old
Control	38.62	34.36	25.00	30.00	24.00	25.00
Water soaking for 48 hrs	46.36	44.73	19.67	25.67	17.33	18.67
Hot water soaking at 70°C for 1 min	42.91	41.59	21.67	27.33	20.00	20.67
Hot water soaking at 80°C for 1 min	44.56	43.89	21.00	27.00	18.67	19.67
10% Sulphuric acid treatment (Quick Dip)	39.89	36.37	24.67	29.00	23.33	23.67
20% Sulphuric acid treatment (Quick Dip)	41.10	39.39	22.67	28.33	21.67	22.00
Soaking in GA ₃ @ 200 ppm for 24 hrs	56.07	49.46	17.67	21.00	14.67	16.67
Soaking in GA ₃ @ 400 ppm for 24 hrs	57.61	50.56	17.00	20.33	14.00	16.00
Soaking in thiourea @ 1000 ppm for 24 hrs	51.96	48.46	18.00	21.67	15.33	17.33
Soaking in thiourea @ 2000 ppm for 24 hrs	49.84	46.77	19.00	24.00	16.33	18.00
C.D. @ 5%	2.38	1.21	0.70	1.04	0.99	0.77

121
122
123
124
125
126
127
128

Seedling survival (%): Seedling survival percentage ranged between 30.0% and 59.7% in fresh seeds, and 24.7% to 54.0% in aged seeds (table. no.2). The maximum survival was noted in GA₃ @ 400 ppm, while the control recorded the minimum. Higher survival in GA₃ - treated seeds could be linked to stronger seedlings with better root systems, enabling higher establishment (Taiz *et al.*, 2015; Hussain *et al.*, 2016).

Table no. 2 Effect of treatments on seedling survival %, seedling length and seedling dry weight .

Treatment details	Survival %		Seedling length		Seedling dry wt.	
	Fresh	1yr old	Fresh	1yr old	Fresh	1yr old
Control	30.00	24.67	4.50	3.80	284.00	257.00
Water soaking for 48 hrs	42.67	43.00	5.33	4.13	316.00	271.67
Hot water soaking at 70°C for 1 min	38.67	36.00	5.14	4.04	311.00	267.33
Hot water soaking at 80°C for 1 min	40.33	40.67	5.26	4.11	313.67	269.33
10% H ₂ SO ₄ treatment (Quick Dip)	33.00	28.00	4.63	3.83	288.00	259.00
20% H ₂ SO ₄ treatment (Quick Dip)	36.00	32.33	4.83	3.95	305.00	264.00
Soaking in GA ₃ @200ppm for 24 hrs	55.67	52.00	5.56	4.18	325.33	279.33
Soaking in GA ₃ @400ppm for 24 hrs	59.67	54.00	5.60	4.20	328.00	281.00
Soaking in thiourea @1000ppm for 24hrs	50.33	49.67	5.44	4.17	321.00	277.67
Soaking in thiourea @2000ppm for 24hrs	45.67	46.67	5.38	4.15	318.33	275.00

C.D. at 5%	1.91	2.34	0.09	0.04	2.43	1.17
-------------------	-------------	-------------	-------------	-------------	-------------	-------------

129 **Seedling length (cm):** Significant variation was observed in seedling length across
130 treatments as presented in table no. 2. Fresh seeds treated with GA₃ @ 400 ppm for 24 h
131 produced the longest seedlings (5.60 cm), followed by GA₃ @ 200 ppm (5.43 cm), while the
132 shortest seedlings (3.03 cm) were in the control. Aged seeds showed similar results. The
133 stimulatory effect of GA₃ on cell elongation and division could explain increased seedling
134 length (Yamaguchi, 2008).

135

136 **Seedling dry weight (mg):** Seedling dry weight ranged from 186.0 mg to 328.0 mg in fresh
137 seeds and 154.0 mg to 296.0 mg in aged seeds. The highest values were recorded in GA₃
138 @ 400 ppm, while control showed the lowest (table no.2). This increase may be attributed to
139 better biomass accumulation through enhanced photosynthetic activity and assimilate
140 distribution (Arteca, 1996; Bewley *et al.*, 2013).

141 **Vigour Index – I:** The results (Table 3) indicated that pre-sowing treatments significantly
142 influenced Vigour Index – I of guava seeds. The highest values were recorded with soaking
143 seeds in GA₃ @ 400 ppm for 24 hrs (205.00 in fresh seeds and 105.67 in one-year-old
144 seeds), which was statistically comparable with soaking in GA₃ @ 200 ppm for 24 hrs and
145 soaking in thiourea @ 1000 ppm for 24 hrs. The lowest values (136.00 and 87.00,
146 respectively) were obtained under the control. These results clearly demonstrate the positive
147 role of GA₃ pre-soaking in enhancing seedling vigour by improving germination and growth
148 performance, even in aged seed lots.

149 **Vigour Index – II:** Data presented in Table 3 showed that Vigour Index – II was also
150 significantly affected by pre-sowing treatments. The maximum values were obtained with
151 soaking seeds in GA₃ @ 400 ppm for 24 hrs (11,923.00 in fresh seeds and 7,493.00 in one-
152 year-old seeds), which remained statistically at par with soaking in GA₃ @ 200 ppm for 24
153 hrs. The lowest vigour indices (7,153.00 and 5,857.33, respectively) were recorded in the
154 control. The superiority of GA₃ treatments can be attributed to improved germination,
155 greater shoot elongation, and enhanced metabolic activity, which facilitated efficient
156 utilization of food reserves (Kucera *et al.*, 2005; Copeland, 2001). These results are
157 consistent with earlier findings in aonla (Khan *et al.*, 2006), papaya (Hassan *et al.*, 2014),
158 Indian gooseberry (Rinku *et al.*, 2019), tamarind (Manoharan, 2018), and khirni (Ratna *et al.*,
159 2018).

160 **Speed of germination:** The speed of germination was improved under GA₃ treatments,
161 with GA₃ @ 400 ppm recording the maximum values (1.88 and 1.78 for fresh and aged
162 seeds, respectively) compared to the control (0.78 and 0.61) as shown in table no.3. This
163 suggests rapid and uniform germination in GA₃ -primed seeds. The faster germination under
164 GA₃ treatment may be attributed to its role in stimulating hydrolytic enzyme activity,
165 particularly amylases, which accelerate the breakdown of stored food reserves into simpler
166 forms readily utilized by the embryo. In addition, GA₃ enhances cell elongation and reduces
167 mechanical resistance of seed coverings, thereby promoting quicker radicle protrusion and
168 uniform seedling emergence.

169 Table no. 3 Effect of treatments on vigour index – I and II and speed of germination.

Treatment details	Vigour Index – I		Vigour Index – II		Speed of germination	
	Fresh	1yr old	Fresh	1yr old	Fresh	1yr old
Control	136.00	87.00	7,153.00	5,857.33	1.72	1.53
Water soaking for 48 hrs	192.33	97.67	10,555.33	6,889.00	1.80	1.54
Hot water soaking at 70°C for 1 min	179.33	94.33	10,036.67	6,421.33	1.75	1.58
Hot water soaking at 80°C for 1 min	187.67	96.33	10,218.67	6,573.00	1.78	1.64
10% H ₂ SO ₄ treatment (Quick Dip)	149.33	88.00	8,274.33	5,937.67	1.73	1.61

20% H ₂ SO ₄ treatment (Quick Dip)	165.00	91.33	9,838.33	6,226.33	1.84	1.57
Soaking in GA ₃ @ 200 ppm for 24 hrs	203.00	103.67	11,615.33	7,376.33	1.86	1.75
Soaking in GA ₃ @ 400 ppm for 24 hrs	205.00	105.67	11,923.00	7,493.00	1.88	1.78
Soaking in thiourea @1000 ppm for 24 hrs	200.33	101.33	11,229.67	7,280.33	1.83	1.72
Soaking in thiourea @2000 ppm for 24 hrs	198.00	100.00	10,990.00	7,154.67	1.81	1.67
C.D. at 5%	6.53	2.66	376.41	104.52	0.03	0.03

170

171

172

4. CONCLUSION

173

174

175

176

177

178

179

180

181

The pre-sowing chemical scarification significantly enhanced germination, seedling growth, and vigour in both fresh and one-year-old guava (*Psidium guajava* L.) seeds. Gibberellic acid (GA₃) at 400 ppm for 24 hours emerged as the most effective treatment, promoting the highest germination percentage, earliest emergence, greatest seedling survival, and superior vigour indices. The treatments with GA₃ at 200 ppm and thiourea at 1000 ppm also improved seed performance, though to a lesser extent. These results underscore the efficacy of GA₃ (400 ppm) as a practical intervention for optimizing seedling establishment in guava nurseries.

182

REFERENCES

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

- Abdul-Baki, A. A., and Anderson, J. D. (1973). Vigour determination in soybean seed by multiple criteria. *Crop Science*, **13**(6): 630–633.
- Arteca, R. N. (1996). *Plant growth substances: Principles and applications*. Chapman & Hall.
- Bewley, J. D., Bradford, K. J., Hilhorst, H. W. M., & Nonogaki, H. (2013). *Seeds: Physiology of development, germination, and dormancy* (3rd ed.). Springer.
- Copeland, L. O., & McDonald, M. B. (2001). *Principles of seed science and technology* (4th ed.). Springer.
- Dinesh, M., and Padmapriya, S. (2022). Role of gibberellic acid in seed germination: Mechanisms and applications. *Journal of Plant Growth Regulation*, **41**(4):1321–1332.
- Finch-Savage, W. E., & Leubner-Metzger, G. (2006). Seed dormancy and control of germination. *New Phytologist*, **171**(3): 501–523.
- Hassan, F. A. S., & Ali, E. F. (2014). Effect of GA₃ on seed germination and growth. *Scientia Horticulturae*, **172**:109–116.
- ISTA. (2001). *International rules for seed testing*. Bassersdorf: International Seed Testing Association.
- Khan, M. M., Iqbal, M. J., Abbas, M., & Usman, M. (2006). Effect of plant growth regulators on seed germination and seedling vigour of fruit crops. *Pakistan Journal of Botany*, **38**(3): 593–600.
- Kucera, B., Cohn, M. A., & Leubner-Metzger, G. (2005). Plant hormone interactions during seed dormancy release and germination. *Seed Science Research*, **15**(4): 281–307.
- Hussain, S., Zheng, M., Khan, F., Khaliq, A., Fahad, S., & Peng, S. (2016). Benefits of seed priming. *Plant Physiology and Biochemistry*, **106**:1–10.
- Manoharan, P., & Sivasubramaniam, K. (2018). Effect of chemical treatments on seed germination of tamarind. *Journal of Applied and Natural Science*, **10**(2): 601–605.
- Naseer, R., Shah, M. A., & Shah, M. I. (2018). Nutritional and medicinal importance of guava (*Psidium guajava* L.). *Journal of Food Science and Technology*, **55**(1): 1–14.
- National Horticulture Board (NHB). (2023). *Horticultural statistics at a glance*. Ministry of Agriculture & Farmers Welfare, Government of India.

212 Nonogaki, H. (2014). Seed dormancy and germination. *Annual Review of Plant Biology*, **65**:
213 447–475.
214 Ratna, R., Samir, M., Srivastava, R., & Uniyal, S. (2018). Improving seed germination and
215 seedling traits by pre-sowing treatments in khirni (*Manilkara hexandra*). *Bulletin of*
216 *Environment, Pharmacology and Life Sciences*, **7**(4): 77–81.
217 Rinku, K., Verma, R., Pandey, C. S., Pandey, S. K., & Sahu, K. (2019). Influence of pre-
218 sowing seed treatment and growing conditions on growth performance of Indian gooseberry
219 seedlings (*Emblca officinalis* Gaertn). *International Journal of Current Microbiology and*
220 *Applied Sciences*, **8**(3): 1936–1948.
221 Ranal, M. A., & Santana, D. G. (2006). How and why to measure germination processes.
222 *Brazilian Journal of Botany*, **29**: 1–11
223 Singh, G. (2011). Guava production in India. *Acta Horticulturae*, **909**: 37–44.
224 Taiz, L., Zeiger, E., Møller, I. M., & Murphy, A. (2015). *Plant physiology and development*
225 (6th ed.). Sinauer.
226 Weitbrecht, K., Müller, K., & Leubner-Metzger, G. (2011). Germination and dormancy
227 mechanisms in seeds. *Plant Biology*, **13**(3): 305–317.
228 Yamaguchi, S. (2008). Gibberellin metabolism and its regulation. *Annual Review of Plant*
229 *Biology*, **59**, 225–251.
230