

Unit Regular Elements in Transformation Semigroups and Continuous Transformation Spaces

Abstract. In the current literature, most works on transformation semigroups have shifted their attention to characterizing regular or unit-regular elements in specialized sub-semigroups, such as those preserving partitions, invariant subspaces, or subspace structures. In this paper, we characterize unit regular elements t of the semigroup $T(X)$ of all transformations defined on an arbitrary set X . An important property of a unit regular element which states that an element t of $T(X)$ is unit regular if and only if there exists a cross section X_0 such that $X - X_0$ and $X - R(t)$ are of the same cardinality has been proved. This result reveals that any transformation t on X having finite range is always unit regular and shows that a transformation t in $T(X)$ which is injective (but not surjective) or surjective (but not injective) can never be a unit regular element of $T(X)$. As a consequence, an alternative proof of the fact that $T(X)$ is a unit regular semigroup if and only if X is finite is given. Analogous characterization of unit regular elements of the semigroup space $CT(X)$ of continuous transformations on a Hausdorff topological space X is also analyzed. The result shows that an element t in $CT(X)$ is unit regular if and only if the range $R(t)$ is a closed subset of X , t is a quotient map onto $R(t)$ with kernel of t possessing continuous cross section X_0 such that the closures of $X - X_0$ and $X - R(t)$ are homeomorphic which coincides with t on the boundary X_0 .

KEY WORDS: Semigroup, Regular element, Monoid, Unit regular transformation.

1 Introduction

The concept of regularity in ring theory was introduced by von Neumann [1] to clarify certain aspects of operator algebras. A stronger form of regularity,

namely unit regularity, was introduced by Ehlich [2] as a finiteness condition on rings. In fact, these concepts are essentially of the domain of semigroup theory, and regularity, especially, is extensively studied as a semigroup theoretic notion [3,4]. In recent years, transformation semigroups have been used to model dynamic systems and non-invertible processes due to the rapid development in computational algebra [5,6]. They are semigroup analogues of permutation groups, representing irreversible state changes in automata, mapping structure in algebras. Nowadays, the focus has been shifted to characterizing the regular and unit regular elements in special sub-semigroups where partitions are preserved [7] or invariant subspaces [8, 9, 10, 11] or subspace structures [12, 13, 14] are connected. In this paper, unit regular elements in the space of transformations defined on an arbitrary set are considered, and a characterization of unit regular elements among the continuous transformations defined on topological spaces is given.

Unlike groups, general semigroups do not possess certain desirable features—most notably, the existence of an identity element and inverses. As a result, semigroup structures are generally less orderly and lack some of the neatness found in group theory. It is in this context that we search for regions in the semigroup space where elements behave nearly like invertible elements, helping us to use group-theoretic tools in semigroups. With this background, regular elements in monoids (semigroups with identity) is defined.

2 Preliminary

An element t in a monoid S is said to be a *regular element* if $t = tt't$ for some $t' \in S$. An element t in S is said to be *unit regular* if $t = tut$ for some invertible element $u \in S$. A monoid in which each element is a regular element (unit regular element) is called *regular semigroup* (*unit regular semigroup*). Let $T(X)$ denote the set of all transformations from the set X to itself and $R(t)$ denote the range of $t \in T(X)$ given by $R(t) = \{y \in X : y = t(x), x \in X\}$. Also, let the kernel of transformation $t \in T(X)$ is defined by $\ker t = \{(x, x') : t(x) = t(x')\}$. The relation ρ_t defined on X by $x \rho_t x'$ if and only if $t(x) = t(x')$ is an equivalence relation on X and it partitions X into disjoint equivalent classes $\rho_t(x) = \{x' \in X : t(x) = t(x')\}$. The family of equivalence classes of the equivalence relation ρ is denoted by X/ρ . By a *cross section* of an equivalence relation ρ we mean a subset X_0 of X which contains exactly one element from each equivalence class of the equivalence relation. That is, $X_0 \cap \rho(x)$ is a singleton set. Note that the cardinality $|X_0|$ of the cross section X_0 is same as the cardinality of the family of equivalence classes X/ρ . The following results will be used as needed.

Theorem 2.1. *Let $t : X \rightarrow Y$ be a mapping, where X and Y are non-empty sets and $\rho = \ker t$. Let $\pi : X \rightarrow X/\rho$ be the natural projection defined by $\pi(x) = \rho(x)$. Then $\phi : X/\rho \rightarrow Y$ defined by $\phi(\rho(x)) = t(x)$ is an injection with $\phi\pi = t$.*

Theorem 2.2. *Let ρ be an equivalence relation on X and X_0 be a cross section of ρ . Let $\phi : X/\rho \rightarrow X_0$ be the associated choice function and $\pi : X \rightarrow X/\rho$ be the projection defined by $\pi(x) = \rho(x)$. Then $\pi\phi$ is the identity function on X/ρ and $e = \phi\pi : X \rightarrow X$ is an idempotent mapping.*

Theorem 2.3. *If $e : X \rightarrow X$ is an idempotent mapping, then range of e , $R(e)$, is a cross section of $\ker e$. Moreover, if $\phi : X/\rho \rightarrow R(e)$ is the associated choice function and $\pi : X \rightarrow X/\rho$ is natural projection, then $\phi\pi = e$.*

Let X be a topological space, Y be a set and $t : X \rightarrow Y$ be a surjective mapping. Then the collection $\{H \subseteq Y : t^{-1}(H) \text{ is open in } X\}$ is a topology on Y called the *quotient topology induced by t* . The space Y with this topology is called *quotient space* and the mapping t is called *quotient mapping* [15]. Let ρ be an equivalence relation on X . By a quotient topology on X/ρ , we mean the quotient topology induced by the natural projection $\pi : X \rightarrow X/\rho$ defined by $\pi(x) = \rho(x)$. A cross section X_0 of an equivalence relation ρ is called a continuous cross section of ρ if the choice function $\phi : X/\rho \rightarrow X_0$ is continuous with respect to the quotient topology on X/ρ . The following theorems are continuous analogues of theorems 2.2 and 2.3.

Theorem 2.4. *Let ρ be an equivalence relation on a topological space X and X_0 be a cross section of ρ . Let $\phi : X/\rho \rightarrow X_0$ be the associated choice function and $\pi : X \rightarrow X/\rho$ be the projection defined by $\pi(x) = \rho(x)$. Then $e = \phi\pi : X \rightarrow X$ is a continuous idempotent mapping.*

Theorem 2.5. *If $e : X \rightarrow X$ be a continuous idempotent mapping, then range of e , $R(e)$, is a continuous cross section of $\ker e$.*

A subspace Y of a topological space X is called a *retract* of X if there exists a continuous function $t : X \rightarrow Y$ whose restriction to Y is the identity function on Y . That is, if a topological space X has a subspace Y , which is a retract of X , then the unique identity mapping $i_Y : Y \rightarrow Y$ can be extended continuously to the function $t : X \rightarrow Y$. In this situation, the extended continuous mapping t is called a *retraction* on X .

3 Unit Regular Transformations

The set $T(X)$ of all transformations from X to itself is a regular monoid [3], where X is non-empty set. But $T(X)$ is not, in general, a unit regular semigroup. We prove the following characterization theorem of unit regular elements of $T(X)$.

Theorem 3.1. *An element $t \in T(X)$ is unit regular if and only if there exists a cross section X_0 of kernel of t ($\ker t$) in X with $|X - X_0| = |X - R(t)|$.*

Proof:

Assume that $t \in T(X)$ is a unit regular element. Then we can find a bijective mapping $u: X \rightarrow X$ with $tut = t$. Here $e = ut$ is an idempotent mapping from X to itself. We claim that $X_0 = R(e)$ is a cross section of $\rho = \ker t$. That is, we will prove that $X \cap \rho(x)$ is a singleton set for each $x \in X$. Let x_0 and y_0 be elements of $X_0 \cap \rho(x)$. Since $\rho(x)$ is an equivalence class from the equivalence relation ρ defined by $t(x) = t(y)$, we get that $t(x_0) = t(x) = t(y_0)$. Since $e = ut$ is an idempotent mapping, e restricted to $X_0 = R(e)$ is the identity mapping and so

$$x_0 = e(x_0) = ut(x_0) = ut(y_0) = e(y_0) = y_0$$

To prove that $|X - X_0| = |X - R(t)|$, first observe that the restriction u_0 of u to $R(t)$ is a bijection of $R(t)$ onto X_0 .

Let $y \in R(t)$. Then $y = t(x)$ for some $x \in X$. Note that

$$u_0(y) = u_0(t(x)) = u_0t(x) = ut(x) = e(x)$$

and so $u_0 : R(t) \rightarrow X_0$.

To prove that u_0 is a surjective mapping, let $x_0 \in X_0$ and $y_0 = t(x_0)$ and so $y_0 \in R(t)$. Since e is identity function on X_0 , we have $u_0(y_0) = u(y_0) = u(t(x_0)) = ut(x_0) = e(x_0) = x_0$. Again, since the function u is an injective function, its restriction u_0 is also an injective function. Hence $u_0 : R(t) \rightarrow X_0$ is a bijective function.

Since u is a bijective function from X onto itself and u_0 is a bijection from $R(t)$ onto X_0 , we see that the function u restricted to $X - R(t)$ is a bijection of $X - R(t)$ onto $X - X_0$. Hence $|X - X_0| = |X - R(t)|$.

Conversely, assume that there exists a cross section X_0 of the equivalence relation $\rho = \ker t$ defined by $x \rho y$ if and only if $t(x) = t(y)$ with $|X - X_0| = |X - R(t)|$ for $t \in T(X)$. We shall prove that t is a unit regular mapping.

Since $|X - X_0| = |X - R(t)|$, we can define a bijective mapping u_0 from $X - X_0$ onto $X - R(t)$. Next, we shall prove that $t_0 = t|_{X_0}$, the restricted map of t to X_0 is a bijective mapping. To prove injection, let x_0 and y_0 be the elements of X_0 with $t(x_0) = t(y_0)$. That is,

$$x_0, y_0 \in X_0 \cap \rho(x_0) = X_0 \cap \rho(y_0)$$

Since X_0 is a cross section of the equivalence relation ρ , we get that $x_0 = y_0$. Hence, t_0 is an injective mapping from X_0 to $R(t)$. To prove that t_0 is a surjective mapping onto $R(t)$, let $y_0 \in R(t)$. Then $y_0 = t(x)$ for some $x \in X$. Since X_0 is a cross section of the equivalence relation ρ , $X_0 \cap \rho(x) = \{x_0\}$, a singleton set. Note that $t(x_0) = t(x)$ and $y_0 = t(x) = t(x_0) = t_0(x_0)$. Thus t_0 is a surjective mapping and hence it is a bijective mapping from X_0 onto $R(t)$.

Now, we define a mapping $u : X \rightarrow X$ by

$$u(x) = \begin{cases} t_0^{-1}(x) & \text{if } x \in R(t) \\ u_0(x) & \text{if } x \notin R(t) \end{cases}$$

By construction, the mapping u is a bijection. Let $x \in X$ and $y = t(x)$. Since $y \in R(t)$, by definition of u , $u(y) = t_0^{-1}(y)$ and $t_0^{-1}(y) \in X_0$ so that

$$tut(x) = tu(t(x)) = tu(y) = t(u(y)) = t(t_0^{-1}(y)) = t_0(t_0^{-1}(y)) = t_0t_0^{-1}(y) = y = t(x)$$

That is, $tut(x) = t(x)$. Hence t is a unit regular mapping. \square

As an immediate consequence, we get the following result.

Corollary 3.1. *An injective (surjective) map which is not surjective (injective) is not unit regular.*

Proof:

In the first case, let $t \in T(X)$ be an injective mapping but not surjection. Since t is not surjective, $R(t) \neq X$ and so $|X - R(t)| \neq 0$.

But, since t is an injective mapping, each equivalence class corresponding to $\ker t$ contains a single element and so the only cross section X_0 of $\ker t$ is the whole set X . Then $|X - X_0| = 0$.

That is, there exists no cross section X_0 of $\ker t$ with $|X - X_0| = |X - R(t)|$. So by theorem 3.1, t is not a unit regular element of $T(X)$.

In the second case, let $t \in T(X)$ be a surjective mapping which is not an injective mapping. Then, $|X - R(t)| = 0$.

Since t is not injective mapping, there will be at least one equivalence class of $\rho = \ker t$ which contains more than one element of X . Note that any cross section X_0 of $\ker t$ contains exactly one element in $X_0 \cap \rho(x)$ for each $x \in X$, it follows that any cross section X_0 of $\ker t$ is a proper subset of X and so $|X - X_0| \neq 0$. Since t is surjective mapping $R(t) = X$ and so $|X - R(t)| = 0$. That is, there exists no cross section X_0 of $\ker t$ with $|X - X_0| = |X - R(t)|$. So, by theorem 3.1, t is not a unit regular element of $T(X)$.

Now we describe a sub-family of elements of $T(X)$ in which each element is a unit regular element but the sub-family is not a unit regular semigroup.

Theorem 3.2. *Let X be a non-empty set and $F = \{t \in T(X) : |R(t)| < \infty\}$. Then each element of F is a unit regular element of $T(X)$.*

Proof:

Let $t \in F$. Then $t \in T(X)$ and $|R(t)| = m < \infty$. The equivalence relation ρ defined by $x \rho y$ if and only if $t(x) = t(y)$ has only a finite number m of equivalence classes. This implies that any cross section X_0 of the equivalence classes contains m number of elements. That is, $|X_0| = m = |R(t)|$. The result is proved in two cases.

In the first case, consider X as a finite set, $|X| = n$. Since $|X_0| = m = |R(t)|$, $|X - X_0| = n - m = |X - R(t)|$. Hence by theorem 3.1, t is a unit regular element.

In the second case, consider X as an infinite set. Since X_0 and $R(t)$ are finite sets and X is an infinite set, $|X - X_0| = |X| = |X - R(t)|$. Hence by theorem 3.1, t is a unit regular element.

Using the above theorem, an alternate proof [16] is provided for the following.

Theorem 3.3. *Let X be non-empty set. The set $T(X)$ of all transformations on X is a unit regular semigroup if and only if X is a finite set.*

Proof:

Assume that X is a finite set and $t \in T(X)$. Then $|R(t)| < \infty$ and so by theorem 3.1, t is unit regular element. Since $T(X)$ is semigroup under composition of functions and also every element of $T(X)$ is unit regular, it is a unit regular semigroup.

Conversely, assume that $T(X)$ is a unit regular semigroup, where X is a non-empty set. If possible, let X be an infinite set and A be a finite non-empty subset of X . Since X is infinite, $|X - A| = |X|$. So we can define a bijective mapping $t : X \rightarrow X - A$. Then $t : X \rightarrow X$ is injective function but not a surjection and so t can not be a unit regular mapping, by corollary 3.1. Hence X must be a finite set.

4 Continuous Unit Regular Transformations

In this section, we provide a characterization of unit regular elements in the semigroup of all continuous transformations from a topological space X to itself. The notation $CT(X)$ denotes the semigroup of all continuous transformations from a topological space X to itself. In general, $CT(X)$ is not a regular semigroup. The following results are needed to characterize unit regular elements of $CT(X)$.

Lemma 4.1. *Every continuous cross section of an equivalence relation on a Hausdorff space is a closed set.*

Proof:

Let X_0 be a continuous cross section of an equivalence relation ρ on Hausdorff space X . Since X_0 is a continuous cross section of ρ , the choice function $\phi : X/\rho \rightarrow X_0$ defined by $\phi(x) = \rho(x)$ is continuous under the quotient topology on X/ρ , $\rho(x)$ is the equivalence class containing x . Let $\pi : X \rightarrow X/\rho$ defined by $\pi(x) = \rho(x)$ be natural projection. Then $e = \phi\pi$ is an idempotent mapping on X by theorem 2.2. Since ϕ and π are continuous $e = \phi\pi$ is also continuous. Observe that

$$e(X) = \phi(\pi(X)) = \phi(X/\rho) = X_0$$

It is claimed that X_0 is a closed set in X . Let $(x_\lambda : \lambda \in \Lambda)$ be a net in X_0 which converges to $x \in X$, where Λ is a directed set. Since $e : X \rightarrow X_0$ is continuous,

the net $(e[x_\lambda] : \lambda \in \Lambda)$ in X_0 converges to $e(x)$ in X . Note that $e(x_\lambda) = x_\lambda$ as e restricted to X_0 is identity mapping on X_0 . Also, note that $e(x) \in X_0$. That is, the net $(x_\lambda : \lambda \in \Lambda) = (e[x_\lambda] : \lambda \in \Lambda)$ converges to $x \in X$ and $e(x) \in X_0 \subset X$. Since X is a Hausdorff space, $e(x) = x$ and so $x \in X_0$. Hence X_0 is closed subset of X as every convergent net in X_0 converges in X_0 itself.

The following theorem is a characterization of regular elements in $CT(X)$ given in [17].

Theorem 4.1. *Let X be a topological space and $t \in CT(X)$. Then t is regular if and only if the following conditions are satisfied:*

1. t is a quotient map onto $R(t)$
2. $R(t)$ is a retract of X
3. $\ker t$ has a continuous cross section

A characterization of unit regular elements of $CT(X)$ is now given.

Theorem 4.2. *Let X be a Hausdorff space and $t \in CT(X)$. Then t is unit regular if and only if the following conditions are satisfied:*

1. $R(t)$ is closed
2. t is a quotient map onto $R(t)$
3. there exists a continuous cross section X_0 of $\ker t$ and a homeomorphism from $\overline{X - X_0}$ to $\overline{X - R(t)}$ which coincides with t on the boundary of X_0 .

Proof:

Assume that $t \in CT(X)$ is a unit regular element. Then t is a regular element and so $R(t)$ is a retract of X and t is a quotient map onto $R(t)$ by theorem 4.1. Since X is a Hausdorff space, $R(t)$ is closed. By definition of unit regularity there exists a homeomorphism $u : X \rightarrow X$ such that $tut = t$. Then $e = ut$ is a continuous idempotent map from X to X and $X_0 = R(e)$ is a continuous cross section of $\ker e$ by theorem 2.5. Also,

$$\begin{aligned} (x, y) \in \ker e &\iff e(x) = e(y) \\ &\iff ut(x) = ut(y) \\ &\iff t(x) = t(y) \\ &\iff (x, y) \in \ker t \end{aligned}$$

This shows that $\ker e = \ker t$ and so X_0 is a continuous cross section of $\ker t$. Since u is a homeomorphism,

$$u(\overline{X - R(t)}) = \overline{u(X - R(t))}$$

$$\begin{aligned}
 &= \overline{(u(X) - u(R(t)))} \\
 &= \overline{(X - R(ut))} \\
 &= \overline{X - R(e)} \\
 &= \overline{X - X_0}
 \end{aligned}$$

Therefore, u maps $\overline{X - R(t)}$ homeomorphically onto $\overline{X - X_0}$. Hence u^{-1} maps $\overline{X - X_0}$ homeomorphically onto $\overline{X - R(t)}$. If $bd(X)$ is the boundary of the set X , then:

$$bd(X_0) = \overline{X_0} \cap \overline{X - X_0} = X_0 \cap \overline{X - X_0}$$

as X_0 is closed, by lemma 4.1. Since $bd(X_0) \subset X_0$, and if $x \in bd(X_0)$, then:

$$ut(x) = e(x) = x$$

So $t(x) = u^{-1}(x)$. That is, the mapping t and the homeomorphism u^{-1} coincides on the boundary of X_0 .

Conversely, assume that t satisfies the conditions (1), (2), and (3) of the theorem 4.2. Let $u_0 : \overline{X - X_0} \rightarrow \overline{X - R(t)}$ which coincides with t on $bd(X_0)$ and $t_0 = t|_{X_0}$. Then t_0 is a bijection from X_0 onto $R(t)$, by theorem 3.1.

Let $\rho = \ker t$ and $\phi : X/\rho \rightarrow X_0$ be the choice function and $\pi : X \rightarrow X/\rho$ be the projection mapping. Then $e = \phi\pi$ is a continuous idempotent map from X to itself. So for each $x \in X$, $e(x) = x_0$, where x_0 is the unique element in $\rho(x) \cap X_0$. Also, we have

$$t_0 e(x) = t_0(x_0) = t(x_0) = t(x)$$

as $x_0 \in \rho(x)$. Thus $e(x) = t_0^{-1}t(x)$ for each $x \in X$ and so $e = t_0^{-1}t$. Since e is continuous and t is a quotient map, it follows that t_0^{-1} is continuous. Hence t_0 is a homeomorphism of X_0 onto $R(t)$.

Define $v : X \rightarrow X$ by

$$v(x) = \begin{cases} u_0(x) & \text{if } x \in \overline{X - X_0} \\ t_0(x) & \text{if } x \in \overline{X_0} = X_0 \end{cases}$$

Since u_0 and t coincide on the $bd(X_0)$, the function v is a bijection and it is continuous by the pasting lemma.

Now we claim that u_0^{-1} coincides with t_0^{-1} on the boundary of $R(t)$.

Let $y \in bd(R(t)) = \overline{R(t)} \cap \overline{X - R(t)}$. Since $R(t)$ is closed, $\overline{R(t)} = R(t)$ and so $y = t_0(x)$ for some $x \in X_0$. Also, as $y \in \overline{X - R(t)}$, $y = u_0(t)$ for some $t \in \overline{X - X_0}$. Now $x \in X_0$ and $t \in \overline{X - X_0}$ implies $v(x) = v(t)$. Since v is an injective function, we get $x = t$. That is, $t_0^{-1}(y) = x = t = u_0^{-1}(y)$. Thus, the claim is proved.

So, the mapping $u : X \rightarrow X$ defined by

$$u(x) = \begin{cases} u_0^{-1}(x) & \text{if } x \in \overline{X - R(t)} \\ t_0^{-1}(x) & \text{if } x \in \overline{R(t)} = R(t) \end{cases}$$

J. Dasan, S. Sajikumar

is a bijection and is continuous by the pasting lemma. Hence u is a homeomorphism. Finally we prove that $tut = t$. Let $x \in X$.

Then, as $y = t(x) \in R(t)$, $tut(x) = tu(y) = tt_0^{-1}(y)$.

Now $tt_0^{-1}(y) = t(t_0^{-1}(y)) = t_0(t_0^{-1}(y)) = y$ as $t_0^{-1}(y) \in X_0$.

Thus $tut(x) = y = t(x)$ for all $x \in X$. That is, t is a unit regular mapping.

5 Acknowledgement

The authors extend their heartfelt thanks to the Editor and the anonymous reviewers for their thoughtful comments, constructive suggestions, and insightful feedback, all of which have significantly enhanced the quality and clarity of this manuscript. The authors also acknowledge the support and encouragement received from Dr. E. Krishnan, Professor (Retired), University College, Thiruvananthapuram.

References

- [1] von Neumann, J.: On regular rings, Proc. Nat. Acad. Sci. USA, 22, 1936, 707-713.
- [2] Ehlich, G.: Unit regular rings, Portugal Math. 27. 209-12, 1968.
- [3] Clifford, A. H., Preston, G. B.: The algebraic Theory of semigroups. Math. Surveys No. 7, Amer. Math. Soc. Providence, R. I. Vol. I 1961, Voll. II 1967.
- [4] Harte, R.: Invertibility and singularity of bounded linear operators, Marcel Dekker, New York 1988.
- [5] Gonca Ayk, Hayrulla Ayk and John M. Howie.: On Factorization and Generators in Transformation semigroups, semigroup Forum OF1-OF13, Springer, 2004.
- [6] Marshal I Sampson, R. B. Abubakar, Reny George, Julie S. George.: semigroup Codes for Diagnostic Sequences, European Journal of Pure and Applied Mathematics, 19(1), 7355-7355, New York Business Global, 2026.
- [7] Mosarof Sarkar, M., Narayan Singh, S. On the Semigroup of partition-preserving transformations whose characters are bijective, Bull. Korean Math. Soc.; 61(1), 117-133, 2024.
- [8] Sawatraksa N, Tantong P., Left and Right Regular elements of Some sub-semigroups of the Linear Transformations Semigroups With invariant Subspace, European Journal of Pure and Applied Mathematics, New York Business Global, 18(1), 5623-5623, 2025.
- [9] Pantarak T, Chaiya Y. Regularity and abundance on semigroups of partial transformations with invariant set. Open Mathematics; 21(1): 20230142, 2023.
- [10] Sarkar M, Singh S. N. On certain semigroups of transformations with an invariant set. Asian-European Journal of Mathematics; 15(11), 2250198, 2022.
- [11] Wijarajak R, Chaiya Y. Green's relations and regularity on semigroups of partial transformations with fixed sets. Communications in Algebra; 50(9), 3827-3839, 2022.
- [12] Sarkar, M., Singh, S. N. Unit-regular and semi-balanced elements in various semigroups of transformations, Semigroup Forum, Volume 106(3), 676-691, 2023.

Unit Regular Elements in Transformation Semigroups and Continuous Transformation Spaces

- [13] Sangkhanan, K. Semigroups of partial linear transformations whose restrictions belong to an injective partial linear transformation semigroup, *AIMS Mathematics*, 10(12), 29470-29487, 2025.
- [14] Sarkar, M., Singh, S. N. On certain semigroups of transformations whose restrictions belong to a given semigroup, *Semigroup Forum*, 108(3), 707–723, 2024.
- [15] Dugundji, J.: *Topology*, Prentice Hall, 1966.
- [16] Hugh D’Alarcao: Factorizable as a finiteness condition, *Semigroup Forum*, 20(3), 281-282, 1980.
- [17] Manoj. S.: von Neumann Regularity of Continuous maps, M. Phil. Dissertation submitted to University of Kerala, 1996.